

Exercises PHY981 Spring 2015

The exercises are available at the beginning of the week and are to be handed in the week thereafter on **Wednesdays at noon**. This can be done electronically by sending your github link or by sending a scan of your notes, a pdf or postscript file, or ipython notebook or any format you prefer, by email to hjensen@nscl.msu.edu.

Exercise 5

This exercise serves to convince you about the relation between two different single-particle bases, where one could be our new Hartree-Fock basis and the other a harmonic oscillator basis.

- a) Consider a Slater determinant built up of single-particle orbitals ψ_λ , with $\lambda = 1, 2, \dots, A$.

The unitary transformation

$$\psi_a = \sum_{\lambda} C_{a\lambda} \phi_{\lambda},$$

brings us into the new basis. The new basis has quantum numbers $a = 1, 2, \dots, A$. Show that the new basis is orthonormal. Show that the new Slater determinant constructed from the new single-particle wave functions can be written as the determinant based on the previous basis and the determinant of the matrix C . Show that the old and the new Slater determinants are equal up to a complex constant with absolute value unity. (Hint, C is a unitary matrix).

- b) Starting with the second quantization representation of the Slater determinant

$$\Phi_0 = \prod_{i=1}^n a_{\alpha_i}^{\dagger} |0\rangle,$$

use Wick's theorem to compute the normalization integral $\langle \Phi_0 | \Phi_0 \rangle$.

Exercise 6

Calculate the matrix elements

$$\langle \alpha_1 \alpha_2 | \hat{F} | \alpha_1 \alpha_2 \rangle$$

and

$$\langle \alpha_1 \alpha_2 | \hat{G} | \alpha_1 \alpha_2 \rangle$$

with

$$|\alpha_1 \alpha_2\rangle = a_{\alpha_1}^{\dagger} a_{\alpha_2}^{\dagger} |0\rangle,$$

$$\hat{F} = \sum_{\alpha\beta} \langle \alpha | f | \beta \rangle a_{\alpha}^{\dagger} a_{\beta},$$

$$\langle \alpha | f | \beta \rangle = \int \psi_{\alpha}^{*}(x) f(x) \psi_{\beta}(x) dx,$$

$$\hat{G} = \frac{1}{2} \sum_{\alpha\beta\gamma\delta} \langle \alpha\beta | g | \gamma\delta \rangle a_{\alpha}^{\dagger} a_{\beta}^{\dagger} a_{\delta} a_{\gamma},$$

and

$$\langle \alpha\beta | g | \gamma\delta \rangle = \int \int \psi_{\alpha}^{*}(x_1) \psi_{\beta}^{*}(x_2) g(x_1, x_2) \psi_{\gamma}(x_1) \psi_{\delta}(x_2) dx_1 dx_2$$

Compare these results with those from exercise 3c).

Exercise 7

Show that the onebody part of the Hamiltonian

$$\hat{H}_0 = \sum_{pq} \langle p | \hat{h}_0 | q \rangle a_p^\dagger a_q$$

can be written, using standard annihilation and creation operators, in normal-ordered form as

$$\begin{aligned} \hat{H}_0 &= \sum_{pq} \langle p | \hat{h}_0 | q \rangle a_p^\dagger a_q \\ &= \sum_{pq} \langle p | \hat{h}_0 | q \rangle \{a_p^\dagger a_q\} + \delta_{pq \in i} \sum_{pq} \langle p | \hat{h}_0 | q \rangle \\ &= \sum_{pq} \langle p | \hat{h}_0 | q \rangle \{a_p^\dagger a_q\} + \sum_i \langle i | \hat{h}_0 | i \rangle \end{aligned}$$

Explain the meaning of the various symbols. Which reference vacuum has been used?

Exercise 8

Show that the twobody part of the Hamiltonian

$$\hat{H}_I = \frac{1}{4} \sum_{pqrs} \langle pq | \hat{v} | rs \rangle a_p^\dagger a_q^\dagger a_s a_r$$

can be written, using standard annihilation and creation operators, in normal-ordered form as

$$\begin{aligned} \hat{H}_I &= \frac{1}{4} \sum_{pqrs} \langle pq | \hat{v} | rs \rangle a_p^\dagger a_q^\dagger a_s a_r \\ &= \frac{1}{4} \sum_{pqrs} \langle pq | \hat{v} | rs \rangle \{a_p^\dagger a_q^\dagger a_s a_r\} + \sum_{pq i} \langle pi | \hat{v} | qi \rangle \{a_p^\dagger a_q\} + \frac{1}{2} \sum_{ij} \langle ij | \hat{v} | ij \rangle \end{aligned}$$

Explain again the meaning of the various symbols.

This exercise is optional: Derive the normal-ordered form of the threebody part of the Hamiltonian.

$$\hat{H}_3 = \frac{1}{36} \sum_{\substack{pqr \\ stu}} \langle pqr | \hat{v}_3 | stu \rangle a_p^\dagger a_q^\dagger a_r^\dagger a_u a_t a_s$$

and specify the contributions to the twobody, onebody and the scalar part.