


# TALENT Course II

## Learning from Data: Bayesian methods and machine learning



### Lecture 23: Model checking II

Daniel Phillips  
Ohio University  
TU Darmstadt  
ExtreMe Matter Institute



OHIO  
UNIVERSITY




TALENT Course II is possible thanks to funding from the STFC

# An EFT expansion in pictures

- General EFT series for observable to order  $k$ :  $y = y_{\text{ref}} \sum_{n=0}^k c_n(x) Q^n$
- In ChiEFT  $Q = \frac{(p, m_\pi)}{\Lambda_b}$ ;  $\Lambda_b \approx 600 \text{ MeV}$ ;  $x = \left\{ \frac{p}{m_\pi}, \cos \theta \right\}$


# An EFT expansion in pictures

- General EFT series for observable to order  $k$ :  $y = y_{\text{ref}} \sum_{n=0}^k c_n(x) Q^n$
- In ChiEFT  $Q = \frac{(p, m_\pi)}{\Lambda_b}$ ;  $\Lambda_b \approx 600 \text{ MeV}$ ;  $x = \left\{ \frac{p}{m_\pi}, \cos \theta \right\}$




# An EFT expansion in pictures

- General EFT series for observable to order  $k$ :  $y = y_{\text{ref}} \sum_{n=0}^k c_n(x) Q^n$
- In ChiEFT  $Q = \frac{(p, m_\pi)}{\Lambda_b}$ ;  $\Lambda_b \approx 600 \text{ MeV}$ ;  $x = \left\{ \frac{p}{m_\pi}, \cos \theta \right\}$




# An EFT expansion in pictures

- General EFT series for observable to order  $k$ :  $y = y_{\text{ref}} \sum_{n=0}^k c_n(x) Q^n$
- In ChiEFT  $Q = \frac{(p, m_\pi)}{\Lambda_b}$ ;  $\Lambda_b \approx 600 \text{ MeV}$ ;  $x = \left\{ \frac{p}{m_\pi}, \cos \theta \right\}$



# An EFT expansion in pictures

- General EFT series for observable to order  $k$ :  $y = y_{\text{ref}} \sum_{n=0}^k c_n(x) Q^n$
- In ChiEFT  $Q = \frac{(p, m_\pi)}{\Lambda_b}$ ;  $\Lambda_b \approx 600 \text{ MeV}$ ;  $x = \left\{ \frac{p}{m_\pi}, \cos \theta \right\}$



**This is what a healthy observable expansion looks like:  
bounded coefficients, that do not grow or shrink with order.**

# Bayesian EFT Parameter Estimation

---

$$X(p) = X_0(p) \sum_{i=0}^k c_i(\{a_j\}) x^i$$

# Bayesian EFT Parameter Estimation

- General EFT series for observable to order k:  $X(p) = X_0(p) \sum_{i=0}^k c_i(\{a_j\}) x^i$
- Truncation error =  $X_0(p) c_{k+1} x^{k+1} + \dots$
- Posterior for parameters  $\mathbf{a} = \{a_0, \dots, a_k\}$  by marginalizing:

$$\text{pr}(\mathbf{a}|D, k, k_{\max}) = \int dc_{k+1} \dots dc_{k_{\max}} \text{pr}(\mathbf{a}, c_{k+1}, \dots, c_{k_{\max}} | D, k, k_{\max})$$

$$\text{and Bayesing} = \int dc_{k+1} \dots dc_{k_{\max}} \frac{\text{pr}(D|\mathbf{a}, c_{k+1}, \dots, c_{k_{\max}}, k, k_{\max}) \text{pr}(\mathbf{a}|\bar{a}_{\text{fix}}) \prod_{j=k+1}^{k_{\max}} \text{pr}(c_j|\bar{c}_{\text{fix}})}{\text{pr}(D|k, k_{\max})}$$

# Bayesian EFT Parameter Estimation

- General EFT series for observable to order k:  $X(p) = X_0(p) \sum_{i=0}^k c_i(\{a_j\}) x^i$
- Truncation error =  $X_0(p) c_{k+1} x^{k+1} + \dots$
- Posterior for parameters  $\mathbf{a} = \{a_0, \dots, a_k\}$  by marginalizing:

$$\text{pr}(\mathbf{a}|D, k, k_{\max}) = \int dc_{k+1} \dots dc_{k_{\max}} \text{pr}(\mathbf{a}, c_{k+1}, \dots, c_{k_{\max}} | D, k, k_{\max})$$

$$\text{and Bayesing} = \int dc_{k+1} \dots dc_{k_{\max}} \frac{\text{pr}(D|\mathbf{a}, c_{k+1}, \dots, c_{k_{\max}}, k, k_{\max}) \text{pr}(\mathbf{a}|\bar{a}_{\text{fix}}) \prod_{j=k+1}^{k_{\max}} \text{pr}(c_j|\bar{c}_{\text{fix}})}{\text{pr}(D|k, k_{\max})}$$

- Marginalization over c's produces revised correlation matrix in standard likelihood, accounts for uncertainties (and correlation structure) induced by omitted terms

$$\text{pr}(\mathbf{a} | D) \propto \exp \left( -\frac{1}{2} \mathbf{r}^T (\Sigma_{\text{exp}} + \Sigma_{\text{th}})^{-1} \mathbf{r} \right) \exp \left( -\frac{\mathbf{a}^2}{2\bar{a}^2} \right) \quad \mathbf{r} \equiv \mathbf{y}_{\text{exp}} - \mathbf{y}_{\text{th}}$$

# Bayesian EFT Parameter Estimation

- General EFT series for observable to order k:  $X(p) = X_0(p) \sum_{i=0}^k c_i(\{a_j\}) x^i$
- Truncation error =  $X_0(p) c_{k+1} x^{k+1} + \dots$
- Posterior for parameters  $\mathbf{a} = \{a_0, \dots, a_k\}$  by marginalizing:

$$\text{pr}(\mathbf{a}|D, k, k_{\max}) = \int dc_{k+1} \dots dc_{k_{\max}} \text{pr}(\mathbf{a}, c_{k+1}, \dots, c_{k_{\max}} | D, k, k_{\max})$$



$$\text{and Bayesing} = \int dc_{k+1} \dots dc_{k_{\max}} \frac{\text{pr}(D|\mathbf{a}, c_{k+1}, \dots, c_{k_{\max}}, k, k_{\max}) \text{pr}(\mathbf{a}|\bar{a}_{\text{fix}}) \prod_{j=k+1}^{k_{\max}} \text{pr}(c_j|\bar{c}_{\text{fix}})}{\text{pr}(D|k, k_{\max})}$$

- Marginalization over c's produces revised correlation matrix in standard likelihood, accounts for uncertainties (and correlation structure) induced by omitted terms

$$\text{pr}(\mathbf{a} | D) \propto \exp \left( -\frac{1}{2} \mathbf{r}^T (\Sigma_{\text{exp}} + \Sigma_{\text{th}})^{-1} \mathbf{r} \right) \exp \left( -\frac{\mathbf{a}^2}{2\bar{a}^2} \right) \quad \mathbf{r} \equiv \mathbf{y}_{\text{exp}} - \mathbf{y}_{\text{th}}$$

- Normal distribution for LECs. Here we take a fixed  $\bar{a}$ , could also marginalize over it.


# Truncation Errors Help



Wesolowski et al., JPG (2016)

$$(\Sigma_{\text{th,corr}})_{ij} = (\mathbf{y}_{\text{ref}})_i (\mathbf{y}_{\text{ref}})_j \bar{c}^2 \sum_{n=k+1}^{k_{\max}} x_i^n x_j^n$$

# Truncation Errors Help




Wesolowski et al., JPG (2016)

$$(\Sigma_{\text{th,corr}})_{ij} = (\mathbf{y}_{\text{ref}})_i (\mathbf{y}_{\text{ref}})_j \bar{c}^2 \sum_{n=k+1}^{k_{\max}} x_i^n x_j^n$$

- Plot results for  $c_0$  and  $c_1$  for fit to data at  $x \in [0, x_{\max}]$  and look for stability as a function of  $x_{\max}$

# Truncation Errors Help




Wesolowski et al., JPG (2016)

$$(\Sigma_{\text{th,corr}})_{ij} = (\mathbf{y}_{\text{ref}})_i (\mathbf{y}_{\text{ref}})_j \bar{c}^2 \sum_{n=k+1}^{k_{\max}} x_i^n x_j^n$$

- Plot results for  $c_0$  and  $c_1$  for fit to data at  $x \in [0, x_{\max}]$  and look for stability as a function of  $x_{\max}$
- Lack of stability for  $k=k_{\max}=1$ . Underfitting.

# Truncation Errors Help



Wesolowski et al., JPG (2016)

$$(\Sigma_{\text{th,corr}})_{ij} = (\mathbf{y}_{\text{ref}})_i (\mathbf{y}_{\text{ref}})_j \bar{c}^2 \sum_{n=k+1}^{k_{\max}} x_i^n x_j^n$$


- Plot results for  $c_0$  and  $c_1$  for fit to data at  $x \in [0, x_{\max}]$  and look for stability as a function of  $x_{\max}$
- Lack of stability for  $k=k_{\max}=1$ . Underfitting.
- $k=1, k_{\max}=3$ : once truncation errors are incorporated correct result is achieved earlier, and with smaller error bars, than for uniform-prior (least-squares) result

# $E_{\max}$ plots for NN

Wesolowski, Furnstahl, Melendez, DP, JPG (2019)

- EKM semi-local potential with R=0.9 fm

Epelbaum, Krebs, Mei $\beta$ nner, EPJA 51, 53 (2015)




# $E_{\max}$ plots for NN

Wesolowski, Furnstahl, Melendez, DP, JPG (2019)

- EKM semi-local potential with R=0.9 fm

Epelbaum, Krebs, Mei  ner, EPJA 51, 53 (2015)




$$(\Sigma_{\text{th,uncorr}})_{ij} = (\mathbf{y}_{\text{ref}})^2 \bar{c}^2 \delta_{ij} \sum_{n=k+1}^{k_{\max}} x_i^{2n}$$

# $E_{\max}$ plots for NN

Wesolowski, Furnstahl, Melendez, DP, JPG (2019)

- EKM semi-local potential with R=0.9 fm

Epelbaum, Krebs, Mei  ner, EPJA 51, 53 (2015)



$$(\Sigma_{\text{th,uncorr}})_{ij} = (\mathbf{y}_{\text{ref}})^2 \bar{c}^2 \delta_{ij} \sum_{n=k+1}^{k_{\max}} x_i^{2n}$$


$$(\Sigma_{\text{th,corr}})_{ij} = (\mathbf{y}_{\text{ref}})_i (\mathbf{y}_{\text{ref}})_j \bar{c}^2 \sum_{n=k+1}^{k_{\max}} x_i^n x_j^n$$

# $E_{\max}$ plots for NN

Wesolowski, Furnstahl, Melendez, DP, JPG (2019)

- EKM semi-local potential with R=0.9 fm

Epelbaum, Krebs, Mei  ner, EPJA 51, 53 (2015)



$$(\Sigma_{\text{th,uncorr}})_{ij} = (\mathbf{y}_{\text{ref}})^2 \bar{c}^2 \delta_{ij} \sum_{n=k+1}^{k_{\max}} x_i^{2n}$$

$$(\Sigma_{\text{th,corr}})_{ij} = (\mathbf{y}_{\text{ref}})_i (\mathbf{y}_{\text{ref}})_j \bar{c}^2 \sum_{n=k+1}^{k_{\max}} x_i^n x_j^n$$


- Can resum truncation error to all orders (under assumptions about its correlation across orders): tests validity of FOTA

# $E_{\max}$ plots for NN

Wesolowski, Furnstahl, Melendez, DP, JPG (2019)

- EKM semi-local potential with  $R=0.9$  fm

Epelbaum, Krebs, Mei  ner, EPJA 51, 53 (2015)





$$(\Sigma_{\text{th,uncorr}})_{ij} = (\mathbf{y}_{\text{ref}})^2 \bar{c}^2 \delta_{ij} \sum_{n=k+1}^{k_{\max}} x_i^{2n}$$

$$(\Sigma_{\text{th,corr}})_{ij} = (\mathbf{y}_{\text{ref}})_i (\mathbf{y}_{\text{ref}})_j \bar{c}^2 \sum_{n=k+1}^{k_{\max}} x_i^n x_j^n$$



- Can resum truncation error to all orders (under assumptions about its correlation across orders): tests validity of FOTA

# $E_{\max}$ plot in the $^1S_0$ at NLO

NLO potential in  $^1S_0$  channel  $V_{1S0}^{(2)}(p', p) = \frac{4\pi}{f_\pi^2} \left( \tilde{C}_{1S0}^{np} + C_{1S0} \frac{p^2 + p'^2}{\Lambda_b^2} \right)$





# Changes in Parameter Estimates





Including truncation errors changes central values and (esp.) errors

# Changes in Parameter Estimates



Including truncation errors changes central values and (esp.) errors

# Changes in Parameter Estimates



Including truncation errors changes central values and (esp.) errors

# NN scattering

Epelbaum, Krebs, Mei  ner, PRC, 2015

Employ “semi-local” potentials of Epelbaum, Krebs, and Mei  ner

$$\chi\text{EFT}: \mathcal{L}(\text{N}, \text{\textpi}) \rightarrow V^{(k)} \rightarrow \delta \rightarrow \sigma_{np}$$

$$\sigma_{np}(E_{\text{lab}}) = \sigma_{\text{LO}} \sum_{n=0}^k c_n(p_{\text{rel}}) \left( \frac{p_{\text{rel}}}{\Lambda_b} \right)^n$$

$$x = \frac{p_{\text{rel}}}{\Lambda_b}$$

# NN scattering

Epelbaum, Krebs, Meißner, PRC, 2015


Employ “semi-local” potentials of Epelbaum, Krebs, and Meißner

$$\chi\text{EFT}: \mathcal{L}(\text{N}, \pi) \rightarrow V^{(k)} \rightarrow \delta \rightarrow \sigma_{np}$$

- NN cross section at  $T_{\text{lab}}=50, 96, 143, 200$  MeV
- Potential regulated by local function, parameterized by  $R$
- EKM identify  $\Lambda_b=600$  MeV for smaller  $R$  values
- Here:  $R=0.9$  fm data
- Results at LO, NLO,  $N^2\text{LO}$ ,  $N^3\text{LO}$ ,  $N^4\text{LO}$  ( $k=0, 2, 3, 4, 5$ )

$$\sigma_{np}(E_{\text{lab}}) = \sigma_{\text{LO}} \sum_{n=0}^k c_n(p_{\text{rel}}) \left( \frac{p_{\text{rel}}}{\Lambda_b} \right)^n$$

$$x = \frac{p_{\text{rel}}}{\Lambda_b}$$




# EFT truncation error model checking

---

# EFT truncation error model checking


Accuracy of three weather forecasting services



Melendez, Furnstahl, Wesolowski, PRC, 2017  
after Furnstahl, Klco, DP, Wesolowski, PRC, 2015  
after Bagnaschi, Cacciari, Guffanti, Jenniches, 2015

- Consider predictions at each order, with their error bars, as data and test them to see if the procedure is consistent
- Fix a given DOB interval: compute success ratio, compare


# EFT truncation error model checking



Melendez, Furnstahl, Wesolowski, PRC, 2017  
after Furnstahl, Klco, DP, Wesolowski, PRC, 2015  
after Bagnaschi, Cacciari, Guffanti, Jenniches, 2015

- Consider predictions at each order, with their error bars, as data and test them to see if the procedure is consistent
- Fix a given DOB interval: compute success ratio, compare
- Look at this for EKM predictions at three different orders and 17 different energies
- Interpret in terms of rescaling of  $\Lambda_b$  by a factor  $\lambda$

# EFT truncation error model checking



Melendez, Furnstahl, Wesolowski, PRC, 2017  
after Furnstahl, Klco, DP, Wesolowski, PRC, 2015  
after Bagnaschi, Cacciari, Guffanti, Jenniches, 2015

- Consider predictions at each order, with their error bars, as data and test them to see if the procedure is consistent
- Fix a given DOB interval: compute success ratio, compare
- Look at this for EKM predictions at three different orders and 17 different energies
- Interpret in terms of rescaling of  $\Lambda_b$  by a factor  $\lambda$

**No evidence for significant rescaling of  $\Lambda_b$**

# Breakdown-scale Inference

---

- $\Lambda_b$  determines the size of the  $c_n$ 's. Choose it too big, and they'll be too big. Choose it too small, they'll be too small. And progressively so as one moves to higher and higher order.
- We have a theory for  $\text{pr}(c_n|c_0, c_1, \dots, c_k)$ : now use Bayes' theorem to see how (im)probable are the  $c_n$ 's that dimensionful EFT coefficients ( $b_n$ 's) produce for a given  $\Lambda_b$ .

# Breakdown-scale Inference

---

- $\Lambda_b$  determines the size of the  $c_n$ 's. Choose it too big, and they'll be too big. Choose it too small, they'll be too small. And progressively so as one moves to higher and higher order.
- We have a theory for  $\text{pr}(c_n|c_0, c_1, \dots, c_k)$ : now use Bayes' theorem to see how (im)probable are the  $c_n$ 's that dimensionful EFT coefficients ( $b_n$ 's) produce for a given  $\Lambda_b$ .

At one energy:

$$\text{pr}(\Lambda_b|b_2, \dots, b_k) \propto \frac{1}{\Lambda_b} \left( \frac{\Lambda_b^{k+2}}{(k+1)\langle b^2 \rangle} \right)^{\frac{k-1}{2}}$$

(NLO: k=2, NNLO: k=3, N<sup>3</sup>LO: k=4, etc.)

# Breakdown-scale Inference


- $\Lambda_b$  determines the size of the  $c_n$ 's. Choose it too big, and they'll be too big. Choose it too small, they'll be too small. And progressively so as one moves to higher and higher order.
- We have a theory for  $\text{pr}(c_n|c_0, c_1, \dots, c_k)$ : now use Bayes' theorem to see how (im)probable are the  $c_n$ 's that dimensionful EFT coefficients ( $b_n$ 's) produce for a given  $\Lambda_b$ .

At one energy:

$$\text{pr}(\Lambda_b|b_2, \dots, b_k) \propto \frac{1}{\Lambda_b} \left( \frac{\Lambda_b^{k+2}}{(k+1)\langle b^2 \rangle} \right)^{\frac{k-1}{2}}$$

(NLO: k=2, NNLO: k=3, N<sup>3</sup>LO: k=4, etc.)

Using 5 energies (and 2 angles):



# Breakdown-scale Inference


- $\Lambda_b$  determines the size of the  $c_n$ 's. Choose it too big, and they'll be too big. Choose it too small, they'll be too small. And progressively so as one moves to higher and higher order.
- We have a theory for  $\text{pr}(c_n|c_0, c_1, \dots, c_k)$ : now use Bayes' theorem to see how (im)probable are the  $c_n$ 's that dimensionful EFT coefficients ( $b_n$ 's) produce for a given  $\Lambda_b$ .

At one energy:

$$\text{pr}(\Lambda_b|b_2, \dots, b_k) \propto \frac{1}{\Lambda_b} \left( \frac{\Lambda_b^{k+2}}{(k+1)\langle b^2 \rangle} \right)^{\frac{k-1}{2}}$$

(NLO: k=2, NNLO: k=3, N<sup>3</sup>LO: k=4, etc.)

Using 17 energies (and 7 angles):



# Breakdown-scale Inference

- $\Lambda_b$  determines the size of the  $c_n$ 's. Choose it too big, and they'll be too big. Choose it too small, they'll be too small. And progressively so as one moves to higher and higher order.
- We have a theory for  $\text{pr}(c_n|c_0, c_1, \dots, c_k)$ : now use Bayes' theorem to see how (im)probable are the  $c_n$ 's that dimensionful EFT coefficients ( $b_n$ 's) produce for a given  $\Lambda_b$ .


At one energy:

$$\text{pr}(\Lambda_b|b_2, \dots, b_k) \propto \frac{1}{\Lambda_b} \left( \frac{\Lambda_b^{k+2}}{(k+1)\langle b^2 \rangle} \right)^{\frac{k-1}{2}}$$

(NLO: k=2, NNLO: k=3, N<sup>3</sup>LO: k=4, etc.)


Using 17 energies (and 7 angles):

R=1.2 fm



# Functional Data


- Truncation-error analysis assumed we have 119 independent data points
- But we don't
- We have a function for each observable at each order
- Can we understand the properties of these functions, and so do  $\Lambda_b$  inference and compute success ratios rigorously?



$$\sigma(E) = \sigma_0(E) [1 + c_2(E)x^2 + c_3(E)x^3 + c_4(E)x^4 + c_5(E)x^5]$$


# Observations and Questions

- $c_n$ 's do not grow or shrink with  $n$ : good  $\Lambda_b$  choice
- Bounded functions, mostly between -2 and 2
- Each “takes a turn” at being largest
- Not oscillating quickly in this energy range




# Observations and Questions

- $c_n$ 's do not grow or shrink with  $n$ : good  $\Lambda_b$  choice
- Bounded functions, mostly between -2 and 2
- Each “takes a turn” at being largest
- Not oscillating quickly in this energy range




# Observations and Questions

- $c_n$ 's do not grow or shrink with  $n$ : good  $\Lambda_b$  choice
- Bounded functions, mostly between -2 and 2
- Each “takes a turn” at being largest
- Not oscillating quickly in this energy range



# Observations and Questions

- $c_n$ 's do not grow or shrink with  $n$ : good  $\Lambda_b$  choice
- Bounded functions, mostly between -2 and 2
- Each “takes a turn” at being largest
- Not oscillating quickly in this energy range




## Physics questions:

- Do curves all fluctuate around zero with some common variance?
- What is the correlation length? Is it different at each order?

# Hypothesis: the $c_n$ 's are a GP

Melendez, Furnstahl, DP, Pratola, Wesolowski (2019)




**GP describes  
“functional data” as**

- Gaussian distribution at each point.
- With specified correlation structure.

# Hypothesis: the $c_n$ 's are a GP

Melendez, Furnstahl, DP, Pratola, Wesolowski (2019)



**GP describes  
“functional data” as**

- Gaussian distribution at each point.
- With specified correlation structure.

**Our hypothesis:**

EFT coefficients at different orders can be modeled as:

- independent but identical realizations of one Gaussian Process;
- with a correlation structure; here we use a “squared exponential” (Gaussian) kernel, but we test it

# Gaussian Process model

---

$$k(c_n(x), c_n(y)) = \bar{c}^2 \exp\left(-\frac{(x-y)^2}{2\ell^2}\right)$$

# Gaussian Process model

---

- Specify correlation matrix of  $c_n$  at  $x$  and  $y$ , e.g.:

$$k(c_n(x), c_n(y)) = \bar{c}^2 \exp\left(-\frac{(x-y)^2}{2\ell^2}\right)$$

# Gaussian Process model

---

- Specify correlation matrix of  $c_n$  at  $x$  and  $y$ , e.g.:

$$k(c_n(x), c_n(y)) = \bar{c}^2 \exp\left(-\frac{(x-y)^2}{2\ell^2}\right)$$

- Assumptions?

# Gaussian Process model

---

- Specify correlation matrix of  $c_n$  at  $x$  and  $y$ , e.g.:

$$k(c_n(x), c_n(y)) = \bar{c}^2 \exp\left(-\frac{(x-y)^2}{2\ell^2}\right)$$

- Assumptions?

- Mean of zero

# Gaussian Process model

---

- Specify correlation matrix of  $c_n$  at  $x$  and  $y$ , e.g.:

$$k(c_n(x), c_n(y)) = \bar{c}^2 \exp\left(-\frac{(x-y)^2}{2\ell^2}\right)$$

- Assumptions?

- Mean of zero
- Stationary

# Gaussian Process model

---

- Specify correlation matrix of  $c_n$  at  $x$  and  $y$ , e.g.:

$$k(c_n(x), c_n(y)) = \bar{c}^2 \exp\left(-\frac{(x-y)^2}{2\ell^2}\right)$$

- Assumptions?
  - Mean of zero
  - Stationary
  - Squared exponential correlation structure

# Gaussian Process model

---


- Specify correlation matrix of  $c_n$  at  $x$  and  $y$ , e.g.:

$$k(c_n(x), c_n(y)) = \bar{c}^2 \exp\left(-\frac{(x-y)^2}{2\ell^2}\right)$$

- Assumptions?
  - Mean of zero
  - Stationary
  - Squared exponential correlation structure
- Need to estimate two parameters:  $\bar{c}$  (variance) and  $\ell$  (correlation length)

# Learning cbar and $\ell$ : Toy Example


- Generate three data sets from an underlying GP with mean 0, variance 1 and length scale 3.



- Fit one GP to “training data” from all three data sets (“three realizations”): estimate  $\mu$ ,  $c_{\bar{a}}$ , and  $\ell$ .

# Learning cbar and $\ell$ : Toy Example


- Generate three data sets from an underlying GP with mean 0, variance  $I$  and length scale 3.



- Fit one GP to “training data” from all three data sets (“three realizations”): estimate  $\mu$ ,  $c_{\bar{a}}$ , and  $\ell$ .

# Learning cbar and $\ell$ : Toy Example

- Generate three data sets from an underlying GP with mean 0, variance 1 and length scale 3.



- Fit one GP to “training data” from all three data sets (“three realizations”): estimate  $\mu$ ,  $c_{\text{bar}}$ , and  $\ell$ .

# What can go wrong

---

Bastos & O'Hagan, Technometrics, 2009

# What can go wrong

---

Bastos & O'Hagan, Technometrics, 2009

- Assumption of GP
  - ⇒ data no more than 2-3  $\sigma$  from mean
  - ⇒ data equally likely above or below

# What can go wrong

---

Bastos & O'Hagan, Technometrics, 2009

- Assumption of GP
  - ⇒ data no more than  $2\text{-}3 \sigma$  from mean
  - ⇒ data equally likely above or below
- Assumption of stationarity

# What can go wrong

Bastos & O'Hagan, Technometrics, 2009

- Assumption of GP
  - ⇒ data no more than  $2-3 \sigma$  from mean
  - ⇒ data equally likely above or below
- Assumption of stationarity
  - ⇒ equal variance at all locations
  - ⇒ correlations depend only on  $x-y$

# What can go wrong

Bastos & O'Hagan, Technometrics, 2009

- Assumption of GP
  - ⇒ data no more than  $2-3 \sigma$  from mean
  - ⇒ data equally likely above or below
- Assumption of stationarity
  - ⇒ equal variance at all locations
  - ⇒ correlations depend only on  $x-y$
- Poor estimate of  $c_{bar}$

# What can go wrong

Bastos & O'Hagan, Technometrics, 2009

- Assumption of GP
  - ⇒ data no more than  $2-3 \sigma$  from mean
  - ⇒ data equally likely above or below
- Assumption of stationarity
  - ⇒ equal variance at all locations
  - ⇒ correlations depend only on  $x-y$
- Poor estimate of  $cbar$ 
  - ⇒ credible intervals too wide or too narrow

# What can go wrong

Bastos & O'Hagan, Technometrics, 2009

- Assumption of GP
  - ⇒ data no more than  $2\text{-}3 \sigma$  from mean
  - ⇒ data equally likely above or below
- Assumption of stationarity
  - ⇒ equal variance at all locations
  - ⇒ correlations depend only on  $x\text{-}y$
- Poor estimate of  $c_{bar}$ 
  - ⇒ credible intervals too wide or too narrow
- Poor estimate of  $\ell$

# What can go wrong

Bastos & O'Hagan, Technometrics, 2009

- Assumption of GP
  - ⇒ data no more than  $2\text{-}3 \sigma$  from mean
  - ⇒ data equally likely above or below
- Assumption of stationarity
  - ⇒ equal variance at all locations
  - ⇒ correlations depend only on  $x\text{-}y$
- Poor estimate of  $c_{bar}$ 
  - ⇒ credible intervals too wide or too narrow
- Poor estimate of  $\ell$ 
  - ⇒ credible intervals too wide or too narrow in vicinity of training points

# GP diagnostics summary

Bastos & O'Hagan, Technometrics, 2009

Melendez, Furnstahl, DP, Pratola, Wesolowski (2019)

# GP diagnostics summary

Bastos & O'Hagan, Technometrics, 2009

Melendez, Furnstahl, DP, Pratola, Wesolowski (2019)

- Assess performance of fitted GP on “testing data” set.

# GP diagnostics summary

Bastos & O'Hagan, Technometrics, 2009

Melendez, Furnstahl, DP, Pratola, Wesolowski (2019)

- Assess performance of fitted GP on “testing data” set.
- Errors are correlated, so can’t just add up number of sigmas and/or compute uncorrelated  $\chi^2$
- But plotting individual errors can still be useful

# GP diagnostics summary

Bastos & O'Hagan, Technometrics, 2009

Melendez, Furnstahl, DP, Pratola, Wesolowski (2019)

- Assess performance of fitted GP on “testing data” set.
- Errors are correlated, so can’t just add up number of sigmas and/or compute uncorrelated  $\chi^2$
- But plotting individual errors can still be useful
- “Consistency plot” has to account for correlations.

# GP diagnostics summary

Bastos & O'Hagan, Technometrics, 2009

Melendez, Furnstahl, DP, Pratola, Wesolowski (2019)

- Assess performance of fitted GP on “testing data” set.
- Errors are correlated, so can’t just add up number of sigmas and/or compute uncorrelated  $\chi^2$
- But plotting individual errors can still be useful
- “Consistency plot” has to account for correlations.
- Define Mahalanobis distance, which does account for correlations

$$MD \equiv (y(x_i) - \mu)^T [k(y(x_i), y(x_j))]^{-1} (y(x_j) - \mu)$$

# GP diagnostics summary

Bastos & O'Hagan, Technometrics, 2009

Melendez, Furnstahl, DP, Pratola, Wesolowski (2019)

- Assess performance of fitted GP on “testing data” set.
- Errors are correlated, so can’t just add up number of sigmas and/or compute uncorrelated  $\chi^2$
- But plotting individual errors can still be useful
- “Consistency plot” has to account for correlations.
- Define Mahalanobis distance, which does account for correlations
$$MD \equiv (y(x_i) - \mu)^T [k(y(x_i), y(x_j))]^{-1} (y(x_j) - \mu)$$
- Write  $k(y(x_i), y(x_j)) = G G^T$  with G from PC or SV decomposition then form “Pivoted Cholesky” or “Eigen” errors from  $G(y(x_i) - \mu)$

# GP diagnostics summary













Bastos & O'Hagan, Technometrics, 2009

Melendez, Furnstahl, DP, Pratola, Wesolowski (2019)

- Assess performance of fitted GP on “testing data” set.
- Errors are correlated, so can’t just add up number of sigmas and/or compute uncorrelated  $\chi^2$
- But plotting individual errors can still be useful
- “Consistency plot” has to account for correlations.
- Define Mahalanobis distance, which does account for correlations
$$MD \equiv (y(x_i) - \mu)^T [k(y(x_i), y(x_j))]^{-1} (y(x_j) - \mu)$$
- Write  $k(y(x_i), y(x_j)) = G G^T$  with G from PC or SV decomposition then form “Pivoted Cholesky” or “Eigen” errors from  $G(y(x_i) - \mu)$
- Can QQ plot any of these individual errors
- Although in the end structure as a function of index may be more revealing



# GP c

- Assess p
- Errors are uncorrelated
- But plots
- “Consistent”
- Define  $\mathbf{M}$
- Write  $k(\cdot)$  “Pivoted”
- Can QQ
- Although





# Model checking a GP

- Fit one GP to “training data” from all three data sets (“three realizations”): estimate  $\mu$ ,  $c_{\bar{a}}$ , and  $\ell$ .





# Model checking a GP

- Fit one GP to “training data” from all three data sets (“three realizations”): estimate  $\mu$ ,  $c_{\bar{a}}$ , and  $\ell$ .



# Model checking a GP

- Fit one GP to “training data” from all three data sets (“three realizations”): estimate  $\mu$ ,  $c_{\bar{a}}$ , and  $\ell$ .




# What can go wrong I: different cbar's

---

- Try to fit a single GP to data generated using different variances


# What can go wrong I: different cbar's

- Try to fit a single GP to data generated using different variances




# What can go wrong I: different cbar's

- Try to fit a single GP to data generated using different variances




# What can go wrong I: different cbar's

- Try to fit a single GP to data generated using different variances




# What can go wrong II: different $\ell$ s

- Try to fit single GP to data with different  $\ell$ s



# What can go wrong II: different $\ell$ s




- Try to fit single GP to data with different  $\ell$ s



# What can go wrong II: different $\ell$ s


- Try to fit single GP to data with different  $\ell$ s

$\ell=3.1$



# $\Lambda_b$ and $\ell$ inference


PRELIMINARY



RKE potential;  $\Lambda=500$  MeV;  
Total cross section only

# $\Lambda_b$ and $\ell$ inference

PRELIMINARY




RKE potential;  $\Lambda=500$  MeV;  
Total cross section only

$\ell \approx 40$  MeV

# $\Lambda_b$ and $\ell$ inference

PRELIMINARY




RKE potential;  $\Lambda=500$  MeV;  
Total cross section only

$\ell \approx 40$  MeV

# $\Lambda_b$ and $\ell$ inference

PRELIMINARY




RKE potential;  $\Lambda=500$  MeV;  
Total cross section only  
Restricted domain

$$\ell \approx 116 \text{ MeV}$$

# $\Lambda_b$ and $\ell$ inference

PRELIMINARY



RKE potential;  $\Lambda=500$  MeV;  
Total cross section only  
Restricted domain

$$\ell \approx 116 \text{ MeV}$$

## To do:

- Model non-stationarity
- Play with “switch over” of expansion parameter at  $p \approx m_\pi$
- Include other observables in analysis