
TALENT COURSE II

LEARNING FROM DATA: BAYESIAN METHODS AND MACHINE LEARNING

Lecture 10: Choosing a prior

Daniel Phillips
Ohio University
TU Darmstadt
ExtreMe Matter Institute



OHIO
UNIVERSITY



TECHNISCHE
UNIVERSITÄT
DARMSTADT

TALENT Course II is possible thanks to funding from the STFC

BAYES' THEOREM

Thomas Bayes (1701?-1761)



$$\text{pr}(A \mid B, I) = \frac{\text{pr}(B \mid A, I)\text{pr}(A \mid I)}{\text{pr}(B \mid I)}$$

BAYES' THEOREM

Thomas Bayes (1701?-1761)



$$\text{pr}(A | B, I) = \frac{\text{pr}(B | A, I)\text{pr}(A | I)}{\text{pr}(B | I)}$$

<http://www.bayesian-inference.com>

BAYES' THEOREM

Thomas Bayes (1701?-1761)



$$\text{pr}(A | B, I) = \frac{\text{pr}(B | A, I)\text{pr}(A | I)}{\text{pr}(B | I)}$$

<http://www.bayesian-inference.com>

$$\text{pr}(\text{model} | \text{data}, I) = \frac{\text{pr}(\text{data} | \text{model}, I)\text{pr}(\text{model} | I)}{\text{pr}(\text{data} | I)}$$

BAYES' THEOREM

Thomas Bayes (1701?-1761)



$$\text{pr}(A | B, I) = \frac{\text{pr}(B | A, I)\text{pr}(A | I)}{\text{pr}(B | I)}$$

<http://www.bayesian-inference.com>

$$\text{pr}(\text{model} | \text{data}, I) = \frac{\text{pr}(\text{data} | \text{model}, I)\text{pr}(\text{model} | I)}{\text{pr}(\text{data} | I)}$$



Posterior

BAYES' THEOREM

Thomas Bayes (1701?-1761)



<http://www.bayesian-inference.com>

$$\text{pr}(A | B, I) = \frac{\text{pr}(B | A, I)\text{pr}(A | I)}{\text{pr}(B | I)}$$

Likelihood



$$\text{pr}(\text{model} | \text{data}, I) = \frac{\text{pr}(\text{data} | \text{model}, I)\text{pr}(\text{model} | I)}{\text{pr}(\text{data} | I)}$$



Posterior

BAYES' THEOREM

Thomas Bayes (1701?-1761)



<http://www.bayesian-inference.com>

$$\text{pr}(A | B, I) = \frac{\text{pr}(B | A, I) \text{pr}(A | I)}{\text{pr}(B | I)}$$

Likelihood

Prior



$$\text{pr}(\text{model} | \text{data}, I) = \frac{\text{pr}(\text{data} | \text{model}, I) \text{pr}(\text{model} | I)}{\text{pr}(\text{data} | I)}$$



Posterior

BAYES' THEOREM

Thomas Bayes (1701?-1761)



<http://www.bayesian-inference.com>

$$\text{pr}(A | B, I) = \frac{\text{pr}(B | A, I) \text{pr}(A | I)}{\text{pr}(B | I)}$$

Likelihood

Prior



$$\text{pr}(\text{model} | \text{data}, I) = \frac{\text{pr}(\text{data} | \text{model}, I) \text{pr}(\text{model} | I)}{\text{pr}(\text{data} | I)}$$

Posterior



Normalization

BAYES' THEOREM

Thomas Bayes (1701?-1761)



<http://www.bayesian-inference.com>

$$\text{pr}(A | B, I) = \frac{\text{pr}(B | A, I) \text{pr}(A | I)}{\text{pr}(B | I)}$$

Likelihood

Prior



$$\text{pr}(\text{model} | \text{data}, I) = \frac{\text{pr}(\text{data} | \text{model}, I) \text{pr}(\text{model} | I)}{\text{pr}(\text{data} | I)}$$

Posterior

Normalization

Probability as degree of belief cf. frequentist view

INDIFFERENCE PRIORS: DISCRETE

INDIFFERENCE PRIORS: DISCRETE

- Consider a six-sided dice

INDIFFERENCE PRIORS: DISCRETE

- Consider a six-sided dice
- How do we assign $\text{pr}(X_i|I)$? ($i=1, 2, 3, 4, 5, 6$)

INDIFFERENCE PRIORS: DISCRETE

- Consider a six-sided dice
- How do we assign $\text{pr}(X_i|I)$? ($i=1, 2, 3, 4, 5, 6$)
- We do know

$$\sum_i \text{pr}(X_i | I) = 1$$

INDIFFERENCE PRIORS: DISCRETE

- Consider a six-sided dice
- How do we assign $\text{pr}(X_i|I)$? ($i=1, 2, 3, 4, 5, 6$)
- We do know

$$\sum_i \text{pr}(X_i | I) = 1$$

- Invariance under labeling $\Rightarrow \text{pr}(X_i|I) = 1/6$.

provided I says nothing that breaks the symmetry

INDIFFERENCE PRIORS: CONTINUOUS

INDIFFERENCE PRIORS: CONTINUOUS

- Location

$$\text{pr}(x | I) dx = \text{pr}(x + x_0 | I) dx$$

- Invariance under origin position $\Rightarrow \text{pr}(x|I) = \text{constant}$

provided I says nothing that breaks the symmetry

INDIFFERENCE PRIORS: CONTINUOUS

- Location

$$\text{pr}(x | I) dx = \text{pr}(x + x_0 | I) dx$$

- Invariance under origin position $\Rightarrow \text{pr}(x|I) = \text{constant}$

provided I says nothing that breaks the symmetry

- Scale

$$\text{pr}(\lambda | I) d\lambda = \text{pr}(\beta\lambda | I) d(\beta\lambda)$$

- Invariance under unit choice $\Rightarrow \text{pr}(\lambda | I) \propto 1/\lambda$

provided I says nothing that breaks the symmetry

- “Jeffreys prior”: uniform prior on $\log(\lambda)$

STRAIGHT LINE APPLICATION

STRAIGHT LINE APPLICATION

- First: rescale to get rid of units, i.e., express plot in terms of “natural units” for the problem

STRAIGHT LINE APPLICATION

- First: rescale to get rid of units, i.e., express plot in terms of “natural units” for the problem
- Intercept: location, scale, or something else?

STRAIGHT LINE APPLICATION

- First: rescale to get rid of units, i.e., express plot in terms of “natural units” for the problem
- Intercept: location, scale, or something else?
- Slope: location, scale, or something else?

STRAIGHT LINE APPLICATION

- First: rescale to get rid of units, i.e., express plot in terms of “natural units” for the problem
- Intercept: location, scale, or something else?
- Slope: location, scale, or something else?

$$\text{pr}(m, b | I) \propto \frac{1}{(1 + m^2)^{3/2}} \quad \text{Jaynes (1967); Jeffreys (1946)}$$

“Symmetric prior”

STRAIGHT LINE APPLICATION

- First: rescale to get rid of units, i.e., express plot in terms of “natural units” for the problem
- Intercept: location, scale, or something else?
- Slope: location, scale, or something else?

$$\text{pr}(m, b | I) \propto \frac{1}{(1 + m^2)^{3/2}} \quad \text{Jaynes (1967); Jeffreys (1946)}$$

“Symmetric prior”

- Standard deviation: location, scale, or something else?
-

STRAIGHT LINE APPLICATION

- First: rescale to get rid of units, i.e., express plot in terms of “natural units” for the problem
- Intercept: location, scale, or something else?
- Slope: location, scale, or something else?

$$\text{pr}(m, b | I) \propto \frac{1}{(1 + m^2)^{3/2}} \quad \text{Jaynes (1967); Jeffreys (1946)}$$

“Symmetric prior”

- Standard deviation: location, scale, or something else?

$$\text{pr}(\sigma | I) \propto \frac{1}{\sigma}$$

PRINCIPLE OF MAXIMUM ENTROPY

$$\sum_{i=1}^6 p_i = 1$$

(something like this is always true)

$$\sum_{i=1}^6 ip_i = 4.5$$

PRINCIPLE OF MAXIMUM ENTROPY

- Suppose we have some information on our dice, e.g.

$$\sum_{i=1}^6 p_i = 1$$

(something like this is always true)

$$\sum_{i=1}^6 ip_i = 4.5$$

PRINCIPLE OF MAXIMUM ENTROPY

- Suppose we have some information on our dice, e.g.

$$\sum_{i=1}^6 p_i = 1 \quad \text{(something like this is always true)}$$

$$\sum_{i=1}^6 ip_i = 4.5$$

- Principle of maximum entropy seeks to maximize

$$S = - \sum_{i=1}^6 p_i \log(p_i) \quad \text{subject to these constraints}$$

PRINCIPLE OF MAXIMUM ENTROPY

- Suppose we have some information on our dice, e.g.

$$\sum_{i=1}^6 p_i = 1 \quad \text{(something like this is always true)}$$

$$\sum_{i=1}^6 ip_i = 4.5$$

- Principle of maximum entropy seeks to maximize

$$S = - \sum_{i=1}^6 p_i \log(p_i) \quad \text{subject to these constraints}$$

- Define $Q(\{p_i\}; \lambda_0, \lambda_1) = - \sum_{i=1}^6 p_i \log(p_i) + \lambda_0 \left(1 - \sum_i p_i \right) + \lambda_1 \left(4.5 - \sum_i ip_i \right)$
-

HAIR COLOR-HEIGHT EXAMPLE REVISITED

- $\frac{1}{3}$ of all Australians are blond, and $\frac{1}{4}$ are tall
- What proportion are blond and tall?

HAIR COLOR-HEIGHT EXAMPLE REVISITED

- 1/3 of all Australians are blond, and 1/4 are tall
- What proportion are blond and tall?

$$Q(\{p_i\}; \lambda_0, \lambda_1, \lambda_2) = - \sum_{i=1}^4 p_i \log(p_i) + \lambda_0 \left(1 - \sum_{i=1}^4 p_i \right) + \lambda_1 \left(\frac{1}{3} - p_1 - p_2 \right) + \lambda_2 \left(\frac{1}{4} - p_1 - p_3 \right)$$

HAIR COLOR-HEIGHT EXAMPLE REVISITED

- 1/3 of all Australians are blond, and 1/4 are tall
- What proportion are blond and tall?

$$Q(\{p_i\}; \lambda_0, \lambda_1, \lambda_2) = - \sum_{i=1}^4 p_i \log(p_i) + \lambda_0 \left(1 - \sum_{i=1}^4 p_i \right) + \lambda_1 \left(\frac{1}{3} - p_1 - p_2 \right) + \lambda_2 \left(\frac{1}{4} - p_1 - p_3 \right)$$

	BLOND	BROWN	SUMS
TALL	x	1/4-x	1/4
SHORT	1/3-x	5/12+x	3/4
SUMS	1/3	2/3	1

MONKEYS ARGUMENT

Thomas Shahan - <https://www.flickr.com/photos/>

- A model for assigning probabilities to M different alternatives: monkeys throwing N balls into M equally sized boxes



MONKEYS ARGUMENT

Thomas Shahan - <https://www.flickr.com/photos/>

- A model for assigning probabilities to M different alternatives: monkeys throwing N balls into M equally sized boxes
- Number of micro-states, W , as a function of $\{p_i\}$ is



MONKEYS ARGUMENT

Thomas Shahan - <https://www.flickr.com/photos/>

- A model for assigning probabilities to M different alternatives: monkeys throwing N balls into M equally sized boxes
- Number of micro-states, W , as a function of $\{p_i\}$ is

$$\log(W(\{n_i\})) = \log(N!) - \sum_{i=1}^M \log(n_i!)$$



MONKEYS ARGUMENT

Thomas Shahan - <https://www.flickr.com/photos/>

- A model for assigning probabilities to M different alternatives: monkeys throwing N balls into M equally sized boxes
- Number of micro-states, W , as a function of $\{p_i\}$ is

$$\log(W(\{n_i\})) = \log(N!) - \sum_{i=1}^M \log(n_i!)$$

$$\log(W(\{n_i\})) \approx -N \sum_{i=1}^M p_i \log(p_i)$$



MONKEYS ARGUMENT

Thomas Shahan - <https://www.flickr.com/photos/>

- A model for assigning probabilities to M different alternatives: monkeys throwing N balls into M equally sized boxes
- Number of micro-states, W , as a function of $\{p_i\}$ is

$$\log(W(\{n_i\})) = \log(N!) - \sum_{i=1}^M \log(n_i!)$$

$$\log(W(\{n_i\})) \approx -N \sum_{i=1}^M p_i \log(p_i)$$

So define $S = - \sum_{i=1}^M p_i \log(p_i)$



MONKEYS ARGUMENT

Thomas Shahan - <https://www.flickr.com/photos/>

- A model for assigning probabilities to M different alternatives: monkeys throwing N balls into M equally sized boxes
- Number of micro-states, W , as a function of $\{p_i\}$ is

$$\log(W(\{n_i\})) = \log(N!) - \sum_{i=1}^M \log(n_i!)$$

$$\log(W(\{n_i\})) \approx -N \sum_{i=1}^M p_i \log(p_i)$$

So define $S = - \sum_{i=1}^M p_i \log(p_i)$ Entropy



MONKEYS ARGUMENT

Thomas Shahan - <https://www.flickr.com/photos/>

- A model for assigning probabilities to M different alternatives: monkeys throwing N balls into M equally sized boxes
- Number of micro-states, W , as a function of $\{p_i\}$ is

$$\log(W(\{n_i\})) = \log(N!) - \sum_{i=1}^M \log(n_i!)$$

$$\log(W(\{n_i\})) \approx -N \sum_{i=1}^M p_i \log(p_i)$$

So define $S = - \sum_{i=1}^M p_i \log(p_i)$ Entropy



Jaynes: derive thermo
using MaxEnt to assign
pdfs subject to macro
constraints

WHY MAXIMIZE THE ENTROPY?

- Information theory: maximum entropy=minimum information (Shannon, 1948)
- Logical consistency (Shore & Johnson, 1960)
- Uncorrelated assignments related monotonically to S (Skilling, 1988)

WHY MAXIMIZE THE ENTROPY?

- Information theory: maximum entropy=minimum information (Shannon, 1948)
- Logical consistency (Shore & Johnson, 1960)
- Uncorrelated assignments related monotonically to S (Skilling, 1988)

VARIATIONAL FUNCTION	OPTIMAL X	IMPLIED CORRELATION
$-\sum p_i \log(p_i)$	0.0833	None
$-\sum p_i^2$	0.0417	Negative
$\sum \log(p_i)$	0.1060	Positive
$\sum \sqrt{p_i}$	0.0967	Positive

CONTINUOUS CASE

- Return to monkeys, but now with different probabilities for each bin. Then

$$S = - \sum_{i=1}^M p_i \log \left(\frac{p_i}{m_i} \right)$$

Shannon-Jaynes entropy
Kullback number
cross entropy

CONTINUOUS CASE

- Return to monkeys, but now with different probabilities for each bin. Then

$$S = - \sum_{i=1}^M p_i \log \left(\frac{p_i}{m_i} \right)$$

Shannon-Jaynes entropy
Kullback number
cross entropy

- Continuous case
-

CONTINUOUS CASE

- Return to monkeys, but now with different probabilities for each bin. Then

$$S = - \sum_{i=1}^M p_i \log \left(\frac{p_i}{m_i} \right)$$

Shannon-Jaynes entropy
Kullback number
cross entropy

- Continuous case

$$S[p] = - \int p(x) \log \left[\frac{p(x)}{m(x)} \right]$$

ONE COMMENT, TWO EXAMPLES

ONE COMMENT, TWO EXAMPLES

- $m(x)$ makes the expression invariant under a change of variables

ONE COMMENT, TWO EXAMPLES

- $m(x)$ makes the expression invariant under a change of variables
- Example 1: maximize S subject only to normalization

ONE COMMENT, TWO EXAMPLES

- $m(x)$ makes the expression invariant under a change of variables
- Example 1: maximize S subject only to normalization

$$Q[p; \lambda] = - \int dx p(x) \log \left[\frac{p(x)}{m(x)} \right] + \lambda \left(1 - \int dx p(x) \right)$$

Solution: $p(x)$ proportional to $m(x)$

ONE COMMENT, TWO EXAMPLES

- $m(x)$ makes the expression invariant under a change of variables
- Example 1: maximize S subject only to normalization

$$Q[p; \lambda] = - \int dx p(x) \log \left[\frac{p(x)}{m(x)} \right] + \lambda \left(1 - \int dx p(x) \right)$$

Solution: $p(x)$ proportional to $m(x)$

- Example 2: maximize $S[p]$ subject to constraint

$$\int dx p(x) (x - \mu)^2 = \sigma^2$$

ONE COMMENT, TWO EXAMPLES

- $m(x)$ makes the expression invariant under a change of variables
- Example 1: maximize S subject only to normalization

$$Q[p; \lambda] = - \int dx p(x) \log \left[\frac{p(x)}{m(x)} \right] + \lambda \left(1 - \int dx p(x) \right)$$

Solution: $p(x)$ proportional to $m(x)$

- Example 2: maximize $S[p]$ subject to constraint

$$\int dx p(x) (x - \mu)^2 = \sigma^2$$

Solution: $p(x)$ proportional to $m(x)$
