

---

# TALENT COURSE II

## LEARNING FROM DATA: BAYESIAN METHODS AND MACHINE LEARNING

---

Lecture 12: More on priors and Maximum Entropy

Daniel Phillips  
Ohio University  
TU Darmstadt  
ExtreMe Matter Institute



OHIO  
UNIVERSITY



TECHNISCHE  
UNIVERSITÄT  
DARMSTADT

TALENT Course II is possible thanks to funding from the STFC

---



---

# BAYES' THEOREM

Thomas Bayes (1701?-1761)

---



$$\text{pr}(A \mid B, I) = \frac{\text{pr}(B \mid A, I)\text{pr}(A \mid I)}{\text{pr}(B \mid I)}$$



# BAYES'THEOREM

Thomas Bayes (1701?-1761)



$$\text{pr}(A | B, I) = \frac{\text{pr}(B | A, I)\text{pr}(A | I)}{\text{pr}(B | I)}$$

<http://www.bayesian-inference.com>



# BAYES' THEOREM

Thomas Bayes (1701?-1761)



$$\text{pr}(A | B, I) = \frac{\text{pr}(B | A, I)\text{pr}(A | I)}{\text{pr}(B | I)}$$

<http://www.bayesian-inference.com>

$$\text{pr}(\text{model} | \text{data}, I) = \frac{\text{pr}(\text{data} | \text{model}, I)\text{pr}(\text{model} | I)}{\text{pr}(\text{data} | I)}$$



# BAYES' THEOREM

Thomas Bayes (1701?-1761)



$$\text{pr}(A | B, I) = \frac{\text{pr}(B | A, I)\text{pr}(A | I)}{\text{pr}(B | I)}$$

<http://www.bayesian-inference.com>

$$\text{pr}(\text{model} | \text{data}, I) = \frac{\text{pr}(\text{data} | \text{model}, I)\text{pr}(\text{model} | I)}{\text{pr}(\text{data} | I)}$$



Posterior



# BAYES' THEOREM

Thomas Bayes (1701?-1761)



<http://www.bayesian-inference.com>

$$\text{pr}(A | B, I) = \frac{\text{pr}(B | A, I)\text{pr}(A | I)}{\text{pr}(B | I)}$$

Likelihood



$$\text{pr}(\text{model} | \text{data}, I) = \frac{\text{pr}(\text{data} | \text{model}, I)\text{pr}(\text{model} | I)}{\text{pr}(\text{data} | I)}$$



Posterior



# BAYES' THEOREM

Thomas Bayes (1701?-1761)



<http://www.bayesian-inference.com>

$$\text{pr}(A | B, I) = \frac{\text{pr}(B | A, I) \text{pr}(A | I)}{\text{pr}(B | I)}$$

Likelihood

Prior



$$\text{pr}(\text{model} | \text{data}, I) = \frac{\text{pr}(\text{data} | \text{model}, I) \text{pr}(\text{model} | I)}{\text{pr}(\text{data} | I)}$$



Posterior



# BAYES' THEOREM

Thomas Bayes (1701?-1761)



<http://www.bayesian-inference.com>

$$\text{pr}(A | B, I) = \frac{\text{pr}(B | A, I) \text{pr}(A | I)}{\text{pr}(B | I)}$$

Likelihood

Prior



$$\text{pr}(\text{model} | \text{data}, I) = \frac{\text{pr}(\text{data} | \text{model}, I) \text{pr}(\text{model} | I)}{\text{pr}(\text{data} | I)}$$

Posterior

Evidence





# BAYES' THEOREM

Thomas Bayes (1701?-1761)



<http://www.bayesian-inference.com>

$$\text{pr}(A | B, I) = \frac{\text{pr}(B | A, I) \text{pr}(A | I)}{\text{pr}(B | I)}$$

Likelihood

Prior



$$\text{pr}(\text{model} | \text{data}, I) = \frac{\text{pr}(\text{data} | \text{model}, I) \text{pr}(\text{model} | I)}{\text{pr}(\text{data} | I)}$$

Posterior

Evidence



**Probability as degree of belief cf. frequentist view**



---

# SUMMARY: PRIOR CHOICE

---



---

# SUMMARY: PRIOR CHOICE

---

- Prior is assigned based on available information



---

# SUMMARY: PRIOR CHOICE

---

- Prior is assigned based on available information
- Invariance arguments/“indifference priors”: priors that are independent of labeling, i.e., uniform.



---

# SUMMARY: PRIOR CHOICE

---

- Prior is assigned based on available information
- Invariance arguments/“indifference priors”: priors that are independent of labeling, i.e., uniform.
- Uniform, but uniform in what?



---

# SUMMARY: PRIOR CHOICE


---

- Prior is assigned based on available information
  - Invariance arguments/“indifference priors”: priors that are independent of labeling, i.e., uniform.
  - Uniform, but uniform in what?
  - Parameterization of priors: independent variables?
-



# SUMMARY: PRIOR CHOICE

- Prior is assigned based on available information
- Invariance arguments/“indifference priors”: priors that are independent of labeling, i.e., uniform.
- Uniform, but uniform in what?
- Parameterization of priors: independence





---

# SUMMARY: PRIOR CHOICE

---

- Prior is assigned based on available information
  - Invariance arguments/“indifference priors”: priors that are independent of labeling, i.e., uniform.
  - Uniform, but uniform in what?
  - Parameterization of priors: independent variables?
-



---

# SUMMARY: PRIOR CHOICE

---

- Prior is assigned based on available information
  - Invariance arguments/“indifference priors”: priors that are independent of labeling, i.e., uniform.
  - Uniform, but uniform in what?
  - Parameterization of priors: independent variables?
  - MaxEnt: allows incorporation of further information, e.g. constraints on mean, variance, etc.
  - MaxEnt: maximizes Shannon information, generates smooth functions
-



---

# SUMMARY: PRIOR CHOICE

---

- Prior is assigned based on available information
- Invariance arguments/“indifference priors”: priors that are independent of labeling, i.e., uniform.
- Uniform, but uniform in what?
- Parameterization of priors: independent variables?
- MaxEnt: allows incorporation of further information, e.g. constraints on mean, variance, etc.
- MaxEnt: maximizes Shannon information, generates smooth functions


Should the likelihood dominate?

---



# PRIOR VS. LIKELIHOOD


- Prior can only be assessed in context of likelihood
- “Robustness” analysis





# BACK TO THE MINI-PROJECT

Schindler, DP (2009); Wesolowski, Klco, Furnstahl, DP, Thapaliya (2016)




$$g(x) = 0.25 + 1.57x + 2.47x^2 + 1.29x^3 + \dots$$



# BACK TO THE MINI-PROJECT

Schindler, DP (2009); Wesolowski, Klco, Furnstahl, DP, Thapaliya (2016)



$$g(x) = 0.25 + 1.57x + 2.47x^2 + 1.29x^3 + \dots$$

- Gaussian prior that encodes naturalness:  $\text{pr}(a_k | \bar{a}, I) \propto \exp\left(-\frac{a_k^2}{2\bar{a}^2}\right)$




---

# RESULTS

---




# RESULTS






# RESULTS

(a) Data set: D1<sub>(5%)</sub>  
Uniform prior



(b) Data set: D1<sub>(5%)</sub>  
Naturalness prior





# RESULTS

**Table 3**

Fit results for Bayesian approach with  $R = 1$ ,  $x_{\max} = 1/\pi$  and  $c = 0.05$

$M$	$\log[\text{pr}(\langle \mathbf{a} \rangle   D_1, M, R)]$	$a_0$	$a_1$	$a_2$
2	12.00	$0.228 \pm 0.018$	$2.06 \pm 0.25$	$1.60 \pm 0.78$
3	11.25	$0.230 \pm 0.018$	$2.04 \pm 0.25$	$1.50 \pm 0.79$
4	10.35	$0.230 \pm 0.018$	$2.04 \pm 0.25$	$1.49 \pm 0.80$
5	9.43	$0.230 \pm 0.018$	$2.04 \pm 0.25$	$1.49 \pm 0.80$
6	8.51	$0.230 \pm 0.018$	$2.04 \pm 0.25$	$1.49 \pm 0.80$
7	7.60	$0.230 \pm 0.018$	$2.04 \pm 0.25$	$1.49 \pm 0.80$


**Table 4**

Fit results for Bayesian approach with  $R = 5$ ,  $x_{\max} = 1/\pi$  and  $c = 0.05$

$M$	$\log[\text{pr}(\langle \mathbf{a} \rangle   D_1, M, R)]$	$a_0$	$a_1$	$a_2$
2	9.62	$0.248 \pm 0.023$	$1.63 \pm 0.39$	$3.15 \pm 1.27$
3	7.10	$0.247 \pm 0.024$	$1.65 \pm 0.45$	$2.98 \pm 2.32$
4	4.57	$0.247 \pm 0.024$	$1.64 \pm 0.46$	$2.98 \pm 2.39$
5	2.04	$0.247 \pm 0.024$	$1.64 \pm 0.46$	$2.98 \pm 2.39$
6	-0.488	$0.247 \pm 0.024$	$1.64 \pm 0.46$	$2.98 \pm 2.39$
7	-3.02	$0.247 \pm 0.024$	$1.64 \pm 0.46$	$2.98 \pm 2.39$



# RELAXATION PLOT





---

# FIVE LEVELS OF PRIOR

<https://github.com/stan-dev/stan/wiki/Prior-Choice-Recommendations>

---

- Flat prior
  - Super-vague, but proper, prior:  $N(0, 1,000,000)$
  - (Very) weakly informative prior:  $N(0, 10)$
  - Weak informative prior:  $N(0, 1)$ : enough information to regularize
  - Specific informative prior:  $N(0.4, .0.2)$
-



---

# FIVE LEVELS OF PRIOR

<https://github.com/stan-dev/stan/wiki/Prior-Choice-Recommendations>

---

- Flat prior
  - Super-vague, but proper, prior:  $N(0, 1,000,000)$
  - (Very) weakly informative prior:  $N(0, 10)$
  - Weak informative prior:  $N(0, 1)$ : enough information to regularize
  - Specific informative prior:  $N(0.4, .0.2)$
  - If you have to choose: weakly informative better than fully informative.
  - Tails: maybe use t-distributions rather than Gaussians
  - When using informative priors, be explicit about every choice!
  - Conjugate priors
-



---

# MOMENTS → FUNCTIONS WITH MAXENT

---



---

# MOMENTS $\rightarrow$ FUNCTIONS WITH MAXENT

---

- Consider a function  $f(x)$  that is positive definite and defined on  $[0,1]$



---

# MOMENTS $\rightarrow$ FUNCTIONS WITH MAXENT

---

- Consider a function  $f(x)$  that is positive definite and defined on  $[0,1]$
- Suppose we know  $N$  moments of the function.



---

# MOMENTS → FUNCTIONS WITH MAXENT

---

- Consider a function  $f(x)$  that is positive definite and defined on  $[0,1]$
- Suppose we know  $N$  moments of the function.

Define

$$S[f] = - \int dx \left( f(x) \log [f(x)] - f(x) \right)$$



---

# MOMENTS $\rightarrow$ FUNCTIONS WITH MAXENT

---

- Consider a function  $f(x)$  that is positive definite and defined on  $[0,1]$
- Suppose we know  $N$  moments of the function.

Define

$$S[f] = - \int dx \left( f(x) \log [f(x)] - f(x) \right)$$



---

# MOMENTS $\rightarrow$ FUNCTIONS WITH MAXENT

---

- Consider a function  $f(x)$  that is positive definite and defined on  $[0,1]$
- Suppose we know  $N$  moments of the function.

Define

$$S[f] = - \int dx \left( f(x) \log [f(x)] - f(x) \right)$$

Maximize  $S$  subject to:

$$\int dx x^j f(x) = \mu_j; j = 0, \dots, N$$

---



---

# MOMENTS → FUNCTIONS WITH MAXENT

---

- Consider a function  $f(x)$  that is positive definite and defined on  $[0,1]$
- Suppose we know  $N$  moments of the function.

Define

$$S[f] = - \int dx \left( f(x) \log [f(x)] - f(x) \right)$$

Maximize  $S$  subject to:

$$\int dx x^j f(x) = \mu_j; j = 0, \dots, N$$

$$Q[f; \{\lambda_j\}] = S[f] + \sum_{j=0}^N \lambda_j \left( \mu_j - \int dx x^j f(x) \right)$$

---



---

# MEAD & PAPANICOLAOU FORMULATION

Mead & Papanicolaou, JMP (1984)

---



---

# MEAD & PAPANICOLAOU FORMULATION

Mead & Papanicolaou, JMP (1984)

---

- Solution:  $f(x) = \exp \left( - \sum_{j=0}^N \lambda_j x^j \right)$



---

# MEAD & PAPANICOLAOU FORMULATION

Mead & Papanicolaou, JMP (1984)

---

- Solution:  $f(x) = \exp \left( - \sum_{j=0}^N \lambda_j x^j \right)$
- Define:  $Z = e^{\lambda_0} = \int dx \exp \left( - \sum_{j=1}^N \lambda_j x^j \right)$



---

# MEAD & PAPANICOLAOU FORMULATION

Mead & Papanicolaou, JMP (1984)

---

- Solution:  $f(x) = \exp \left( - \sum_{j=0}^N \lambda_j x^j \right)$
- Define:  $Z = e^{\lambda_0} = \int dx \exp \left( - \sum_{j=1}^N \lambda_j x^j \right)$
- Introduce:  $\Gamma(\lambda_1, \dots, \lambda_N) = \log Z + \sum_{j=1}^N \mu_j \lambda_j$



---

# MEAD & PAPANICOLAOU FORMULATION

Mead & Papanicolaou, JMP (1984)

---

- Solution:  $f(x) = \exp \left( - \sum_{j=0}^N \lambda_j x^j \right)$
  - Define:  $Z = e^{\lambda_0} = \int dx \exp \left( - \sum_{j=1}^N \lambda_j x^j \right)$
  - Introduce:  $\Gamma(\lambda_1, \dots, \lambda_N) = \log Z + \sum_{j=1}^N \mu_j \lambda_j$
  - $\Gamma$  is everywhere convex, and its stationary points are solutions of the moment equations.
-



# MEAD & PAPANICOLAOU FORMULATION

Mead & Papanicolaou, JMP (1984)

- Solution:  $f(x) = \exp \left( - \sum_{j=0}^N \lambda_j x^j \right)$
- Define:  $Z = e^{\lambda_0} = \int dx \exp \left( - \sum_{j=1}^N \lambda_j x^j \right)$
- Introduce:  $\Gamma(\lambda_1, \dots, \lambda_N) = \log Z + \sum_{j=1}^N \mu_j \lambda_j$
- $\Gamma$  is everywhere convex, and its stationary points are solutions of the moment equations.
- $N=1: \Gamma = \ln[(1 - e^{-\lambda_1})/\lambda_1] + \mu_1 \lambda_1$



# MEAD & PAPANICOLAOU FORMULATION

Mead & Papanicolaou, JMP (1984)

- Solution:  $f(x) = \exp \left( - \sum_{j=0}^N \lambda_j x^j \right)$
- Define:  $Z = e^{\lambda_0} = \int dx \exp \left( - \sum_{j=1}^N \lambda_j x^j \right)$
- Introduce:  $\Gamma(\lambda_1, \dots, \lambda_N) = \log Z + \sum_{j=1}^N \mu_j \lambda_j$
- $\Gamma$  is everywhere convex, and its stationary points are solutions of the moment equations.
- $N=1: \Gamma = \ln[(1 - e^{-\lambda_1})/\lambda_1] + \mu_1 \lambda_1$
- Only possesses a minimum if  $\mu_1 < 1 = \mu_0$ .



# MEAD & PAPANICOLAOU FORMULATION

Mead & Papanicolaou, JMP (1984)

- Solution:  $f(x) = \exp \left( - \sum_{j=0}^N \lambda_j x^j \right)$
  - Define:  $Z = e^{\lambda_0} = \int dx \exp \left( - \sum_{j=1}^N \lambda_j x^j \right)$
  - Introduce:  $\Gamma(\lambda_1, \dots, \lambda_N) = \log Z + \sum_{j=1}^N \mu_j \lambda_j$
  - $\Gamma$  is everywhere convex, and its stationary points are solutions of the moment equations.
  - $N=1: \Gamma = \ln[(1 - e^{-\lambda_1})/\lambda_1] + \mu_1 \lambda_1$
  - Only possesses a minimum if  $\mu_1 < 1 = \mu_0$ .
- Moment  
conditions:  
 $\{\mu_j: j=0, \dots, N\}$   
must be completely  
monotonic



---

# LIKELIHOOD + MAXENT

---



---

# LIKELIHOOD + MAXENT

---

- Suppose we have data  $d_j$ , with errors  $\sigma_j$ , on the function at points  $x_j$ .



---

# LIKELIHOOD + MAXENT

---

- Suppose we have data  $d_j$ , with errors  $\sigma_j$ , on the function at points  $x_j$ .
- We can formulate this as a likelihood for  $f(x_j)$ 's



---

# LIKELIHOOD + MAXENT

---

- Suppose we have data  $d_j$ , with errors  $\sigma_j$ , on the function at points  $x_j$ .
- We can formulate this as a likelihood for  $f(x_j)$ 's
- Combine it with MaxEnt prior: maximize  $\alpha S - \chi^2/2$



---

# LIKELIHOOD + MAXENT

---

- Suppose we have data  $d_j$ , with errors  $\sigma_j$ , on the function at points  $x_j$ .
  - We can formulate this as a likelihood for  $f(x_j)$ 's
  - Combine it with MaxEnt prior: maximize  $\alpha S - \chi^2/2$
  - Non-parametric estimation technique
-



---

# LIKELIHOOD + MAXENT

---

- Suppose we have data  $d_j$ , with errors  $\sigma_j$ , on the function at points  $x_j$ .
  - We can formulate this as a likelihood for  $f(x_j)$ 's
  - Combine it with MaxEnt prior: maximize  $\alpha S - \chi^2/2$
  - Non-parametric estimation technique
  - Likelihood prefers something with lots of wiggles
-



---

# LIKELIHOOD + MAXENT

---

- Suppose we have data  $d_j$ , with errors  $\sigma_j$ , on the function at points  $x_j$ .
  - We can formulate this as a likelihood for  $f(x_j)$ 's
  - Combine it with MaxEnt prior: maximize  $\alpha S - \chi^2/2$
  - Non-parametric estimation technique
  - Likelihood prefers something with lots of wiggles
  - MaxEnt prefers something smooth
-



---

# LIKELIHOOD + MAXENT

---

- Suppose we have data  $d_j$ , with errors  $\sigma_j$ , on the function at points  $x_j$ .
  - We can formulate this as a likelihood for  $f(x_j)$ 's
  - Combine it with MaxEnt prior: maximize  $\alpha S - \chi^2/2$
  - Non-parametric estimation technique
  - Likelihood prefers something with lots of wiggles
  - MaxEnt prefers something smooth
  - Discussion in literature over how Bayesian this is
-



---

# LIKELIHOOD + MAXENT

---

- Suppose we have data  $d_j$ , with errors  $\sigma_j$ , on the function at points  $x_j$ .
  - We can formulate this as a likelihood for  $f(x_j)$ 's
  - Combine it with MaxEnt prior: maximize  $\alpha S - \chi^2/2$
  - Non-parametric estimation technique
  - Likelihood prefers something with lots of wiggles
  - MaxEnt prefers something smooth
  - Discussion in literature over how Bayesian this is
  - Choice of  $\alpha$ : classic maximum entropy, historic maximum entropy, Bryan's method
-



---

# IMAGE RECONSTRUCTION

Grotenhus, Masters' thesis

---



---

# IMAGE RECONSTRUCTION

---

Grotenhus, Masters' thesis






---

# IMAGE RECONSTRUCTION

---

Grotenhus, Masters' thesis





---

# IMAGE RECONSTRUCTION

Grotenhus, Masters' thesis

---




---

# IMAGE RECONSTRUCTION

---

Grotenhus, Masters' thesis





# IMAGE RECONSTRUCTION

Grotenhus, Masters' thesis





---

# INVERTING THE EUCLIDEAN RESPONSE

---

Lovato, Gandolfi, Carlson, Pieper, Schiavilla, PRC (2015)

Jarrell & Gubernatis, Phys. Rep. (1996)

---



---

# INVERTING THE EUCLIDEAN RESPONSE

---

Lovato, Gandolfi, Carlson, Pieper, Schiavilla, PRC (2015)

Jarrell & Gubernatis, Phys. Rep. (1996)

- Theorist calculates: 
$$\frac{E_{\alpha\beta}(q, \tau)}{C_{\alpha\beta}(q)} = \frac{\langle 0 | O_{\alpha}^{\dagger}(\mathbf{q}) e^{-(H-E_0)\tau} O_{\beta}(\mathbf{q}) | 0 \rangle}{\langle 0 | e^{-(H-E_0)\tau} | 0 \rangle}$$



# INVERTING THE EUCLIDEAN RESPONSE

Lovato, Gandolfi, Carlson, Pieper, Schiavilla, PRC (2015)

Jarrell & Gubernatis, Phys. Rep. (1996)

- Theorist calculates:  $\frac{E_{\alpha\beta}(q, \tau)}{C_{\alpha\beta}(q)} = \frac{\langle 0 | O_{\alpha}^{\dagger}(\mathbf{q}) e^{-(H-E_0)\tau} O_{\beta}(\mathbf{q}) | 0 \rangle}{\langle 0 | e^{-(H-E_0)\tau} | 0 \rangle}$
- Experimentalist measures:

$$R_{\alpha\beta}(q, \omega) \propto \sum_f \delta(\omega + E_0 - E_f) \langle 0 | O_{\alpha}(\mathbf{q}) | f \rangle \langle f | O_{\beta}(\mathbf{q}) | 0 \rangle$$



# INVERTING THE EUCLIDEAN RESPONSE

Lovato, Gandolfi, Carlson, Pieper, Schiavilla, PRC (2015)

Jarrell & Gubernatis, Phys. Rep. (1996)

- Theorist calculates: 
$$\frac{E_{\alpha\beta}(q, \tau)}{C_{\alpha\beta}(q)} = \frac{\langle 0 | O_{\alpha}^{\dagger}(\mathbf{q}) e^{-(H-E_0)\tau} O_{\beta}(\mathbf{q}) | 0 \rangle}{\langle 0 | e^{-(H-E_0)\tau} | 0 \rangle}$$

- Experimentalist measures:

$$R_{\alpha\beta}(q, \omega) \propto \sum_f \delta(\omega + E_0 - E_f) \langle 0 | O_{\alpha}(\mathbf{q}) | f \rangle \langle f | O_{\beta}(\mathbf{q}) | 0 \rangle$$

- Relationship: 
$$E_{\alpha\beta}(q, \tau) = C_{\alpha\beta}(\tau) \int_{\omega_{th}}^{\infty} d\omega e^{-\tau\omega} R_{\alpha\beta}(q, \omega)$$



# INVERTING THE EUCLIDEAN RESPONSE

Lovato, Gandolfi, Carlson, Pieper, Schiavilla, PRC (2015)

Jarrell & Gubernatis, Phys. Rep. (1996)

- Theorist calculates:  $\frac{E_{\alpha\beta}(q, \tau)}{C_{\alpha\beta}(q)} = \frac{\langle 0 | O_{\alpha}^{\dagger}(\mathbf{q}) e^{-(H-E_0)\tau} O_{\beta}(\mathbf{q}) | 0 \rangle}{\langle 0 | e^{-(H-E_0)\tau} | 0 \rangle}$
- Experimentalist measures:

$$R_{\alpha\beta}(q, \omega) \propto \sum_f \delta(\omega + E_0 - E_f) \langle 0 | O_{\alpha}(\mathbf{q}) | f \rangle \langle f | O_{\beta}(\mathbf{q}) | 0 \rangle$$

- Relationship:  $E_{\alpha\beta}(q, \tau) = C_{\alpha\beta}(\tau) \int_{\omega_{th}}^{\infty} d\omega e^{-\tau\omega} R_{\alpha\beta}(q, \omega)$
- Why not just invert the Laplace transform?



# INVERTING THE EUCLIDEAN RESPONSE

Lovato, Gandolfi, Carlson, Pieper, Schiavilla, PRC (2015)

Jarrell & Gubernatis, Phys. Rep. (1996)

- Theorist calculates: 
$$\frac{E_{\alpha\beta}(q, \tau)}{C_{\alpha\beta}(q)} = \frac{\langle 0 | O_{\alpha}^{\dagger}(\mathbf{q}) e^{-(H-E_0)\tau} O_{\beta}(\mathbf{q}) | 0 \rangle}{\langle 0 | e^{-(H-E_0)\tau} | 0 \rangle}$$

- Experimentalist measures:

$$R_{\alpha\beta}(q, \omega) \propto \sum_f \delta(\omega + E_0 - E_f) \langle 0 | O_{\alpha}(\mathbf{q}) | f \rangle \langle f | O_{\beta}(\mathbf{q}) | 0 \rangle$$

- Relationship: 
$$E_{\alpha\beta}(q, \tau) = C_{\alpha\beta}(\tau) \int_{\omega_{th}}^{\infty} d\omega e^{-\tau\omega} R_{\alpha\beta}(q, \omega)$$

- Why not just invert the Laplace transform?

- At large positive frequencies the kernel is exponentially small, so large  $\omega$  features of  $R(\omega)$  depend on subtle features of  $E(\tau)$ .



# RESULTS

