

ON THE JAMES AND HILTON–MILNOR SPLITTINGS, & THE METASTABLE EHP SEQUENCE

SANATH DEVALAPURKAR AND PETER HAINE

ABSTRACT. This note provides modern proofs of some classical results in algebraic topology, such as the James Splitting, the Hilton–Milnor Splitting, and the metastable EHP sequence. We prove fundamental splitting results

$$\Sigma\Omega\Sigma X \simeq \Sigma X \vee (X \wedge \Sigma\Omega\Sigma X) \quad \text{and} \quad \Omega(X \vee Y) \simeq \Omega X \times \Omega Y \times \Omega\Sigma(\Omega X \wedge \Omega Y)$$

in the maximal generality of an ∞ -category with finite limits and pushouts in which pushouts squares remain pushouts after basechange along an arbitrary morphism (i.e., Mather’s Second Cube Lemma holds). For connected objects, these imply the classical James and Hilton–Milnor Splittings. Moreover, working in this generality shows that the James and Hilton–Milnor splittings hold in many new contexts, for example in: elementary ∞ -topoi, profinite spaces, and motivic spaces over arbitrary base schemes. The splitting result in this last context extend Wickelgren and Williams’ splitting result for motivic spaces over a perfect field. We also give two proofs of the metastable EHP sequence in the setting of ∞ -topoi: the first is a new, non-computational proof that only utilizes basic connectedness estimates involving the James filtration and the Blakers–Massey Theorem, while the second reduces to the classical computational proof.

CONTENTS

1. Introduction	1
1.1. Linear overview	3
1.2. Notation & background	4
2. The James Splitting	5
2.1. Universal pushouts and Mather’s Second Cube Lemma	5
2.2. Statement of the James Splitting & Consequences	7
2.3. Proof of the James Splitting	8
2.4. Ganea’s Lemma	11
3. The Hilton–Milnor Splitting	12
3.1. Reminder on group objects & the Splitting Lemma	13
4. Connectedness & the James Splitting	14
4.1. Connectedness and the Blakers–Massey Theorem	15
4.2. The infinite James and Hilton–Milnor Splittings	18
5. The James filtration & metastable EHP sequence	21
5.1. The James filtration	21
5.2. Splitting the James filtration	22
5.3. Proofs of the metastable EHP sequence	25
References	27

1. INTRODUCTION

A classical result of James shows that given a pointed connected space X , the homotopy type $\Sigma\Omega\Sigma X$ given by suspending the loop space on the suspension of X splits as a wedge sum

$$(1.1) \quad \Sigma\Omega\Sigma X \simeq \bigvee_{i \geq 1} \Sigma X^{\wedge i}$$

of suspensions of smash powers of X [7; 19]. Hilton and Milnor proved a related splitting result [15; 16; 24, Theorem 3]: given pointed connected spaces X and Y , they showed that there is a homotopy equivalence

$$(1.2) \quad \Omega\Sigma(X \vee Y) \simeq \Omega\Sigma X \times \Omega\Sigma Y \times \Omega\Sigma \left(\bigvee_{i,j \geq 1} X^{\wedge i} \wedge Y^{\wedge j} \right).$$

In the classical setting, these splitting results follow from combining a connectedness argument using the hypothesis that X and Y are connected with the following more fundamental splittings: given any pointed spaces X and Y , there are natural equivalences

$$(1.3) \quad \Sigma\Omega\Sigma X \simeq \Sigma X \vee (X \wedge \Sigma\Omega\Sigma X) \quad \text{and} \quad \Omega(X \vee Y) \simeq \Omega X \times \Omega Y \times \Omega\Sigma(\Omega X \wedge \Omega Y).$$

The first objective of this note is to provide clear, modern, and non-computational proofs of these fundamental splittings (1.3). The only property particular to the ∞ -category of spaces that our proofs utilize is *Mather's Second Cube Lemma* [22]; this asserts that pushout squares remain pushouts after basechange along an arbitrary morphism (see §2.1). Hence these ‘fundamental’ James and Hilton–Milnor Splittings hold in any ∞ -category where we can make sense of suspensions, loops, wedge sums, and smash products, and have access to Mather's Second Cube Lemma:

Theorem 1.4 (Fundamental James Splitting; Theorem 2.10). *Let \mathcal{X} be an ∞ -category with finite limits and pushouts, and assume that Mather's Second Cube Lemma holds in \mathcal{X} . Then for every pointed object $X \in \mathcal{X}_*$, there is a natural equivalence*

$$\Sigma\Omega\Sigma X \simeq \Sigma X \vee (X \wedge \Sigma\Omega\Sigma X).$$

Moreover, for each integer $n \geq 1$ then there is a natural equivalence

$$\Sigma\Omega\Sigma X \simeq \left(\bigvee_{1 \leq i \leq n} \Sigma X^{\wedge i} \right) \vee (X^{\wedge n} \wedge \Sigma\Omega\Sigma X).$$

In general, the infinite splitting $\Sigma\Omega\Sigma X \simeq \bigvee_{i \geq 1} \Sigma X^{\wedge i}$ need not hold; roughly speaking, the problem is that if X is not connected, then $(X^{\wedge n} \wedge \Sigma\Omega\Sigma X)$ need not vanish as $n \rightarrow \infty$. However, there is always a natural map $\bigvee_{i \geq 1} \Sigma X^{\wedge i} \rightarrow \Sigma\Omega\Sigma X$.

Theorem 1.5 (Fundamental Hilton–Milnor Splitting; Theorem 3.1). *Let \mathcal{X} be an ∞ -category with finite limits and pushouts, and assume that Mather's Second Cube Lemma holds in \mathcal{X} . Then for every pair of pointed objects $X, Y \in \mathcal{X}_*$ there is an natural equivalence*

$$\Omega(X \vee Y) \simeq \Omega X \times \Omega Y \times \Omega\Sigma(\Omega X \wedge \Omega Y).$$

In §4.2 we explain in what generality the the infinite James Splitting (1.1) and Hilton–Milnor Splitting (1.2) hold, and how to deduce them from Theorems 1.4 and 1.5.

It might seem that knowing that the James and Hilton–Milnor Splittings in this level of generality is of dubious advantage; the settings in which one is most likely to want to apply these splittings are the ∞ -category **Spc** of spaces (where the results are already known), or an ∞ -topos (where the results follows immediately from the results for **Spc**; see §4.2). However, algebraic geometry provides an example that does not immediately follow from the result for spaces: motivic spaces. The obstruction is that the ∞ -category of motivic spaces over a scheme is not an ∞ -topos; since motivic localization almost never commutes with taking loops, knowing the James and Hilton–Milnor Splittings in the ∞ -topos of Nisnevich sheaves does not allow one to deduce that they hold in motivic spaces.

Wickelgren and Williams used the *James filtration* to prove that the infinite James Splitting (1.1) holds for \mathbf{A}^1 -connected motivic spaces over a perfect field [35, Theorem 1.5]. The reason for the restriction on the base is because their proof relies on Morel's unstable \mathbf{A}^1 -connectivity Theorem [25, Theorems 5.46 and 6.1], which implies that motivic localization commutes with loops [3, Theorem 2.4.1; 25, Theorem 6.46]. However, the unstable \mathbf{A}^1 -connectivity property does not hold for higher-dimensional bases [3, Remark 3.3.5; 4], so a different method is needed if one wants to prove James and Hilton–Milnor Splittings for motivic spaces over more general bases. This is where our generalization pays off: work of Hoyois [18,

Proposition 3.15] shows that, in particular, Mather’s Second Cube Lemma holds in motivic spaces over an *arbitrary* base scheme. Therefore, [Theorems 1.4](#) and [1.5](#) apply in this setting. We use these splittings to give a description of the motivic space $\Sigma\Omega\mathbf{P}^1$ in terms of wedges of motivic spheres $S^{i+1,i}$ ([Example 2.14](#)), and also give a new description of $\Omega\Sigma(\mathbf{P}^1 \setminus \{0, 1, \infty\})$ ([Example 3.3](#)). Over a perfect field, we give a new decomposition of $\Omega\Sigma^2(\mathbf{P}^1 \setminus \{0, 1, \infty\})$ in terms of motivic spheres of the form $S^{2m+1,m}$ ([Example 4.25](#)).

The second goal of this note is to give a modern construction of the metastable EHP sequence in an ∞ -topos \mathcal{X} . For every pointed connected object $X \in \mathcal{X}_*$, the James Splitting provides *Hopf maps*

$$h_n : \Omega\Sigma X \rightarrow \Omega\Sigma(X^{\wedge n}).$$

There is also a *James filtration* $\{J_m(X)\}_{m \geq 0}$ on $\Omega\Sigma X$, and, moreover, the composite

$$(1.6) \quad J_{n-1}(X) \longrightarrow \Omega\Sigma X \xrightarrow{h_n} \Omega\Sigma X^{\wedge n}$$

is trivial. The sequence (1.6) is not a fiber sequence in general¹, but is in the *metastable range*:

Theorem 1.7 (metastable EHP sequence; [Theorem 5.19](#)). *Let \mathcal{X} be an ∞ -topos, $k \geq 0$ an integer, and $X \in \mathcal{X}_*$ a pointed k -connected object. Then for each integer $n \geq 1$, the morphism $J_{n-1}(X) \rightarrow \text{fib}(h_n)$ is $((n+1)(k+1)-3)$ -connected.*

We note here that a morphism is m -connected in our terminology if and only if it is $(m+1)$ -connected in the classical terminology (see [Warning 4.9](#)).

We provide two proofs of [Theorem 1.7](#). The first proof is new and non-computational; it only makes use of some basic connectedness estimates involving the James filtration and the Blakers–Massey Theorem. In the second proof we simply note that [Theorem 1.7](#) for a general ∞ -topos follows immediately from the claim for the ∞ -topos of spaces. In the case of spaces, we provide a computational proof; we include this second proof because we were unable to find the computational proof we were familiar with in the literature.

1.1. Linear overview. We have written this note with two audiences in mind: the student interested in seeing proofs of [Theorems 1.4](#), [1.5](#) and [1.7](#) in the classical setting of spaces, and the expert homotopy theorist interested in applying these results to more general contexts such as motivic spaces or profinite spaces. The student can always take \mathcal{X} to be the ∞ -category of spaces, and the expert can safely skip the background sections provided for the student. We also note that this text should still be accessible to the reader familiar with homotopy (co)limits but unfamiliar with higher categories, since all we use in our proofs are basic manipulations of homotopy (co)limits.

[Section 2](#) is dedicated to proving [Theorem 1.4](#). In [§2.1](#), we provide background on Mather’s Second Cube Lemma and the universality of pushouts. In [§2.2](#), we provide a proof of the James Splitting. Our proof is roughly the same as proofs presented elsewhere [[17](#); [32](#), §17.2; [36](#)], but it seems that the generality of the argument we present here is not very well-known.

[Section 3](#) provides a quick proof of [Theorem 1.5](#). Again, shadows of the proof we provide appear in the literature [[11](#); [12](#), §2 & 3; [32](#), §17.8], but it seems that the generality of the proof has not been completely internalized by the community. As an application, we use work of Wickelgren [[34](#), Corollary 3.2] to give a new description of the motivic space $\Omega\Sigma(\mathbf{P}^1 \setminus \{0, 1, \infty\})$ ([Example 3.2](#)).

[Section 4](#) explains how to use a connectedness argument to show prove the infinite James Splitting (1.1) and Hilton–Milnor Splitting (1.2) for pointed connected objects of an ∞ -topos, and for pointed \mathbf{A}^1 -connected motivic spaces over a perfect field. In [§4.1](#), we begin by recalling the basics of connectedness and the Blakers–Massey Theorem in an ∞ -topos; this material is also instrumental in our proof of [Theorem 1.7](#). [Subsection 4.2](#) presents the connectedness argument to deduce the infinite splittings and defines the Hopf maps appearing in [Theorem 1.7](#). As an application we give a new description of the motivic space $\Omega\Sigma^2(\mathbf{P}^1 \setminus \{0, 1, \infty\})$ over a perfect field ([Example 4.25](#)).

[Section 5](#) is dedicated to proving [Theorem 1.7](#). In [§5.1](#), we provide the background on the James filtration needed to understand the statement of [Theorem 1.7](#), as well as some connectedness estimates we need to prove [Theorem 1.7](#). In [§5.2](#), we give a refinement of the James Splitting in terms of the James filtration.

¹When \mathcal{X} is the ∞ -category of spaces and X is a sphere, James and Toda proved that, roughly, the sequence (1.6) becomes a fiber sequence after p -localization. See [[20](#); [21](#); [33](#)] for a precise statement.

In § 5.3, we first provide a proof of [Theorem 1.7](#) using the Blakers–Massey Theorem (which we have not seen elsewhere), and then record for posterity what we imagine is the standard computational proof of [Theorem 1.7](#).

Acknowledgements. We are grateful to Tom Bachmann for pointing out that colimits are universal in motivic spaces. We are indebted to André Joyal for noticing some errors in an earlier version as well as alerting us to our misuse of terminology regarding connectedness. We are grateful to Tomer Schlank for explaining why the infinite version of the James Splitting requires X to be connected, and for explaining some subtleties around [Lemma 2.17](#). We thank Elden Elmanto, Marc Hoyois, and Dylan Wilson for helpful conversations. The second-named author gratefully acknowledges support from both the MIT Dean of Science Fellowship and the National Science Foundation Graduate Research Fellowship under Grant #112237.

1.2. Notation & background. In this subsection we set the basic notational conventions that we use throughout this note as well as recall a bit of relevant background.

Notation 1.8. Let \mathcal{X} be an ∞ -category. If \mathcal{X} has a terminal object, we write $*$ $\in \mathcal{X}$ for the terminal object and \mathcal{X}_* for the ∞ -category of pointed objects in \mathcal{X} . If \mathcal{X}_* has coproducts and $X, Y \in \mathcal{X}_*$, we write $X \vee Y$ for the coproduct of X and Y in \mathcal{X}_* . If \mathcal{X}_* has coproducts and products, note that there is a natural comparison morphism $X \vee Y \rightarrow X \times Y$ induced by the morphisms

$$(\mathrm{id}_X, *) : X \rightarrow X \times Y \quad \text{and} \quad (*, \mathrm{id}_Y) : Y \rightarrow X \times Y.$$

We say that a morphism $f : X \rightarrow Y$ in \mathcal{X}_* is *null* if f factors through the zero object $*$.

Recollection 1.9. Let \mathcal{X} be an ∞ -category with finite products and pushouts, and $X, Y \in \mathcal{X}_*$ pointed objects of \mathcal{X} . The *smash product* $X \wedge Y$ of X and Y is the cofiber

$$\begin{array}{ccc} X \vee Y & \longrightarrow & X \times Y \\ \downarrow & \lrcorner & \downarrow \\ * & \longrightarrow & X \wedge Y \end{array}$$

of the comparison morphism $X \vee Y \rightarrow X \times Y$.

Recollection 1.10. Let \mathcal{X} be an ∞ -category with pushouts and a terminal object. The *suspension* of an object $X \in \mathcal{X}$ is the pushout

$$\begin{array}{ccc} X & \longrightarrow & * \\ \downarrow & \lrcorner & \downarrow \\ * & \longrightarrow & \Sigma X. \end{array}$$

Recollection 1.11. Let \mathcal{X} be an ∞ -category with finite limits. The *loop object* of a pointed object $X \in \mathcal{X}_*$ is the pullback

$$\begin{array}{ccc} \Omega X & \longrightarrow & * \\ \downarrow \lrcorner & & \downarrow \\ * & \longrightarrow & X \end{array}$$

in \mathcal{X}_* .

We also make repeated use of the following easy fact. The unfamiliar reader should consult [5, §2; 27].

Lemma 1.12. *Let \mathcal{X} be an ∞ -category with pushouts and*

$$(1.13) \quad \begin{array}{ccccc} X_1 & \longleftarrow & X_0 & \longrightarrow & X_2 \\ \uparrow & & \uparrow & & \uparrow \\ W_1 & \longleftarrow & W_0 & \longrightarrow & W_2 \\ \downarrow & & \downarrow & & \downarrow \\ Y_1 & \longleftarrow & Y_0 & \longrightarrow & Y_2 \end{array}$$

a commutative diagram in \mathcal{X} . Then the colimit of the diagram (1.13) exists and is equivalent to both of the following two iterated pushouts:

(1.12.1) *Form the pushout of the rows of (1.13), then take the pushout of the resulting span*

$$X_1 \overset{X_0}{\sqcup} X_2 \longleftarrow W_1 \overset{W_0}{\sqcup} W_2 \longrightarrow Y_1 \overset{Y_0}{\sqcup} Y_2.$$

(1.12.2) *Form the pushout of the columns of (1.13), then take the pushout of the resulting span*

$$X_1 \overset{W_1}{\sqcup} Y_1 \longleftarrow X_0 \overset{W_0}{\sqcup} Y_0 \longrightarrow X_2 \overset{W_2}{\sqcup} Y_2.$$

2. THE JAMES SPLITTING

In this section, we present a proof of the James Splitting which holds in any ∞ -category with finite limits and pushouts, where pushout squares remain pushouts after basechange along an arbitrary morphism. The argument we give roughly follows the argument Hopkins gave in his course on stable homotopy theory in the setting of spaces [17, Lecture 4, §3]; Hopkins attributes this proof to James [19; 20; 21] and Ganea [10].

2.1. Universal pushouts and Mather’s Second Cube Lemma. The key property utilized in the proofs we present of the James and Hilton–Milnor Splittings is that pushout squares are preserved by arbitrary basechange. This implies that, in particular, the James and Hilton–Milnor Splittings hold in any ∞ -topos, but also in other situations (such as motivic spaces). In this subsection, we provide the categorical context that we work in for the rest of the paper and give a convenient reformulation of the stability of pullbacks under basechange in terms of Mather’s Second Cube Lemma (Lemma 2.6).

Recollection 2.1. Let \mathcal{I} be an ∞ -category and let \mathcal{X} be an ∞ -category with pullbacks and all \mathcal{I} -shaped colimits. We say that *\mathcal{I} -shaped colimits in \mathcal{X} are universal* if \mathcal{I} -shaped colimits in \mathcal{X} are stable under pullback along any morphism. That is, for every diagram $F : \mathcal{I} \rightarrow \mathcal{X}$ and pair of morphisms $\text{colim}_{i \in \mathcal{I}} F(i) \rightarrow Z$ and $Y \rightarrow Z$ in \mathcal{X} , the natural morphism

$$\text{colim}_{i \in \mathcal{I}} (F(i) \times_Z Y) \rightarrow \left(\text{colim}_{i \in \mathcal{I}} F(i) \right) \times_Z Y$$

is an equivalence.

Example 2.2. Let $0 \leq n \leq \infty$, and let \mathcal{X} be an n -topos. One of the Giraud–Lurie axioms for n -topoi guarantees that all small colimits in \mathcal{X} are universal [HTT, Theorem 6.1.0.6 & Proposition 6.4.1.5]. In particular, small colimits in the category **Set** of sets and the ∞ -category **Spc** of spaces are universal.

Example 2.3. Let \mathcal{X} be an *elementary* ∞ -topos in the sense of [29, Definition 3.5]. Then all colimits that exist in \mathcal{X} are universal [29, Corollary 3.12]. In particular, finite colimits are universal in \mathcal{X} .

Example 2.4 (motivic spaces). Let S be a scheme. The ∞ -category $\mathbf{H}(S)$ of *motivic spaces over S* is defined as the \mathbf{A}^1 -localization of the ∞ -topos $\text{Sh}_{\text{nis}}(\mathbf{Sm}_S)$ of sheaves of spaces on the category \mathbf{Sm}_S of smooth schemes of finite type over S equipped with the Nisnevich topology. Concretely, $\mathbf{H}(S)$ is the full subcategory of $\text{Sh}_{\text{nis}}(\mathbf{Sm}_S)$ spanned by those Nisnevich sheaves \mathcal{F} on \mathbf{Sm}_S with the property that for every smooth S -scheme X , the projection $\text{pr}_1 : X \times_S \mathbf{A}_S^1 \rightarrow X$ induces an equivalence

$$\text{pr}_1^* : \mathcal{F}(X) \xrightarrow{\sim} \mathcal{F}(X \times_S \mathbf{A}_S^1).$$

The inclusion $\mathbf{H}(S) \subset \mathbf{Sh}_{\text{nis}}(\mathbf{Sm}_S)$ admits a left adjoint $L_{\text{mot}} : \mathbf{Sh}_{\text{nis}}(\mathbf{Sm}_S) \rightarrow \mathbf{H}(S)$ called *motivic localization*. Motivic localization preserves finite products, but not all finite limits. Moreover, the ∞ -category $\mathbf{H}(S)$ is not an ∞ -topos (see [31, Remark 3.5; 28, §4.3]), and it is not immediately clear from the construction if any colimits are universal in $\mathbf{H}(S)$. Nonetheless, Hoyois has shown that all small colimits are universal in $\mathbf{H}(S)$ [18, Proposition 3.15].

Example 2.5 (profinite spaces). We say that a space X is π -finite if X is truncated, has finitely many connected components, and $\pi_i(X, x)$ is finite for every integer $i \geq 1$ and point $x \in X$. Write $\mathbf{Spc}_\pi \subset \mathbf{Spc}$ for the full subcategory spanned by the π -finite spaces and $\text{Pro}(\mathbf{Spc}_\pi)$ for the ∞ -category of *profinite spaces*. Infinite coproducts in $\text{Pro}(\mathbf{Spc}_\pi)$ are not universal [SAG, Warning E.6.0.9], however, finite colimits and geometric realizations of simplicial objects are universal in $\text{Pro}(\mathbf{Spc}_\pi)$ [SAG, Theorem E.6.3.1 & Corollary E.6.3.2].

The following result gives a reformulation of what it means for pushouts to be universal in terms of Mather’s Second Cube Lemma, which Mather originally proved in the ∞ -category of spaces [22, Theorem 25]. See [Ker, Tag 011H] for an elegant proof of Mather’s Second Cube Lemma in \mathbf{Spc} .

Lemma 2.6. *Let \mathcal{X} be an ∞ -category with pullbacks and pushouts. The following conditions are equivalent:*

(2.6.1) *Pushouts in \mathcal{X} are universal.*

(2.6.2) *Mather’s Second Cube Lemma holds in \mathcal{X} : Given a commutative cube*

$$\begin{array}{ccccc}
 A_0 & \longrightarrow & A_2 & & \\
 \downarrow & \searrow & \downarrow & \searrow & \\
 & A_1 & \longrightarrow & A_3 & \\
 \downarrow & \downarrow & \downarrow & \downarrow & \\
 B_0 & \longrightarrow & B_2 & & \\
 & \searrow & \downarrow & \searrow & \\
 & B_1 & \longrightarrow & B_3 &
 \end{array}$$

in \mathcal{X} where the bottom horizontal face is a pushout square and all vertical faces are pullback squares, then the top horizontal square is a pushout square.

Proof. The implication (2.6.1) \Rightarrow (2.6.2) is immediate. To see that (2.6.2) \Rightarrow (2.6.1), suppose that we are given a pushout square

$$(2.7) \quad \begin{array}{ccc}
 B_0 & \longrightarrow & B_2 \\
 \downarrow & & \downarrow \\
 B_1 & \longrightarrow & B_3
 \end{array}$$

in \mathcal{X} and morphisms $f : B_3 \rightarrow Z$ and $g : Y \rightarrow Z$ in \mathcal{X} . For each $i \in \{0, 1, 2, 3\}$, define $A_i := B_i \times_Z Y$, so that all the vertical squares in the diagram

$$(2.8) \quad \begin{array}{ccccccc}
 A_0 & \longrightarrow & A_2 & & & & \\
 \downarrow & \searrow & \downarrow & \searrow & & & \\
 & A_1 & \longrightarrow & A_3 & \longrightarrow & Y & \\
 \downarrow & \downarrow & \downarrow & \downarrow & & \downarrow g & \\
 B_0 & \longrightarrow & B_2 & & & & \\
 & \searrow & \downarrow & \searrow & & & \\
 & B_1 & \longrightarrow & B_3 & \xrightarrow{f} & Z &
 \end{array}$$

are pullbacks. Since the bottom horizontal square of the cube in (2.8) is a pushout, (2.6.2) implies that the top horizontal square is also a pushout. Thus the pushout square (2.7) remains a pushout after base change along an arbitrary morphism, as desired. \square

Since the main results of this note are about pointed objects, we make the following mildly abusive convention:

Convention 2.9. We say that an ∞ -category \mathcal{X} has *universal pushouts* if \mathcal{X} has finite limits and pushouts, and pushouts in \mathcal{X} are universal.

2.2. Statement of the James Splitting & Consequences. The James Splitting, originally proven in [19], provides a splitting of the space $\Omega\Sigma X$ after a single suspension. The goal of this subsection is to provide a proof of the James Splitting that only relies on the universality of pushouts and a few elementary computations involving the interaction between forming suspensions, loop objects, and smash products.

Theorem 2.10 (Fundamental James Splitting). *Let \mathcal{X} be an ∞ -category with universal pushouts. For every pointed object $X \in \mathcal{X}_*$, there is a natural equivalence*

$$\Sigma\Omega\Sigma X \simeq \Sigma X \vee \Sigma(X \wedge \Omega\Sigma X).$$

Using the fact that $\Sigma(X \wedge \Omega\Sigma X) \simeq X \wedge \Sigma\Omega\Sigma X$ (Lemma 2.26) and iterating the equivalence of Theorem 2.10, we see:

Corollary 2.11 (Fundamental James Splitting, redux). *Let \mathcal{X} be an ∞ -category with universal pushouts. For each pointed object $X \in \mathcal{X}_*$ and integer $n \geq 1$, then there is a natural equivalence*

$$\Sigma\Omega\Sigma X \simeq \left(\bigvee_{1 \leq i \leq n} \Sigma X^{\wedge i} \right) \vee (X^{\wedge n} \wedge \Sigma\Omega\Sigma X).$$

Notation 2.12. Let \mathcal{X} be an ∞ -category with universal pushouts and $X \in \mathcal{X}_*$ a pointed object. Assume that \mathcal{X}_* has countable coproducts. Passing to the colimit as $n \rightarrow \infty$, the coproduct insertions

$$\bigvee_{i=1}^n \Sigma X^{\wedge i} \rightarrow \Sigma\Omega\Sigma X$$

provided by Corollary 2.11 provide a natural comparison morphism

$$\bigvee_{i=1}^{\infty} \Sigma X^{\wedge i} \rightarrow \Sigma\Omega\Sigma X,$$

which we denote by $c_X : \bigvee_{i=1}^{\infty} \Sigma X^{\wedge i} \rightarrow \Sigma\Omega\Sigma X$.

Warning 2.13. The comparison morphism c_X need not be an equivalence. For example, if $\mathcal{X} = \mathbf{Spc}$ and $X = S^0$ is the 0-sphere, then the map

$$c_{S^0} : \bigvee_{i=1}^{\infty} S^1 \rightarrow \Sigma\Omega\Sigma S^0 \simeq \bigvee_{i \in \mathbf{Z}} S^1$$

is not an equivalence. Even though both the source and target of c_{S^0} are both countable wedges of copies of S^1 , the map c_{S^0} is the summand inclusion induced by the inclusion $\mathbf{Z}_{\geq 1} \subset \mathbf{Z}$.

We analyze when the comparison morphism c_X is an equivalence in §4.2.

Example 2.14. Let S be a scheme. Since colimits are universal in the ∞ -category $\mathbf{H}(S)$ of motivic spaces over S (Example 2.4), Theorem 2.10 implies that for any pointed motivic space $X \in \mathbf{H}(S)_*$ and integer $n \geq 1$, we have S^1 -James Splittings

$$\Sigma\Omega\Sigma X \simeq \left(\bigvee_{1 \leq i \leq n} \Sigma X^{\wedge i} \right) \vee (X^{\wedge n} \wedge \Sigma\Omega\Sigma X).$$

Write \mathbf{G}_m for the multiplicative group scheme over S . Since $\Sigma\mathbf{G}_m \simeq \mathbf{P}_S^1$, setting $X = \mathbf{G}_m$ we see that

$$(2.15) \quad \Sigma\Omega\mathbf{P}_S^1 \simeq \left(\bigvee_{1 \leq i \leq n} \Sigma\mathbf{G}_m^{\wedge i} \right) \vee (\mathbf{G}_m^{\wedge n} \wedge \Sigma\Omega\mathbf{P}_S^1).$$

Using the grading convention

$$S^{a,b} := \mathbf{G}_m^{\wedge b} \wedge (S^1)^{\wedge(a-b)}$$

for motivic spheres, we can rewrite the equivalence (2.15) as

$$\Sigma \Omega \mathbf{P}_S^1 \simeq \left(\bigvee_{1 \leq i \leq n} S^{i+1,i} \right) \vee (S^{n+1,n} \wedge \Omega \mathbf{P}_S^1).$$

Remark 2.16. There is another suspension in motivic homotopy theory, given by smashing with the multiplicative group scheme \mathbf{G}_m . One would like an analogue of the James Splitting in $\mathbf{H}(S)_*$ for \mathbf{G}_m -suspensions. For $S = \text{Spec}(\mathbf{R})$, Betti realization defines a functor

$$\mathbf{H}(\text{Spec}(\mathbf{R})) \rightarrow \mathbf{Spc}_{C_2}$$

to C_2 -spaces which sends S^1 to the circle with trivial C_2 -action, and \mathbf{G}_m to the sign representation circle S^σ . Even though Betti realization is not an equivalence, it closely ties \mathbf{R} -motivic homotopy theory with C_2 -equivariant homotopy theory. In [14], Hill studies the *signed James construction* in C_2 -equivariant unstable homotopy theory, and shows that an analogue of the James Splitting holds for $\Omega^\sigma \Sigma^\sigma X$ after suspending by the regular representation sphere $S^\rho = S^1 \wedge S^\sigma$. This might lead one to hope that there is an analogue of Hill's result in motivic homotopy theory which proves the James Splitting for $\Omega_{\mathbf{G}_m} \Sigma_{\mathbf{G}_m} X$ after \mathbf{P}^1 -suspension; at the moment, we are not aware of such a result.

2.3. Proof of the James Splitting. Before we prove Theorem 2.10, we need a few preliminary results. First, we give a convenient expression for $\Sigma \Omega \Sigma X$ as the cofiber of the projection $\text{pr}_2 : X \times \Omega \Sigma X \rightarrow \Omega \Sigma X$. This expression for $\Sigma \Omega \Sigma X$ is an immediate consequence of the following:

Lemma 2.17. *Let \mathcal{X} be an ∞ -category with universal pushouts. For every pointed object $X \in \mathcal{X}_*$, there exists a natural morphism $\alpha_X : X \times \Omega \Sigma X \rightarrow \Omega \Sigma X$ and a pushout square*

$$\begin{array}{ccc} X \times \Omega \Sigma X & \xrightarrow{\text{pr}_2} & \Omega \Sigma X \\ \alpha_X \downarrow & & \downarrow \ulcorner \\ \Omega \Sigma X & \xrightarrow{\quad} & * \end{array}$$

Proof. Write

$$\begin{array}{ccc} X & \longrightarrow & * \\ \downarrow & & \downarrow i_2 \\ * & \xrightarrow{i_1} & \Sigma X \end{array}$$

for the pushout square defining the suspension ΣX . The definition of ΣX provides an equivalence between the points $i_1, i_2 : * \rightarrow \Sigma X$, hence there are natural pullback squares

$$\begin{array}{ccccc} X \times \Omega \Sigma X & \xrightarrow{\alpha_X} & \Omega X & \longrightarrow & * \\ \downarrow \lrcorner & & \downarrow \lrcorner & & \downarrow i_2 \\ X & \longrightarrow & * & \xrightarrow{i_1} & X \end{array}$$

The claim now follows from Mather's Second Cube Lemma applied to the cube

$$\begin{array}{ccccc}
 X \times \Omega \Sigma X & \xrightarrow{\text{pr}_2} & \Omega \Sigma X & & \\
 \downarrow a_X & \searrow & \downarrow & \searrow & \\
 & \Omega \Sigma X & \xrightarrow{\quad} & * & \\
 \downarrow & \downarrow & \downarrow & \downarrow i_2 & \\
 X & \xrightarrow{\quad} & * & \searrow i_2 & \\
 & \downarrow & \downarrow i_1 & \downarrow & \\
 & * & \xrightarrow{\quad} & \Sigma X & .
 \end{array}$$

□

Warning 2.18. The morphism $a_X : X \times \Omega \Sigma X \rightarrow \Omega \Sigma X$ in Lemma 2.17 cannot generally be identified with the second projection $\text{pr}_2 : X \times \Omega \Sigma X \rightarrow \Omega \Sigma X$. Indeed, since \mathcal{X} is assumed to have universal pushouts, there is a natural pushout square

$$\begin{array}{ccc}
 X \times \Omega \Sigma X & \xrightarrow{\text{pr}_2} & \Omega \Sigma X \\
 \text{pr}_2 \downarrow & \lrcorner & \downarrow \\
 \Omega \Sigma X & \longrightarrow & \Sigma X \times \Omega \Sigma X .
 \end{array}$$

Moreover, the object $\Sigma X \times \Omega \Sigma X$ is not generally terminal in \mathcal{X} .

Corollary 2.19. Let \mathcal{X} be an ∞ -category with universal pushouts. For every pointed object $X \in \mathcal{X}_*$, there is a natural equivalence

$$\text{cofib}(\text{pr}_2 : X \times \Omega \Sigma X \rightarrow \Omega \Sigma X) \simeq \Sigma \Omega \Sigma X .$$

Next, we give a convenient expression for the term $\Sigma(X \wedge \Omega \Sigma X)$ in the James Splitting as the pushout of the span

$$X \xleftarrow{\text{pr}_1} X \times \Omega \Sigma X \xrightarrow{\text{pr}_2} \Omega \Sigma X .$$

Our proof of this appeals to the following fact, which follows immediately from the definitions.

Lemma 2.20. Let \mathcal{X} be an ∞ -category with pushouts and a terminal object, and let $X, Y \in \mathcal{X}_*$ be pointed objects of \mathcal{X} . Then the square

$$\begin{array}{ccc}
 X \vee Y & \xrightarrow{(*, \text{id}_Y)} & Y \\
 (\text{id}_X, *) \downarrow & & \downarrow \\
 X & \longrightarrow & *
 \end{array}$$

is a pushout square.

Proposition 2.21. Let \mathcal{X} be an ∞ -category with finite limits and pushouts. Then for every pair of pointed objects $X, Y \in \mathcal{X}$, there is a pushout square

$$\begin{array}{ccc}
 X \times Y & \xrightarrow{\text{pr}_2} & Y \\
 \text{pr}_1 \downarrow & & \downarrow \\
 X & \longrightarrow & \Sigma(X \wedge Y) ,
 \end{array}$$

where the morphisms $X \rightarrow \Sigma(X \wedge Y)$ and $Y \rightarrow \Sigma(X \wedge Y)$ are null.

Proof. Let C denote the pushout $X \sqcup^{X \times Y} Y$; we desire to show that $C \simeq \Sigma(X \wedge Y)$. We apply [Lemma 1.12](#) to the commutative diagram

$$(2.22) \quad \begin{array}{ccccc} & * & \longleftarrow & * & \longrightarrow & * \\ \uparrow & & & \uparrow & & \uparrow \\ X & \longleftarrow & X \vee Y & \longrightarrow & Y \\ \parallel & & \downarrow & & \parallel \\ X & \longleftarrow & X \times Y & \longrightarrow & Y. \end{array}$$

Appealing to [Lemma 2.20](#), taking pushouts of the rows of (2.22) results in the span

$$C \longleftarrow * \longrightarrow *,$$

which has pushout C . Alternatively, since the smash product $X \wedge Y$ is the cofiber of the comparison morphism $X \vee Y \rightarrow X \times Y$, taking pushouts of the columns of (2.22) results in the span

$$(2.23) \quad * \longleftarrow X \wedge Y \longrightarrow *.$$

By definition, the pushout of the span (2.23) is the suspension $\Sigma(X \wedge Y)$, so [Lemma 1.12](#) shows that

$$C \simeq \Sigma(X \wedge Y).$$

To conclude the proof, note that it follows from the definitions that the induced morphisms

$$X \rightarrow \Sigma(X \wedge Y) \quad \text{and} \quad Y \rightarrow \Sigma(X \wedge Y)$$

factor through the zero object $* \in \mathcal{X}_*$. □

[Proposition 2.21](#) also provides a general formula for the cofiber $\text{cofib}(\text{pr}_2 : X \times Y \rightarrow Y)$ that allows us to relate the expressions for $\Sigma\Omega\Sigma X$ and $\Sigma(X \wedge \Omega\Sigma X)$ from [Corollary 2.19](#) and [Proposition 2.21](#), respectively.

Corollary 2.24. *Let \mathcal{X} be an ∞ -category with finite limits and pushouts. Then, for every pair of pointed objects $X, Y \in \mathcal{X}_*$:*

(2.24.1) *There is a natural equivalence $\text{cofib}(\text{pr}_2 : X \times Y \rightarrow Y) \simeq \Sigma X \vee \Sigma(X \wedge Y)$.*

(2.24.2) *There a natural equivalence $\Sigma(X \times Y) \simeq \Sigma(X \wedge Y) \vee \Sigma X \vee \Sigma Y$.*

Proof. Consider the diagram

$$(2.25) \quad \begin{array}{ccccccc} X \times Y & \xrightarrow{\text{pr}_2} & Y & \xrightarrow{\quad} & * \\ \text{pr}_1 \downarrow & & \downarrow & & \downarrow \\ X & \xrightarrow{\quad} & \Sigma(X \wedge Y) & \xrightarrow{\quad} & \Sigma Y \vee \Sigma(X \wedge Y) \\ \downarrow & & \downarrow & & \downarrow \\ * & \xrightarrow{\quad} & \Sigma X \vee \Sigma(X \wedge Y) & \xrightarrow{\quad} & \Sigma X \vee \Sigma Y \vee \Sigma(X \wedge Y), \end{array}$$

where the top-left square is the pushout square of [Proposition 2.21](#) and all of the morphisms in the bottom-right square are coproduct insertions. Since the maps $X \rightarrow \Sigma(X \wedge Y)$ and $Y \rightarrow \Sigma(X \wedge Y)$ are null, the diagram (2.25) commutes and the bottom-left and top-right squares of (2.25) are pushout squares. This proves (2.24.1). To prove (2.24.2), note that the bottom-right square in the diagram (2.25) is a pushout. □

[Corollaries 2.19](#) and [2.24](#) now combine to give the James Splitting.

Proof of Theorem 2.10. Combining [Corollary 2.19](#) with [Corollary 2.24](#) in the case that $Y = \Omega\Sigma X$ we see that there are natural equivalences

$$\begin{aligned} \Sigma\Omega\Sigma X &\simeq \text{cofib}(\text{pr}_2 : X \times \Omega\Sigma X \rightarrow \Omega\Sigma X) \\ &\simeq \Sigma X \vee \Sigma(X \wedge \Omega\Sigma X). \end{aligned}$$

□

The splitting

$$\Sigma\Omega\Sigma X \simeq \left(\bigvee_{1 \leq i \leq n} \Sigma X^{\wedge i} \right) \vee (X^{\wedge n} \wedge \Sigma\Omega\Sigma X)$$

of [Corollary 2.11](#) is immediate from [Theorem 2.10](#) combined with the following elementary fact:

Lemma 2.26. *Let \mathcal{X} be an ∞ -category with universal pushouts. For every pair of pointed objects $X, Y \in \mathcal{X}_*$, there is a natural equivalence*

$$\Sigma(X \wedge Y) \simeq X \wedge \Sigma Y.$$

Proof. Since pushouts in \mathcal{X} are universal and colimits commute, the squares

$$\begin{array}{ccc} X \times Y & \xrightarrow{\text{id}_X \times *} & X \times * \\ \text{id}_X \times * \downarrow & & \downarrow \text{id}_X \times * \\ X \times * & \xrightarrow{\text{id}_X \times *} & X \times \Sigma Y \end{array} \quad \text{and} \quad \begin{array}{ccc} X \vee Y & \xrightarrow{\text{id}_X \vee *} & X \vee * \\ \text{id}_X \vee * \downarrow & & \downarrow \text{id}_X \vee * \\ X \vee * & \xrightarrow{\text{id}_X \vee *} & X \vee \Sigma Y \end{array}$$

are both pushouts in \mathcal{X}_* . By the definition of the smash product and the facts that colimits commute and $X \wedge * \simeq *$, we see that

$$\begin{aligned} X \wedge \Sigma Y &= \text{cofib}(X \vee \Sigma Y \rightarrow X \times \Sigma Y) \\ &\simeq \text{cofib} \left((X \vee *) \sqcup^{X \vee Y} (X \vee *) \rightarrow (X \times *) \sqcup^{X \times Y} (X \times *) \right) \\ &\simeq (X \wedge *) \sqcup^{X \wedge Y} (X \wedge *) \\ &\simeq \Sigma(X \wedge Y). \end{aligned}$$

□

2.4. Ganea's Lemma. Since the method of proof is similar to the arguments in this section, we close with the following lemma of Ganea [9, Theorem 1.1]. This will not be used in the sequel.

Lemma 2.27. *Let \mathcal{X} be an ∞ -category with universal pushouts. Let $f : X \rightarrow Y$ be a morphism in \mathcal{X}_* , and write $i : \text{fib}(f) \rightarrow X$ for the induced morphism from the fiber of f . Then there is a natural equivalence*

$$\text{fib}(\text{cofib}(i) \rightarrow Y) \simeq \Sigma(\Omega Y \wedge \text{fib}(f)).$$

Proof. By [Proposition 2.21](#), it suffices to show that the square

$$\begin{array}{ccc} \Omega Y \times \text{fib}(f) & \xrightarrow{\text{pr}_2} & \text{fib}(f) \\ \text{pr}_1 \downarrow & & \downarrow \\ \Omega Y & \longrightarrow & \text{fib}(\text{cofib}(i) \rightarrow Y) \end{array}$$

is a pushout. Consider the diagram

$$(2.28) \quad \begin{array}{ccccccc} \Omega Y \times \text{fib}(f) & \xrightarrow{\text{pr}_2} & \text{fib}(f) & & & & \\ \text{pr}_2 \downarrow & \searrow \text{pr}_1 & \downarrow & \searrow & & & \\ \Omega Y & \xrightarrow{\quad} & \text{fib}(\text{cofib}(i) \rightarrow Y) & \xrightarrow{\quad} & * & & \\ \downarrow & \downarrow & \downarrow & & \downarrow & & \\ \text{fib}(f) & \xrightarrow{\quad} & X & \xrightarrow{\quad} & \text{cofib}(i) & \xrightarrow{\quad} & Y \\ \downarrow & \downarrow & \downarrow & & \downarrow & & \\ * & \xrightarrow{\quad} & * & \xrightarrow{\quad} & \text{cofib}(i) & \xrightarrow{\quad} & Y \end{array}$$

and note that each vertical square is a pullback square. The bottom horizontal square in (2.28) is a pushout square by definition, so the assumption that pushouts in \mathcal{X} are universal implies that the top horizontal square is a pushout as well. \square

3. THE HILTON–MILNOR SPLITTING

The main result of this section is the following:

Theorem 3.1 (Fundamental Hilton–Milnor Splitting). *Let \mathcal{X} be an ∞ -category with universal pushouts and $X, Y \in \mathcal{X}_*$. Then there is a natural equivalence*

$$\Omega(X \vee Y) \simeq \Omega X \times \Omega Y \times \Omega\Sigma(\Omega X \wedge \Omega Y).$$

Before giving the proof of [Theorem 3.1](#), we discuss some applications.

Example 3.2. Let S be a scheme. Since colimits are universal in the ∞ -category $\mathbf{H}(S)$ of motivic spaces over S ([Example 2.4](#)), [Theorem 3.1](#) implies that for any pointed motivic spaces $X, Y \in \mathbf{H}(S)_*$ we have an equivalence

$$\Omega(X \vee Y) \simeq \Omega X \times \Omega Y \times \Omega\Sigma(\Omega X \wedge \Omega Y).$$

The Hilton–Milnor Splitting allows us to give a new description of the motivic space $\Omega\Sigma(\mathbf{P}^1 \setminus \{0, 1, \infty\})$:

Example 3.3. Let S be a quasicompact quasiseparated scheme. Write $\{0, 1, \infty\} \subset \mathbf{P}_S^1$ for the closed subscheme given by the union of the images of the closed embeddings $S \hookrightarrow \mathbf{P}_S^1$ at 0, 1, and ∞ , respectively. Using the Morel–Voevodsky motivic purity Theorem [[2](#); [18](#), §7.5; [26](#), §3, Theorem 2.23;] Wickelgren [[34](#), Corollary 3.2] showed that there is an equivalence

$$\Sigma(\mathbf{P}_S^1 \setminus \{0, 1, \infty\}) \simeq \Sigma(\mathbf{G}_m \vee \mathbf{G}_m)$$

in the ∞ -category $\mathbf{H}(S)_*$. Since $\Sigma\mathbf{G}_m \simeq \mathbf{P}_S^1$, setting $X = Y = \Sigma\mathbf{G}_m$ in [Example 3.2](#) we see that there are equivalences

$$\begin{aligned} \Omega\Sigma(\mathbf{P}_S^1 \setminus \{0, 1, \infty\}) &\simeq \Omega\Sigma(\mathbf{G}_m \vee \mathbf{G}_m) \\ &\simeq \Omega\Sigma\mathbf{G}_m \times \Omega\Sigma\mathbf{G}_m \times \Omega\Sigma(\Omega\Sigma\mathbf{G}_m \wedge \Omega\Sigma\mathbf{G}_m) \\ &\simeq \Omega\mathbf{P}_S^1 \times \Omega\mathbf{P}_S^1 \times \Omega\Sigma(\Omega\mathbf{P}_S^1 \wedge \Omega\mathbf{P}_S^1). \end{aligned}$$

Applying the James Splitting of [Example 2.14](#) we see that for each integer $n \geq 1$, we can also express $\Omega\Sigma(\mathbf{P}_S^1 \setminus \{0, 1, \infty\})$ as

$$\Omega\Sigma(\mathbf{P}_S^1 \setminus \{0, 1, \infty\}) \simeq \Omega\mathbf{P}_S^1 \times \Omega\mathbf{P}_S^1 \times \Omega\left(\left(\bigvee_{1 \leq i \leq n} \Omega\mathbf{P}_S^1 \wedge S^{i+1, i}\right) \vee (S^{n+1, n} \wedge (\Omega\mathbf{P}_S^1)^{\wedge 2})\right).$$

We now turn to the proof of the Hilton–Milnor Splitting. We first show that there is a fiber sequence

$$(3.4) \quad \Sigma(\Omega Y \wedge \Omega X) \longrightarrow X \vee Y \longrightarrow X \times Y.$$

We then show that the sequence (3.4) splits after taking loops. To do this, we construct a section

$$\Omega(X \times Y) \rightarrow \Omega(X \vee Y),$$

and use the fact that a fiber sequence of group objects with a section splits on the level of underlying objects. After proving that (3.4) is a fiber sequence we give a quick review of group objects and deduce [Theorem 3.1](#) from the Splitting Lemma ([Lemma 3.11](#)).

We start with the following observation:

Lemma 3.5. *Let \mathcal{X} be an ∞ -category with finite limits and $X, Y \in \mathcal{X}_*$. Then there is a natural equivalence*

$$\mathrm{fib}((\mathrm{id}_X, *): X \rightarrow X \times Y) \simeq \Omega Y.$$

Next, we prove the existence of the fiber sequence (3.4).

Lemma 3.6. Let \mathcal{X} be an ∞ -category with universal pushouts and $X, Y \in \mathcal{X}_*$. Then there is a natural equivalence

$$\mathrm{fib}(X \vee Y \rightarrow X \times Y) \simeq \Sigma(\Omega Y \wedge \Omega X).$$

Proof. Write $F := \mathrm{fib}(X \vee Y \rightarrow X \times Y)$. By [Proposition 2.21](#), it suffices to show that there is a pushout square

$$\begin{array}{ccc} \Omega Y \times \Omega X & \xrightarrow{\mathrm{pr}_2} & \Omega X \\ \mathrm{pr}_1 \downarrow & & \downarrow \\ \Omega Y & \longrightarrow & F. \end{array}$$

Consider the diagram

$$(3.7) \quad \begin{array}{ccccccc} \Omega Y \times \Omega X & \xrightarrow{\mathrm{pr}_2} & \Omega X & & & & \\ \downarrow \mathrm{pr}_1 & \searrow & \downarrow & \searrow & & & \\ & \Omega Y & \longrightarrow & F & \longrightarrow & * & \\ & \downarrow & & \downarrow & & \downarrow & \\ * & \longrightarrow & Y & \longrightarrow & X \vee Y & \longrightarrow & X \times Y \\ & \searrow & \searrow & & & & \\ & X & & & & & \end{array}$$

The right-most vertical square in (3.7) is a pullback by definition, and the front and right vertical squares in the cube appearing in (3.7) are pullback squares by [Lemma 3.5](#). The back and left vertical squares in the cube appearing in (3.7) are pullback squares by the Gluing Lemma for pullback squares. The bottom horizontal square in (3.7) is a pushout square by definition, so the assumption that pushouts in \mathcal{X} are universal implies that the top horizontal square is a pushout as well. \square

3.1. Reminder on group objects & the Splitting Lemma. In order to split the fiber sequence (3.4) after taking loops, we need a few basic facts about group objects, which we now review.

Definition 3.8. Let \mathcal{X} be an ∞ -category. A *group object* in \mathcal{X} is a simplicial object $G_\bullet : \Delta^{\mathrm{op}} \rightarrow \mathcal{X}$ such that:

(3.8.1) For each integer $n \geq 0$ and partition $[n] = S \cup S'$ such that $S \cap S' = \{s\}$ consists of a single element, the induced square

$$\begin{array}{ccc} G_n & \longrightarrow & G_\bullet(S') \\ \downarrow & & \downarrow \\ G_\bullet(S) & \longrightarrow & G_\bullet(\{s\}) \end{array}$$

is a pullback square in \mathcal{X} .

(3.8.2) The object G_0 is a terminal object of \mathcal{X} .

In this case, we call $G_1 \in \mathcal{X}$ the *underlying object* of G_\bullet . The face map $d_1 : G_1 \times G_1 \simeq G_2 \rightarrow G_1$ provides a multiplication on G_1 with unit given by the degeneracy map $s_0 : * \simeq G_0 \rightarrow G_1$.

We write $\mathrm{Grp}(\mathcal{X}) \subset \mathrm{Fun}(\Delta^{\mathrm{op}}, \mathcal{X})$ for the full subcategory spanned by the group objects.

The key example of a group object is loops on a pointed object. As a simplicial object, ΩX can be written as the Čech nerve of the basepoint $* \rightarrow X$; since we use Čech nerves in §4.1, we recall the definition here.

Recollection 3.9. Let \mathcal{X} be an ∞ -category with pullbacks, and let $e : W \rightarrow X$ be a morphism in \mathcal{X} . The Čech nerve $\check{C}_\bullet(e)$ of e is the simplicial object

$$\cdots \rightrightarrows W \times_X W \times_X W \rightrightarrows W \times_X W \rightrightarrows W$$

in \mathcal{X} , where $\check{C}_n(e)$ is the $(n+1)$ -fold fiber product of W over X , each degeneracy map is a diagonal morphism, and each face map is a projection. Note that the morphism $e : W \rightarrow X$ defines a natural augmentation $\check{C}_\bullet(e) \rightarrow X$.

Lemma 3.10. *Let \mathcal{X} be an ∞ -category with finite limits and $X \in \mathcal{X}_*$. Then ΩX naturally admits the structure of a group object of \mathcal{X} .*

Proof. Let $U_\bullet(X)$ denote the Čech nerve of the basepoint $* \rightarrow X$. Since $U_0(X) \simeq *$, [HTT, Proposition 6.1.2.11] shows that the Čech nerve $U_\bullet(X)$ is a group object of \mathcal{X} . Since $U_1(X) \simeq \Omega X$, it follows that the loop functor $\Omega : \mathcal{X}_* \rightarrow \mathcal{X}_*$ factors as the composite

$$\mathcal{X}_* \xrightarrow{U_\bullet} \mathrm{Grp}(\mathcal{X}) \rightarrow \mathcal{X}_*$$

of the functor given by the assignment $X \mapsto U_\bullet(X)$ followed by the forgetful functor $\mathrm{Grp}(\mathcal{X}) \rightarrow \mathcal{X}_*$. \square

We leave the following Splitting Lemma as an amusing exercise for the reader.

Lemma 3.11 (Splitting Lemma). *Let \mathcal{X} be an ∞ -category with finite limits, let*

$$A \xrightarrow{i} B \xrightarrow{p} C$$

be a fiber sequence of group objects in \mathcal{X} , and write $m : B \times B \rightarrow B$ for the multiplication on B . For any section $s : C \rightarrow B$ of p on the level of underlying pointed objects of \mathcal{X} , the composite

$$A \times C \xrightarrow{i \times s} B \times B \xrightarrow{m} B$$

is an equivalence in \mathcal{X}_ .*

We can now prove [Theorem 3.1](#).

Proof of Theorem 3.1. By [Lemmas 3.6](#) and [3.10](#), there is a fiber sequence

$$(3.12) \quad \Omega \mathrm{fib}(X \vee Y \rightarrow X \times Y) \rightarrow \Omega(X \vee Y) \rightarrow \Omega X \times \Omega Y$$

of group objects of \mathcal{X} . Note that the map $\Omega(X \vee Y) \rightarrow \Omega X \times \Omega Y$ has a section defined by the composite

$$\Omega X \times \Omega Y \xrightarrow{\Omega i_1 \times \Omega i_2} \Omega(X \vee Y) \times \Omega(X \vee Y) \xrightarrow{m} \Omega(X \vee Y),$$

where $i_1 : X \rightarrow X \vee Y$ and $i_2 : Y \rightarrow X \vee Y$ are the coproduct insertions, and m is the multiplication coming from the group structure on $\Omega(X \vee Y)$. By [Lemma 3.11](#) the fiber sequence (3.12) splits, so applying [Lemma 3.6](#) we see that there are equivalences

$$\begin{aligned} \Omega(X \vee Y) &\simeq \Omega X \times \Omega Y \times \Omega \mathrm{fib}(X \vee Y \rightarrow X \times Y) \\ &\simeq \Omega X \times \Omega Y \times \Omega \Sigma(\Omega X \wedge \Omega Y). \end{aligned}$$

\square

4. CONNECTEDNESS & THE JAMES SPLITTING

The purpose of this section is explain how to use a connectedness argument to show that if X is a pointed connected object of an ∞ -topos, then the comparison morphism

$$c_X : \bigvee_{i=1}^{\infty} \Sigma X^{\wedge i} \rightarrow \Sigma \Omega \Sigma X$$

is an equivalence. To do this, we start by reviewing the basics of connectedness in an ∞ -topos ([§4.1](#)). We also prove some basic connectedness results that we need in our proof of the metastable EHP sequence in [Section 5](#). [Subsection 4.2](#) proves the infinite James and Hilton–Milnor Splittings for connected objects and explains how to deduce Wickelgren and William’s James Splitting in motivic spaces over a perfect field from these results.

4.1. Connectedness and the Blakers–Massey Theorem. In this subsection, we review the basic properties of k -truncated and k -connected morphisms in an ∞ -topos. We also recall the Blakers–Massey Theorem ([Theorem 4.15](#)) and Freudenthal Suspension Theorem ([Corollary 4.16](#)) in an ∞ -topos, since our proof of the metastable EHP sequence in [§ 5](#) relies on these results.

The reader interested in the details of the results reviewed here should consult [[HTT](#), §6.5.1; [1](#), §3.3] for connectedness results, and [[1](#)] for the Blakers–Massey Theorem.

Definition 4.1. Let \mathcal{X} be an ∞ -topos. For each integer $k \geq -2$, define k -truncatedness for morphisms in \mathcal{X} recursively as follows.

(4.1.1) A morphism f is (-2) -truncated if f is an equivalence.

(4.1.2) For $k \geq -1$, a morphism $f : X \rightarrow Y$ is k -truncated if the diagonal $\Delta_f : X \rightarrow X \times_Y X$ is $(k - 1)$ -truncated.

An object $X \in \mathcal{X}$ is k -truncated if the unique morphism $X \rightarrow *$ is k -truncated.

Write $\mathcal{X}_{\leq k} \subset \mathcal{X}$ for the full subcategory spanned by the k -truncated objects. The inclusion $\mathcal{X}_{\leq k} \subset \mathcal{X}$ admits a left adjoint which we denote by $\tau_{\leq k} : \mathcal{X} \rightarrow \mathcal{X}_{\leq k}$.

Example 4.2. Let \mathcal{C} be a small ∞ -category equipped with a Grothendieck topology τ , and let $k \geq -2$ be an integer. Then a sheaf $\mathcal{F} \in \mathrm{Sh}_\tau(\mathcal{C})$ of spaces on \mathcal{C} with respect to τ is k -truncated if and only if $\mathcal{F}(c)$ is a k -truncated space for every $c \in \mathcal{C}$. That is, \mathcal{F} is k -truncated if and only if \mathcal{F} is a sheaf of k -truncated spaces.

Remark 4.3. If \mathcal{X} is an ∞ -topos, then the full subcategory $\mathcal{X}_{\leq 0}$ spanned by the 0-truncated objects is an ordinary topos, i.e., a category of sheaves of sets on a Grothendieck site.

Recollection 4.4. Let \mathcal{X} be an ∞ -topos. A morphism $f : X \rightarrow Y$ in \mathcal{X} is an *effective epimorphism* if the augmentation $\check{C}_\bullet(f) \rightarrow Y$ exhibits Y as the colimit of the Čech nerve of f (see [Recollection 3.9](#)). Equivalently, f is an effective epimorphism if and only if $\tau_{\leq 0}(f) : \tau_{\leq 0}(X) \rightarrow \tau_{\leq 0}(Y)$ is an effective epimorphism in the ordinary topos $\mathcal{X}_{\leq 0}$ of 0-truncated objects of \mathcal{X} [[HTT](#), Proposition 7.2.1.14].

Example 4.5. A morphism $f : X \rightarrow Y$ in the ∞ -topos **Spc** of spaces is an effective epimorphism if and only if $\pi_0(f) : \pi_0(X) \rightarrow \pi_0(Y)$ is a surjection of sets.

Definition 4.6. Let \mathcal{X} be an ∞ -topos. For each integer $k \geq -2$, define k -connectedness for morphisms in \mathcal{X} recursively as follows.

(4.6.1) Every morphism is (-2) -connected.

(4.6.2) For $k \geq -1$, a morphism $f : X \rightarrow Y$ is k -connected if f is an effective epimorphism and the diagonal $\Delta_f : X \rightarrow X \times_Y X$ is $(k - 1)$ -connected.

A morphism f is ∞ -connected if f is k -connected for all $k \geq -2$.

An object $X \in \mathcal{X}$ is k -connected if the unique morphism $X \rightarrow *$ is k -connected.

Remark 4.7. A morphism f is (-1) -connected if and only if f is an effective epimorphism.

Definition 4.8. An ∞ -topos \mathcal{X} is *hypercomplete* if every ∞ -connected morphism in \mathcal{X} is an equivalence.

Warning 4.9. Our conventions for connectedness follow those of Anel, Biedermann, Finster, and Joyal [[1](#), §3.3]. For $k \geq 0$, a homotopy type X is k -connected in our sense if and only if X is k -connected in the classical terminology [[8](#), §6.7; [13](#), p. 346; [23](#), Chapter 10, §4]. In particular, X is 0-connected if and only if X is path-connected. This convention *differs* from the classical one for maps: an n -connected map in our sense is an $(n + 1)$ -connected map in the classical sense.

Comparing to Lurie’s terminology [[HTT](#), §6.5.1], an object or morphism is n -connected in our sense if and only if it is $(n + 1)$ -connective in Lurie’s sense. One of the benefits of our terminological choice is that the constant factors in many connectedness estimates are eliminated (see [Theorem 4.15](#) and [Corollary 4.16](#)).

The following basic properties of k -connected morphisms are proven in [[HTT](#), §6.5.1].

Proposition 4.10. Let \mathcal{X} be an ∞ -topos and $k \geq -2$ be an integer. Then:

(4.10.1) The class of k -connected morphisms in \mathcal{X} is stable under composition.

(4.10.2) The class of k -connected morphisms in \mathcal{X} is stable under pushout along any morphism.

- (4.10.3) The class of k -connected morphisms in \mathcal{X} is stable under pullback along any morphism.
- (4.10.4) The class of k -connected objects in \mathcal{X} is stable under finite products.
- (4.10.5) Given morphisms $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ in \mathcal{X} where f is k -connected, the morphism g is k -connected if and only if gf is k -connected.
- (4.10.6) Given a morphism $f : X \rightarrow Y$ in \mathcal{X} with a section $s : Y \rightarrow X$, the morphism f is $(k + 1)$ -connected if and only if the section s is k -connected.
- (4.10.7) An object $X \in \mathcal{X}$ is k -connected if and only if the k -truncation $\tau_{\leq k}(X)$ of X is terminal in \mathcal{X} .

Since the k -truncation functor $\tau_{\leq k} : \mathcal{X} \rightarrow \mathcal{X}$ preserves filtered colimits, from (4.10.7) we deduce:

Corollary 4.11. *Let \mathcal{X} be an ∞ -topos and $k \geq -2$ be an integer. Then the class of k -connected objects of \mathcal{X} is stable under filtered colimits.*

In the ∞ -topos of spaces, the following connectedness estimates are usually done by appealing to cell structures. Such arguments are unavailable in an arbitrary ∞ -topos, so we deduce these connectedness estimates from [Proposition 4.10](#).

Proposition 4.12. *Let \mathcal{X} be an ∞ -topos, $X, Y \in \mathcal{X}_*$ pointed objects, and $k, \ell \geq 0$ integers. If X is k -connected and Y is ℓ -connected, then:*

- (4.12.1) The suspension ΣX is $(k + 1)$ -connected.
- (4.12.2) The coproduct insertion $X \rightarrow X \vee Y$ is $(\ell - 1)$ -connected.
- (4.12.3) The natural morphism $X \vee Y \rightarrow X \times Y$ is $(k + \ell)$ -connected.
- (4.12.4) The smash product $X \wedge Y$ is $(k + \ell + 1)$ -connected.
- (4.12.5) For each positive integer n , the n -fold smash product $X^{\wedge n}$ is $(n(k + 1) - 1)$ -connected.

Proof. For (4.12.1) note that since k -connected morphisms are stable under pushout (4.10.2), the definition of the suspension ΣX as a pushout shows that the basepoint $* \rightarrow \Sigma X$ is k -connected. Hence ΣX is $(k + 1)$ -connected (4.10.6).

Second, (4.12.2) follows from the fact that the basepoint $* \rightarrow Y$ is $(\ell - 1)$ -connected (4.10.6), combined with the fact that $(\ell - 1)$ -connected morphisms are stable under pushout (4.10.2). Third, (4.12.3) follows from the fact that the basepoints $* \rightarrow X$ and $* \rightarrow Y$ are $(k - 1)$ -connected and $(\ell - 1)$ -connected (4.10.6), respectively, and a general fact about pushout-products of connected morphisms [1, Corollary 3.3.7(4)].

Now we prove (4.12.4). Since $(k + \ell)$ -connected morphisms are stable under pushout (4.10.2), by (4.12.1) and the pushout square

$$\begin{array}{ccc} X \vee Y & \longrightarrow & X \times Y \\ \downarrow & \lrcorner & \downarrow \\ * & \longrightarrow & X \wedge Y \end{array}$$

defining the smash product $X \wedge Y$, we see that the basepoint $* \rightarrow X \wedge Y$ is $(k + \ell)$ -connected. Hence $X \wedge Y$ is $(k + \ell + 1)$ -connected (4.10.6).

Finally, (4.12.5) follows from (4.12.4) by induction. \square

Now we record a convenient fact about the interaction between connectedness and pullbacks that we need in our proof of the metastable EHP sequence.

Proposition 4.13. *Let \mathcal{X} be an ∞ -topos, $\ell \geq -2$ be an integer, and*

$$\begin{array}{ccccc} A & \xrightarrow{f} & C & \xleftarrow{g} & B \\ a \downarrow & & \downarrow c & & \downarrow b \\ A' & \xrightarrow{f'} & C' & \xleftarrow{g'} & B' \end{array}$$

be a commutative diagram in \mathcal{X} . If a and b are ℓ -connected and c is $(\ell + 1)$ -connected, then the induced morphism on pullbacks $a \times_c b : A \times_C B \rightarrow A' \times_{C'} B'$ is ℓ -connected.

Proof. Since ℓ -connected morphisms are stable under composition, by factoring the induced morphism $A \times_C B \rightarrow A' \times_{C'} B'$ as a composite of induced morphisms

$$A \times_C B \xrightarrow{a \times_c \text{id}} A' \times_{C'} B \xrightarrow{\text{id} \times_{\text{id}} b} A' \times_{C'} B',$$

it suffices to prove the claim in the special case $B = B'$ and the morphism $b : B \rightarrow B'$ is the identity. To prove the claim when b is the identity, first write $(\text{id}_A, f) : A \rightarrow A \times_{C'} C$ for the section of the projection $A \times_{C'} C \rightarrow A$ induced by the commutative square

$$\begin{array}{ccc} A & \xrightarrow{f} & C \\ \parallel & & \downarrow c \\ A & \xrightarrow{f'a} & C' \end{array}.$$

Consider the following commutative diagram of pullback squares

$$\begin{array}{ccccccc} A \times_C B & \longrightarrow & A \times_{C'} B & \longrightarrow & A' \times_{C'} B & \longrightarrow & B \\ \downarrow \lrcorner & & \downarrow \lrcorner & & \downarrow \lrcorner & & \downarrow g \\ A & \xrightarrow{(\text{id}_A, f)} & A \times_{C'} C & \longrightarrow & A' \times_{C'} C & \longrightarrow & C \\ & & \downarrow \lrcorner & & \downarrow \lrcorner & & \downarrow c \\ & & A & \xrightarrow{a} & A' & \xrightarrow{f'} & C' \end{array},$$

and notice that the composite middle horizontal map $A \rightarrow C$ is f . Since $c : C \rightarrow C'$ is $(\ell + 1)$ -connected, the projection $A \times_{C'} C \rightarrow A$ is $(\ell + 1)$ -connected (4.10.3). Hence the section $(\text{id}_A, f) : A \rightarrow A \times_{C'} C$ is ℓ -connected (4.10.6). Consequently, the induced morphism $A \times_C B \rightarrow A \times_{C'} B$ is ℓ -connected. Now note that since $a : A \rightarrow A'$ is ℓ -connected, the induced morphism $A \times_{C'} B \rightarrow A' \times_{C'} B$ is ℓ -connected. Hence the composite morphism $A \times_C B \rightarrow A' \times_{C'} B$ is ℓ -connected, as desired. \square

In particular, [Proposition 4.13](#) shows that the class of ℓ -connected morphisms in an ∞ -topos is closed under finite products. Setting $B = B' = *$ in [Proposition 4.13](#) we deduce:

Corollary 4.14. *Let \mathcal{X} be an ∞ -topos, $\ell \geq -2$ be an integer, and*

$$\begin{array}{ccc} A & \xrightarrow{f} & C \\ a \downarrow & & \downarrow c \\ A' & \xrightarrow{f'} & C' \end{array}$$

*be a commutative square in \mathcal{X} . If a is ℓ -connected and c is $(\ell + 1)$ -connected, then for every point $x : * \rightarrow C$, the induced morphism $\text{fib}_x(f) \rightarrow \text{fib}_{cx}(f')$ on fibers is ℓ -connected.*

We conclude this subsection by recalling the Blakers–Massey and Freudenthal Suspension Theorems in the setting of ∞ -topoi.

Theorem 4.15 (Blakers–Massey [1, Corollary 4.3.1]). *Let \mathcal{X} be an ∞ -topos and let*

$$\begin{array}{ccc} A & \xrightarrow{g} & C \\ f \downarrow & & \downarrow \\ B & \longrightarrow & D \end{array}$$

be a pushout square in \mathcal{X} . If f is k -connected and g is ℓ -connected, then the induced morphism $A \rightarrow B \times_D C$ is $(k + \ell)$ -connected.

As in the classical setting, applying the Blakers–Massey Theorem to the pushout defining the suspension immediately implies the Freudenthal Suspension Theorem.

Corollary 4.16 (Freudenthal Suspension Theorem). *Let \mathcal{X} be an ∞ -topos, and $X \in \mathcal{X}_*$ a pointed k -connected object. Then the unit morphism $X \rightarrow \Omega\Sigma X$ is $2k$ -connected.*

4.2. The infinite James and Hilton–Milnor Splittings. We now explain how to use a connectedness argument to show that if X is a pointed 0-connected object of an ∞ -topos, then the comparison morphism

$$c_X : \bigvee_{i=1}^{\infty} \Sigma X^{\wedge i} \rightarrow \Sigma\Omega\Sigma X$$

introduced in [Notation 2.12](#) is an equivalence ([Proposition 4.18](#)). The James Splitting gives us access to generalized Hopf invariants in this very general setting, and implies the stable Snaith splitting for $\Omega\Sigma X$ [\[30\]](#). We also prove the infinite version of the Hilton–Milnor Splitting ([Example 4.22](#)). Using Morel’s unstable \mathbf{A}^1 -connectivity Theorem, we explain why these infinite splittings hold in motivic spaces over a perfect field ([Corollary 4.24](#)). As an application, we give a new description of $\Omega\Sigma^2(\mathbf{P}^1 \setminus \{0, 1, \infty\})$ ([Example 4.25](#)).

The key tool to prove all of these results is the following lemma about how infinite James Splittings interact with localizations.

Lemma 4.17. *Let \mathcal{Y} be an ∞ -category, $j : \mathcal{X} \hookrightarrow \mathcal{Y}$ a full subcategory, and assume that the inclusion j admits a left adjoint $L : \mathcal{Y} \rightarrow \mathcal{X}$. Assume that:*

- (4.17.1) *The ∞ -categories \mathcal{X} and \mathcal{Y} have universal pushouts and the ∞ -category \mathcal{Y}_* has countable coproducts.*
- (4.17.2) *The functor $L : \mathcal{Y} \rightarrow \mathcal{X}$ commutes with finite products and the formation of loop objects (e.g. L is left exact).*

If $X \in \mathcal{X}_$ is a pointed object with the property that the comparison morphism*

$$c_{j(X)} : \bigvee_{i \geq 1} \Sigma_{\mathcal{Y}} j(X)^{\wedge i} \rightarrow \Sigma_{\mathcal{Y}} \Omega_{\mathcal{Y}} \Sigma_{\mathcal{Y}} j(X)$$

is an equivalence in \mathcal{Y}_ , then the comparison morphism*

$$c_X : \bigvee_{i \geq 1} \Sigma_{\mathcal{X}} X^{\wedge i} \rightarrow \Sigma_{\mathcal{X}} \Omega_{\mathcal{X}} \Sigma_{\mathcal{X}} X$$

is an equivalence in \mathcal{X}_ .*

Proof. Since the localization $L : \mathcal{Y} \rightarrow \mathcal{X}$ preserves the terminal object, the functor L descends to the level of pointed objects. Moreover, since L preserves finite products, the functor $L : \mathcal{Y}_* \rightarrow \mathcal{X}_*$ commutes with smash products. Since the functor $L : \mathcal{Y}_* \rightarrow \mathcal{X}_*$ also commutes with the formation of loop objects, we see that the morphism c_X is equivalent to $L(c_{j(X)})$. The assumption that the morphism $c_{j(X)}$ is an equivalence completes the proof. \square

Proposition 4.18 (James Splitting). *Let \mathcal{X} be an ∞ -topos. Then for each 0-connected pointed object $X \in \mathcal{X}_*$, the natural comparison morphism*

$$c_X : \bigvee_{i \geq 1} \Sigma X^{\wedge i} \rightarrow \Sigma\Omega\Sigma X$$

is an equivalence in \mathcal{X}_ .*

Proof. Since every ∞ -topos is a left exact localization of a presheaf ∞ -topos and presheaf ∞ -topoi are hypercomplete (see [Definition 4.8](#)), by [Lemma 4.17](#) we are reduced to the case that \mathcal{X} is hypercomplete. That is, it suffices to show that the morphism c_X is ∞ -connected.

To see this, notice that for each integer $n \geq 1$, the summand inclusions of $\bigvee_{i=1}^n \Sigma X^{\wedge i}$ into $\bigvee_{i=1}^{\infty} \Sigma X^{\wedge i}$ and

$$\Sigma\Omega\Sigma X \simeq \left(\bigvee_{1 \leq i \leq n} \Sigma X^{\wedge i} \right) \vee (X^{\wedge n} \wedge \Sigma\Omega\Sigma X)$$

(Corollary 2.11) fit into a commutative triangle

$$(4.19) \quad \begin{array}{ccc} \bigvee_{i=1}^n \Sigma X^{\wedge i} & \longrightarrow & \bigvee_{i=1}^{\infty} \Sigma X^{\wedge i} \\ & \searrow & \downarrow c_X \\ & & \Sigma \Omega \Sigma X. \end{array}$$

Since X is 0-connected, combining (4.12.1), (4.12.4), (4.12.5) and Corollary 4.16 we see that $X^{\wedge n} \wedge \Sigma \Omega \Sigma X$ is n -connected. Hence (4.12.2) shows that the summand inclusion

$$\bigvee_{i=1}^n \Sigma X^{\wedge i} \rightarrow \Sigma \Omega \Sigma X$$

is $(n-1)$ -connected. Similarly, since X is 0-connected, combining (4.12.1), (4.12.5), and Corollary 4.11 we see that the object $\bigvee_{i \geq n+1}^{\infty} \Sigma X^{\wedge i}$ is $(n+1)$ -connected. Again applying (4.12.2), we see that the summand inclusion

$$\bigvee_{i=1}^n \Sigma X^{\wedge i} \rightarrow \bigvee_{i=1}^{\infty} \Sigma X^{\wedge i}$$

is n -connected. The commutativity of the triangle (4.19) combined with (4.10.5) show that the morphism

$$c_X : \bigvee_{i \geq 1} \Sigma X^{\wedge i} \rightarrow \Sigma \Omega \Sigma X$$

is $(n-1)$ -connected. Since this is true for each integer $n \geq 1$, we have shown that morphism c_X is ∞ -connected, as desired. \square

Remark 4.20. Of course, in the proof of Proposition 4.18 we can further reduce to the case $\mathcal{X} = \mathbf{Spc}$ and the claim follows from the classical James Splitting. The purpose of our proof is to provide an explanation that does not appeal to the classical result but, rather, only uses basic manipulations available in an ∞ -topos.

Proposition 4.21 (Hilton–Milnor Splitting, general version). *Let \mathcal{X} be an ∞ -category with universal pushouts and countable coproducts, and let $X, Y \in \mathcal{X}_*$ be pointed objects.*

(4.21.1) *If the comparison morphism c_Y is an equivalence, then there is a natural equivalence of pointed objects*

$$\Omega X \times \Omega \Sigma Y \times \Omega \left(\Sigma \Omega X \wedge \bigvee_{j \geq 1} Y^{\wedge j} \right) \simeq \Omega(X \vee \Sigma Y).$$

(4.21.2) *If the comparison morphisms c_X and c_Y are equivalences, then there is a natural equivalence of pointed objects*

$$\Omega \Sigma X \times \Omega \Sigma Y \times \Omega \Sigma \left(\bigvee_{i, j \geq 1} X^{\wedge i} \wedge Y^{\wedge j} \right) \simeq \Omega \Sigma(X \vee Y).$$

Proof. For (4.21.1), first note that the Hilton–Milnor Splitting of Theorem 3.1 provides an equivalence.

$$\Omega(X \vee \Sigma Y) \simeq \Omega X \times \Omega \Sigma Y \times \Omega(\Omega X \wedge \Omega \Sigma Y)$$

The claim now follows from the natural natural equivalences

$$\begin{aligned} \Omega \left(\Sigma \Omega X \wedge \bigvee_{j \geq 1} Y^{\wedge j} \right) &\simeq \Omega \left(\Omega X \wedge \bigvee_{j \geq 1} \Sigma Y^{\wedge j} \right) && \text{(Lemma 2.26)} \\ &\simeq \Omega(\Omega X \wedge \Sigma \Omega \Sigma Y) && (c_Y \text{ is an equivalence}) \\ &\simeq \Omega \Sigma(\Omega X \wedge \Omega \Sigma Y) && \text{(Lemma 2.26)} \end{aligned}$$

Similarly, (4.21.2) follows from Lemma 2.26, (4.21.1), and the assumption that $c_X : \bigvee_{i \geq 1} \Sigma X^{\wedge i} \rightarrow \Sigma \Omega \Sigma X$ is an equivalence \square

Example 4.22 (Hilton–Milnor Splitting). Let \mathcal{X} be an ∞ -topos and let $X, Y \in \mathcal{X}_*$ be pointed objects. By [Proposition 4.18](#), the hypotheses of [\(4.21.1\)](#) are satisfied if Y is 0-connected, and the hypotheses of [\(4.21.2\)](#) are satisfied if both X and Y are 0-connected.

The next application is that the infinite James and Hilton–Milnor Splittings hold for \mathbf{A}^1 -0-connected motivic spaces over a perfect field. The infinite James Splitting was first proven by Wickelgren and Williams using the *James filtration* [\[35, Theorem 1.5\]](#); the infinite Hilton–Milnor Splittings in this context is new.

Recollection 4.23. Let S be a scheme and $n \geq 0$ an integer. A motivic space $X \in \mathbf{H}(S) \subset \mathrm{Sh}_{\mathrm{nis}}(S)$ is \mathbf{A}^1 - n -connected if X is an n -connected object of the Nisnevich ∞ -topos $\mathrm{Sh}_{\mathrm{nis}}(S)$.

Corollary 4.24. Let k be a perfect field and let X and Y be pointed motivic spaces over k .

(4.24.1) If X is \mathbf{A}^1 -0-connected, then the natural comparison morphism

$$c_X : \bigvee_{i \geq 1} \Sigma X^{\wedge i} \rightarrow \Sigma \Omega \Sigma X$$

is an equivalence in $\mathbf{H}(\mathrm{Spec}(k))_*$.

(4.24.2) If Y is \mathbf{A}^1 -0-connected, then there is a natural equivalence of pointed motivic spaces

$$\Omega X \times \Omega \Sigma Y \times \Omega \left(\Sigma \Omega X \wedge \bigvee_{j \geq 1} Y^{\wedge j} \right) \simeq \Omega(X \vee \Sigma Y).$$

(4.24.3) If X and Y are \mathbf{A}^1 -0-connected, then there is a natural equivalence of pointed motivic spaces

$$\Omega \Sigma X \times \Omega \Sigma Y \times \Omega \Sigma \left(\bigvee_{i, j \geq 1} X^{\wedge i} \wedge Y^{\wedge j} \right) \simeq \Omega \Sigma(X \vee Y).$$

Proof. First note that [\(4.24.2\)](#) and [\(4.24.3\)](#) follow from [Proposition 4.21](#) and [\(4.24.1\)](#). To verify [\(4.24.1\)](#), note that since pushouts are universal in the ∞ -category $\mathbf{H}(\mathrm{Spec}(k))$ of motivic spaces over k ([Example 2.4](#)) and $\mathbf{H}(\mathrm{Spec}(k))$ is a localization of the ∞ -topos $\mathrm{Sh}_{\mathrm{nis}}(\mathrm{Spec}(k))$ of Nisnevich sheaves on $\mathrm{Spec}(k)$, it suffices to check that the motivic localization functor

$$L_{\mathrm{mot}} : \mathrm{Sh}_{\mathrm{nis}}(\mathrm{Spec}(k)) \rightarrow \mathbf{H}(\mathrm{Spec}(k))$$

satisfies the hypotheses of [Lemma 4.17](#). To see this, first note that over any base scheme the motivic localization commutes with finite products [\[18, Corollary 3.5\]](#). Second, since the field k is perfect, Morel’s unstable \mathbf{A}^1 -connectivity Theorem [\[25, Theorems 5.46 and 6.1\]](#) implies that motivic localization commutes with the formation of loop objects [\[3, Theorem 2.4.1; 25, Theorem 6.46\]](#). \square

Example 4.25. Let k be a perfect field. [Corollary 4.24](#) gives the following variant on [Example 3.3](#). Since $\Sigma \mathbf{G}_m \simeq \mathbf{P}^1$ is \mathbf{A}^1 -0-connected, [\(4.24.3\)](#) provides equivalences of pointed motivic spaces over k

$$\begin{aligned} \Omega \Sigma^2(\mathbf{P}_k^1 \setminus \{0, 1, \infty\}) &\simeq \Omega \Sigma(\mathbf{P}_k^1 \vee \mathbf{P}_k^1) \\ &\simeq \Omega \Sigma \mathbf{P}_k^1 \times \Omega \Sigma \mathbf{P}_k^1 \times \Omega \Sigma \left(\bigvee_{i, j \geq 1} (\mathbf{P}_k^1)^{\wedge(i+j)} \right) \\ &\simeq \Omega S^{3,1} \times \Omega S^{3,1} \times \Omega \left(\bigvee_{i, j \geq 1} S^{2(i+j)+1, i+j} \right). \end{aligned}$$

We complete this section by constructing the *Hopf maps* that appear in the metastable EHP sequence.

Construction 4.26 (Hopf maps). Let \mathcal{X} be an ∞ -topos and X a pointed 0-connected object of \mathcal{X} . For each integer $n \geq 1$, we define the *Hopf map* $h_n : \Omega \Sigma X \rightarrow \Omega \Sigma X^{\wedge n}$ as the adjoint to the collapse map

$$\Sigma \Omega \Sigma X \simeq \bigvee_{i \geq 1} \Sigma X^{\wedge i} \rightarrow \Sigma X^{\wedge n}$$

induced by the James Splitting of [Proposition 4.18](#).

5. THE JAMES FILTRATION & METASTABLE EHP SEQUENCE

In classical algebraic topology, the *metastable EHP sequence* is the statement that the composite

$$X \longrightarrow \Omega\Sigma X \xrightarrow{h_2} \Omega\Sigma X^{\wedge 2}$$

is a fiber sequence in a range depending on the connectedness of X , known as the *metastable range*. Here the first map $X \rightarrow \Omega\Sigma X$ is the unit and h_2 is the Hopf map (Construction 4.26). For the higher Hopf maps $h_n : \Omega\Sigma X \rightarrow \Omega\Sigma X^{\wedge n}$, there is an analogous fiber sequence in a range

$$J_{n-1}(X) \longrightarrow \Omega\Sigma X \xrightarrow{h_n} \Omega\Sigma X^{\wedge n},$$

where $J_{n-1}(X)$ is the $(n-1)^{\text{st}}$ piece of the *James filtration* on $\Omega\Sigma X$.

This section is dedicated to a non-computational proof of the metastable EHP sequence in an ∞ -topos that only makes use of the Blakers–Massey Theorem and some basic connectedness results (see Theorem 5.19). In §5.1 we review the James filtration. In §5.2 we refine the James Splitting to a splitting

$$\Sigma J_n(X) \simeq \bigvee_{i=1}^n \Sigma X^{\wedge i}.$$

In §5.3, we give our non-computational proof of the metastable EHP sequence via the Blakers–Massey Theorem, and also record a computational proof for posterity.

5.1. The James filtration. Classically, the James filtration $\{J_n(X)\}_{n \geq 0}$ provides a multiplicative filtration on the free monoid $J(X)$ on a pointed space X , in the homotopical sense. At the point-set level, $J(X)$ can be presented as the free topological monoid on X , and $J_n(X)$ can be identified the subspace of words of length at most n in $J(X)$. Concatenation of words then supplies $\{J_n(X)\}_{n \geq 0}$ with the structure of a filtered monoid. Since the trivial monoid and trivial group coincide, if X is connected, then the free monoid $J(X)$ on X coincides with the free group $\Omega\Sigma X$ on X .

In a general ∞ -category, we can define the James filtration as follows. This definition is provided in [6, Section 3] in the context of homotopy type theory; the arguments made in [6, Section 3] are formal and valid in any ∞ -topos.

Construction 5.1 (James filtration). Let \mathcal{X} be an ∞ -category with finite products and pushouts, and let $X \in \mathcal{X}_*$ be a pointed object. For each integer $n \geq 0$ we define a pointed object $J_n(X) \in \mathcal{X}_*$ as well as morphisms

$$i_n : J_n(X) \rightarrow J_{n+1}(X) \quad \text{and} \quad \alpha_n : X \times J_n(X) \rightarrow J_{n+1}(X)$$

in \mathcal{X}_* recursively as follows.

(5.1.1) We define $J_0(X) := *$, $J_1(X) := X$, the morphism $i_0 : * \rightarrow X$ is the basepoint, and the morphism $\alpha_0 : X \times * \rightarrow X$ is the projection $\text{pr}_1 : X \times * \simeq X$.

(5.1.2) For $n \geq 2$, we define $J_n(X)$, i_{n-1} , and α_{n-1} by the pushout square

$$(5.2) \quad \begin{array}{ccc} X \times J_{n-2}(X) & \xrightarrow{J_{n-2}(X)} & J_{n-1}(X) \\ \downarrow & \lrcorner & \downarrow i_{n-1} \\ X \times J_{n-1}(X) & \xrightarrow{\alpha_{n-1}} & J_n(X), \end{array}$$

where: the top horizontal morphism is induced by the universal property of the pushout by the commutative square

$$\begin{array}{ccc} J_{n-2}(X) & \xrightarrow{i_{n-2}} & J_{n-1}(X) \\ (*, \text{id}) \downarrow & & \parallel \\ X \times J_{n-2}(X) & \xrightarrow{\alpha_{n-2}} & J_{n-1}(X), \end{array}$$

and the left vertical morphism is induced by the universal property of the pushout by the commutative square

$$\begin{array}{ccc} J_{n-2}(X) & \xrightarrow{i_{n-2}} & J_{n-1}(X) \\ (*, \text{id}) \downarrow & & \downarrow (*, \text{id}) \\ X \times J_{n-2}(X) & \xrightarrow{\text{id} \times i_{n-2}} & X \times J_{n-1}(X) \end{array}$$

For each positive integer n , define a morphism $a_n : X^{\times n} \rightarrow J_n(X)$ as the composite

$$X^{\times n} \simeq X^{\times n-1} \times J_1(X) \xrightarrow{\text{id} \times \alpha_1} X^{\times n-2} \times J_2(X) \xrightarrow{\text{id} \times \alpha_2} \dots \longrightarrow X \times J_{n-1}(X) \xrightarrow{\alpha_{n-1}} J_n(X).$$

Finally, define $J(X) := \text{colim}_{n \geq 0} J_n(X)$.

Definition 5.3. Let \mathcal{X} be an ∞ -category with finite products and pushouts, and let $X \in \mathcal{X}_*$ be a pointed object. For each integer $n \geq 0$ define a morphism $u_n : J_n(X) \rightarrow \Omega\Sigma X$ recursively as follows. The morphism u_0 is the basepoint, and the morphism $u_1 : X \rightarrow \Omega\Sigma X$ is the unit. For $n \geq 2$, the morphism u_n is induced by the commutative square

$$\begin{array}{ccc} X \times J_{n-2}(X) & \xrightarrow{J_{n-2}(X)} J_{n-1}(X) & \longrightarrow J_{n-1}(X) \\ \downarrow & \downarrow & \downarrow u_{n-1} \\ X \times J_{n-1}(X) & \xrightarrow{m \circ (u_1 \times u_{n-1})} & \Omega\Sigma X \end{array}$$

where $m : \Omega\Sigma X \times \Omega\Sigma X \rightarrow \Omega\Sigma X$ is the group multiplication.

The morphisms u_n induce a morphism $u : J(X) \rightarrow \Omega\Sigma X$.

Theorem 5.4 ([6, Section 6]). *Let \mathcal{X} be an ∞ -topos and $X \in \mathcal{X}_*$ a pointed object. If X is 0-connected, then the morphism $u : J(X) \rightarrow \Omega\Sigma X$ is an equivalence.*

Brunerie [6] gives an elementary proof of the following connectedness estimate.

Lemma 5.5 ([6, Proposition 4]). *Let \mathcal{X} be an ∞ -topos, $k \geq 0$ an integer, and $X \in \mathcal{X}_*$ a pointed k -connected object. Then the morphism $i_{n-1} : J_{n-1}(X) \rightarrow J_n(X)$ is $(n(k+1) - 2)$ -connected.*

Corollary 5.6. *Let \mathcal{X} be an ∞ -topos, $k \geq 0$ an integer, and $X \in \mathcal{X}_*$ a pointed k -connected object. Then for each integer $n \geq 0$, the object $J_n(X)$ is k -connected.*

Proof. If $n = 0$, then the claim is clear since $J_0(X) = *$. If $n > 0$, then by Lemma 5.5 the morphisms i_0, \dots, i_{n-1} are all $(k-1)$ -connected. Hence the basepoint

$$i_{n-1} \cdots i_0 : * \rightarrow J_n(X)$$

is $(k-1)$ -connected; equivalently, $J_n(X)$ is k -connected (4.10.6). \square

Lemma 5.7 ([6, Proposition 6]). *Let \mathcal{X} be an ∞ -topos, $k \geq 0$ an integer, and $X \in \mathcal{X}_*$ a pointed k -connected object. Then the morphism $u_n : J_n(X) \rightarrow \Omega\Sigma X$ is $((n+1)(k+1) - 2)$ -connected.*

5.2. Splitting the James filtration. The purpose of this subsection is to prove the following splitting of the James filtration, which we use in our proof of the metastable EHP sequence (Theorem 5.19).

Proposition 5.8. *Let \mathcal{X} be an ∞ -category with universal pushouts, and let $X \in \mathcal{X}_*$. Then there is a splitting*

$$(5.9) \quad \Sigma J_n(X) \simeq \bigvee_{1 \leq i \leq n} \Sigma X^{\wedge i}.$$

If \mathcal{X} is an ∞ -topos and X is 0-connected, then under the map $\Sigma u_n : \Sigma J_n(X) \rightarrow \Sigma \Omega\Sigma X$, the splitting (5.9) is an equivalence onto the first n factors of the splitting $\Sigma \Omega\Sigma X \simeq \bigvee_{i \geq 1} \Sigma X^{\wedge i}$ of Proposition 4.18.

The proof of Proposition 5.8 requires some preliminaries. We need to relate the cofiber of i_n to smash powers of X ; before doing so we need some preparatory lemmas.

Lemma 5.10. *Let \mathcal{X} be an ∞ -category with finite products and pushouts, and let $X, Y \in \mathcal{X}_*$. Then there is a cofiber sequence*

$$Y \longrightarrow \mathrm{cofib}((\mathrm{id}_X, *) : X \rightarrow X \times Y) \longrightarrow X \wedge Y.$$

Proof. There is a map of cofiber sequences

$$(5.11) \quad \begin{array}{ccccc} X & \xlongequal{\quad} & X & \longrightarrow & * \\ \downarrow & & \downarrow (\mathrm{id}_X, *) & & \downarrow \\ X \vee Y & \longrightarrow & X \times Y & \longrightarrow & X \wedge Y, \end{array}$$

where the leftmost vertical map is the coproduct insertion. The cofiber of the coproduct insertion $X \rightarrow X \vee Y$ is Y , and the cofiber of the basepoint $* \rightarrow X \wedge Y$ is $X \wedge Y$. To conclude, note that taking vertical cofibers in (5.11) results in a cofiber sequence. \square

The following is a straightforward application of [Lemma 1.12](#).

Lemma 5.12. *Let \mathcal{X} be an ∞ -category with pushouts and a terminal object and let*

$$\begin{array}{ccc} A & \longrightarrow & B \\ \downarrow & & \downarrow \\ C & \longrightarrow & D \end{array}$$

be a commutative square in \mathcal{X}_ . Then there is a natural equivalence*

$$\mathrm{cofib}(B \sqcup^A C \rightarrow D) \simeq \mathrm{cofib}(\mathrm{cofib}(A \rightarrow C) \rightarrow \mathrm{cofib}(B \rightarrow D)).$$

We are now ready to show that $\mathrm{cofib}(i_n) \simeq X^{\wedge n+1}$.

Proposition 5.13. *Let \mathcal{X} be an ∞ -category with universal pushouts, and let $X \in \mathcal{X}_*$. Then for each integer $n \geq 0$, there is a natural equivalence*

$$\mathrm{cofib}(i_n : J_n(X) \rightarrow J_{n+1}(X)) \simeq X^{\wedge n+1}.$$

Moreover, the composite

$$X^{\times n+1} \xrightarrow{a_n} J_{n+1}(X) \longrightarrow X^{\wedge n+1}$$

is equivalent to the canonical map $X^{\times n+1} \rightarrow X^{\wedge n+1}$.

Proof. We prove the claim by induction on n . For the base case, note that since the morphism i_0 is the basepoint $* \rightarrow X$, the cofiber of i_0 is X . For the inductive step we assume that $\mathrm{cofib}(i_n) \simeq X^{\wedge n+1}$ and show that $\mathrm{cofib}(i_{n+1}) \simeq X^{\wedge n+2}$. From the defining pushout square (5.2), we see that

$$\mathrm{cofib}(i_{n+1}) \simeq \mathrm{cofib}(X \times J_n(X) \sqcup^{J_n(X)} J_{n+1}(X) \rightarrow X \times J_{n+1}(X)).$$

Applying [Lemma 5.12](#) shows that

$$\mathrm{cofib}(i_{n+1}) \simeq \mathrm{cofib}(\mathrm{cofib}(J_n(X) \rightarrow X \times J_n(X)) \rightarrow \mathrm{cofib}(J_{n+1}(X) \rightarrow X \times J_{n+1}(X))),$$

where the map of cofibers is induced by the map $i_n : J_n(X) \rightarrow J_{n+1}(X)$. By [Lemma 5.10](#), there is a cofiber sequence

$$X \rightarrow \mathrm{cofib}(J_n(X) \rightarrow X \times J_n(X)) \rightarrow X \wedge J_n(X);$$

moreover, the map $i_n : J_n(X) \rightarrow J_{n+1}(X)$ induces a map of cofiber sequences

$$(5.14) \quad \begin{array}{ccccc} X & \longrightarrow & \mathrm{cofib}(J_n(X) \rightarrow X \times J_n(X)) & \longrightarrow & X \wedge J_n(X) \\ \parallel & & \downarrow & & \downarrow \mathrm{id}_X \wedge i_n \\ X & \longrightarrow & \mathrm{cofib}(J_{n+1}(X) \rightarrow X \times J_{n+1}(X)) & \longrightarrow & X \wedge J_{n+1}(X). \end{array}$$

Since the leftmost vertical map is the identity, taking vertical cofibers in the map of cofiber sequences (5.14) produces an equivalence between the vertical cofibers of the middle and right vertical maps. Since the cofiber of the middle vertical map is $\text{cofib}(i_{n+1})$, we find that

$$\text{cofib}(i_{n+1}) \simeq \text{cofib}(\text{id}_X \wedge i_n : X \wedge J_n(X) \rightarrow X \wedge J_{n+1}(X)).$$

Since pushouts in \mathcal{X} are universal we have a natural equivalence

$$\text{cofib}(\text{id}_X \wedge i_n : X \wedge J_n(X) \rightarrow X \wedge J_{n+1}(X)) \simeq X \wedge \text{cofib}(i_n : J_n(X) \rightarrow J_{n+1}(X))$$

By the inductive hypothesis, $\text{cofib}(i_n) \simeq X^{\wedge n+1}$, so $\text{cofib}(i_{n+1}) \simeq X^{\wedge n+2}$, as desired. \square

Next we split the term

$$\Sigma(X \times J_{n-1}(X) \overset{J_{n-1}(X)}{\sqcup} J_n(X))$$

in the pushout square (5.2) defining $\Sigma J_{n+1}(X)$ and prove **Proposition 5.8**.

Lemma 5.15. *Let \mathcal{X} be an ∞ -category with universal pushouts, $X \in \mathcal{X}_*$, and $n \geq 1$ an integer. Then there is a natural equivalence.*

$$\Sigma(X \times J_{n-1}(X) \overset{J_{n-1}(X)}{\sqcup} J_n(X)) \simeq \Sigma(X \wedge J_{n-1}(X)) \vee \Sigma X \vee \Sigma J_n(X).$$

Proof. Since suspension preserves pushouts, we have a pushout square

$$(5.16) \quad \begin{array}{ccc} \Sigma J_{n-1}(X) & \longrightarrow & \Sigma J_n(X) \\ \downarrow & \ulcorner & \downarrow \\ \Sigma(X \times J_{n-1}(X)) & \longrightarrow & \Sigma(X \times J_{n-1}(X) \overset{J_{n-1}(X)}{\sqcup} J_n(X)). \end{array}$$

Under the equivalence

$$\Sigma(X \times J_{n-1}(X)) \simeq \Sigma(X \wedge J_{n-1}(X)) \vee \Sigma X \vee \Sigma J_{n-1}(X)$$

of **Corollary 2.24**, the left vertical map in (5.16) is the coproduct insertion. Hence on pushouts we see that

$$\Sigma(X \times J_{n-1}(X) \overset{J_{n-1}(X)}{\sqcup} J_n(X)) \simeq \Sigma(X \wedge J_{n-1}(X)) \vee \Sigma X \vee \Sigma J_n(X). \quad \square$$

Proof of Proposition 5.8. We prove the claim by induction on n . The base case where $n = 1$ is obvious. For the inductive step, assume that $n \geq 1$ and $\Sigma J_n(X) \simeq \bigvee_{i=1}^n \Sigma X^{\wedge i}$. By **Proposition 5.13** we have a cofiber sequence

$$J_n(X) \xrightarrow{i_n} J_{n+1}(X) \longrightarrow X^{\wedge n+1},$$

so the inductive hypothesis and the duals of **Lemmas 3.10** and **3.11**, it suffices to define a retraction

$$r_n : \Sigma J_{n+1}(X) \rightarrow \Sigma J_n(X)$$

of the map Σi_n .

We construct the retractions $r_n : \Sigma J_{n+1}(X) \rightarrow \Sigma J_n(X)$ inductively. For the base case, the retraction $r_0 : \Sigma X \rightarrow *$ of Σi_0 is the unique morphism. For the inductive step, assume that $n \geq 1$ and we have constructed a retraction $r_{n-1} : \Sigma J_n(X) \rightarrow \Sigma J_{n-1}(X)$ of Σi_{n-1} ; we use this to construct a retraction r_n of Σi_n . Since suspension preserves pushouts, suspending the defining pushout square (5.2) yields a pushout square

$$(5.17) \quad \begin{array}{ccc} \Sigma(X \times J_{n-1}(X) \overset{J_{n-1}(X)}{\sqcup} J_n(X)) & \longrightarrow & \Sigma J_n(X) \\ \downarrow & \ulcorner & \downarrow \Sigma i_n \\ \Sigma(X \times J_n(X)) & \xrightarrow{\Sigma \alpha_n} & \Sigma J_{n+1}(X). \end{array}$$

In order to define a retraction of Σi_n , it suffices to define a retraction of the left vertical map in (5.17), i.e., it suffices to define a retraction

$$\Sigma(X \times J_n(X)) \rightarrow \Sigma\left(X \times J_{n-1}(X) \overset{J_{n-1}(X)}{\sqcup} J_n(X)\right).$$

By [Corollary 2.24](#) and [Lemma 5.15](#), we have equivalences

$$\Sigma(X \times J_n(X)) \simeq \Sigma(X \wedge J_n(X)) \vee \Sigma X \vee \Sigma J_n(X)$$

and

$$\Sigma\left(X \times J_{n-1}(X) \overset{J_{n-1}(X)}{\sqcup} J_n(X)\right) \simeq \Sigma(X \wedge J_{n-1}(X)) \vee \Sigma X \vee \Sigma J_n(X).$$

Moreover, the left vertical map in (5.17) is induced by the suspensions of the identity on X , identity on $J_n(X)$, and the map $i_n : J_{n-1}(X) \rightarrow J_n(X)$. Under the identifications

$$\Sigma(X \wedge J_{n-1}(X)) \simeq X \wedge \Sigma J_{n-1}(X) \quad \text{and} \quad \Sigma(X \wedge J_n(X)) \simeq X \wedge \Sigma J_n(X)$$

of [Lemma 2.26](#), we see that the map

$$\text{id}_X \wedge r_{n-1} : \Sigma(X \wedge J_{n-1}(X)) \simeq X \wedge \Sigma J_{n-1}(X) \longrightarrow X \wedge \Sigma J_n(X) \simeq \Sigma(X \wedge J_n(X))$$

is a retraction of $\Sigma(\text{id}_X \wedge i_{n-1})$. Hence the map

$$(\text{id}_X \wedge r_{n-1}) \vee \text{id} \vee \text{id} : \Sigma(X \wedge J_n(X)) \vee \Sigma X \vee \Sigma J_n(X) \longrightarrow \Sigma(X \wedge J_{n-1}(X)) \vee \Sigma X \vee \Sigma J_n(X)$$

supplies the desired retraction of the left vertical map in (5.17). \square

5.3. Proofs of the metastable EHP sequence. In this subsection, we present two proofs of the metastable EHP sequence in the setting of ∞ -topoi. Before making a precise statement of the main result, we need the following easy lemma.

Lemma 5.18. *Let \mathcal{X} be an ∞ -topos, $X \in \mathcal{X}_*$ be a pointed 0-connected object, and $n \geq 1$ an integer. Then the composite*

$$J_{n-1}(X) \xrightarrow{u_n} \Omega \Sigma X \xrightarrow{h_n} \Omega \Sigma X^{\wedge n}$$

is null. Here h_n is the Hopf map of [Construction 4.26](#).

Proof. It suffices to prove the corresponding statement on adjoints: in other words, we need to show that the composite

$$\Sigma J_{n-1}(X) \xrightarrow{\Sigma u_n} \Sigma \Omega \Sigma X \longrightarrow \Sigma X^{\wedge n}$$

is null. This is an immediate consequence of [Proposition 5.8](#). \square

We can now state the metastable EHP sequence.

Theorem 5.19 (metastable EHP sequence). *Let \mathcal{X} be an ∞ -topos, $k \geq 0$ an integer, and $X \in \mathcal{X}_*$ a pointed k -connected object. Then for every integer $n \geq 1$, the morphism $J_{n-1}(X) \rightarrow \text{fib}(h_n)$ induced by [Lemma 5.18](#) is $((n+1)(k+1)-3)$ -connected.*

Remark 5.20. [Theorem 5.19](#) implies the metastable EHP sequence of Asok–Wickelgren–Williams for ∞ -topoi of hypersheaves on an ordinary site with enough points [[3](#), [Proposition 3.1.4](#)].

The first proof of [Theorem 5.19](#) we present is *internal* to ∞ -topoi, and only uses basic facts about connectedness and the James construction, as well as the Blakers–Massey Theorem. The second reduces to the ∞ -topos **Spc** of spaces, then uses the homology Whitehead Theorem and Serre spectral sequence to give a calculational proof of the metastable EHP sequence in the classical setting. Both perspectives are valuable, and we present the second here in part because the calculational proof of the metastable EHP sequence does not seem to be easy to locate in the literature.

Internal proof of Theorem 5.19. First we show that it suffices to prove the claim where we replace $\text{fib}(h_n)$ by the fiber of the natural morphism $J_n(X) \rightarrow X^{\wedge n}$. Observe that we have a commutative square

$$(5.21) \quad \begin{array}{ccc} J_n(X) & \longrightarrow & X^{\wedge n} \\ u_n \downarrow & & \downarrow \\ \Omega\Sigma X & \xrightarrow{h_n} & \Omega\Sigma X^{\wedge n}, \end{array}$$

where the right vertical morphism is the unit. Since X is k -connected, the morphism

$$u_n : J_n(X) \rightarrow \Omega\Sigma X$$

is $((n+1)(k+1)-2)$ -connected (Lemma 5.7) and $X^{\wedge n}$ is $(n(k+1)-1)$ -connected (4.12.5). By the Freudenthal Suspension Theorem (Corollary 4.16) the unit morphism $X^{\wedge n} \rightarrow \Omega\Sigma X^{\wedge n}$ is $2(n(k+1)-1)$ -connected. Since $n \geq 1$, we have that

$$2(n(k+1)-1) \geq (n+1)(k+1)-2,$$

so that both of the vertical morphisms in (5.21) are $((n+1)(k+1)-2)$ -connected. Applying Corollary 4.14 to the square (5.21), we see that the induced morphism on horizontal fibers

$$\text{fib}(J_n(X) \rightarrow X^{\wedge n}) \rightarrow \text{fib}(h_n)$$

is $((n+1)(k+1)-3)$ -connected. Therefore, to prove that the morphism $J_{n-1}(X) \rightarrow \text{fib}(h_n)$ is $((n+1)(k+1)-3)$ -connected, it suffices to show that the induced morphism

$$(5.22) \quad J_{n-1}(X) \rightarrow \text{fib}(J_n(X) \rightarrow X^{\wedge n})$$

is $((n+1)(k+1)-3)$ -connected.

Since X is k -connected, $J_{n-1}(X)$ is k -connected (Corollary 5.6) and the morphism

$$i_{n-1} : J_{n-1}(X) \rightarrow J_n(X)$$

is $(n(k+1)-2)$ -connected (Lemma 5.5). Applying the Blakers–Massey Theorem (Theorem 4.15) to the cofiber sequence

$$\begin{array}{ccc} J_{n-1}(X) & \xrightarrow{i_{n-1}} & J_n(X) \\ \downarrow & & \downarrow \\ * & \longrightarrow & X^{\wedge n} \end{array}$$

provided by Proposition 5.13, we see that morphism (5.22) is $((n+1)(k+1)-3)$ -connected. \square

Computational proof of Theorem 5.19. Let $m := (n+1)(k+1)-3$; we need to show that the map

$$J_{n-1}(X) \rightarrow \text{fib}(h_n)$$

is m -connected. The following two facts allow us to reduce to proving the claim in the case that $\mathcal{X} = \mathbf{Spc}$.

- (1) If the conclusion of Theorem 5.19 holds for the ∞ -topos \mathbf{Spc} , then it holds for any presheaf ∞ -topos.
- (2) If $L : \mathcal{Y} \rightarrow \mathcal{X}$ is a left exact left adjoint between ∞ -topoi, then L preserves: suspensions, loops, smash products, fibers, the James filtration, and m -connectedness (this last fact is [HTT, Proposition 6.5.1.16]).

Now we prove the claim for $\mathcal{X} = \mathbf{Spc}$. The claim is trivial if $n = 1$, so assume that $n \geq 2$. Since X is k -connected by assumption, the smash power $X^{\wedge n}$ is $(nk + n - 1)$ -connected (4.12.5). Since $\Omega\Sigma X^{\wedge n}$ is simply-connected, the Serre spectral sequence for (integral) homology has E^2 -page

$$E_{p,q}^2 = H_p(\Omega\Sigma X^{\wedge n}; H_q(\text{fib}(h_n))) \cong H_p(\Omega\Sigma X^{\wedge n}) \otimes H_q(\text{fib}(h_n)).$$

Since

$$H_p(\Omega\Sigma X^{\wedge n}) \cong \bigoplus_{i \geq 0} \tilde{H}_p(X^{\wedge n})^{\otimes i},$$

and $\tilde{H}_p(X^{\wedge n})^{\otimes i}$ becomes nontrivial in degree $i(nk + n + 2)$, we find that

$$H_p(\Omega\Sigma X^{\wedge n}) \cong H_p(X^{\wedge n}) \quad \text{for} \quad p < 2(nk + n + 2).$$

In particular, $E_{p,0}^2 = H_p(X^{\wedge n})$ for $p < 2(nk + n + 2)$. Consequently, the Serre spectral sequence has no nontrivial differentials off bidegrees $(p, 0)$ with $p < 2(nk + n + 2)$.

The E^2 -page of this spectral sequence is very simple if $p + q < (n + 1)(k + 1)$: in this range, $E_{p,q}^2$ vanishes unless one of p or q is zero, in which case

$$E_{p,0}^2 = H_p(X^{\wedge n}) \quad \text{and} \quad E_{0,q}^2 = H_q(\text{fib}(h_n))$$

(note that $(n + 1)(k + 1) \leq 2(nk + n + 2)$). Recall that for $p < 2(nk + n + 2)$, the Serre spectral sequence has no nontrivial differentials off bidegrees $(p, 0)$. Moreover, for $q < (n + 1)(k + 1) - 1$, are also no nontrivial differentials with target in bidegree $(0, q)$. Consequently, for

$$p + q < (n + 1)(k + 1) - 2,$$

we find that the Serre spectral sequence collapses at the E^2 -page, and therefore that

$$\tilde{H}_*(\Omega \Sigma X) \cong \tilde{H}_*(\text{fib}(h_n)) \oplus \tilde{H}_*(\Omega \Sigma X^{\wedge n}) \quad \text{for} \quad * < (n + 1)(k + 1) - 2.$$

The map $J_{n-1}(X) \rightarrow \text{fib}(h_n)$ then induces a homology equivalence in degrees $< (n + 1)(k + 1) - 2$. We conclude by the homology Whitehead Theorem. \square

Remark 5.23. In the case that $n = 2$ and $\mathcal{X} = \mathbf{Spc}$, the computational proof of the metastable EHP sequence given here reduces to the proof presented in [3, Proposition 3.1.2].

REFERENCES

- HTT J. Lurie, *Higher topos theory*, Annals of Mathematics Studies. Princeton, NJ: Princeton University Press, 2009, vol. 170, pp. xviii+925, ISBN: 978-0-691-14049-0; 0-691-14049-9.
- SAG _____, *Spectral algebraic geometry*, Preprint available at math.ias.edu/~lurie/papers/SAG-rootfile.pdf, Feb. 2018.
- Ker _____, *Kerodon*, kerodon.net, 2020.
- 1. M. Anel, G. Biedermann, E. Finster, and A. Joyal, *A generalized Blakers–Massey Theorem*, Preprint available at [arXiv:1703.09050](https://arxiv.org/abs/1703.09050), Mar. 2017.
- 2. B. Antieau and E. Elmanto, *A primer for unstable motivic homotopy theory*, in *Surveys on recent developments in algebraic geometry*, Proc. Sympos. Pure Math. Vol. 95, Amer. Math. Soc., Providence, RI, 2017, pp. 305–370.
- 3. A. Asok, K. Wickelgren, and B. Williams, *The simplicial suspension sequence in \mathbf{A}^1 -homotopy*, Geom. Topol., vol. 21, no. 4, pp. 2093–2160, 2017. DOI: 10.2140/gt.2017.21.2093.
- 4. J. Ayoub, *Un contre-exemple à la conjecture de \mathbf{A}^1 -connexité de F. Morel*, C. R. Math. Acad. Sci. Paris, vol. 342, no. 12, pp. 943–948, 2006. DOI: 10.1016/j.crma.2006.04.017.
- 5. T. Barthel and O. Antolín-Camarena, *Chromatic fracture cubes*, Preprint [arXiv:1410.7271](https://arxiv.org/abs/1410.7271), Oct. 2014.
- 6. G. Brunerie, *The James Construction and $\pi_4(S^3)$ in Homotopy Type Theory*, Journal of Automated Reasoning, vol. 63, no. 2, pp. 255–284, Aug. 2019. DOI: 10.1007/s10817-018-9468-2.
- 7. F. R. Cohen, J. P. May, and L. R. Taylor, *Splitting of certain spaces CX* , Math. Proc. Cambridge Philos. Soc., vol. 84, no. 3, pp. 465–496, 1978. DOI: 10.1017/S0305004100055298.
- 8. T. tom Dieck, *Algebraic topology*, EMS Textbooks in Mathematics. European Mathematical Society (EMS), Zürich, 2008, pp. xii+567, ISBN: 978-3-03719-048-7. DOI: 10.4171/048.
- 9. T. Ganea, *A generalization of the homology and homotopy suspension*, Comment. Math. Helv., vol. 39, pp. 295–322, 1965. DOI: 10.1007/BF02566956.
- 10. _____, *On the homotopy suspension*, Comment. Math. Helv., vol. 43, pp. 225–234, 1968. DOI: 10.1007/BF02564393.
- 11. B. Gray, *A note on the Hilton–Milnor theorem*, Topology, vol. 10, pp. 199–201, 1971. DOI: 10.1016/0040-9383(71)90004-8.
- 12. J. Grbić, *Homotopy theory and the complement of a coordinate subspace arrangement*, in *Toric topology*, Contemp. Math. Vol. 460, Amer. Math. Soc., Providence, RI, 2008, pp. 111–130. DOI: 10.1090/conm/460/09014.
- 13. A. Hatcher, *Algebraic topology*. Cambridge University Press, Cambridge, 2002, pp. xii+544, ISBN: 0-521-79160-X; 0-521-79540-0.
- 14. M. A. Hill, *On the algebras over equivariant little disks*, Preprint available at [arXiv:1709.02005](https://arxiv.org/abs/1709.02005), Sep. 2017.
- 15. P. J. Hilton, *On the homotopy groups of the union of spheres*, J. London Math. Soc., vol. 30, pp. 154–172, 1955. DOI: 10.1112/jlms/s1-30.2.154.

16. _____, *On the homotopy groups of unions of spaces*, Comment. Math. Helv., vol. 29, pp. 59–92, 1955. DOI: 10.1007/BF02564271.
17. M. Hopkins, Course notes for *Spectra and stable homotopy theory* taken by Akhil Mathew. Available at math.uchicago.edu/~amathew/256y.pdf, Fall 2012.
18. M. Hovey, *The six operations in equivariant motivic homotopy theory*, Adv. Math., vol. 305, pp. 197–279, 2017. DOI: 10.1016/j.aim.2016.09.031.
19. I. M. James, *Reduced product spaces*, Ann. of Math. (2), vol. 62, pp. 170–197, 1955. DOI: 10.2307/2007107.
20. _____, *On the suspension triad*, Ann. of Math. (2), vol. 63, pp. 191–247, 1956. DOI: 10.2307/1969607.
21. _____, *The suspension triad of a sphere*, Ann. of Math. (2), vol. 63, pp. 407–429, 1956. DOI: 10.2307/1970011.
22. M. Mather, *Pull-backs in homotopy theory*, Canadian J. Math., vol. 28, no. 2, pp. 225–263, 1976. DOI: 10.4153/CJM-1976-029-0.
23. J. P. May, *A concise course in algebraic topology*, Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 1999, pp. x+243, ISBN: 0-226-51182-0; 0-226-51183-9.
24. J. W. Milnor, *On the construction FK*, in *Algebraic Topology: A Student's Guide*, J. W. Milnor, J. F. Adams, and G. C. Shepherd, Eds., London Mathematical Society Lecture Note Series. Cambridge University Press, 1972, pp. 118–136. DOI: 10.1017/CB09780511662584.011.
25. F. Morel, *\mathbf{A}^1 -algebraic topology over a field*, Lecture Notes in Mathematics. Springer, Heidelberg, 2012, vol. 2052, pp. x+259, ISBN: 978-3-642-29513-3. DOI: 10.1007/978-3-642-29514-0.
26. F. Morel and V. Voevodsky, *\mathbf{A}^1 -homotopy theory of schemes*, Inst. Hautes Études Sci. Publ. Math., no. 90, 45–143 (2001), 1999.
27. D. Nardin, *Answer to Math Overflow Question 333239: Describing fiber products in stable ∞ -categories*, MO : 333239, Jun. 2019.
28. G. Raptis and F. Strunk, *Model topoi and motivic homotopy theory*, Doc. Math., vol. 23, pp. 1757–1797, 2018.
29. N. Rasekh, *Truncations and blakers–massey in an elementary higher topos*, Preprint available at [arXiv:1812.10527](https://arxiv.org/abs/1812.10527), Dec. 2018.
30. V. P. Snaith, *A stable decomposition of $\Omega^n S^n X$* , J. London Math. Soc. (2), vol. 7, pp. 577–583, 1974. DOI: 10.1112/jlms/s2-7.4.577.
31. M. Spitzweck and P. A. Østvær, *Motivic twisted K-theory*, Algebr. Geom. Topol., vol. 12, no. 1, pp. 565–599, 2012. DOI: 10.2140/agt.2012.12.565.
32. J. Strom, *Modern classical homotopy theory*, Graduate Studies in Mathematics. American Mathematical Society, Providence, RI, 2011, vol. 127, pp. xxii+835, ISBN: 978-0-8218-5286-6. DOI: 10.1090/gsm/127.
33. H. Toda, *Composition methods in homotopy groups of spheres*, Annals of Mathematics Studies, No. 49. Princeton University Press, Princeton, N.J., 1962, pp. v+193.
34. K. Wickelgren, *Desuspensions of $S^1 \wedge (\mathbf{P}_Q^1 \setminus \{0, 1, \infty\})$* , Internat. J. Math., vol. 27, no. 7, pp. 1640010, 18, 2016. DOI: 10.1142/S0129167X16400103.
35. K. Wickelgren and B. Williams, *The simplicial EHP sequence in \mathbf{A}^1 -algebraic topology*, Geom. Topol., vol. 23, no. 4, pp. 1691–1777, 2019. DOI: 10.2140/gt.2019.23.1691.
36. D. Wilson, *James construction*, Notes available at www.math.uchicago.edu/~dwilson/pretalbot2017/james-construction.pdf, Feb. 2017.