

$$t \in < 0, 0,5 >$$

$$h = 0,5$$

| x_i | 0 | 0,5 |
|-----------|-----|-----|
| $y_{1,i}$ | 400 | 520 |
| $y_{2,i}$ | 80 | 80 |

$$\begin{cases} \frac{dy_1}{dt} = c_1 y_1 - a_{12} y_1 y_2 \\ \frac{dy_2}{dt} = -c_2 y_2 + a_{21} y_1 y_2 \end{cases}$$

$$\begin{cases} t_0 = 0 \\ y(0) = \begin{bmatrix} y_1(0) \\ y_2(0) \end{bmatrix} = \begin{bmatrix} 400 \\ 80 \end{bmatrix} \end{cases}$$

$$f_1(400, 80) = 1 \cdot 400 - 0,005 \cdot 400 \cdot 80 = 240$$

$$f_2(400, 80) = -1 \cdot 80 + 0,0025 \cdot 400 \cdot 80 = 0$$

$$t_1 = 0,5$$

~~8/8~~
$$y_{1,1} = y_{1,0} + h \cdot f_1(400, 80) = 400 + 0,5 \cdot 240 = 520$$

~~8/8~~
$$y_{2,1} = y_{2,0} + h \cdot f_2(400, 80) = 80 + 0,5 \cdot 0 = 80$$

$$x_0 = \frac{b-a}{2} \cdot t_0 + \frac{b+a}{2} =$$

$$= \frac{0,1 - (-0,1)}{2} \cdot (-0,5774) + \frac{0,1 + (-0,1)}{2} =$$

$$= \frac{0,2}{2} \cdot (-0,5774) + \frac{0}{2} =$$

$$= 0,1 \cdot (-0,5774) = -0,05774$$

$$f(x_0) = -0,1486$$

$$x_1 = 0,05774$$

$$f(x_1) = 0,3772$$

$$S(f) = 0,1 \cdot (-0,1486 + 0,3772) = 0,02286$$

$$n=2$$

$$t_0 = -0,7746$$

$$t_1 = 0$$

$$t_2 = 0,7746$$

$$A_0 = \frac{5}{9}$$

$$A_1 = \frac{8}{9}$$

$$A_2 = \frac{5}{9}$$

$$S(f) = \frac{b-a}{2} \left(A_0 \cdot f(x_0) + A_1 \cdot f(x_1) + A_2 \cdot f(x_2) \right)$$

$$x_0 = \frac{b-a}{2} \cdot t_0 + \frac{b+a}{2} = 0,1 \cdot (-0,7746) = -0,07746$$

$$f(x_0) = -0,1721$$

$$x_1 = 0,1 \cdot 0 = 0$$

$$f(x_1) = 0$$

$$x_2 = 0,1 \cdot 0,7746 = 0,07746$$

$$f(x_2) = 0,5995$$

$$\begin{aligned} S(f) &= 0,1 \cdot \left(\frac{5}{9} \cdot (-0,1721) + \frac{8}{9} \cdot 0 + \frac{5}{9} \cdot 0,5995 \right) = \\ &= 0,02374 \end{aligned}$$

| Metoda | Stopień wielomianu (Liczba węzłów) | Wartość całki |
|------------------------------|---------------------------------------|---------------|
| Węzły Legendre'a | 1 (2) | 0,02286 |
| | 2 (3) | 0,02374 |
| | 3 (4) | 0,0237513 |
| | 4 (5) | 0,0237512 |
| Metoda złożona parabol | ~ (1000) | 0,04693 |
| Wolfram | — | 0,0237525 |