

Centro de Investigación y de Estudios Avanzados del Instituto
Politécnico Nacional

DEPARTAMENTO DE FÍSICA

TRABAJO FIN DE GRADO


Sensibilidades a radio de carga de neutrinos en experimentos de neutrinos de reactor involucrando $CE\nu NS$

Realizado por
Mario Antonio Guerra Ruiz

Para la obtención del título de
Maestra en Ciencias
en la Especialidad de Física

Dirigido por
Dr. Omar Gustavo Miranda Romagnoli

Ciudad de México, Noviembre de 2026



Centro de Investigación y de Estudios Avanzados del Instituto
Politécnico Nacional

PHYSICS DEPARTMENT

**Sensitivities to neutrino charge radius at reactor
neutrino experiments involving CE ν NS**

Presented by
Mario Antonio Guerra Ruiz

In order to obtain the
Master of Science degree
speciality in Physics

Under the supervision of
PhD Omar Gustavo Miranda Romagnoli

Mexico City, November of 2026

Agradecimientos

agradecimiento aqui.

Resumen

resumen aqui

Abstract

resumen en ingles

Contents

1	Introduction	1
2	Electroweak theory in the Standard Model of particles	2
2.1	Introduction	2
2.2	Symmetries and fermion content	2
2.2.1	Neutrinos	2
2.2.2	Weak Nuclear Force	3
2.3	Massive and Massless Mediators	5
2.4	Electroweak mixing $\sin^2 \theta_W$	5
3	Radiative Corrections and Neutrino Charge Radius	6
3.1	Introduction	6
3.2	Neutrino Charge Radius	6
3.3	One-loop Corrections to the $\nu\nu\gamma$ vertex	6
3.4	NCR as a radiative correction of Electroweak Mixing	6
4	Coherent Elastic Neutrino-Nucleus Scattering	7
4.1	Introduction	7
4.2	Cross Section	7
4.3	Experiments	7
4.4	Quenching Factors	7
4.5	Neutrino Sources	7
4.6	Form Factors	7
5	SBC-ININ-LAr	8
5.1	Results	8
6	Summary and conclusions	10
	Bibliography	12

1. Introduction

intro

2. Electroweak theory in the Standard Model of particles

2.1. Introduction

The Standard Model of particles (SM) is the most currently utilized theory of interactions of subatomic particles. It describes all matter particles and forces between them, excluding gravity, as interactions between 12 fermions (6 quarks and 6 leptons) with their corresponding anti-fermions, 4 spin-1 force-carrying gauge bosons and 1 scalar boson as shown in Fig. 2.1. Within this framework exist 3 distinct forces: strong nuclear force, mediated by the gluon (g) boson; electromagnetic (EM) force, mediated by the photon (γ) boson; and weak nuclear force, mediated by the $W^\pm$ and Z bosons. This work pertains to the description of the properties of the only family of particles that present no electric or color charge known as neutrinos (ν), wherein the SM exist three known flavors (ν_e , ν_μ and ν_τ) corresponding to the latter half of (electro)weak isospin doublets formed with the known charged leptons (e^- , μ^- and τ^-). This chapter aims to describe neutrinos in the SM, the weak force and its origin as a part of a larger broken symmetry known as "Electroweak Theory" (EW).

2.2. Symmetries and fermion content

2.2.1 Neutrinos

Neutrinos are charge-less, color-less, spin- $\frac{1}{2}$ particles with negligible masses compared to other fundamental particles first proposed in 1930 by Pauli W. [1], and later detected by C. L. Cowan et al. in 1956 [2], to solve known apparent contradictions in conserved quantities of beta decays. Beta decay was known to be as the emission of beta particles (electrons) by the nuclei of unstable atoms, wherein the difference in mass of the original and resulting nuclei of the decay should fix the energy at which the emitted particles are detected. Also of note, due to conservation of angular momentum the sum of the spin (S_e) of the electron, the spin of the resulting nucleus (S_{N^*}) and the angular momentum between them (L) should equal the original nucleus' spin (S_N). Despite this, it was known experimentally that the resulting electrons were emitted in a spectrum of energies less than expected and at $L = 0$, making the sum of spins impossible to match. Pauli's solution to the apparent violation of the conservation of these quantities was to hypothesize the existence of a second neutral spin- $\frac{1}{2}$ with very small mass compared to the electron that allows for both the distribution of energy between both emitted particles, thus letting the electron be emitted at different energies depending on the kinematics, and the conservation of




Figure 2.1: Particles in the SM

angular momentum.

$$\begin{array}{ccc}
 N \rightarrow N^* + e^- & \rightarrow & N \rightarrow N^* + e^- + \nu \\
 m_N = m_{N^*} + E_{e^-} & \rightarrow & m_N = m_{N^*} + E_{e^-} + E_\nu \\
 S_z N = S_z N^* + S_z e^- & \rightarrow & S_z N = S_z N^* + S_z e^- + S_z \nu
 \end{array}$$

Later refinement of the atomic model found that nuclei were made up of positively charged elements named protons (p) and neutrally charged elements known as neutrons¹ (n), leaving the small hypothesized particle to be named neutrino (ν) later by Enrico Fermi. It is well understood now that the beta decay is a process in which a neutron within an unstable nucleus decays into a proton, an electron and a(n) (anti)neutrino.

$$n \rightarrow p + e^- + \bar{\nu}$$

Subsequent experiments exploiting cross symmetry of the above interaction [2], accelerator-produced pion decay [3], and charged-current tau neutrino interactions in nuclear emulsion [4, 5] confirmed the existence of three distinct active neutrino flavors and the conservation of lepton family number in the interactions that involved these particles. Given that neutrinos don't interact via electromagnetic or strong forces a theory of a different force that mediated these interactions was developed into what is now known as the "weak nuclear force".

2.2.2 Weak Nuclear Force


Developments in quantum mechanics (QM) and special relativity (SR) led up to the first theories that could describe the behavior of some subatomic particles and their

¹this was the name first proposed for the particle hypothesized by Pauli, later changed

interactions through excitations of quantized fields called quantum field theories (QFTs) obtained by promoting classical fields to operator-valued quantities satisfying canonical (anti)commutation relations and deriving their dynamics from a Lorentz-invariant action principle. One of the earliest and most successful QFTs was Quantum Electro Dynamics (QED), describing how spin-1/2 (anti)particles ($\bar{\psi}$) interacted with an EM 4-potential (A^μ) by minimizing the action given by the lagrangian density:

$$\mathcal{L}_{\text{QED}} = \underbrace{\bar{\psi}(i\gamma^\mu\partial_\mu - m)\psi}_{\text{free Dirac field}} - \underbrace{\frac{1}{4}F_{\mu\nu}F^{\mu\nu}}_{\text{free EM field}} - \underbrace{e\bar{\psi}\gamma^\mu\psi A_\mu}_{\text{interaction}}. \quad (2.1)$$

While the ‘free’ terms in (2.1) describe the propagation of Dirac fermions and photons in free Minkowski space, the ‘interaction’ term describes a vertex that couples the (anti)fermion current with the electromagnetic four-potential, representing the elementary process of a(n) (anti)fermion emitting or absorbing a photon. At zeroth-order $\hbar$ perturbative terms, QED had shown predictive power for processes like Compton ($\gamma + e^- \rightarrow \gamma + e^-$) [6] and Møller ($e^- + e^- \rightarrow e^- + e^-$) [7] scattering by the time Fermi first proposed an interaction inspired by the QED interaction term, where instead of a fermion current emits a photon, two fermion currents meet at a vertex such that a coupling constant (now called Fermi’s constant G_F) quantified the intensity of the interaction, and thus can be estimated experimentally.



$$\sim e\bar{\psi}_f\gamma^\mu\psi_f A_\mu$$

$$\sim \frac{G_F}{\sqrt{2}}(\bar{\psi}_n\gamma^\mu\psi_p)(\bar{\psi}_v\gamma_\mu\psi_{e^-})$$

Subsequent discoveries in particle physics revealed that charged-current weak interactions violate parity symmetry maximally. First theorized by Lee and Yang in 1956 [8] to resolve the then-known $\tau - \theta$ puzzle, and later verified experimentally by Wu et al. in 1957 [9], this motivated a modification of the Fermi interaction in the form of a combination of the two bilinear Lorentz-invariant structures that admit such violation [10, 11].

$$V : j_V^\mu = \bar{\psi}\gamma^\mu\psi \quad \xrightarrow{\mathcal{P}} \quad j'_{V\mu} = \bar{\psi}\gamma_\mu\psi \quad (2.2)$$

$$A : j_A^\mu = \bar{\psi}\gamma^\mu\gamma^5\psi \quad \xrightarrow{\mathcal{P}} \quad j'_{A\mu} = -\bar{\psi}\gamma_\mu\gamma^5\psi \quad (2.3)$$

The expressions (2.2) and (2.3) show how different bilinear structures change under parity ($\mathcal{P}$) and how pure vector (axial) currents are $\mathcal{P}$ – even ($\mathcal{P}$ – odd). Given a general current structure, $j^\mu = g_V j_V^\mu + g_A j_A^\mu$, observables that depend on the inner product of two such currents transform under parity as:

$$|\mathcal{M}|^2 = (j_1 \cdot j_2) = g_V^2 (j_{1V} \cdot j_{2V}) + g_A^2 (j_{1A} \cdot j_{2A}) + g_V g_A (j_{1V} \cdot j_{2A} + j_{1A} \cdot j_{2V}) \quad (2.4)$$

$$\xrightarrow{\mathcal{P}} \quad (2.5)$$

$$|\mathcal{M}'|^2 = (j'_1 \cdot j'_2) = g_V^2 (j'_{1V} \cdot j'_{2V}) + g_A^2 (j'_{1A} \cdot j'_{2A}) - g_V g_A (j'_{1V} \cdot j'_{2A} + j'_{1A} \cdot j'_{2V}).$$

The only difference between 2.4 and ?? lies in the sign of the interference term, such that the degree of $\mathcal{P}$ -symmetry violation is given by the coefficient $\frac{g_V g_A}{g_V^2 + g_A^2}$. Normalizing this there are two options for maximally violated $\mathcal{P}$ -symmetry given by $\frac{g_V}{g_A} = \pm 1$, also known as $V \pm A$ currents. Given that in the Wu experiment it was determined that (anti)neutrinos are emitted exclusively as left(right)-handed fermions the appropriate structure to use for weak interactions was determined to be $V - A$. Modifying the Fermi lagrangian density for beta decay to include this gives the following expression:

$$\mathcal{L}_{CC} = \frac{G_F}{\sqrt{2}} \left[\bar{\psi}_{\nu_e} \gamma^\mu (1 - \gamma^5) \psi_e \right] \left[\bar{\psi}_p \gamma_\mu (1 - \gamma^5) \psi_n \right] + \text{h.c.} \quad (2.6)$$

Later refinements of this theory succeeded in demonstrating the universality of the weak interaction across leptonic decays and strangeness-changing meson decays through Cabibbo mixing [12]. This not only established that all such processes are mediated by the same underlying force, but also motivated the prediction and subsequent discovery of the charm quark [13]. With these improvements on the theory thus far, the point-like structure of the interaction still led to scattering amplitudes that were proportional to center-of-mass energy $\sqrt{s}$ and to probability violation at higher energies. In direct analogy with QED, where the point-like Coulomb interaction is resolved into the exchange of a massless photon, the Fermi vertex can be interpreted as the low-energy limit of the exchange of a charged spin-1 $W^\pm$, wherein a mass is considered for such a mediator in order to explain the short range of weak interactions.

2.3. Massive and Massless Mediators

2.4. Electroweak mixing $\sin^2 \theta_W$

3. Radiative Corrections and Neutrino Charge Radius

3.1. Introduction

3.2. Neutrino Charge Radius

3.3. One-loop Corrections to the $\nu\nu\gamma$ vertex

3.4. NCR as a radiative correction of Electroweak Mixing

4. Coherent Elastic Neutrino-Nucleus Scattering

4.1. Introduction

4.2. Cross Section

4.3. Experiments

4.4. Quenching Factors

4.5. Neutrino Sources

4.6. Form Factors

5. SBC-ININ-LAr

5.1. Results

6. Summary and conclusions

Bibliography

- [1] Wolfgang Pauli. “Letter to the Physical Institute of the Federal Institute of Technology, Zürich”. Open letter to the Gauverein meeting in Tübingen. Reproduced in L. M. Brown, “The idea of the neutrino,” *Physics Today*, vol. 31, no. 9, pp. 23–28, 1978. Also in R. Kronig and V. F. Weisskopf (eds.), *Collected Scientific Papers by Wolfgang Pauli*, vol. 2, Interscience, New York, 1964. 1930. URL: <http://www.math.utah.edu/~beebe/talks/2015/qtm/pdf/pauli-1930-ltc.pdf>.
- [2] C. L. Cowan et al. “Detection of the Free Neutrino: a Confirmation”. In: *Science* 124.3212 (1956), pp. 103–104. DOI: [10.1126/science.124.3212.103](https://doi.org/10.1126/science.124.3212.103). eprint: <https://www.science.org/doi/pdf/10.1126/science.124.3212.103>. URL: <https://www.science.org/doi/abs/10.1126/science.124.3212.103>.
- [3] G. Danby et al. “Observation of High-Energy Neutrino Reactions and the Existence of Two Kinds of Neutrinos”. In: *Physical Review Letters* 9.1 (1962), pp. 36–44. DOI: [10.1103/PhysRevLett.9.36](https://doi.org/10.1103/PhysRevLett.9.36).
- [4] K. Kodama et al. “Observation of Tau Neutrino Interactions”. In: *Physics Letters B* 504.3 (2001), pp. 218–224. DOI: [10.1016/S0370-2693\(01\)00307-0](https://doi.org/10.1016/S0370-2693(01)00307-0). arXiv: [hep-ex/0012035](https://arxiv.org/abs/hep-ex/0012035).
- [5] K. Kodama et al. “Final tau-neutrino results from the DONuT experiment”. In: *Physical Review D* 78 (2008), p. 052002. DOI: [10.1103/PhysRevD.78.052002](https://doi.org/10.1103/PhysRevD.78.052002).
- [6] Oskar Klein and Yoshio Nishina. “Über die Streuung von Strahlung durch freie Elektronen nach der neuen relativistischen Quantendynamik von Dirac”. In: *Zeitschrift für Physik* 52 (1929), pp. 853–868. DOI: [10.1007/BF01366453](https://doi.org/10.1007/BF01366453).
- [7] Christian Møller. “Zur Theorie des Durchgangs schneller Elektronen durch Materie”. In: *Annalen der Physik* 406 (1932), pp. 531–585. DOI: [10.1002/andp.19324060506](https://doi.org/10.1002/andp.19324060506).
- [8] T. D. Lee and C. N. Yang. “Question of Parity Conservation in Weak Interactions”. In: *Phys. Rev.* 104 (1956), pp. 254–258. DOI: [10.1103/PhysRev.104.254](https://doi.org/10.1103/PhysRev.104.254).
- [9] C. S. Wu et al. “Experimental Test of Parity Conservation in Beta Decay”. In: *Phys. Rev.* 105 (1957), pp. 1413–1415. DOI: [10.1103/PhysRev.105.1413](https://doi.org/10.1103/PhysRev.105.1413).
- [10] E. C. G. Sudarshan and R. E. Marshak. “Chirality Invariance and the Universal Fermi Interaction”. In: *Phys. Rev.* 109 (1958), pp. 1860–1862. DOI: [10.1103/PhysRev.109.1860](https://doi.org/10.1103/PhysRev.109.1860).
- [11] R. P. Feynman and M. Gell-Mann. “Theory of the Fermi Interaction”. In: *Phys. Rev.* 109 (1958), pp. 193–198. DOI: [10.1103/PhysRev.109.193](https://doi.org/10.1103/PhysRev.109.193).
- [12] N. Cabibbo. “Unitary Symmetry and Leptonic Decays”. In: *Phys. Rev. Lett.* 10 (1963), pp. 531–533. DOI: [10.1103/PhysRevLett.10.531](https://doi.org/10.1103/PhysRevLett.10.531).
- [13] S. L. Glashow, J. Iliopoulos, and L. Maiani. “Weak Interactions with Lepton-Hadron Symmetry”. In: *Phys. Rev. D* 2 (1970), pp. 1285–1292. DOI: [10.1103/PhysRevD.2.1285](https://doi.org/10.1103/PhysRevD.2.1285).