

PESTICIDES IN FRUITS AND VEGETABLES



PESTICIDES IN FRUITS AND VEGETABLES

A PROJECT REPORT

Submitted by

G. Divya

203307137396

B. Durga Maha Lakshmi

203307137400

M. Sandhya Rani

203307137404

N. Thanusha Rani

203307137411

K. Ishwariya

203307137398

K. Anitha

203307137402

M. Sai Prasanna

203307137409

P. Naga Durga Mounika

203307137412

P. Hima Bindu

203307137415

K. Sruthi

203307137401

R. Likitha

203307137403

M. Sai Jyothi

203307137410

P. Kanaka sri

203307137413

In partially fulfilment for the award of the degree of

Bachelor of Science

Under the guidance of

Sri O.G.V. SAI DIVYA



DEPARTMENT OF MATHMATICS

SIR C. R. REDDY COLLEGE FOR WOMEN

(Affiliated to Adikavi Nannaya University, Rajamahendravaram)

ELURU, W.G.Dist., (A.P)

2021-2022

SIR C. R. REDDY COLLEGE FOR WOMEN

(Affiliated to Adikavi Nannaya University, Rajamahendravaram)

ELURU, W.G.DIST., (A.P)



CERTIFICATE

This is to certify that this is a bonafide record of the project work entitled "**Pesticides in Fruits and Vegetables**" done by _____
bearing Regd. No _____, a student of I B.Sc., 1st Semester in the
Department of Mathematics of **Sir C. R. Reddy College for Women**, Eluru, during the
academic year 2021-2022

Signature of the Lecturer

Signature of
Head of the Department

Signature of the Examiners

DECLARATION

We hereby declare that the project entitled “*Pesticides in Fruits and Vegetables*” authentic work carried out by us under the esteemed guidance **Sri O. G. V. Sai Divya** for the partially fulfilment of the award of Bachelor of Science (B.Sc) degree and we further admit that this has not been submitted by us anywhere else for the award of any other degree.

Name:

Regd. No:

ACKNOWLEDGEMENT

We feel highly grateful to Smt. P.Sailaja, Principal, Sir C.R.R College for Women, Eluru, for being a motivating factor throughout the course of the project.

We are thankful to **Smt. S.S.L. Sabari Kumari**, Head of the Department, Department of Chemistry, Sir C R Reddy College for Women, Eluru for providing necessary facilities during our work

We take this opportunity to express my deep sense of gratitude to our project guide, **Sri. O.G.V. Sai Divya**, for her keen interest and valuable guidance right from the inception of the project till the complete execution of the same. We also thank other staff for their help.

I am using this opportunity to express my gratitude to all the staff who supported me throughout the course of this project. I am thankful for their aspiring guidance, invaluable constructive criticism and friendly advice during the project work.

We will be failing in our duty if we do not acknowledge the continued encouragement and support we have had from all our family members, without whose help this project would not have been completed.

CONTENTS

Chapter No	Title	Page No
1	Introduction	1
	Scope	4
	Objectives	8
2	Methodology	11
	Primary Data	12
	Secondary Data	23
3	Questionnaire	24
4	Data Analysis	28
	Problem Identification	29
	Data Tables	33
5	Conclusion and Recommendation	44
6	References	49

INTRODUCTION

Chapter 1

INTRODUCTION

Pesticides are a numerous and diverse group of chemical compounds. Which are used to eliminate pests in agriculture and house holds. They enable the quantities and the quality of crops and food to be controlled and help to limit the many human diseases transmitted by insect or rodent vectors. However despite their many merits, pesticides are some of the most toxic environmentally stable and mobiles substances in the Environment their excessive use has a deleterious effect on humans and the environment their presence in food is particularly dangerous with their Environmental stability, ability to and toxicity, pesticides may places the human body at greater risk of disease and poisoning.

Pesticides enter the environment in various forms (e. g: powders, moistened powders, for preparing aqueous solutions and concentrates for making up emulsion or sprays) pesticides are of enormous importance increasing the yields and quality of agriculture products they are used to:

- ⇒ Control the numbers of pests destroying whole plants or their parts;
- ⇒ Increase the production of animal and plant biomass
- ⇒ Combat microorganisms causing farm produce to rot and to decay
- ⇒ Combat algal, bacteria, fungi, and weeds;
- ⇒ Combat animal pests damaging crops (e. g: mites, aphids, insects, larval and nematodes) ;
- ⇒ Stimulate or inhibit plant-growth processes (e. g: remove Excess flowers, destroy foliage or dry out plants) ;
- ⇒ Make possible the action of other substances
- ⇒ Counteract growths on boats and ships
- ⇒ Kill harmful organisms in the farm buildings the home, hospitals , stores and vehicles



The widespread use of pesticides not only contaminates water, soil, and air but also causes their accumulation in crops (e. g :fruits and vegetables). Pesticides are transported mainly by rain and wind from their points of application to crops and land. Where their presence be undesirable or harmful.

A pesticides can be defined as any chemical used to directly control pest populations or to prevent or reduce pest damage. Pesticides also include compounds intended for use as plant growth regulators, defoliants, or desiccants even though they are not normall used as pest control agents, nor are they usually effective as such. It is important to remember that the"-cide" in pesticides means "to kill", these products can be dangerous if not used properly.

Pesticides use has increased agricultural production worldwide and there by contributed to foodsecurity . With the expect 30% increase of world population more than 1100 pesticides have been widely used in various combinations at different stages of cultivation and postharvest storage. Despite their Usefulness' pesticides could potential risks to food safety and the environment. Pesticides could possible remain in foodstuff, such as fruits and vegetables which pose a potential threat to human and animal the use of pesticides during production of ten leads to the presence of pesticides in fram .



SCOPE

SCOPE

Pesticides use with the expected 30% increase of world population 9.2 billion by 2050, there is a projected demand to increase food production by 70% according to the calculation by population, there will be a continuing need for pesticides based solution. For pest control and food security in the future to increase food product to meet the increasing food supply more than 1100 pesticides have been widely used in various combinations at different stages of cultivation and postharvest storage. Despite their usefulness, pesticides could pose potential risks to food safety and the environment pesticides residues could possibly remain in food stuff.

Such as fruits and vegetables, which pose a potential threat to human and animals health for this reason, most countries or regions have drafted rigorous regulations to levels for compliance. However, pesticides in crop production and to monitor their residues are one of the most frequently detected xenobiotic in food analysis, especially in enormously consumed fruits and vegetables all over the world.

The EPA PGP will cover pesticides authorized for use under FIFRA, and generally includes pesticides application covered by the 2006 rule.

1. The EPA PGP will cover the following pesticides use pattern for control of.
2. Mosquitoes and other flying insect pesticides
3. Weed and algae
4. Animals pests
5. Forest canopy pest

Pesticides are among leading causes of death by self poisoning, in particular in low and middle income countries as they are intrinsically toxic and deliberately spread in the environment, the production regulations and use of pesticides require strict regulation and control. Regular monitoring of residues in food and the environment is also required.

Wide scope screening of pesticides fruits and vegetables using information dependent aca vision employing 41+PLC-TOF-MS and automated Ms/Ms library for simultaneous screening and identification of 427 pesticides in showed that 97.4% of the pesticides in fresh fruits and vegetables had the screening detection limit less than 50mg/kg and more than 86.7% pesticides two objectives in relation pesticide.

1. To ban pesticides that are most toxic to human, as well as pesticides that remain for the longest time in the environment.

2. To protect public health by setting maximum limits for pesticides residues in food and water.

Scope of pesticides in fruits and vegetables:

- ❖ Pesticides use with the expected 30% increase of world population 9.2 billion by 2050, there is a projected demand to increase food production by 70% according to the calculation by Population, there will be a continuing need for pesticides based solution.
- ❖ For pest control and food security in the future to increase food product to meet the increasing food supply more than 1100 pesticides have been widely used in various combinations at different stages of cultivation and postharvest storage.
- ❖ Despite their usefulness. Pesticides could pose potential risks to food safety and the environment pesticides residues could possibly remain in food stuff.
- ❖ Such as fruits and vegetables, which pose a potential threat to human and animals health for this reason, most countries or regions have drafted rigorous regulations to levels for compliance.
- ❖ However, pesticides in crop production and to monitor their residues are one of the most frequently detected xenobiotic in food analysis, especially in enormously consumed fruits and vegetables all over the world.
- ❖ The EPA PGP will cover pesticides authorized for use under FIFRA, and generally includes pesticides application covered by the 2006 rule.

The EPA PGP will cover the following pesticides use pattern for control of.

1. Mosquitoes and other flying insect pesticides
2. Weed and algae
3. Animals pests
4. Forest canopy pest

- ❖ Pesticides are among leading causes of death by self poisoning, in particular in low and middle
- ❖ Income countries as they are intrinsically toxic and deliberately spread in the environment, the
- ❖ Production regulations and use of pesticides require strict regulation and control. Regular
- ❖ Monitoring of residues in food and the environment is also required.
- ❖ Wide scope screening of pesticides fruits and vegetables using information dependent HPLC-TOF-MS and automated MS/MS library for simultaneous screening and identification of 427 pesticides in showed

that 97.4% of the pesticides in fresh fruits and Vegetables had the screening detection limit less than 50mg/kg and more than 86.7% pesticides

❖ Two objectives in relation pesticide.

1. To ban pesticides that are most toxic to human, as well as pesticides that remain for the longest time in the environment.
2. To protect public health by setting maximum limits for pesticides residues in food and water.
3. A total of 111 pesticides were analyzed in vegetable samples. Out of a total of 111 pesticides, 85 pesticides were analyzed on LCMS/MS and remainder 26 pesticides were on GC-ECD. GC-MS was used for ion confirmation for 26 analytes.

2.4.1 Gas chromatography-electron capture detector (GC-ECD) analysis

4. A total of 26 compounds were analyzed with the help of a Thermo Scientific Trace 1310 gas chromatograph with an auto-sampler (AS 1310), coupled with an Electron Capture Detector (ECD).
5. The separation was performed on a Restek- Rxi-5ms capillary column (30 m long \times 0.25 mm internal diameter \times 0.25 μ m film thickness).
6. Sample injection was operated in the splitting mode 1:10, with an injector temperature of 250°C and the detector temperature was 300°C.
7. The oven temperature was held initially at 110°C for 1 minute, programmed from 110 to 210°C at 10°C/min. and held for 1 minute at 210°C, further programmed from 210 to 300°C at 5°C/min. and finally held at 300°C for 6 minutes.
8. Makeup gas with a flow of 25.0 mL/min.
9. The Thermo Scientific Chromeleon 7.2 version software was used for system control and acquisition.
10. 2.4.2 Liquid chromatography-tandem mass spectrometry analysis total of 85 pesticides were analyzed by LCMS.

OBJECTIVES

OBJECTIVES

Analyse food and chemical issues determine how science affected food through production, packing and health. Debate and benefits and risks of pesticides use Calculate the standard liquid measurements for mixing of agricultural chemicals.

To develop a method of accessing imports on human health and ecosystems considering the most used pesticides to make the general method operable for any kind of crops growth in

Mediterranean green house

1. Understand pesticides.
2. Identify common four misted chemicals.
3. Determine the deference between chronic and acute toxicity.
4. Discuss explosive routes of chemicals.
5. Identify personal protective equipment (PPE).
6. Examine signs of pesticides poisoning.
7. Look at first and treatment.
8. Recognize proper storage and disposal practices.
9. To describe occurrence in the environment & sampling regime of atrazine in public water suppliers.
10. To assess Kentucky water systems complete with the EPA regulations identity the health implications & public health role.
11. To determine if the current monitoring regions is adequate in detecting atrazine levels of comer with focus on the critical time of the exposure the atrazine.
12. To investigate pesticides abuse on fruits and vegetables in Tianjin area to detect pesticide residues in fruits and vegetables.
13. To study the methods for reducing pesticides residues in fruits and vegetables

METHODS

- ❖ A questionnaire on the pesticides application during growing in fruits and vegetables was administered to 185 farmers Tianjin area.
- ❖ According to the information from the questionnaire survives fruits and vegetables samples was collected in four seasons around the year and measured for organophosphorus pesticides residues by gas chromatography.
- ❖ Fruits and vegetables samples contained pesticides residue were treated by scald immersion in 0.15 % and 0.30 % detergent solution immersion impure water peeling cutting root and pesticides residues were measured before and after the treatment.
- ❖ To ban pesticides that most toxic to humans, as well as pesticides that remain for the longest time in the environment.
- ❖ To protect public health by setting maximum limits for pesticides residues in food and water.

METHODOLOGY

Chapter 2

METHODOLOGY

In addition, fruit also we have found in 20 types of pesticides residue in mango trees found 21 types of in orange trees. In the method of minimizing pesticides residue that the washing with water first soak in water 10 min flow

STEP 1 : collection of respective samples.

STEP 2 : Isolation of residue extraction .

STEP 3 : Separation of co-extracted matrix components clean up.

STEP 4 : Identification and quantification of far get analysis.

STEP 5 : confirmation of results by an additional analysis.

Methodology are two types

1. primary method
2. Secondary method.

Primary method

Some primary methods are

- *Dusting
- * Sparing
- * Granular
- * Speed pelleting/ seed dressing
- * Seedling root dip
- * Sett treatment
- * Trunk/ steam injection
- * Padding

Dusting :

Dusting In carried out in the morning hours and during very light air stream. It can be done manually or by using dusters. Some times dust can be applied in soil for cheaper and suited for dry land crop pest control.

Spraying :

Spraying is normally carried out by mixing EC or WP formulations in water.



Grnular:

Highly toxic pesticides are handled safely in the form of granules. Granules can be applied

directly on the soil or in the plant part

- Broadcasting
- In furrow apply
- Seed dressing
- Spot application
- Ring
- Root zone
- Pralinage
- Broadcasting :

Granules are mixed with equal quantity of sand broadcasted directly on the or in the film of standing water rice nursery carbonfuran 3%G applied 1.45 kg/8 cent rice nursery in this film of water and impound water for 3 days.

• In furrow application:

Granules are applied at the time of sowing in furrow applied@3g sorghum shoottly.

- **Seed dressing :**

After the establishment of the plants the granules are applied a little away from the plant (10-15cm) in a furrow.



- **Spot application :**

Granules are applied @ 5cm away and 5 cm on the sides of plant. This reduces the quantity of insecticide required.

- **Ring :**

Granules are applied in a ring form around the trees.

- **Root zone :**

Granules are encapsulated and placed in the root zone of the plant by mixing it with equal quantity of sand in the central whorl of crops like sorghum maize fruit , vegetables ,sugar cane to control internal borders.

- **Pralinage :**

This surface of banana sucker intended for planting is trimmed . The sucker is dipped in wet clay slurry and carbon 3G is sprinkled to control burning nematode

- **Speed pelleting/speed dressing :**

The insecticide mixed with seed before sowing sorghum seed are treated with chlorpyrifos to control shootfly. The carbendazim 50 sp and imidacloprid is directly used as day seed dressing insecticide against cotton sucking pests.

Seedling root dip :

It is followed to control early stage pests in fruits to control sucking pests and stem borer in early transplanted crop a shallow pit lined with polythene sheet is prepared in the field. To this 0.5 kg urea in 2.5 litres of water prepared separately are poured. The solution is made up to 50ml with water and the root of seedlings in bundles are dipped for 20min before transplanting

- **Sett treatment:**

Treat the sugar cane and fruits setts in 0.05% malathion for 15 min to protect them from scale. Treat the fruits setts in 0.05% Imidacloprid 70ws @ 175g/ha for 15min to protect them from termites.

- **Trunk/steam injection :**

This method is used for the control of coconut pests like black headed caterpillar mite etc., Drill a downward slanting hole of 1.25cm diameter to a depth of 5cm at a height of about 1.5 m above ground level and inject 5ml of monocrotophos 36 WSC into the stem and plug the hole with cement or clay mixed with a fungicide. For stem injection of banana an injecting gun or hypodermic syringe is used for the control of banana aphid vector of bunchy top diseases.

- **Padding :**

St borers of mango, silk and cotton and cashew can be controlled by this method. Bark of infected tree (5*5 cm) is removed on three sides leaving bottom as a flap. Small quantity of absorbent cotton is placed in exposed area and 5-10ml of monocrotophos 36 WSP is added using an inc filler, close the flap and cover with clay mixed with fungicide.



Name : P . Kanaka Sri

Roll Number: 203732

Farmer
name: Venkatalakshmi

Age: 45

Type of farming:
Tamota

Area : Vatluru

Name : M . Sai Jyothi

Roll Number:
203729

Farmer name :
Ramarao

Age: 37

Type of farming:
Banana

Area : Nagavaram





Name:M . Sai Prasana

Roll Number:203728

Farmer name: Lakshmi

Age: 47

Type of farming:
Brinjal

Area : Denduluru

Name: P . Naga Durga
Mounika

Roll Number: 203731

Farmer name: p.
Venkateswararao

Age: 40

Type of farming:
banana

Area :
Ramaraogudem





Name:K . Ishwariya

Roll Number:
203723

Farmer name: Prasad

Age: 49

Type of farming: corn

Area : Jogannapalem

Name: K . Anitha

Roll
Number:203725

Farmer name:
Nirmala

Age: 50

Type of farming:
Mirchi

Area:
Bommuluru





Name: N.Thanusha
Rani

Roll Number: 203730

Farmer name:
M.srinivasa

Age: 50

Type of farming:
Guava

Area: Thotagudem

Name: G . Divya

Roll
Number:203721

Farmer name : ck
sindhu

Age: 29

Type of farming:
Apple

Area:Bommuluru





Name: M . Sandhya

Roll Number:203704

Farmer name: Yesu babu
. madasi

Age:42

Type of farming: corn

Area: Krishnapuram

Name:K . Sruthi

Roll
Number:203724

Farmer name:
Baburao

Age:40

Type of farming:
Mirchi

Area: Thotagudem





Name:Likitha.R

Roll Number: 203726

Farmer name:
R.Koteswara Rao

Age:45

Type of farming:
Coconut

Area: Pedavegi

Name:Durga
mahalakshmi

Roll Number: 203703

Farmer name:
Koteswarao

Age:48

Type of farming:
sugar cane

Area: Vatuluru





Name:P . Hima Bindhu

Roll Number: 203734

Farmer name: Y . Chanti

Age: 55

Type of farming: Mango

Area: Hanuman
Junction

Secondary Methodology For Pesticides

Both EPA'S National Pesticides and National water programs assess the effect of pesticides on aquatic ecosystems using approaches were developed with high quality data using rigorously peer reviewed assessment methodology. In characterizing pesticides effects as well as chronic and sub-lethal effects on growth survival and reproduction in their assessments.

- *Are legally defense under our statutory mandates.

- *Are implementable at the federal and state level.

- *Are developed as quickly and efficiently as possible.

- *Reflect stakeholders input and comments.

- *Multi residue determination of Pesticides in wine by electrospray ionization liquid chromatography tandem mass spectrometry (Ls/Ms/Ma).

- *Multi residue determination of Pesticides in wine by GL/MS triple quad (GL/MS/MS).

- *Limit of quantification:0.010ppm(10ppb) consider the lowest worldwide standard for most agrochemicals (MRLS).

- *Values <0.01ppm consider non-defectable.

- *Includes fungicides, insecticides,aracnicides and herbicides.

- *Lab values stored in LCMS database

The water flow third wash are at a time for 20mins.It can reduce the pesticides vesidue up to 60-70% well get fruits and vegetables that have less pesticides residue vinegal baking soda.Turmeric powered is also used to remove Pesticides and help to kill germs in fruits and vegetables.

Current Activities For Developing Common Effects Methodology For Pesticides:

A FIFRA (Federal Insecticides, Fungicides and Rodenticide Act) Scientific Aduisory pane(SAP) meeting was held in late January 2012.The FIFRA SAP is composed of biological statisticians toxicologists and other experts who provide independent scientific advice to EPA on a wide range of health and safety issues related to pesticides a common effects methodology.

The SAP review was highly favorable and the SAP made a number of recommendations for EPA to move forward with their approach in a report issued in April 2012.

QUESTIONNAIRE

Chapter 4
QUESTIONNAIRE

SIR CRR COLLEGE FOR WOMEN

TITLE : PESTICIDES IN FRUITS AND VEGETABLES

QUESTIONNAIRE

NAME:

What type of crops are _____

1.Experiences in vegetables producing ?

(a)< 5 years (b) 5-10 years (c) > 10 years

2.Protection measure during spraying ?

(a) no gloves and masks (b) gloves & masks

3. where do you store pesticides?

(a) store room (b) garden,openfield

(c) coolstorages,refrigerator

4. Do you follow instructions on the label

(a) yes (b) No

5. How many hours do you work in the field

(a) 3-4hrs (b) 5-7hrs (c) >7 hrs

6. How many pesticides do you apply per month

(a) 1 (b) 2 (c) none of these

7. which type of field you have
(a) vegetables (b) fruits
8. pesticides application timing
(a) morning (b) afternoon (c) none
9. positions of spray head
(a) 20cm above plants tops
(b) at plant tops
(c) targeted insects & diseases
10. pesticide spraying techniques in field
(a) spray against the wind (b) spray with the wind
(c) walk forward (d) walk backward
11. Number of spray application per crop
(a) <3 (b) 3-4 (c) 5-7 (d) >7
12. pesticides are poisonous
(a) agree (b) natural (c) disagree
13. sign of poisoning with pesticides
(a) burning eyes (b) cough, headache
(c) skin inflammation (d) others

14. Type of pesticides applicator

- (a) knapsack sprayer (b) sprayer hose
- (c) motorise sprayer (d) none of the above

15. Do you think that pesticides affect the environment

- (a) Strongly agree (b) agree
- (c) Strongly disagree (d) disagree

16. which age is harmful to pesticides

- (a) Below 20 (b) 25
- (c) Above 50 (d) all of the above

17. which disease to causes the pesticides

- (a) T.B (b) cancer
- (c) sugar (d) none of the above

18. Insecticides kill

- (a) Both harmful and useful insects (b) Specific insects
- (c) Harmful insects (d) None

19. The most common pesticides used in india is

- (a) DDT (b) Endrin
- (c) BHC (d) both a&b

20. Pesticides generally attack

- (a) Muscular system (b) Respiratory system
- (c) Nervous system (d) None

PROBLEMS IDENTIFICATION

Chapter 4

PROBLEM IDENTIFICATION


- pesticides are among leading causes of death by self-poisoning in particular in low and middle income countries as they are intrinsically toxic and deliberately spread in the environment, the production, distribution, and the use of pesticides require strict regulation and control. regular monitoring of residue in food and the environment is also required.
- The EPA PGP will cover pesticides authorized for use under RIERA, and generally includes pesticide applications covered by the 2006 rule.
- The EPA PGP will cover the following pesticide use patterns with discharges to waters of the u.s for the control of:
 - Mosquitoes and other flying insect pests
 - Weed and algae
 - Animal pests
 - Forest canopy pest
 - Direct impact on humans
 - Possible carcinogenicity, neurotoxicity, reurotoxic reproductive and metabolic toxicity of pesticide accumulating in the body
 - Impact on environment soil contamination Surface water contamination
 - Ground water contamination
 - Effect on soil fertility, air contamination
 - Effect and destruction of useful organisms
 - Non-target vegetation contamination
 - Increase the production of animal and plants biomass
 - Combal micro-organisms causing from produce and to delay
 - combal algae,bacteria,fungi and weeds
- There are numerous health hazards linked to the use of pesticides . pesticides can have grace effects on health of any body consuming fruits or vegetables highly contaminated with pesticides

- High levels of pesticides in food can lead to development of diseases such as cancer, kidney and lung ailments .
- childrens have developing organs, prone to catching infections and diseases .
- Any exposure to these high chemical residues can lead to childhood cancers, mental health problems such as autism and attention deficit hyperactivity disorder .
- pesticide may also be transported by rain and wind from the points of application to adjacent areas where their presence may be undesirable or harmful .
- Endocrine disruptor
- Thyroid disruption properties in birds, amphibians and fish
- carcinogen
- susceptibility in fungal infection
- Earthworms became infected with monoxystid gregarines
- Interact with vertebrate immune systems
- Animal infections, disease outbreaks and higher mortality
- Immunotoxicity primarily caused by the inhibition of serine hydrolases or esterases
- oxidative damage
- Impaired metabolic functions such as thermoregulation
- Respiratory, cardiovascular, neurological and immunological toxicity in rats and humans
- Effect various physiological and behavioural traits of beneficial orthoptods, particularly hymenopterans
- Reduced food availability and adverse secondary effects on soil invertebrates and butterflies
- Decreased species abundance and diversity in small mammals
- Pesticides have some direct harmful effect on plant including poor root hair development, shoot yellowing and reduced plant growth
- The use of pesticides is that their residues persist in water and other components of the environment as they are not easily degraded in the environment

- pesticides also have the potential to harm the nervous system ,the reproductive system,and the endocrine system
- pesticides can contaminate soil,water,turf,and other vegetation
- pesticides can reduce populations of helpful insects
- pesticides may lead to polination problems
- Adverse health effects on farmers
- crops may get contaminated with harmful substances
- pest resistance in the long run
- pesticide poisoning
- problematic for flora and fauna
- Ecological imbalance
- Rather unclear long-term effects

Problem Analysis

1



Analysis of the Problems:

The Method has been widely used for the Analysis GCMS Analyze, Triple Guadnepole. Traditional techniques are laborious, costly and consuming. They are several advanced techniques for the determination of pesticides. Pesticide toxicity and available extraction methods any chemical, biological substances or mixture of substances intended for preventing, destroying, attracting, repelling or controlling pest. Reduction of beneficial species resistance ground water contamination Drift of sprays and Vapours residues in food. Extraction is the Process by which toxicant is transformed from the treated bulky biological material into solvent clean up isolation of toxicant from interfering substance or solvent.


Three kinds of Spectrophotometric Methods in pesticide residue analysis. ultraviolet visible and in formal methods

- Various techniques of Chromatography
 1. Column chromatography gas chromatography liquid chromatography
 2. Affinity chromatography
 3. Ion exchange chromatography

DATA TABLES


1 . Experine in Vegetables Producing

Options	Experience	
A	5 Years	35
B	10 Years	52
C	Both A and B	37
D	15 Years	0




2 . Pestiside sparying Technique Used in field

opitions	techniques	
A	Spray against the world	49
B	Spray with the wind	30
C	Walk forward	21
D	Walk bckward	04




3.Number sign of passing with Pesticide

Options	Effects	
A	Burning gas	39
B	Cough and Head Ache	52
C	Skin inflowbin	32
D	Others	09




4 . Pesticides are poisonous

Opitions		
A	Agree	93
B	Natural	16
C	Disagree	21
D	Both A and B	00




5 . Do you think pesticides affect the Environment

Opitions		
A	Strongly Agree	60
B	Agree	55
C	Storngly Disagree	07
D	Disagree	03




1 . How many hours do you work in the field

Options		
A	3.4 Hours	23
B	5.7 Hours	62
C	➤ 7 Hours	49
D	Non of These	00




2 . How many pesticides do you apply per month

Opitions		
A	1	33
B	2	12
C	None of These	99
D	More Than 2	0




3 . Which type of field you have

Options		
A	Vegetables	77
B	Fruits	52
C	Both A and B	00
D	None of These	00




4 . Where do you store pesticides

options		
A	Store room	82
B	Garden,Open field	34
C	Cool storge, Refrigerator	13
D	outside	0




5. Positions of spray head

Options		
A	20cm above plants top	29
B	At plant tops	49
C	Targeted Insectse diseases	49
D	None of these	0



Analysis

- The main stages in analytical procedures for determination pesticides in samples of fruits & vegetables
- 2 ➤ The analysis of pesticides in biological samples continues to present challenges to analysts
- The number of problems crop up in the analysis of pesticides.
- The complexity and the diversity of matrices in biological



- The less additional methods are required to all analytes
- The more efficient and economical the analysis
- Less time, personal, materials.....

Some of the most Commonly used solvents for pesticide residue in fruits and vegetables and acetoethyl acetate, dichloromethane, methanol and toluence. In certain cases, the mixtures of solvents are used to improve the recovery of the methods.

RECOMMENDATIONS

RECOMMENDATIONS

- Collaborate with the Rotterdam convention to strengthen capacity building programs and the use of the knowledge base maintained by the Convention.
- Fulfill a role in supporting collaboration among developing countries to strengthen pesticide risk regulation.
- Explore the options to make regulatory risk data more transparent and publicly accessible. This could require long-term changes in the organization and funding of the underlying risk research and the structure of the intellectual property rights of risk data.
- Strengthen research and Extension in the fields of agroecology, organic farming and IPM, in particular supporting network initiatives on these themes and local universities and farmer associations in developing countries.
- Stop all exports of crop control products banned in the EU.
- Only allow the export of several restricted pesticides if these are regulated accordingly and used properly in the importing country.
- Support developing countries in developing an efficient process of re-evaluating pesticide registrations according to contemporary good regulatory practices in line with the FAO/WHO Code of Conduct.
- Buy only original products, with authorized sellers.
- Buy the pesticides according to your needs to reduce storage time and Surplus.
- Comply with all local legislations concerning storage.
- Pesticides should be stored Safely, locked, out of the reach of Children and unauthorized persons. They must also be out of the reach of animals.
- The Storage place should dry and have a good ventilation.
 - In case of spillage follow the instructions on the label for cleaning and proper disposal.
 - More dilute then on the label.

- At the lower rate than on the label
- Less frequently than on the label.
- For pests not on the label, as Long as the Site or cmp is listed and other restrictions are observed.
- It is illegal to apply pesticides in the following ways:
 - Using less diluent (water) than on the label, thus increasing the concentration.
 - At a higher rate per acre than on the label.
 - Shortening the specified interval between applications.

CONCLUSION

CONCLUSION

- Pesticides are an increasingly important part of
- Agricultural production technology but...
- Their inherent toxicity means they will continue to face stringent Regulation.
- Difference in access to pesticides, or in their cost will affect Production opportunities.
- Farmers appear to be successful in adapting their pest management Strategies to difference in prices in the cases where a number of Substitutes exist.
- Pesticides are a very diverse group of chemical hence it is Difficult to explore all the ramifications .
- Microbial and faunal population in time get to tolerant to Pesticides though there is considerable stress on the ecosystemic.
- Also major processes such as enzyme activity, respirations, Carbon and nitrogen mineralization are majorly affected.
- The widespread use of pesticides is ineffective and economically Harmful in the long run.
- Their detrimental effect on health and environment make Them an inadequate long term solution.
- In addition, most synthetic and natural pesticides are susceptible To ineffectiveness due to resistance build up in insects.
- Thus the only viable solutions for the future is IPM.
- The economic benefits and reduced social costs of these Systems present a logical answer to the pest control problem.
- These products can be dangerous if not used properly.
- Parasitoids will help in selecting which pesticides to use.
- The developed method allows qualitative and quantitative
- Analysis of 427 pesticides in fruits and vegetables samples in UHPLC- QTOF-MS

REFERENCES

REFERENCES

Some directions are used that pesticides users must obey are contained in documents that are only referred to on the product labelling.

These references to other documents is a new practice it is necessary because there is no longer room in the traditional pesticide label to explain the requirement of all laws and regulations that may apply to the user for example EPA has adopted or is considering new requirements

☞ ground water protection

☞ pesticides transportation, storage and disposal worker protection

☞ Endangered species protection

1.WWWEncyclopedia.com

2.WWWveckypad.com

3.Practical manual in chemistry for class×11

AUTHORS

1.Scientific committee on health and environment risks , toxicity and assessment of chemical mixture, European union scientific committee Brussels 2012, <https://ec.europa.eu>

2.F. Plaršić, temeljni pojmovi i tokovi psihologije i one koje rade otrovima koriste
d.o.o. Zagreb, 1999

3.S. Koo, R. Raroui, W. Bracke et al. current eu research activities on combined exposure to Multiple chemical, "environment international, vol." 120, pp. 544-562, 2018.

4.K.B. Nargaard and N. Cedergreen, "pesticides users can interact synergistically on aquatic" Environmental science and pollution research , vol. 17, pp. 957-967, 2010.

5.G.Glavan and J.Bozici"The synergy of cenobiotics in honey bee *Apis mellifera* mechanism and effects "Acta Biologica slovenica ,vol.56,no1pp,11-25,2013.

6.F.Sgolastra,p.medrzycki,L.Bortolotti "synergistic mortality between a neonicotinoid Insecticide and an ergosterol-biosynthesis-inhibiting fungicideinpp1236-1243.2017.

7.R.Raimets,Rkarise,mmandetal,"synergistic inter action between a variety of insecticides and an ergosterol biosynthesis inhibitors fungicide in dietary exposure of bumble bees(*Bombus terrestris*)"pest management science,vol74,no3,pp.541-546,2018.

8."Regulation(EC)no396/2005 of the European parliament and of the council of 23 February 2005 on maximum residue levels of pesticides in or on food and feed of plant and animal Origin and amending council directive 91/414/EEC , pp116,official journal of the Europe