

Document Management System

	Sultanate of Oman Ministry of Health The Royal Hospital Department of Child Health
Code: CH-MG-PRT-4-Vers.1.0	
Effective Date: 19/02/2024	
Target Review Date: 20/03/2027	
Title: Cerliponase Alfa Treatment	
Recombinant human tripeptidyl peptidase 1 (Cerliponase Alfa) is an enzyme-replacement therapy that has been developed to treat neuronal ceroid lipofuscinosis type 2 (CLN2) disease, a rare lysosomal disorder that causes progressive dementia in children. The Hamburg Scale (the Clinical Scoring System for Late Infantile Neuronal Ceroid Lipofuscinoses) is the most commonly used system for patient assessment and follow up. Indication to start the treatment: 1.CLN2 diagnosis by TPP1 genotype analysis and enzyme activity. 2.If the Hamburg motor scoring system prior to the initiation of t the treatment 2 and above the patient will be eligible for treatment. 3.Score between 3 and 6 out of 6 points on a motor-language scale • 2 and above on motor scale • ≥ 1 on language scale (Up to 15 years of age) 4.If there is family history to start the treatment in pre-symptomatic patients at age of 18 months and above unless there is new evidence to be started earlier. 5. Consent form for the treatment protocol including the termination police should be discussed with family and signed. 6. Therapy should be initiated with agreement of two consultants (metabolic and neurology) Surveillance: 1. Three monthly assessment using the Hamburg Scale (the Clinical Scoring System for Late Infantile Neuronal Ceroid Lipofuscinoses) by consultant 2. Ophthalmology assessment every 6 months 3. Swallowing assessment every 6 months 4. Brain MRI assessment and EEG	

Termination of Treatment:

1. The 2-point decline in the score on the motor and language domains of the CLN2 Clinical Rating Scale on the dose 24th of the treatment, the decision to stop the treatment should be discussed.
2. Generalized motor status epilepticus or severe infections within four weeks of first drug dose
3. If there is an evidence of the compliance of less than 85% based on the doses consumed and not institutional related

Hamburg Scoring System for Neuronal Ceroid Lipofuscinosis

Function	Ability	Score
Motor function	Walks normally	3
	Frequent falls, clumsiness	2
	Can't walk without assistance or crawling only	1
	Immobile, bedridden	0
Language	Normal	3
	Recognizable abnormal	2
	Hardly understandable	1
	Unintelligible or no language	0
Vision	Recognizes objects and grabs them	3
	Attempts to grab objects are uncoordinated	2
	Reacts to visual light	1
	No reaction to visual stimuli	0
Seizures (grand mal)	No seizures in the past 3 months	3
	1-2 seizures in the past 3 months	2

	>1 seizure per month	0

Reference:

1. Clinical Review Report: Cerliponase Alfa (Brineura): (Biomarin Pharmaceutical (Canada) Inc.): **Indication:** For the treatment of neuronal ceroid lipofuscinosis type 2 (CLN2) disease, also known as tripeptidyl peptidase 1 (TPP1) deficiency

2-Study of Intraventricular Cerliponase Alfa for CLN2 Disease

Angela Schulz, M.D., Temitayo Ajayi, M.D., Nicola Specchio, M.D., Ph.D., Emily de Los Reyes, M.D., Paul Gissen, M.B., Ch.B., Ph.D., Douglas Ballon, Ph.D., Jonathan P. Dyke, Ph.D., Heather Cahan, M.D., Peter Slasor, Sc.D., David Jacoby, M.D., Ph.D., and Alfried Kohlschütter, M.D. for the CLN2 Study Group

Written By: Dr.Aisha Al Balushi

Checked By: Dr.Nadia Al Hashmi

Authorized by: KHOULA JULENDA AL SAID

[Back](#)