



FALL PREVENTION POLICY –RH

- ▶ **DEFINITION** –A fall is defined as an event that results in a person coming to rest inadvertently on the ground or floor or other lower level ,with or without injury .



SCOPE OF FALL PREVENTION

- ▶ **All staff members are responsible for**
- ▶ **Implementing the intent and directives contained within this policy ,and for creating a safe environment of care .Any**
- ▶ **Staff member ,physician,or family member may request that be placed on**
- ▶ **Fall precautions**



GOALS

- ▶ **Decrease incidence of falls**
- ▶ **Decrease severity of falls**
- ▶ **Increase mobility and function**
- ▶ **Improve environmental safety**
- ▶ **Provide comprehensive assessment**
- ▶ **Enhance staff knowledge on falls reduction prevention strategy**
- ▶ **Improve the patients confidence**

PURPOSE

The fall prevention policy is intended to objectively and systematically identify patients at risk for fall

To provide a standardized approach to prevent client falls and injury from falls through the use of a standardized fall prevention strategy



Cont—PURPOSE

To protect patients and promote patient safety

To effectively identify and intervene with patients who are at for falling

To educate patients ,families,and staff members on measures to prevent falls and promot identified as a higher level of risk. All patients will be assessed for fall during the in -patient and ambulatory service



❖ **FALL RISK WILL BE ASSESSED DURING ANY TYPE OF VISIT AND WILL BE REASSESSED**

Within 24 hours of admission to acute or long-term care

At change in patients status

Following a fall

Once a week



FALL RISK ASSESSMENT TEAM

APPROACH

Identify risk of fall

John Hopkins fall Scale Communicate

Risk /Humpty Dumpty

Bedside signage

Fall Risk ID band

Nursing Documentation .

Review john Hopkins Fall Scale Risk

Factors

Implement Standard interventions

Conduct HighRisk Assessment

Implement High Risk nursing

interventions

provide patient /family education



Patient Name:

Date:

Fall Risks *(Check all that apply)*



History of Falls

☐


Medication
Side Effects

☐


Walking Aid

☐


IV Pole or Equipment

☐


Unsteady Walk

☐


May Forget or
Choose Not to Call

☐

You are at high risk for injury if you fall:

- ☐ Age: 85 years old or older, frailty
- ☐ Bones: osteoporosis, risk or history of fracture, etc
- ☐ Coagulation: risk for bleeding, low platelet counts or taking anticoagulation
- ☐ Surgery (recent): lower limb amputation, major abdominal, or thoracic surgery

☐

Fall Interventions *(Circle selection based on color)*

Communicate
Recent Falls and/or
Risk of Harm



Walking Aids



Crutches



Cane



Walker

IV Assistance
When Walking



Toileting Schedule: Every _____ hours



Bed Pan



Assist to
Commode



Assist to
Bathroom

Bed Alarm On



Assistance Out of Bed



1 person



2 people

RESPONSIBILITIES OF STAFF

**RESPONSIBILITIES OF THE FOLLOWING STAFF
ARE DELINEATED**

A. Director of Nursing

B. Unit Nurses and Ward Nurses

C. Admission Nurses

F. Pharmacists

G. Physical and Occupational

Therapists

H. Biomedical

I. Nursing Quality

J. General Service Management Staff

K. Education Service

L. Fall Team

How Can We Improve?



IMPLEMENTATION OF INTERVENTION

Apply a yellow armband Implement appropriate standard safety measures

Escort or offer wheelchair assistance to a patient who is at risk for falls

provide adequate lighting

Bed/Stretcher should be on the lowest position with bed lock

While on stretcher all rails should be raised for patient safety

If high risk ,do not place patient on examination table without attendants

Toilet patient regularly ,especially prior to procedure .

Instruct patient to use emergency call bell in restroom and keep path to bathroom clear

While in recovery area –patient to be visible and not left alone

All moderate sedation patients must be monitored according to hospital policy

Educate patient if on medications that increased risk for falls.provide pamphlet on fall safety

 Attachment: 1

Fall Assessment Tool Humpty Dumpty Scale
(Paediatric)

Parameter	Criteria	Score
Age		
Less than 3 years old		4
3 to less than 7 years old		3
7 to less than 13 years old		2
13 years old and above		1
Gender		
Male		2
Female		1

Neurological Diagnosis	4
Alterations in Oxygenation (Respiratory Diagnosis, Dehydration, Anemia, Anorexia, Syncope/Dizziness, etc.)	3
Psych/Behavioral Disorders	2
Other Diagnosis	1
Cognitive Impairments	
Not Aware of Limitations	3
Forget Limitations	2
Oriented to own Ability	1
Environmental Factors	
History of Falls or Infant-Toddler Placed in Bed	4
Patient uses assistive devices or Infant Toddler in Crib or Furniture/Lighting (Tripled Room)	3
Patient Placed in Bed	2
Outpatient Area	1
Response to Surgery/Sedation/ Anesthesia	
Within 24 hours	3
Within 48 hours	2



Thank you