

ROP- Medication use

Ensure the safe use of high-risk medications.

ACI medication management ROPs

- Antimicrobial stewardship.
- Concentrated electrolytes.
- Heparin safety.
- High alert medications.
- Infusion pump safety.[infusion pump training]
- Narcotic safety.

Antimicrobial stewardship program {ASP}

Consequences of unintended use of antimicrobial agents include.

1. Toxicity
2. Selection of pathogenic organism
3. Development of organisms resistant to antimicrobial agents.

ASP is an activity that includes.

- Appropriate selection
- Appropriate dosing.
- Appropriate route
- Appropriate duration of antimicrobial therapy.

A successful ASP requires an interdisciplinary approach.

- Pharmacy { clinical pharmacist, hospital pharmacist }
- Infection control team
- Support from hospital administration
- Medical staff leadership { ID team }
- Other health care providers { nurses, medical team, lab personnel. }

Concentrated electrolytes

- Calcium : concentration $>0.9\%$
 - Calcium chloride 10%
 - Calcium gluconate 10%
- . Magnesium : concentration $>20\%$.
 - Magnesium sulfate 50%.
- . Potassium : concentration $>2\text{mmol}$
 - Potassium chloride 15%.
 - Potassium phosphate {3mmol +4.4mEqK/ml}
- . Sodium chloride- concentration $>0.9\%$
 - sodium chloride 3% { 500ml}

Heparin safety

- Heparin is HAM, limiting its availability and ensuring that high dose formats are not stocked in a client service areas are effective strategies to minimize the risk of death or disability injury associated with these agents.
- For flushing intravenous line, organizations are encouraged to consult best practice guidelines to explore options other than heparin.
- LMWH [clexane]
- Heparin 1000 units
- Heparin 5000 units

HIGH ALERT MEDICATIONS

- High alert medications are drugs that bear a heightened risk of causing significant patient harm when they are used in error.

Why are HAMS a safety concern ?

. When used improperly, will produce more serious harm to patients than other medications.

Infusion Pump Safety

A documented and co ordinated approach for infusion pump safety that includes training , evaluation of competence and a process to report problem with infusion pump use is implemented.

Narcotic safety

The organization evaluates and limits the availability of narcotic products and removes high- dose, high- potency formats from patient care areas.

- . Morphine ampoules with concentration >10mg/ml and >2mg/ml.
- . High potency Fentanyl ampoules ; >500mcg and >100mcg.

Summary

- Medication use process is complex.
- Medication errors are preventable.
- Employ safety measures.

THANK YOU

Medication group

