

Calculus on Manifolds

Nutan Nepal

January 31, 2019

1 1-30

1. Let $f : [a, b] \rightarrow \mathbf{R}$ be an increasing function. If $x_1, \dots, x_n \in [a, b]$ are distinct, show that

$$\sum_{i=1}^n o(f, x_i) \leq f(b) - f(a).$$

Solution: Let order be defined in $\{x_1, \dots, x_n\}$ such that $x_1 < \dots < x_n$, then since f is an increasing function, we get $f(x_1) \leq \dots \leq f(x_n)$. We have

$$o(f, x_i) = \lim_{\delta \rightarrow 0} (M(x_i, f, \delta) - m(x_i, f, \delta))$$

where,

$$\begin{aligned} M(x_i, f, \delta) &= \sup \{f(x) : x \in [a, b] \text{ and } \|x - x_i\| < \delta\}, \\ m(x_i, f, \delta) &= \inf \{f(x) : x \in [a, b] \text{ and } \|x - x_i\| < \delta\}. \end{aligned}$$

If we denote a δ -neighborhood of some $x_i \in [a, b]$ by $N_\delta(x_i)$, then since $\delta \rightarrow 0$, we can choose a sufficiently small $\delta > 0$ such that

$$\bigcap_{i=1}^n N_\delta(x_i) = \emptyset.$$

Then for all such δ we have,

$$\begin{aligned} f(x_{i+1}) &\geq M(x_i, f, \delta) \geq f(x_{i-1}), \text{ and} \\ f(x_{i+1}) &\geq m(x_i, f, \delta) \geq f(x_{i-1}). \end{aligned}$$

We simplify the given summation as

$$\begin{aligned} \sum_{i=1}^n o(f, x_i) &= \lim_{\delta \rightarrow 0} \sum_{i=1}^n (M(x_i, f, \delta) - m(x_i, f, \delta)) \\ &\leq \lim_{\delta \rightarrow 0} \sum_{i=1}^n (f(x_{i+1}) - f(x_{i-1})) \\ &= f(x_{n+1}) - f(x_0), \\ &\leq f(b) - f(a). \end{aligned}$$

where the last statement follows from the fact that the max and min $f(x)$ can get is $f(b)$ and $f(a)$. ■

2 1-25

1. Prove that a linear transformation $T : \mathbf{R}^n \rightarrow \mathbf{R}^m$ is continuous.

Solution:

Proof. We need to show that $T : \mathbf{R}^n \rightarrow \mathbf{R}^m$ is continuous at all $a \in \mathbf{R}^n$. That is, for every $\varepsilon > 0$ we can find a $\delta > 0$ such that $0 < \|x - a\| < \delta \implies \|T(x) - T(a)\| < \varepsilon$, where $x \in \mathbf{R}^n$. But we have,

$$\|T(x) - T(a)\| = \|T(x - a)\| \leq M\|x - a\|$$

for some $M \in \mathbf{R}$. So for any given $\varepsilon > 0$, we can choose $\delta = \varepsilon/M$. Then certainly, if $0 < \|x - a\| < \delta$, then

$$\|T(x) - T(a)\| \leq M\|x - a\| < M\delta = \varepsilon.$$

So it follows that the linear transformation is continuous. ■

3 1-29

1. If A is compact, prove that every continuous function $f : A \rightarrow \mathbf{R}$ takes on a maximum and a minimum value.

Solution:

Proof. Since A is compact and f is continuous, we know that the image of A under f is compact in \mathbf{R} . Hence, it follows from 3.1, that f takes on a maximum and a minimum value. ■

Lemma 3.1. *A compact set in \mathbf{R} has a maximum and a minimum value.*

Proof. We know that a compact set is closed and bounded and in \mathbf{R} , a compact set is in the form $[a, b]$. And since $a, b \in [a, b]$, all we need to show is that a and b are infimum and supremum, respectively, of the given interval. ■

4 2-4

Let g be a continuous real-valued function on the unit circle $x \in \mathbf{R}^2 : \|x\| = 1$ such that $g(0, 1) = g(1, 0) = 0$ and $g(-x) = -g(x)$. Define $f : \mathbf{R}^2 \rightarrow \mathbf{R}$ by

$$f(x) = \begin{cases} \|x\| \cdot g\left(\frac{x}{\|x\|}\right) & x \neq 0, \\ 0 & x = 0. \end{cases}$$

1. If $x \in \mathbf{R}^2$ and $h : \mathbf{R} \rightarrow \mathbf{R}$ is defined by $h(t) = f(tx)$, show that h is differentiable.

Solution: We need to show that for every $a \in \mathbf{R}$, there exists a $\lambda : \mathbf{R} \rightarrow \mathbf{R}$ such that

$$\lim_{t \rightarrow 0} \frac{h(a+t) - h(a) - \lambda(t)}{t} = 0. \tag{1}$$

We see that

$$h(a+t) = f(ax+tx) = \begin{cases} \|(a+t)x\| \cdot g\left(\frac{x}{\|x\|}\right) & (a+t)x \neq 0, \\ 0 & (a+t)x = 0. \end{cases}$$

This follows from the fact that $g\left(\frac{(a+t)x}{\|(a+t)x\|}\right) = g\left(\frac{x}{\|x\|}\right) = g\left(\frac{x}{\|x\|}\right)$. We have,

$$\frac{h(a+t) - h(a) - \lambda(t)}{t} = \frac{\|(a+t)x\| \cdot g(\hat{x}) - \|a\|g(\hat{x}) - \lambda(t)}{t}$$

$$=$$

5

Theorem 5.1 (Heine-Borel Theorem). *The closed interval $[a, b]$ is compact.*

Proof. If \mathcal{O} is an open cover of $[a, b]$, let

$$A = \{x : a \leq x \leq b \text{ and } [a, x] \text{ is covered by some finite number of open sets in } \mathcal{O}\}$$

■

6 2-1

1. Prove that if $f : \mathbf{R}^n \rightarrow \mathbf{R}^m$ is differentiable at $a \in \mathbf{R}^n$, then it is continuous at a .

Solution: Hi.