Notes: Chow Theory for Matroid Deletion

Nutan Nepal

October 15, 2025

1 Algebraic Preliminaries

1.1 The Module of Flats and Perverse Subgroups

Let $R := \mathbb{Z}[x, x^{-1}]$ be the ring of Laurent polynomials with integer coefficients. For any integer n, we denote by $\mathrm{Pal}(n)$ the additive group of polynomials $f(x) \in R$ satisfying $f(x) = x^n f(x^{-1})$. While $\mathrm{Pal}(0)$ is a subring of R, we note that $\mathrm{Pal}(n)$ for $n \neq 0$ is not, as $\mathrm{Pal}(n) \cdot \mathrm{Pal}(m) \subseteq \mathrm{Pal}(n+m)$.

Let M be a matroid with lattice of flats $\mathcal{L}(\mathrm{M})$.

Definition 1. *The module $\mathcal{H}(\mathrm{M})$ is the free R-module with a standard basis $\{\mathrm{F}\}_{F \in \mathcal{L}(\mathrm{M})}$. An element $\alpha \in \mathcal{H}(\mathrm{M})$ is a formal sum*

$$\alpha = \sum_{F \in \mathcal{L}(\mathrm{M})} \alpha_F \cdot \mathrm{F}, \quad \text{where } \alpha_F \in R.$$

We also define the R-linear maps $S_F : \mathcal{H}(\mathrm{M}) \to R$ for any flat $F \in \mathcal{L}(\mathrm{M})$ by

$$S_F(\alpha) := \sum_{G \geq F} x^{\mathrm{rk}(F) - \mathrm{rk}(G)} \alpha_G.$$

Following Braden-Vysogorets, there is an important abelian subgroup of $\mathcal{H}(\mathrm{M})$.

Definition 2. *The subgroup of **perverse elements** $\mathcal{H}_p(\mathrm{M}) \subset \mathcal{H}(\mathrm{M})$ consists of all α such that for every flat $F \in \mathcal{L}(\mathrm{M})$:*

1. *The coefficient α_F is a polynomial, i.e., $\alpha_F \in \mathbb{Z}[x]$.*
2. *The Laurent polynomial $S_F(\alpha)$ is in $\mathrm{Pal}(0)$.*

Definition 3. *We relax the first condition to define a larger subgroup $\mathcal{H}'_p(\mathrm{M}) \subseteq \mathcal{H}(\mathrm{M})$ consisting of all elements α that satisfy $S_F(\alpha) \in \mathrm{Pal}(0)$ for all $F \in \mathcal{L}(\mathrm{M})$.*

1.2 The Chow and Kazhdan-Lusztig Bases

The module $\mathcal{H}(\mathrm{M})$ has natural bases constructed from the Kazhdan-Lusztig and Chow polynomials.

Definition 4. *The **Kazhdan-Lusztig basis** $\{\zeta^F\}$ is defined by*

$$\zeta^F = \sum_{G \leq F} \zeta^F_G \cdot \mathrm{G} := \sum_{G \leq F} x^{\mathrm{rk}(F) - \mathrm{rk}(G)} \mathrm{P}_{\mathrm{M}^F_G}(x^{-2}) \cdot \mathrm{G},$$

where $\mathrm{P_M}$ *is the Kazhdan-Lusztig polynomial of* M.

Remark. Braden-Vysogorets prove that the elements $\{\zeta^F\}$ form a $\mathbb{Z}$-basis for $\mathcal{H}_p(\mathrm{M})$. In particular, if $\alpha = \sum_F \alpha_F \cdot \mathrm{F} \in \mathcal{H}_p(\mathrm{M})$, then

$$\alpha = \sum_F \alpha_F(0) \cdot \zeta^F.$$

Definition 5. *The* ***Chow basis*** $\{\mathrm{c}^F\}$ *is defined by*

$$\mathrm{c}^F = \sum_{G \leq F} \mathrm{c}^F_G \cdot \mathrm{G} := \sum_{G \leq F} x^{\mathrm{rk}(F)-\mathrm{rk}(G)} \underline{\mathrm{H}}_{\mathrm{M}^F_G}(x^{-2}) \cdot \mathrm{G},$$

where $\underline{\mathrm{H}}_{\mathrm{M}}$ *is the Chow polynomial of* M*, which is palindromic of degree* $\mathrm{rk}(\mathrm{M}) - 1$.

Remark. The coefficients of the Chow basis have a special structure:

$$\mathrm{c}^F_F = 1 \in \mathrm{Pal}(0), \quad \text{and} \quad \mathrm{c}^F_G \in \mathrm{Pal}(2) \text{ for all } G < F.$$

Definition 6. *We define the module* $\mathcal{S}(\mathrm{M})$ *to be the free* $\mathrm{Pal}(0)$*-module with basis* $\{\mathrm{c}^F\}$.

Definition 7. *For an element* e *in the ground set of* M*, the* ***deletion operator*** *is an R-module homomorphism* $\Delta : \mathcal{H}(\mathrm{M}) \to \mathcal{H}(\mathrm{M} \setminus e)$ *defined by*

$$\Delta(\mathrm{F}) = x^{-\delta(F)} \cdot \mathrm{F} \setminus \mathrm{e}, \quad \textit{where } \delta(F) := \mathrm{rk}_{\mathrm{M}}(F) - \mathrm{rk}_{\mathrm{M}\setminus e}(F \setminus e).$$

Definition 8. *For any element* e *in the ground set of* M*, we define the set* $S_e(\mathrm{M}) := \{F \in \mathcal{L}(\mathrm{M}) \mid e \notin F \text{ and } F \cup e \in \mathcal{L}(\mathrm{M})\}$. *As demonstrated in Braden-Vysogorets,* $S_e(\mathrm{M})$ *plays a crucial role in the deletion formula.*

2 Main Propositions

Proposition 9. *For all* $F \in \mathcal{L}(\mathrm{M})$, $\mathrm{c}^F \in \mathcal{H}'_p(\mathrm{M})$*, i.e., it satisfies* $S_G(\mathrm{c}^F) \in \mathrm{Pal}(0)$ *for all* $G \in \mathcal{L}(\mathrm{M})$.

Proof. We see that, for any $G \in \mathcal{L}(\mathrm{M})$,

$$\begin{aligned} S_G(\mathrm{c}^F) = \sum_{H:G\leq H\leq F} x^{\mathrm{rk}(G)-\mathrm{rk}(H)} \cdot \mathrm{c}^F_H &= \sum_{H:G\leq H\leq F} x^{\mathrm{rk}(G)-\mathrm{rk}(H)} \left(x^{\mathrm{rk}(F)-\mathrm{rk}(H)} \underline{\mathrm{H}}_{\mathrm{M}^F_H}(x^{-2}) \right) \\ &= x^{\mathrm{rk}(F)-\mathrm{rk}(G)} \sum_{H:G\leq H\leq F} x^{-2(\mathrm{rk}(H)-\mathrm{rk}(G))} \underline{\mathrm{H}}_{\mathrm{M}^F_H}(x^{-2}). \end{aligned}$$

This sum is precisely the definition of the *augmented* Chow polynomial, $\mathrm{H}_{\mathrm{M}^F_G}(x^{-2})$. So, $S_G(c^F) = x^{\mathrm{rk}(F)-\mathrm{rk}(G)} \mathrm{H}_{\mathrm{M}^F_G}(x^{-2})$. Let $d = \mathrm{rk}(F) - \mathrm{rk}(G)$. The term x^d is in $\mathrm{Pal}(2d)$. The term $\mathrm{H}_{\mathrm{M}^F_G}(x^{-2})$ is in $\mathrm{Pal}(-2d)$. The product is in $\mathrm{Pal}(2d - 2d) = \mathrm{Pal}(0)$ as required. □

Proposition 10. *The subgroup* $\mathcal{H}'_p(\mathrm{M})$ *is a* $\mathrm{Pal}(0)$*-module.*

Proof. Let $\alpha = \sum_F p_F \cdot \mathrm{F} \in \mathcal{H}'_p(\mathrm{M})$ and $q \in \mathrm{Pal}(0)$.

We see that

$$S_G(q \cdot \alpha) = \sum_{H \geq G} x^{\mathrm{rk}(G)-\mathrm{rk}(H)} (q \cdot p_H) = q \cdot S_G(\alpha) \in \mathrm{Pal}(0).$$

Therefore, $q \cdot \alpha \in \mathcal{H}'_p(\mathrm{M})$. □

Proposition 11. *Any Laurent polynomial f can be uniquely written as a sum $f = f_0 + f_2$ where $f_0 \in \mathrm{Pal}(0)$ and $f_2 \in \mathrm{Pal}(2)$. The projections are given by*

$$f_0(x) = \frac{x^2 f(x^{-1}) - f(x)}{x^2 - 1} \quad \textit{and} \quad f_2(x) = \frac{f(x) - f(x^{-1})}{1 - x^{-2}}.$$

Proposition 12. $\mathcal{S}(\mathrm{M}) = \mathcal{H}'_p(\mathrm{M})$ *as* $\mathrm{Pal}(0)$*-modules.*

Proof. It is enough to show that $\mathcal{H}'_p(\mathrm{M}) \subseteq \mathcal{S}(\mathrm{M})$.

Let $\alpha = \sum_F \alpha_F \cdot \mathrm{F}$ be an arbitrary element of $\mathcal{H}'_p(\mathrm{M})$. Since $\{\mathrm{c}^F\}$ is a basis for the entire module $\mathcal{H}(\mathrm{M})$, α has a unique expansion in this basis:

$$\alpha = \sum_{F \in \mathcal{L}(\mathrm{M})} q_F \cdot \mathrm{c}^F \quad \text{for } q_F \in \mathbb{Z}[x, x^{-1}].$$

To show that $\alpha \in \mathcal{S}(\mathrm{M})$, we must prove that each of these coefficients q_F is actually in $\mathrm{Pal}(0)$. We prove this by induction on the corank of F.

Base Case: The relationship between the coefficients is given by the change-of-basis formula:

$$\alpha_F = \sum_{G \geq F} q_G \cdot \mathrm{c}^G_F = q_F + \sum_{G > F} q_G \cdot \mathrm{c}^G_F.$$

For $F = E$, this simplifies to $q_E = \alpha_E$. Since $\alpha \in \mathcal{H}'_p(\mathrm{M})$, we have $S_E(\alpha) = \alpha_E \in \mathrm{Pal}(0)$. This implies that $q_E \in \mathrm{Pal}(0)$, establishing the base case.

Inductive Step: Assume that for all flats G with $\mathrm{rk}(G) > \mathrm{rk}(F)$, the coefficient q_G is in $\mathrm{Pal}(0)$.

Since $\alpha \in \mathcal{H}'_p(\mathrm{M})$, it satisfies:

$$S_F(\alpha) = \sum_{H \geq F} x^{\mathrm{rk}(F) - \mathrm{rk}(H)} \alpha_H \in \mathrm{Pal}(0).$$

We substitute the change-of-basis formula $\alpha_H = \sum_{L \geq H} q_L \cdot \mathrm{c}^L_H$ into this expression:

$$S_F(\alpha) = \sum_{H \geq F} x^{\mathrm{rk}(F) - \mathrm{rk}(H)} \left(\sum_{L \geq H} q_L \cdot \mathrm{c}^L_H \right) = \sum_{L \geq F} q_L \left(\sum_{H : F \leq H \leq L} x^{\mathrm{rk}(F) - \mathrm{rk}(H)} \mathrm{c}^L_H \right) = \sum_{L \geq F} q_L \cdot S_F(\mathrm{c}^L).$$

We isolate the $L = F$ term and rearrange the sum:

$$q_F = S_F(\alpha) - \sum_{L > F} q_L \cdot S_F(\mathrm{c}^L).$$

By the inductive hypothesis, each q_L in the sum is in $\mathrm{Pal}(0)$. Since $\mathrm{Pal}(0)$ is a ring, the difference on the right is in $\mathrm{Pal}(0)$. Thus, q_F must also be in $\mathrm{Pal}(0)$. □

Proposition 13 (Change of Basis Formula)**.** *If $\alpha = \sum_F \alpha_F \cdot \mathrm{F} \in \mathcal{S}(\mathrm{M})$, then its expansion in the c^F basis is $\alpha = \sum_F \overline{\alpha_F} \cdot \mathrm{c}^F$, where $\overline{\alpha_F}$ is the unique $\mathrm{Pal}(0)$ component of α_F from the decomposition in Proposition 11.*

Proof. Let $\alpha \in \mathcal{S}(\mathrm{M})$. By the definition of $\mathcal{S}(\mathrm{M})$, α can be uniquely expressed as a $\mathrm{Pal}(0)$-linear combination of the c^F basis vectors:

$$\alpha = \sum_{F \in \mathcal{L}(\mathrm{M})} q_F \cdot \mathrm{c}^F, \quad \text{where each } q_F \in \mathrm{Pal}(0).$$

We can also express α in the standard basis as $\alpha = \sum_F \alpha_F \cdot \mathrm{F}$. As in the proof of proposition 12, the change-of-basis formula relating the two sets of coefficients is:

$$q_F = \alpha_F - \sum_{G > F} q_G \cdot \mathrm{c}_F^G. \tag{1}$$

The goal is to show that $q_F = \overline{\alpha_F}$. We prove this by induction on the corank of F as in the previous proposition.

Base Case: For the base case, we see that $q_E = \alpha_E = S_E(\alpha) \in \mathrm{Pal}(0)$. This implies that $q_E = \overline{\alpha_E}$.

Inductive Step: Assume that for all flats G with $\mathrm{rk}(G) > \mathrm{rk}(F)$, the coefficient $q_G = \overline{\alpha_G}$.

Then the entire sum $\sum_{G>F} q_G \cdot \mathrm{c}_F^G$ is in $\mathrm{Pal}(2)$, since for any $G > F$, the coefficient q_G is in $\mathrm{Pal}(0)$ and the basis coefficient c_F^G is in $\mathrm{Pal}(2)$, as noted in a previous remark. This shows that α_F is the sum of q_F (which, by definition, is in $\mathrm{Pal}(0)$) and another term that is in $\mathrm{Pal}(2)$.

By Proposition 11, this decomposition is unique. Therefore, the coefficient q_F must be the unique $\mathrm{Pal}(0)$ component of the α_F. □

Proposition 14. *For any flat $F \in \mathcal{L}(\mathrm{M})$, the element $\Delta(\mathrm{c}^F)$ is in $\mathcal{S}(\mathrm{M} \setminus e)$.*

Proof. Let $F \in \mathcal{L}(\mathrm{M})$ be an arbitrary flat. We need to show that $S_H^{\mathrm{M}\setminus e}(\Delta(\mathrm{c}^F))$ is in $\mathrm{Pal}(0)$ for all $H \in \mathcal{L}(\mathrm{M} \setminus e)$.

Let $\beta = \Delta(\mathrm{c}^F) = \sum_{G \in \mathcal{L}(\mathrm{M}\setminus e)} \beta_G \cdot \mathrm{G}$.

By definition, $\mathrm{c}^F = \sum_{L \le F} \mathrm{c}_L^F \cdot \mathrm{L}$. Applying the Δ operator gives:

$$\beta = \Delta(\mathrm{c}^F) = \sum_{L \le F} \mathrm{c}_L^F x^{-\delta(L)} \cdot \mathrm{L} \setminus \mathrm{e}.$$

The coefficient β_G is the sum over all flats $L \le F$ such that $L \setminus e = G$. Thus, we have:

$$\begin{aligned} S_H^{\mathrm{M}\setminus e}(\Delta(\mathrm{c}^F)) = \sum_{G \ge H} x^{\mathrm{rk}(H)-\mathrm{rk}(G)} \beta_G &= \sum_{G \ge H} x^{\mathrm{rk}(H)-\mathrm{rk}(G)} \left(\sum_{L : L \le F, L\setminus e = G} \mathrm{c}_L^F x^{-\delta(L)} \right) \\ &= \sum_{L : L \le F, L \setminus e \ge H} x^{\mathrm{rk}(H)-\mathrm{rk}(L\setminus e)} \mathrm{c}_L^F x^{-\delta(L)} \end{aligned}$$

The exponent on x simplifies to $\mathrm{rk}(H) - \mathrm{rk}(L)$ using the definition of discrepancy.

$$S_H^{\mathrm{M}\setminus e}(\Delta(\mathrm{c}^F)) = \sum_{L : L \le F, L \setminus e \ge H} x^{\mathrm{rk}(H)-\mathrm{rk}(L)} \mathrm{c}_L^F.$$

Now we apply Lemma 2.15 from Braden-Vysogorets. The condition $L \setminus e \ge H$ is equivalent to $L \ge \overline{H}$, where $\overline{H}$ is the closure of H in M. Furthermore, $\mathrm{rk}_{\mathrm{M}\setminus e}(H) = \mathrm{rk}_{\mathrm{M}}(\overline{H})$.

Substituting these two facts, the summation range becomes $L \in [\overline{H}, F]$ and the expression is:

$$S_H^{\mathrm{M}\setminus e}(\Delta(\mathrm{c}^F)) = \sum_{L:\overline{H}\le L\le F} x^{\mathrm{rk}(\overline{H})-\mathrm{rk}(L)}\mathrm{c}_L^F = S_{\overline{H}}(\mathrm{c}^F) \in \mathrm{Pal}(0).$$

Since the palindromic condition holds for any flat H, we have proven that $\Delta(\mathrm{c}^F) \in \mathcal{S}(\mathrm{M}\setminus e)$. □

Remark. The map $\Delta : \mathcal{S}(\mathrm{M}) \to \mathcal{S}(\mathrm{M}\setminus e)$ is a $\mathrm{Pal}(0)$-module homomorphism that restricts from the $\mathbb{Z}[x, x^{-1}]$-module homomorphism $\Delta : \mathcal{H}(\mathrm{M}) \to \mathcal{H}(\mathrm{M}\setminus e)$.

Proposition 15. *For any flat $F \in \mathcal{L}(\mathrm{M})$, the following identities hold in $\mathcal{S}(\mathrm{M}\setminus e)$:*

1. *If e is not a coloop in M^F, then*

$$\Delta(\mathrm{c}^F) = \mathrm{c}^{F\setminus e} + x^{-1}\sum_{G\in S_e(\mathrm{M}^F)} \mathrm{c}^F_{G\cup e}\cdot \mathrm{c}^G \tag{2}$$

2. *If e is a coloop in M^F, then*

$$\Delta(\mathrm{c}^F) = (x+x^{-1})\mathrm{c}^{F\setminus e} + x^{-1}\sum_{G\in S_e(\mathrm{M}^F), G\neq F\setminus e} \mathrm{c}^F_{G\cup e}\cdot \mathrm{c}^G \tag{3}$$

Proof. By Proposition 13, the coefficients q_G in the expansion $\Delta(\mathrm{c}^F) = \sum q_G \cdot \mathrm{c}^G$ are the unique $\mathrm{Pal}(0)$ components of the standard-basis coefficients,

$$\beta_G = \sum_{L:L\le F, L\setminus e=G} \mathrm{c}^F_L x^{-\delta(L)}.$$

Thus we analyze q_G by considering different cases for the flat G.

Coefficients for $G \neq F\setminus e$: For any flat G not equal to $F\setminus e$, its preimages $L \le F$ are a subset of $\{\overline{G}, G\cup e\}$ where $\overline{G}$ is the closure of G in M.

- If $G \in S_e(\mathrm{M}^F)$, $\overline{G} = G$ and the standard-basis coefficient is $\beta_G = \mathrm{c}^F_G + x^{-1}\mathrm{c}^F_{G\cup e}$. This is a sum of a $\mathrm{Pal}(2)$ term and a $\mathrm{Pal}(0)$ term. Its unique $\mathrm{Pal}(0)$ part is $q_G = x^{-1}\mathrm{c}^F_{G\cup e}$.
- If $G \notin S_e(\mathrm{M}^F)$, the only possible preimage is $L = \overline{G}$. The standard-basis coefficient is $\beta_G = \mathrm{c}^F_{\overline{G}}$, which is in $\mathrm{Pal}(2)$. Its $\mathrm{Pal}(0)$ part is $q_G = 0$.

These coefficients account for the summation in both formulas of the proposition.

Coefficient for $G = F\setminus e$: Here we analyze $\beta_{F\setminus e}$ by considering the the following two cases:

- If e is **not a coloop** in M^F, the only flat $L \le F$ that deletes to $F\setminus e$ is $L = F$. The standard-basis coefficient is $\beta_{F\setminus e} = \mathrm{c}^F_F x^{-0} = 1 \in \mathrm{Pal}(0)$. So $q_{F\setminus e} = 1$, proving the first part of the proposition.
- If e is **a coloop** in M^F, the preimages are $L = F\setminus e$ and $L = F$. The standard-basis coefficient is $\beta_{F\setminus e} = \mathrm{c}^F_{F\setminus e} + x^{-1}\mathrm{c}^F_F = x + x^{-1} \in \mathrm{Pal}(0)$. So $q_{F\setminus e} = x + x^{-1}$. This proves the second part of the proposition.

□

3 Recovery of the Semi-Small Decomposition Formulas

We now show how the formulas in Proposition 15 recover the known semi-small decomposition formulas for the Chow polynomial $\underline{\mathrm{H}}$ from [FMSV24]. This is achieved by equating the coefficients of the empty flat $\emptyset$ on both sides of the identities.

Let $f = \mathrm{rk}(F)$. The coefficient of $\emptyset$ in $\Delta(\mathrm{c}^F)$ is:

$$\beta_\emptyset = \mathrm{c}^F_\emptyset x^{-\delta(\emptyset)} + \mathrm{c}^F_{\{e\}} x^{-\delta(\{e\})} = x^f \underline{\mathrm{H}}_{\mathrm{M}^F}(x^{-2}) + x^{f-2}\underline{\mathrm{H}}_{\mathrm{M}^F_e}(x^{-2})$$

Case 1: e is not a coloop in M^F. We take the coefficient of $\emptyset$ in the formula (2).

$$x^f \underline{\mathrm{H}}_{\mathrm{M}^F}(x^{-2}) + x^{f-2}\underline{\mathrm{H}}_{\mathrm{M}^F_e}(x^{-2}) = x^f \underline{\mathrm{H}}_{\mathrm{M}^F\setminus e}(x^{-2}) + x^{-1} \sum_{G \in S_e(\mathrm{M}^F)} x^{f-g-1} \underline{\mathrm{H}}_{\mathrm{M}^F_{G\cup e}}(x^{-2}) \cdot x^g \underline{\mathrm{H}}_{\mathrm{M}^G}(x^{-2}).$$

Dividing by x^f and setting $y = x^{-2}$, we get:

$$\underline{\mathrm{H}}_{\mathrm{M}^F}(y) + y\underline{\mathrm{H}}_{\mathrm{M}^F_e}(y) = \underline{\mathrm{H}}_{\mathrm{M}^F\setminus e}(y) + y \sum_{G \in S_e(\mathrm{M}^F)} \underline{\mathrm{H}}_{\mathrm{M}^F_{G\cup e}}(y) \cdot \underline{\mathrm{H}}_{\mathrm{M}^G}(y)$$

The sum on the right includes the term for $G = \emptyset$, which is $\underline{\mathrm{H}}_{\mathrm{M}^F_e}(y) \cdot \underline{\mathrm{H}}_{U_{0,0}}(y) = \underline{\mathrm{H}}_{\mathrm{M}^F_e}(y)$. Separating this term and canceling it with the $y\underline{\mathrm{H}}_{\mathrm{M}^F_e}(y)$ from the LHS yields the semi-small decomposition formula:

When e is not a coloop in M^F,

$$\underline{\mathrm{H}}_{\mathrm{M}^F}(y) = \underline{\mathrm{H}}_{\mathrm{M}^F\setminus e}(y) + \sum_{G \in S_e(\mathrm{M}^F)\setminus\{\emptyset\}} y\underline{\mathrm{H}}_{\mathrm{M}^F_{G\cup e}}(y) \cdot \underline{\mathrm{H}}_{\mathrm{M}^G}(y). \tag{4}$$

Case 2: e is a coloop in M^F. We take the coefficient of $\emptyset$ in the formula (3). Equating the coefficients gives:

$$x^f \underline{\mathrm{H}}_{\mathrm{M}^F}(x^{-2}) + x^{f-2}\underline{\mathrm{H}}_{\mathrm{M}^F_e}(x^{-2}) = (x + x^{-1})x^{f-1}\underline{\mathrm{H}}_{\mathrm{M}^F\setminus e}(x^{-2}) + x^{f-2}\sum_{\dots} \dots$$

Dividing by x^f and setting $y = x^{-2}$ leads to the known formula for adding a coloop:

When e is a coloop in M^F,

$$\underline{\mathrm{H}}_{\mathrm{M}^F}(y) = (1+y)\underline{\mathrm{H}}_{\mathrm{M}^F\setminus e}(y) + \sum_{G \in \mathcal{L}(\mathrm{M}^F\setminus e)\setminus\{\emptyset\}} y\underline{\mathrm{H}}_{\mathrm{M}^F_{G\cup e}}(y) \cdot \underline{\mathrm{H}}_{\mathrm{M}^G}(y). \tag{5}$$

This completes the recovery of the known decomposition formulas.

3.1 Recovery of the Augmented Chow Polynomial Decomposition Formulas

We consider another $\mathbb{Z}[x, x^{-1}]$-module homomorphism $\Phi_{\mathrm{M}} : \mathcal{H}(\mathrm{M}) \to \mathbb{Z}[x, x^{-1}]$ defined by

$$\Phi_{\mathrm{M}}(\alpha) = \sum_{F \in \mathcal{L}(\mathrm{M})} x^{-\mathrm{rk}(F)} \alpha_F,$$

which satisfies $\Phi_{\mathrm{M}\setminus e} \circ \Delta = \Phi_{\mathrm{M}}$. Applying this map to c^F gives

$$\Phi_{\mathrm{M}}(\mathrm{c}^F) = \sum_{G \leq F} x^{-\mathrm{rk}(G)} \mathrm{c}^F_G = x^{\mathrm{rk}(F)} \sum_{G \leq F} x^{-2\mathrm{rk}(G)} \underline{\mathrm{H}}_{\mathrm{M}^F_G}(x^{-2}) = x^{\mathrm{rk}(F)} \mathrm{H}_{\mathrm{M}^F}(x^{-2}) \in \mathrm{Pal}(0).$$

Remark. We see that the factorization $\Phi_{\mathrm{M}} = \Phi_{\mathrm{M}\setminus e} \circ \Delta$ as $\mathbb{Z}[x, x^{-1}]$-module homomorphisms also holds as $\mathrm{Pal}(0)$-module homomorphisms $\mathcal{S}(\mathrm{M}) \to \mathrm{Pal}(0)$.

Applying the $\Phi_{\mathrm{M}\setminus e}$ map to (2) yields the following identity:

$$x^f \mathrm{H}_{\mathrm{M}^F}(x^{-2}) = x^f \mathrm{H}_{\mathrm{M}^F\setminus e}(x^{-2}) + x^{-1} \sum_{G \in S_e(\mathrm{M}^F)} x^{f-g-1} \underline{\mathrm{H}}_{\mathrm{M}^F_{G\cup e}}(x^{-2}) \cdot x^g \mathrm{H}_{\mathrm{M}^G}(x^{-2}).$$

Simplifying the exponents and setting $y = x^{-2}$, we get:

When e is not a coloop in M^F,

$$\mathrm{H}_{\mathrm{M}^F}(y) = \mathrm{H}_{\mathrm{M}^F\setminus e}(y) + \sum_{G \in S_e(\mathrm{M}^F)} y \underline{\mathrm{H}}_{\mathrm{M}^F_{G\cup e}}(y) \cdot \mathrm{H}_{\mathrm{M}^G}(y). \tag{6}$$

Proceeding similarly for the coloop case, we get the following identity:

$$x^f \mathrm{H}_{\mathrm{M}^F}(x^{-2}) = (x + x^{-1}) x^{f-1} \mathrm{H}_{\mathrm{M}^F\setminus e}(x^{-2}) + x^{-1} \sum_{G \in S_e(\mathrm{M}^F)} x^{f-g-1} \underline{\mathrm{H}}_{\mathrm{M}^F_{G\cup e}}(x^{-2}) \cdot x^g \mathrm{H}_{\mathrm{M}^G}(x^{-2}).$$

Simplifying the exponents and setting $y = x^{-2}$, we get:

When e is a coloop in M^F,

$$\mathrm{H}_{\mathrm{M}^F}(y) = (1 + y) \mathrm{H}_{\mathrm{M}^F\setminus e}(y) + \sum_{G \in S_e(\mathrm{M}^F)} y \underline{\mathrm{H}}_{\mathrm{M}^F_{G\cup e}}(y) \cdot \mathrm{H}_{\mathrm{M}^G}(y). \tag{7}$$

4 Relating the ζ^F and c^F bases

Braden-Vysogorets define a basis $\{\zeta^F\}_{F \in \mathcal{L}(\mathrm{M})}$ of $\mathcal{H}(\mathrm{M})$ as

$$\zeta^F = \sum_{G \leq F} \zeta^F_G \cdot \mathrm{G} := \sum_{G \leq F} x^{\mathrm{rk}(F)-\mathrm{rk}(G)} \mathrm{P}_{\mathrm{M}^F_G}(x^{-2}) \cdot \mathrm{G}. \tag{8}$$

Using the fact that the inverse of the Chow function, $\underline{\mathrm{H}}$, in the incidence algebra of M is negative the reduced characteristic function, $\overline{\chi}$, and that the inverse of the Kazhdan-Lusztig function, P, is the inverse Kazhdan-Lusztig function, $\hat{\mathrm{Q}}$, of the matroid, we have the following change of basis formulas:

$$\mathrm{F} = \sum_{G \leq F} \mathfrak{f}^F_G \cdot \mathrm{c}^G = \sum_{G \leq F} x^{\mathrm{rk}(F)-\mathrm{rk}(G)} \overline{\chi}_{\mathrm{M}^F_G}(x^{-2}) \cdot \mathrm{c}^G. \tag{9}$$

$$\mathrm{F} = \sum_{G \leq F} \mathfrak{q}^F_G \cdot \zeta^G = \sum_{G \leq F} x^{\mathrm{rk}(F)-\mathrm{rk}(G)} \hat{\mathrm{Q}}_{\mathrm{M}^F_G}(x^{-2}) \cdot \zeta^G. \tag{10}$$

Applying the Φ_{M} map to (9) and (10) yields the following identities:

$$x^{-f} = -\sum_{G \leq F} x^{f-g} \overline{\chi}_{\mathrm{M}_G^F}(x^{-2}) \cdot x^g \mathrm{H}_{\mathrm{M}^G}(x^{-2}) \implies 1 + \sum_{G \leq F} x^{2f} \overline{\chi}_{\mathrm{M}_G^F}(x^{-2}) \cdot \mathrm{H}_{\mathrm{M}^G}(x^{-2}) = 0. \tag{11}$$

$$x^{-f} = \sum_{G \leq F} x^{f-g} \hat{\mathrm{Q}}_{\mathrm{M}_G^F}(x^{-2}) \cdot x^g \mathrm{Z}_{\mathrm{M}^G}(x^{-2}) \implies 1 - \sum_{G \leq F} x^{2f} \hat{\mathrm{Q}}_{\mathrm{M}_G^F}(x^{-2}) \cdot \mathrm{Z}_{\mathrm{M}^G}(x^{-2}) = 0. \tag{12}$$

Thus we have the following two identities in the incidence algebra of M:

$$\overline{\chi} \cdot \mathrm{H} = -x^r \text{ and } \hat{\mathrm{Q}} \cdot Z = x^r. \tag{13}$$

We define a new family of elements in $\mathcal{H}(\mathrm{M})$:

$$\xi^F := \sum_{G \leq F} x^{\mathrm{rk}(G)} \mu(\mathrm{M}_G^F) \cdot \mathrm{G}.$$

We have the following lemma from the previous paper:

Lemma 16. *Let* M *be a matroid on* E*, and let* $i \in E$ *be an element of* M*.*

1. *If* e *is a coloop of* M^F*, then* $\Delta(\xi^F) = 0$*.*
2. *If* e *is not a coloop of* M^F*, then* $\Delta(\xi^F) = \xi^{F \setminus e}$*.*

Since the $\zeta^F \in \mathcal{H}_p(\mathrm{M}) \subset \mathcal{S}(\mathrm{M})$, we can express it as:

$$\zeta^F = \sum_{G \in \mathcal{L}(\mathrm{M})} \mathfrak{a}_G^F \cdot \mathrm{c}^G \quad \text{where } \mathfrak{a}_G^F = \overline{\zeta_G^F} \in \mathrm{Pal}(0).$$

Example—Incorrect Simplification

For the boolean matroid B_n, the KL polynomial is $P_{B_k}(x) = 1$, so $\zeta_G^F = x^{\mathrm{rk}(F)-\mathrm{rk}(G)} = x^{f-g}$. Using the projection formula, we have that

$$\mathfrak{a}_G^F = \overline{\zeta_G^F} = \frac{x^{g-f+2} - x^{f-g}}{x^2-1} = \sum_{i=0}^{f-g} x^{f-g-2i}. \qquad \text{(the product telescopes)}$$

Thus, for the boolean matroid B_n,

$$\zeta^F = \sum_{G \le F} \left(\sum_{i=0}^{f-g} x^{f-g-2i} \right) \cdot \mathrm{c}^G.$$

The previous simplification to a single sum was incorrect. The correct coefficients are:

$$\begin{aligned}
f-g=0: &\quad \mathfrak{a}_G^F = \overline{x^0} = \overline{1} = 1 \\
f-g=1: &\quad \mathfrak{a}_G^F = \overline{x^1} = \frac{x^2(x^{-1}) - x}{x^2-1} = 0 \\
f-g=2: &\quad \mathfrak{a}_G^F = \overline{x^2} = \frac{x^2(x^{-2}) - x^2}{x^2-1} = \frac{1-x^2}{x^2-1} = -1 \\
f-g=3: &\quad \mathfrak{a}_G^F = \overline{x^3} = \frac{x^2(x^{-3}) - x^3}{x^2-1} = \frac{x^{-1}-x^3}{x^2-1} = -x - x^{-1} \\
f-g=4: &\quad \mathfrak{a}_G^F = \overline{x^4} = \frac{x^2(x^{-4}) - x^4}{x^2-1} = \frac{x^{-2}-x^4}{x^2-1} = -x^2 - 1 - x^{-2}
\end{aligned}$$

and so on.

Remark. Computing these coefficients in general does not seem to be easy.

We have that

$$\sum_{H \le F} \mathfrak{f}_H^F \cdot \mathrm{c}^H = \mathrm{F} = \sum_{G \le F} \mathfrak{q}_G^F \cdot \zeta^G = \sum_{G \le F} \mathfrak{q}_G^F \cdot \sum_{H \le G} \overline{\zeta_H^G} \cdot \mathrm{c}^H = \sum_{H \le F} \left(\sum_{G : H \le G \le F} \mathfrak{q}_G^F \cdot \overline{\zeta_H^G} \right) \cdot \mathrm{c}^H.$$

Thus $\mathfrak{f}_H^F = \sum_{G:H\le G\le F} \mathfrak{q}_G^F \cdot \overline{\zeta_H^G} \implies \mathfrak{f} = \mathfrak{q} \cdot \overline{\zeta}$. Multiplying both sides by *zeta* on the left yields:

$$\zeta \cdot \mathfrak{f} = \zeta \cdot \mathfrak{q} \cdot \overline{\zeta} = \overline{\zeta}.$$

5 The incidence algebra of M

Let $\mathrm{Int}(\mathrm{M})$ be the set of all intervals in $\mathcal{L}(\mathrm{M})$. In our study, **the incidence algebra** $\mathcal{I}(\mathrm{M})$ of M is the $\mathbb{Z}[x, x^{-1}]$-algebra with basis $\{\mathfrak{b}_G^F\}_{[G,F] \in \mathrm{Int}(\mathrm{M})}$. The algebra structure is given by

$$\mathfrak{b}_G^F \cdot \mathfrak{b}_H^K = \begin{cases} \mathfrak{b}_H^F & \text{if } G = K, \\ 0 & \text{otherwise.} \end{cases}$$

for all $F, G, H, K \in \mathcal{L}(\mathrm{M})$. The elements of $\mathcal{I}(\mathrm{M})$ are formal sums of the form

$$\alpha = \sum_{[G,F] \in \mathrm{Int}(\mathrm{M})} \alpha_{G,F} \cdot \mathfrak{b}_G^F \quad \text{where } \alpha_{G,F} \in \mathbb{Z}[x, x^{-1}].$$

6 todo

- Find the correct formula for the coefficients $\mathfrak{a}_G^F$ in the change of basis formula of the Boolean matroid.
- Define similar polynomial as Y for the reduced characteristic polynomial and see if it is palindromic.
- Use the definition of the mobius basis as in previous paper and see if it gives similar identities as (13).

Proposition 17. *Let $\alpha \in \mathcal{S}(\mathrm{M})$ and let p_F be the coefficients of its expansion in the $\{\zeta^F\}$ basis. These coefficients must satisfy the global condition:*

$$\sum_{F \in \mathcal{L}(\mathrm{M})} p_F(x) \cdot x^{\mathrm{rk}(F)} Z_{\mathrm{M}^F}(x^{-2}) \in \mathrm{Pal}(0)$$

Proof.

$$\mathrm{Pal}(0) \ni \Phi_{\mathrm{M}}(\alpha) = \sum_{F \in \mathcal{L}(\mathrm{M})} p_F(x) \cdot x^{\mathrm{rk}(F)} Z_{\mathrm{M}^F}(x^{-2})$$

□

$$\begin{aligned}
(\Delta_e \circ \Delta_f)(\mathrm{F}) &= \Delta_e(x^{-\delta_f(F)}\mathrm{F} \setminus \mathrm{f}) \\
&= x^{-\delta_f(F)} \cdot \Delta_e(\mathrm{F} \setminus \mathrm{f}) \\
&= x^{-\delta_f(F)} \cdot x^{-\delta_e(F \setminus f)}(\mathrm{F} \setminus \mathrm{f}) \setminus \mathrm{e} \\
&= x^{-(\delta_f(F)+\delta_e(F \setminus f))}\mathrm{F} \setminus \mathrm{e},\mathrm{f}
\end{aligned}$$

The exponent is the sum of the discrepancies. Let's expand it:

$$\begin{aligned}
\delta_f(F) + \delta_e(F \setminus f) &= \Big(\mathrm{rk}_{\mathrm{M}}(F) - \mathrm{rk}_{\mathrm{M} \setminus f}(F \setminus f)\Big) \\
&\quad + \Big(\mathrm{rk}_{\mathrm{M} \setminus f}(F \setminus f) - \mathrm{rk}_{\mathrm{M} \setminus e,f}(F \setminus e, f)\Big) \\
&= \mathrm{rk}_{\mathrm{M}}(F) - \mathrm{rk}_{\mathrm{M} \setminus e,f}(F \setminus e, f)
\end{aligned}$$

Notice this is a telescoping sum. The total exponent just measures the total rank drop from M to the minor $\mathrm{M} \setminus \{e, f\}$.

7 Commutativity of Deletion Operators

A fundamental property of the deletion operator is that the order in which elements are deleted does not matter. This combinatorial fact is reflected in the commutativity of the corresponding algebraic operators acting on our modules. Let $e, f \in E$ be distinct elements of the ground set. We have two deletion operators:

$$\begin{aligned}
\Delta_e &: \mathcal{H}(\mathrm{M}) \to \mathcal{H}(\mathrm{M} \setminus e) \\
\Delta_f &: \mathcal{H}(\mathrm{M}) \to \mathcal{H}(\mathrm{M} \setminus f)
\end{aligned}$$

We can compose them to form maps from $\mathcal{H}(\mathrm{M})$ to $\mathcal{H}(\mathrm{M} \setminus \{e, f\})$.

Proposition 18. *The deletion operators commute. That is, as maps from $\mathcal{H}(\mathrm{M})$ to $\mathcal{H}(\mathrm{M} \setminus \{e, f\})$, we have the equality:*

$$\Delta_f \circ \Delta_e = \Delta_e \circ \Delta_f.$$

Proof. Since the operators are $\mathbb{Z}[x, x^{-1}]$-module homomorphisms, it is sufficient to check that they agree on the standard basis vectors $\{\mathrm{H}\}_{H \in \mathcal{L}(\mathrm{M})}$. Let $\delta_e(H) := \mathrm{rk}_{\mathrm{M}}(H) - \mathrm{rk}_{\mathrm{M}\setminus e}(H \setminus e)$ denote the discrepancy of H when deleting e.

First, we compute the composition $\Delta_f \circ \Delta_e$.

$$\begin{aligned}
(\Delta_f \circ \Delta_e)(\mathrm{H}) &= \Delta_f\left(\Delta_e(\mathrm{H})\right) \\
&= \Delta_f\left(x^{-\delta_e(H)} \cdot \mathrm{H} \setminus \mathrm{e}\right) \\
&= x^{-\delta_e(H)} \cdot \Delta_f(\mathrm{H} \setminus \mathrm{e}) \\
&= x^{-\delta_e(H)} \cdot x^{-\delta_f(H\setminus e)} \cdot (\mathrm{H} \setminus \mathrm{e}) \setminus \mathrm{f} \\
&= x^{-(\delta_e(H)+\delta_f(H\setminus e))} \cdot \mathrm{H} \setminus \{\mathrm{e}, \mathrm{f}\}
\end{aligned}$$

The total exponent is the sum of the discrepancies:

$$\begin{aligned}
\delta_e(H) + \delta_f(H \setminus e) &= \left(\mathrm{rk}_{\mathrm{M}}(H) - \mathrm{rk}_{\mathrm{M}\setminus e}(H \setminus e)\right) + \left(\mathrm{rk}_{\mathrm{M}\setminus e}(H \setminus e) - \mathrm{rk}_{\mathrm{M}\setminus\{e,f\}}(H \setminus \{e, f\})\right) \\
&= \mathrm{rk}_{\mathrm{M}}(H) - \mathrm{rk}_{\mathrm{M}\setminus\{e,f\}}(H \setminus \{e, f\}).
\end{aligned}$$

This final expression is the total discrepancy from deleting both elements. It is symmetric with respect to e and f. If we were to compute the composition $\Delta_e \circ \Delta_f$, we would arrive at the same exponent and the same final basis vector $\mathrm{H} \setminus \{\mathrm{e}, \mathrm{f}\}$. Therefore, the operators commute. □

Remark. Since Δ_e and Δ_f are also $\mathrm{Pal}(0)$-module homomorphisms that preserve the submodules $\mathcal{S}(\mathrm{M})$, their composition is also a $\mathrm{Pal}(0)$-module homomorphism $\mathcal{S}(\mathrm{M}) \to \mathcal{S}(\mathrm{M} \setminus \{e, f\})$ that is independent of the order of application. This commutativity is essential for building chain complexes using sequences of deletion operators, a potential direction for future research.

8 A Cohomology Complex for $\mathcal{S}(\mathrm{M})$

The commutativity of the deletion operators allows us to construct a cochain complex whose properties could reveal deep information about the structure of the $\mathcal{S}(\mathrm{M})$ modules. This formalizes the idea of using the Boolean lattice of operators to deduce properties of $\mathcal{S}(\mathrm{M})$.

Definition 19 (The Cochain Complex of $\mathcal{S}$)**.** *Let* M *be a matroid on a ground set E.*

1. *For each integer $k \geq 0$, we define the **k-th cochain group** as the* $\mathrm{Pal}(0)$*-module*

$$C^k(\mathcal{S}, \mathrm{M}) := \bigoplus_{A \subseteq E, |A|=k} \mathcal{S}(\mathrm{M} \setminus A).$$

By definition, $C^0(\mathcal{S}, \mathrm{M}) = \mathcal{S}(\mathrm{M})$.

2. *Let us fix a total ordering on the ground set E. For $k \geq 0$, we define the* ***k-th coboundary map*** $d^k : C^k(\mathcal{S}, \mathrm{M}) \to C^{k+1}(\mathcal{S}, \mathrm{M})$ *by its action on each summand. For an element $\alpha \in \mathcal{S}(\mathrm{M} \setminus A)$ with $|A| = k$, we define*

$$d^k(\alpha) := \sum_{e \notin A} (-1)^{pos(e, E \setminus A)} \Delta_e(\alpha) \in \bigoplus_{e \notin A} \mathcal{S}(\mathrm{M} \setminus (A \cup \{e\})) \subseteq C^{k+1}(\mathcal{S}, \mathrm{M}),$$

where $pos(e, E \setminus A)$ is the position of the element e in the ordered set $E \setminus A$.

Proposition 20. *The pair $(C^\bullet(\mathcal{S}, \mathrm{M}), d^\bullet)$ is a cochain complex. That is, the composition of any two consecutive coboundary maps is zero: $d^{k+1} \circ d^k = 0$.*

Proof Sketch. We need to show that applying the map d twice results in the zero map. An element in the image of d^k is a sum of elements of the form $\Delta_e(\alpha)$. Applying d^{k+1} to such an element gives a sum of terms like $\Delta_f(\Delta_e(\alpha))$.

The full expression for $(d^{k+1} \circ d^k)(\alpha)$ will be a double summation over pairs of distinct elements $\{e_1, e_2\}$ not in the original deletion set. Each pair will appear twice, for example as $\Delta_{e_2} \circ \Delta_{e_1}$ and $\Delta_{e_1} \circ \Delta_{e_2}$. The alternating signs in the definition of the coboundary map are chosen precisely so that these two terms cancel each other out. This cancellation is guaranteed because the deletion operators commute, as shown in Proposition 18 ($\Delta_{e_1} \circ \Delta_{e_2} = \Delta_{e_2} \circ \Delta_{e_1}$). Thus, the entire sum is zero. □

Remark. Having defined a cochain complex, one can now study its **cohomology groups**, $H^k(\mathcal{S}, \mathrm{M}) := \ker(d^k)/\mathrm{im}(d^{k-1})$. These groups are new, potentially very rich, algebraic invariants of the matroid M. The "pipe dream" would be to compute these groups and discover that they have a simple structure (for example, that they vanish in most degrees). Such a result would imply profound new identities among the Chow polynomials of the matroid and its minors.

Proposition 21 (Expansion in the KL Basis)**.** *If $\alpha = \sum_F \alpha_F \cdot \mathrm{F}$ is an element of $\mathcal{S}(\mathrm{M})$, its expansion in the Kazhdan-Lusztig basis is $\alpha = \sum_F p_F \cdot \zeta^F$, where the coefficients $p_F \in \mathbb{Z}[x, x^{-1}]$ are determined by the recursive formula:*

$$p_F = \alpha_F - \sum_{G > F} p_G \cdot \zeta^G_F.$$

Proof. Let the expansion of α in the ζ^F basis be $\alpha = \sum_F p_F \cdot \zeta^F$. We can expand this in the standard basis:

$$\alpha = \sum_F p_F \left(\sum_{H \leq F} \zeta^F_H \cdot \mathrm{H} \right) = \sum_H \left(\sum_{F \geq H} p_F \cdot \zeta^F_H \right) \cdot \mathrm{H}.$$

By comparing the coefficient of H with the original expansion of α, we get:

$$\alpha_H = \sum_{F \geq H} p_F \cdot \zeta^F_H.$$

Since $\zeta^H_H = 1$, we can isolate the p_H term and solve for it, which yields the recursive formula in the proposition. The coefficients p_F for $F > H$ are computed first in an induction on decreasing rank. □

Remark (Why there is no simple projection). It's important to understand why this result is a recursion and not a simple projection like the other two cases.

- For $\alpha \in \mathcal{H}_p(\mathrm{M})$, the coefficients α_F are polynomials and the basis coefficients ζ_F^G are divisible by x for $G < F$. This special structure allows an inductive proof that the ζ^F-basis coefficients are simply $\alpha_F(0)$.
- For $\alpha \in \mathcal{S}(\mathrm{M})$ expanded in the c^F basis, the basis coefficients c_F^G have the convenient Pal(2) property. This forces the standard-basis coefficient α_F to have a unique 'Pal(0) + Pal(2)' structure, allowing you to find the c^F-basis coefficient by projection.

In the present case—expanding $\alpha \in \mathcal{S}(\mathrm{M})$ in the ζ^F basis—neither of these special conditions applies. The coefficients α_F are Laurent polynomials and the basis coefficients ζ_F^G have no special palindromic structure. Therefore, the full recursive formula is the only way to describe the relationship. The resulting coefficients p_F will, in general, be complex Laurent polynomials, not simply constants or 'Pal(0)' elements.

9 Relating the ζ^F and c^F Bases

The two special bases for $\mathcal{H}(\mathrm{M})$ can be expressed in terms of each other. The formulas for this change of basis reveal a deep connection between the polynomial invariants that define them.

9.1 Expressing c^F in the ζ^F Basis

We can find the coefficients for expanding c^F in the ζ^F basis by composing the known change-of-basis maps through the standard basis $\{\mathrm{G}\}$. In the language of the incidence algebra, this amounts to a convolution product.

Proposition 22. *The expansion of the Chow basis element c^F in the Kazhdan-Lusztig basis $\{\zeta^G\}$ is given by:*

$$\mathrm{c}^F = \sum_{G \leq F} B_{FG}(x) \cdot \zeta^G,$$

where the coefficient $B_{FG}(x)$ is given by the convolution of the Chow polynomial and the inverse Kazhdan-Lusztig polynomial:

$$B_{FG}(x) = \sum_{K : G \leq K \leq F} \left(x^{\mathrm{rk}(F)-\mathrm{rk}(K)} \underline{\mathrm{H}}_{\mathrm{M}_K^F}(x^{-2}) \right) \cdot \left(x^{\mathrm{rk}(K)-\mathrm{rk}(G)} Q_{\mathrm{M}_G^K}(x^{-2}) \right).$$

Proof. In the incidence algebra, the relationships between the bases are given by convolutions. The definition of the c^F basis can be written as $\mathrm{c} = * \mathbf{c}$, where $\mathbf{c}(G, F) = \mathrm{c}_G^F$. Similarly, we have $\zeta = * \mathbf{z}$.

From [**gx-inversekazhdanlustig**], we know that the inverse of the KL polynomial function $\mathbf{p}$ is the inverse KL polynomial function $\mathbf{q}$. Since $\mathbf{z}$ is defined using $\mathbf{p}$, its inverse in the incidence algebra is the function for Q. Therefore, we can express the standard basis as $= \zeta * \mathbf{q}$, where $\mathbf{q}(G, F) = x^{\mathrm{rk}(F)-\mathrm{rk}(G)} Q_{\mathrm{M}_G^F}(x^{-2})$.

Substituting this into the definition of c^F gives:

$$\mathrm{c} = * \mathbf{c} = (\zeta * \mathbf{q}) * \mathbf{c} = \zeta * (\mathbf{q} * \mathbf{c}).$$

The formula in the proposition is the explicit expression of the convolution product $(\mathbf{q} * \mathbf{c})(G, F)$. □

Remark. This proposition provides a direct formula to translate from the Chow basis to the Kazhdan-Lusztig basis. The coefficients $B_{FG}(x)$ are, in general, complex Laurent polynomials whose structure is determined by the interplay between the Chow polynomials and the inverse Kazhdan-Lusztig polynomials of the matroid's minors.

$$\mathfrak{b}_G^F(x) = x^{\mathrm{rk}(F)-\mathrm{rk}(G)} \left(\underline{\mathrm{H}}_{\mathrm{M}_G^F}(x^{-2}) - \mathrm{P}_{\mathrm{M}_G^F}(x^{-2}) \right) - \sum_{K:G<K<F} \mathfrak{b}_K^F(x) \cdot x^{\mathrm{rk}(K)-\mathrm{rk}(G)} \mathrm{P}_{\mathrm{M}_G^K}(x^{-2}) \tag{14}$$

Expressing the Chow Basis in the KL Basis

We want to find the coefficients $\mathfrak{b}_G^F(x) \in \mathrm{Pal}(0)$ in the expansion:

$$\mathrm{c}^F = \sum_{G \leq F} \mathfrak{b}_G^F(x) \cdot \zeta^G$$

Since both bases are lower-triangular with respect to the standard basis $\{\mathrm{L}\}$, we can solve for the coefficients $\mathfrak{b}_G^F(x)$ recursively.

By expanding both sides in the standard basis and equating the coefficients of a given basis vector L where $L \leq F$, we get:

$$\mathrm{c}_L^F = \sum_{G:L \leq G \leq F} \mathfrak{b}_G^F(x) \cdot \zeta_L^G$$

Using the definitions of the basis coefficients from your notes, this equation becomes:

$$x^{\mathrm{rk}(F)-\mathrm{rk}(L)} \underline{\mathrm{H}}_{\mathrm{M}_L^F}(x^{-2}) = \sum_{G:L \leq G \leq F} \mathfrak{b}_G^F(x) \cdot x^{\mathrm{rk}(G)-\mathrm{rk}(L)} \mathrm{P}_{\mathrm{M}_L^G}(x^{-2})$$

A Recursive Formula for the Coefficients

We can rearrange this equation to get a recursive formula for the coefficients $\mathfrak{b}_G^F(x)$. We solve for the coefficients by induction on the corank of G within the interval $[\emptyset, F]$.

The relationship is defined by the following recursive formula:

1. **Base Case ($G = F$):**
$$\mathfrak{b}_F^F(x) = 1$$
2. **Recursive Step (for $G < F$):**
$$\mathfrak{b}_G^F(x) = x^{\mathrm{rk}(F)-\mathrm{rk}(G)} \left(\underline{\mathrm{H}}_{\mathrm{M}_G^F}(x^{-2}) - \mathrm{P}_{\mathrm{M}_G^F}(x^{-2}) \right) - \sum_{K:G<K<F} \mathfrak{b}_K^F(x) \cdot x^{\mathrm{rk}(K)-\mathrm{rk}(G)} \mathrm{P}_{\mathrm{M}_G^K}(x^{-2})$$

You can use this formula to compute the coefficients starting from the top of the lattice (where $\mathfrak{b}_F^F = 1$) and working your way down. The fact that both bases belong to the module $\mathcal{S}(\mathrm{M})$ guarantees that each coefficient $\mathfrak{b}_G^F(x)$ you compute will be in $\mathrm{Pal}(0)$.

Expressing the Chow Basis in the KL Basis

We want to find the coefficients $\mathfrak{b}_G^F(x) \in \mathrm{Pal}(0)$ in the expansion:

$$\mathrm{c}^F = \sum_{G \leq F} \mathfrak{b}_G^F(x) \cdot \zeta^G$$

By expanding both sides in the standard basis and equating the coefficients of a given basis vector L where $L \leq F$, we get:

$$\mathrm{c}_L^F = \sum_{G: L \leq G \leq F} \mathfrak{b}_G^F(x) \cdot \zeta_L^G$$

Using the definitions of the basis coefficients from your notes, this equation becomes:

$$x^{\mathrm{rk}(F)-\mathrm{rk}(L)} \underline{\mathrm{H}}_{\mathrm{M}_L^F}(x^{-2}) = \sum_{G: L \leq G \leq F} \mathfrak{b}_G^F(x) \cdot x^{\mathrm{rk}(G)-\mathrm{rk}(L)} \mathrm{P}_{\mathrm{M}_L^G}(x^{-2})$$

10 Simplification by Palindromicity

Your observations are correct, and they lead to a significant simplification of the change-of-basis formula. By using the properties of palindromicity, we can decompose the single equation into two separate, simpler identities.

10.1 The Decomposition Principle

The key here is the unique decomposition of any Laurent polynomial f into a $\mathrm{Pal}(0)$ part and a $\mathrm{Pal}(2)$ part, as stated in Proposition 11. We will denote the $\mathrm{Pal}(0)$ component of a polynomial f as $(f)_0$ and the $\mathrm{Pal}(2)$ component as $(f)_2$, such that $f = (f)_0 + (f)_2$. We apply this principle to the equation:

$$\mathrm{c}_L^F = \sum_{G: L \leq G \leq F} \mathfrak{b}_G^F(x) \cdot \zeta_L^G$$

10.2 Applying the Decomposition

1. **Analyze the Left-Hand Side (LHS):** As you noted, the Chow basis coefficient c_L^F is in $\mathrm{Pal}(2)$ for all $L < F$. This means its decomposition is trivial:
   - $(\mathrm{c}_L^F)_0 = 0$
   - $(\mathrm{c}_L^F)_2 = \mathrm{c}_L^F$
2. **Analyze the Right-Hand Side (RHS):** Each term in the sum is a product of $\mathfrak{b}_G^F(x)$ and ζ_L^G.
   - We know that the coefficient $\mathfrak{b}_G^F(x)$ must be in $\mathrm{Pal}(0)$.
   - The term ζ_L^G is a polynomial but has no inherent palindromic symmetry. We can decompose it into its parts: $\zeta_L^G = (\zeta_L^G)_0 + (\zeta_L^G)_2$.

   Therefore, each term in the sum is $\mathfrak{b}_G^F \cdot ((\zeta_L^G)_0 + (\zeta_L^G)_2)$. Since multiplying a $\mathrm{Pal}(0)$ term by a $\mathrm{Pal}(n)$ term results in a $\mathrm{Pal}(n)$ term, we can separate the entire sum into its $\mathrm{Pal}(0)$ and $\mathrm{Pal}(2)$ components:
   - $(\mathrm{RHS})_0 = \sum_{G: L \leq G \leq F} \mathfrak{b}_G^F(x) \cdot (\zeta_L^G)_0$

- $(\text{RHS})_2 = \sum_{G:L\le G\le F} \mathfrak{b}_G^F(x)\cdot\left(\zeta_L^G\right)_2$

10.3 The Simplified Equations

By equating the corresponding palindromic components of the LHS and RHS, we replace the original mixed equation with two "purer" identities.

Simplified Identities (for $L < F$)

1. Equating the $\text{Pal}(0)$ components gives a set of linear constraints on the coefficients:
$$\sum_{G:L\le G\le F} \mathfrak{b}_G^F(x)\cdot\left(\zeta_L^G\right)_0 = 0$$
2. Equating the $\text{Pal}(2)$ components gives a new formula for the Chow coefficients:
$$\mathrm{c}_L^F = \sum_{G:L\le G\le F} \mathfrak{b}_G^F(x)\cdot\left(\zeta_L^G\right)_2$$

This is a powerful simplification. The original equation has been split into two distinct relationships that separate the symmetric and anti-symmetric aspects of the connection between the Chow and Kazhdan-Lusztig bases.

We know that
$$\zeta_L^F = x^{\mathrm{rk}(F)-\mathrm{rk}(L)}\mathrm{P}_{\mathrm{M}_L^F}(x^{-2})$$
Let
$$\zeta^F = \sum_{G\le F}\zeta_G^F\cdot\mathrm{G} = \sum_{H\le F}\mathfrak{a}_H^F\cdot\mathrm{c}^H = \sum_{H\le F}\mathfrak{a}_H^F\cdot\sum_{G\le H}\mathrm{c}_G^H\cdot\mathrm{G} = \sum_{G\le F}\left(\sum_{H:G\le H\le F}\mathfrak{a}_H^F\cdot\mathrm{c}_G^H\right)\cdot\mathrm{G}$$
We know that
$$\mathfrak{a}_H^F = \overline{\zeta_H^F} = \frac{x^{h-f+2}\cdot\mathrm{P}_{\mathrm{M}_H^F}(x^2) - x^{f-h}\cdot\mathrm{P}_{\mathrm{M}_H^F}(x^{-2})}{x^2-1}$$
So,
$$\zeta_G^F = \sum_{H:G\le H\le F}\frac{x^{h-f+2}\cdot\mathrm{P}_{\mathrm{M}_H^F}(x^2) - x^{f-h}\cdot\mathrm{P}_{\mathrm{M}_H^F}(x^{-2})}{x^2-1}x^{h-g}\cdot\underline{\mathrm{H}}_{\mathrm{M}_G^H}(x^{-2})$$
$$x^{f-g}\cdot\mathrm{P}_{\mathrm{M}_G^F}(x^{-2}) = \sum_{H\in[G,F]}\frac{x^{2h-f+2}\cdot\mathrm{P}_{\mathrm{M}_H^F}(x^2) - x^{f}\cdot\mathrm{P}_{\mathrm{M}_H^F}(x^{-2})}{x^2-1}\cdot x^{-g}\underline{\mathrm{H}}_{\mathrm{M}_G^H}(x^{-2})$$
$$\mathrm{P}_{\mathrm{M}_G^F}(x^{-2}) = \sum_{H\in[G,F]}\frac{x^{2h-2f+2}\cdot\mathrm{P}_{\mathrm{M}_H^F}(x^2) - \mathrm{P}_{\mathrm{M}_H^F}(x^{-2})}{x^2-1}\cdot\underline{\mathrm{H}}_{\mathrm{M}_G^H}(x^{-2})$$

Setting $y = x^{-2}$, we get:
$$(y^{-1}-1)\cdot\mathrm{P}_{\mathrm{M}_G^F}(y) = \sum_{H\in[G,F]}\left(y^{\mathrm{rk}(\mathrm{M}_H^F)-1}\cdot\mathrm{P}_{\mathrm{M}_H^F}(y^{-1}) - \mathrm{P}_{\mathrm{M}_H^F}(y)\right)\cdot\underline{\mathrm{H}}_{\mathrm{M}_G^H}(y)$$

$$(1-y)\cdot \mathrm{P}_{\mathrm{M}_G^F}(y) = \sum_{H\in[G,F]} \left(y^{\mathrm{rk}(\mathrm{M}_H^F)}\cdot \mathrm{P}_{\mathrm{M}_H^F}(y^{-1}) - y\cdot \mathrm{P}_{\mathrm{M}_H^F}(y)\right)\cdot \underline{\mathrm{H}}_{\mathrm{M}_G^H}(y)$$

Taking $F = \mathrm{M}$ and $G = \emptyset$, we get:

$$(1-y)\cdot \mathrm{P}_{\mathrm{M}}(y) = \sum_{H\in\mathcal{L}(\mathrm{M})} \left(y^{\mathrm{crk}(H)}\cdot \mathrm{P}_{\mathrm{M}_H}(y^{-1}) - y\cdot \mathrm{P}_{\mathrm{M}_H}(y)\right)\cdot \underline{\mathrm{H}}_{\mathrm{M}^H}(y)$$

$$(1-y)\cdot \mathrm{P}_{\mathrm{M}}(y) = y^r\cdot \mathrm{P}_{\mathrm{M}}(y^{-1}) - y\cdot \mathrm{P}_{\mathrm{M}}(y) + \sum_{H\neq\emptyset} \left(y^{\mathrm{crk}(H)}\cdot \mathrm{P}_{\mathrm{M}_H}(y^{-1}) - y\cdot \mathrm{P}_{\mathrm{M}_H}(y)\right)\cdot \underline{\mathrm{H}}_{\mathrm{M}^H}(y)$$

$$\mathrm{P}_{\mathrm{M}}(y) - y^r\cdot \mathrm{P}_{\mathrm{M}}(y^{-1}) = \sum_{H\neq\emptyset} \left(y^{\mathrm{crk}(H)}\cdot \mathrm{P}_{\mathrm{M}_H}(y^{-1}) - y\cdot \mathrm{P}_{\mathrm{M}_H}(y)\right)\cdot \underline{\mathrm{H}}_{\mathrm{M}^H}(y)$$

Can we use the following identity to simplify the sum:

$$y^r\cdot \mathrm{P}_{\mathrm{M}}(y^{-1}) - \mathrm{P}_{\mathrm{M}}(y) = \sum_{F\neq\emptyset} \chi_{\mathrm{M}^F}(y)\cdot \mathrm{P}_{\mathrm{M}_F}(y) = R_{\mathrm{M}}(y)$$

$$\sum_{F\neq\emptyset} \chi_{\mathrm{M}^F}(y)\cdot \mathrm{P}_{\mathrm{M}_F}(y) = \sum_{H\neq\emptyset} \left(y\cdot \mathrm{P}_{\mathrm{M}_H}(y) - y^{\mathrm{crk}(H)}\cdot \mathrm{P}_{\mathrm{M}_H}(y^{-1})\right)\cdot \underline{\mathrm{H}}_{\mathrm{M}^H}(y)$$

$$R_{\mathrm{M}}(y) = \sum_{H\neq\emptyset} \left(y\cdot \mathrm{P}_{\mathrm{M}_H}(y) - R_{\mathrm{M}_H}(y) - \mathrm{P}_{\mathrm{M}_H}(y)\right)\cdot \underline{\mathrm{H}}_{\mathrm{M}^H}(y)$$

$$R_{\mathrm{M}}(y) = (y-1)\cdot \sum_{H\neq\emptyset} \mathrm{P}_{\mathrm{M}_H}(y)\cdot \underline{\mathrm{H}}_{\mathrm{M}^H}(y) - \sum_{H\neq\emptyset} R_{\mathrm{M}_H}(y)\cdot \underline{\mathrm{H}}_{\mathrm{M}^H}(y)$$

$$\sum_{H\in\mathcal{L}(\mathrm{M})} R_{\mathrm{M}_H}(y)\cdot \underline{\mathrm{H}}_{\mathrm{M}^H}(y) = (y-1)\sum_{H\neq\emptyset} \mathrm{P}_{\mathrm{M}_H}(y)\cdot \underline{\mathrm{H}}_{\mathrm{M}^H}(y)$$

$$\sum_{H\in\mathcal{L}(\mathrm{M})}\sum_{H<F} \chi_{\mathrm{M}_H^F}(y)\cdot \mathrm{P}_{\mathrm{M}_F}(y)\cdot \underline{\mathrm{H}}_{\mathrm{M}^H}(y) = (y-1)\sum_{H\neq\emptyset} \mathrm{P}_{\mathrm{M}_H}(y)\cdot \underline{\mathrm{H}}_{\mathrm{M}^H}(y)$$

$$\sum_{\emptyset\neq F\in\mathcal{L}(\mathrm{M})} \mathrm{P}_{\mathrm{M}_F}(y)\cdot \sum_{H<F} \underline{\mathrm{H}}_{\mathrm{M}^H}\cdot \chi_{\mathrm{M}_H^F}(y) = (y-1)\sum_{H\neq\emptyset} \mathrm{P}_{\mathrm{M}_H}(y)\cdot \underline{\mathrm{H}}_{\mathrm{M}^H}(y)$$