

1) If $t_1(n) \in o(g_1(n))$ and $t_2(n) \in o(g_2(n))$, then $t_1(n) + t_2(n) \in o(\max\{g_1(n), g_2(n)\})$. prove the assertions.

Sol: we need to show that $t_1(n) + t_2(n) \in o(\max\{g_1(n), g_2(n)\})$. This means there exists a positive constant c and n_0 such that $t_1(n) + t_2(n) \leq c$.

$$t_1(n) \leq c_1 g_1(n) \text{ for all } n \geq n_1$$

$$t_2(n) \leq c_2 g_2(n) \text{ for all } n \geq n_2$$

$$\text{let } n_0 = \max\{n_1, n_2\} \text{ for all } n \geq n_0$$

consider $t_1(n) + t_2(n)$ for all $n \geq n_0$

$$t_1(n) + t_2(n) \leq c_1 g_1(n) + c_2 g_2(n)$$

we need to relate $g_1(n)$ and $g_2(n)$ to $\max\{g_1(n), g_2(n)\}$:

$$g_1(n) \leq \max\{g_1(n), g_2(n)\} \text{ and } g_2(n) \leq \max\{g_1(n), g_2(n)\}$$

thus,

$$c_1 g_1(n) \leq c_1 \max\{g_1(n), g_2(n)\}$$

$$c_2 g_2(n) \leq c_2 \max\{g_1(n), g_2(n)\}$$

$$c_1 g_1(n) + c_2 g_2(n) \leq c_1 \max\{g_1(n), g_2(n)\} + c_2 \max\{g_1(n), g_2(n)\}$$

$$c_1 g_1(n) + c_2 g_2(n) \leq (c_1 + c_2) \max\{g_1(n), g_2(n)\}$$

$$t_1(n) + t_2(n) \leq (c_1 + c_2) \max\{g_1(n), g_2(n)\} \text{ for all } n \geq n_0$$

By the definition of Big-O notation.

$$t_1(n) + t_2(n) \in o(\max\{g_1(n), g_2(n)\})$$

$$t_1(n) + t_2(n) \in o(\max\{g_1(n), g_2(n)\})$$

thus, the assertion is proved.

2) Find the time complexity of the Recurrence equation.

Sol: let us consider such that Recurrence for merge sort.

$$T(n) = 2T(n/2) + n$$

By using master theorem

$$T(n) = aT(n/b) + f(n)$$

where $a \geq 1$, $b \geq 1$ and $f(n)$ is positive function.

Ex:- $T(n) = 2T(n/2) + n$

$$a = 2, b = 2, f(n) = n$$

By comparing of $f(n)$ with $n^{\log_b a}$

$$\log_b a = \log_2 2 = 1$$

Compare $f(n)$ with $n^{\log_b a}$:

$$f(n) = n$$

$$n^{\log_b a} = n^1 = n$$

$$* f(n) = O(n^{\log_b a}), \text{ then } T(n) = O(n^{\log_b a} \log n)$$

In our case:

$$\log_b a = 1$$

$$T(n) = O(n^1 \log n) = O(n \log n)$$

Then time complexity of Recurrence Relation is $T(n) = 2T(n/2) + n$ is

$$O(n \log n)$$

$$T(n) = \begin{cases} 2T(n/2) + 1 & \text{if } n > 1 \\ 1 & \text{otherwise} \end{cases}$$

By applying of master theorem

$$T(n) = aT(n/b) + f(n) \text{ where } a \geq 1, b > 1$$

$$T(n) = 2T(n/2) + 1$$

$$\text{here } a=2, b=2, f(n)=1$$

By comparison of $f(n)$ and $n^{\log_b a}$

If $f(n) = O(n^c)$ where $c < \log_b a$, then $T(n) = O(n^{\log_b a})$.

If $f(n) = O(n^{\log_b a})$, then $T(n) = O(n^{\log_b a} \log n)$.

If $f(n) = \Omega(n^c)$ where $c > \log_b a$ then $T(n) = O(f(n))$

Let's calculate $\log_b a$:

$$\log_b a = \log_2 2 = 1$$

$$f(n) = 1$$

$$n^{\log_b a} = n^1 = n$$

$f(n) = O(n^c)$ with $c < \log_b a$ (case 1)

In this case $c=0$ and $\log_b a = 1$

$$c < 1, \text{ so } T(n) = O(n^{\log_b a}) = O(n^1) = O(n)$$

Time complexity of Recurrence Relation.

$$T(n) = 2T(n/2) + 1 \text{ is } O(n)$$

$$4) T(n) = \begin{cases} 2T(n-1) & \text{if } n > 0 \\ 1 & \text{otherwise} \end{cases}$$

Sol: Here, where $n=0$

$$T(0) = 1$$

Recurrence Relation Analysis

for $n > 0$:

$$T(n) = 2T(n-1)$$

$$T(n) = 2T(n-1)$$

$$T(n-1) = 2T(n-2)$$

$$T(n-2) = 2T(n-3)$$

$$T(1) = 2T(0)$$

from this pattern

$$T(n) = 2 \cdot 2 \cdot 2 \dots 2 T(0) = 2^n T(0)$$

Since $T(0) = 1$, we have

$$T(n) = 2^n$$

The Recurrence Relation is

$$T(n) = 2T(n-1) \text{ for } n > 0 \text{ and } T(0) = 1 \text{ is } T(n) = 2^n$$

5) Big O notation: Show that $f(n) = n^2 + 3n + 5$ is $O(n^2)$.

Sol: $f(n) = O(g(n))$ means $c > 0$ and $n_0 \geq 0$

$$f(n) \leq c \cdot g(n) \text{ for all } n \geq n_0$$

$$\text{Given is } f(n) = n^2 + 3n + 5$$

$$c > 0, n_0 \geq 0 \text{ such that } f(n) \leq c \cdot n^2$$

$$f(n) = n^2 + 3n + 5$$

Let's choose $c = 2$

$$f(n) \leq 2 \cdot n^2$$

$$f(n) = n^2 + 3n + 5 \leq n^2 + 3n^2 + 5n^2 = 9n^2$$

$$\text{So, } c = 9, n_0 = 1 \text{ } f(n) \leq 9n^2 \text{ for all } n \geq 1$$

$$f(n) = n^2 + 3n + 5 \text{ is } O(n^2).$$