

Lecture notes

Einführung in die Logik 2024W

This is a summary of the material discussed in the lecture "Mathematische Logik". It is still a work in progress and there **may be mistakes** in this work. If you find any, feel free to let me know and I will correct them

The content of this script relies on [EE01], [Van98] and [Kri98] Dieses Skript ist noch nicht vollständig und wird regelmäßig aktualisiert.

Contents

1	Propositional logic	2
1.1	Truth assignments	3
1.2	A parsing algorithm	4
1.3	Induction and recursion	6
1.4	Sentential connectives	8
1.5	Compactness Theorem	10
2	Predicate - / first order logic	11
2.1	Formulas	12
2.2	Semantics of first order logic	13
2.3	definability in a structure	16
2.4	Homomorphisms of structures	17
2.5	Unique readability for terms	19
2.6	A parsing algorithm for first order logic	20
2.7	Deductions (formal proofs)	20
2.8	Generalization and deduction theorem	22
2.9	Soundness and completeness	25
2.9.1	Sizes of models	28
2.9.2	Completeness for uncountable languages	29
3	Model Theory	30
3.1	Löwenheim-Skolem-Theorem	30
3.2	Theories and completeness	30
3.3	Theory of algebraic closed fields	32
3.4	Nonstandard Analysis	33
3.5	o-minimality	36
3.6	o-minimal ordered groups and rings	36
4	Boolean Algebra	43
4.1	Stone Representation Theorem	44
4.2	Lindenbaum-Tarski Algebras	45
5	Set Theory of ZFC	49
5.1	First axioms of ZFC	49
5.1.1	Classes and functions	50
5.2	Ordinals	52
5.2.1	Transfinite induction	54
5.2.2	Axiom of Choice	55
5.2.3	Ordinal arithmetic	56
5.2.4	Cardinals	58
6	Outernotes, Abbreviations, Overview	65

CHAPTER 1

Propositional logic

The definitions, lemmata, propositions and theorems as well as the notes in this chapter are sourced from [EE01, chapter 1].

Definition 1.1. Language of PL: The Language of Propositional logic is a set containing

Language

- logical symbols: consisting of the **sentential connective** symbols $\neg, \wedge, \vee, \rightarrow, \leftrightarrow$ and parenthesis $(,)$
- non-logical symbols: A_1, A_2, A_3, \dots (also called sentential atoms, variables)

from which we assume (for unique readability) that no symbol is a finite sequence of any other symbols.

Note :

1. The role of the logical symbols doesn't change, the sentential atoms we see as variables, they function as placeholders or variables.
2. we assumed the set of non-logical symbols is countable, for most of our conclusions you could use any set of prop. atoms of any size

Definition 1.2. Expression / prop. sentence: An **expression** is a any finite sequence of symbols We define **grammatically correct exp.** recursively

expression


1. every prop. atom is a prop. sentence
2. if α, β are prop. sentences, then also $(\neg\alpha), (\alpha \wedge \beta), (\alpha \vee \beta), (\alpha \rightarrow \beta), (\alpha \leftrightarrow \beta)$
3. nothing else (in particular \emptyset is not a prop. fla.)

and call them **prop. sentences** or **prop. fla.** Equivalently stated every prop. sentence is built up by applying finitely many formula building operations on atoms and the prop. sent. returned from building operations.

prop. fla.
prop. sentence

$$\mathcal{E}_{\neg}, \mathcal{E}_{\neg}(\alpha) := (\neg\alpha) \text{ for any prop. fla. } \alpha \text{ and similarly for } \mathcal{E}_{\wedge}, \mathcal{E}_{\vee}, \mathcal{E}_{\rightarrow}, \mathcal{E}_{\leftrightarrow}$$

This allows us to symbolize the **expression tree** (Here for example for $((\neg(A_1 \rightarrow A_2)) \vee A_3)$)



We will return to these construction trees in 1.2, where we answer the question of what truth value a given prop. sentence might have.

Definition 1.3. Construction sequence: Given a prop. sentence α a **construction sequence** of α is a finite sequence $\langle \alpha_1, \dots, \alpha_{n-1}, \alpha \rangle$ such that for all $i \leq n$ the following holds

construction
sequence

- α_i is a sentential atom
- or $\alpha_i = \mathcal{E}_{\neg}(\alpha_j)$ for some $j < i$
- or $\alpha_i = \mathcal{E}_{\Box}(\alpha_j, \alpha_k)$ for some $j, k < i$ and $\Box \in \{\wedge, \vee, \rightarrow, \leftrightarrow\}$

Definition 1.4. Closedness of a set: Let S be a set. We say S is **closed** under an n -ary operational symbol f iff for all $s_1, s_2, \dots, s_n \in S$ it holds $f(s_1, s_2, \dots, s_n) \in S$

closure

Induction principle: Suppose S is a set of prop. sentences containing all prop. atoms and closed under the 5 formula building operations, then S is the set of all prop. sentences.

Proof. let PS = set of all prop. sent.

$S \subseteq PS$: is clear

$S \supseteq PS$: let $\alpha \in PS$ then α has a construction seq. $\langle \alpha_1, \dots, \alpha_{n-1}, \alpha \rangle$ and $\alpha_1 \in S$. Let's assume that for $i \leq k < n$ each α_i is in S . Then α_{k+1} is either an atom and therefore in S or its obtained by one of the formula building operations and therefore $\alpha_{k+1} \in S$

□

1.1 TRUTH ASSIGNMENTS

The interpretation of a prop. atom is either true or false, denoted by 0/1 or T/F or \top/\perp . A truth assignment is simply any map $\nu : S \mapsto \{0, 1\}$, where S is a map of propositional atoms. Our goal is going to be to extend any truth assignment ν to a function $\bar{\nu} : \bar{S} \mapsto \{0, 1\}$, where \bar{S} is the closure of S under the 5 fla. building operations.

Definition 1.5. Truth assignment: Let $\{0, 1\}$ be the set of truth values. A truth assignment (TA) for a set S of prop. atoms is a map $\nu : S \rightarrow \{0, 1\}$

Truth assignment
TA

We now want to extend ν to $\bar{\nu} : \bar{S} \rightarrow \{0, 1\}$, where \bar{S} is the closure of S under the 5 fla. building operations such that for all propositional atoms $A \in S$ and propositional formulas α, β in \bar{S}

1. $\bar{\nu}(A) = \nu(A)$
2. $\bar{\nu}(\neg\alpha) = 1 - \nu(\alpha)$
3. $\bar{\nu}(\alpha \wedge \beta) = \begin{cases} 1 & \text{iff } \bar{\nu}(\alpha) = 1 = \bar{\nu}(\beta) \\ 0 & \text{otherwise} \end{cases}$
4. $\bar{\nu}(\alpha \vee \beta) = \begin{cases} 1 & \text{iff } \bar{\nu}(\alpha) = 1 \text{ or } \bar{\nu}(\beta) = 1 \\ 0 & \text{otherwise} \end{cases}$
5. $\bar{\nu}(\alpha \rightarrow \beta) = \begin{cases} 1 & \text{iff } \bar{\nu}(\alpha) = 0 \text{ or } \bar{\nu}(\beta) = 1 \\ 0 & \text{otherwise} \end{cases}$
6. $\bar{\nu}(\alpha \leftrightarrow \beta) = \begin{cases} 1 & \text{iff } \bar{\nu}(\alpha) = \bar{\nu}(\beta) \\ 0 & \text{otherwise} \end{cases}$

We also want the extension to be unique, that is

Theorem 1.1. Unique readability: For all TA ν for a set S $\exists! \bar{\nu} : \bar{S} \rightarrow \{0, 1\}$ satisfying the above properties

We will prove this later

Definition 1.6. Satisfaction: A TA ν satisfies a prop. sent. α if $\bar{\nu}(\alpha) = 1$ (that is, provided that every atom of α is in the domain of ν). We call α **satisfiable** if there exists a TA that satisfies it.

satisfy
satisfiable

Definition 1.7. Tautological implication: Let Σ be a set of prop. sent. and α a prop. sent. then we say: Σ tautologically implies α if for all TA that satisfy Σ , α is also satisfied and we write $\Sigma \models \alpha$. If $\Sigma = \{\beta\}$, we simply write $\beta \models \alpha$. If $\Sigma = \emptyset$ then α is called a **tautology** and we write $\models \alpha$ instead of $\emptyset \models \alpha$. α, β are called **tautologically equivalent** iff $\alpha \models \beta$ and $\beta \models \alpha$, we then write $\alpha \models \beta$.

taut. implication
 \models

Note : In other words, tautological implication $\Sigma \models \alpha$ means that you can not find a TA, that satisfy all members of Σ but not α . A tautology is satisfied by every TA. Suppose there is no TA that satisfies Σ , then we have $\Sigma \models \alpha$ for every prop. sent. α .

Example 1.1. $\{\neg A \vee B\} \models A \rightarrow B$

Note : In order to check if a prop. sent. is satisfiable we need to check 2^N TAs, where $N = \#$ of atoms. It is unknown if this can be done by an algorithm in polynomial time. Answering this would settle the debate whether $P = NP$.

However we can find a way to reduce satisfiability of an infinite set Σ of prop. sent. to all finite subsets of Σ . There later will be a more elementary proof of the compactness theorem, this proof is not part of the exam.

Theorem 1.2. Compactness theorem: Let Σ be an infinite set of prop. sent. such that

$$\forall \Sigma_0 \subseteq \Sigma, \Sigma_0 \text{ finite} \exists \text{ TA satisfying every member of } \Sigma_0 \quad (\text{finite satisfiability})$$

then there is a TA satisfying every member of Σ .

Proof. using topology: We have our infinite set of prop. sent. which satisfies above condition. One way to look at TA is as a sequence of 0 and 1. Let $\mathcal{A} = \{A_0, A_1, \dots\}$ be the set of all prop. atoms. We are going to identify the truth assignments on \mathcal{A} with elements in $\{0, 1\}^{\mathcal{A}} := \{f : \mathcal{A} \rightarrow \{0, 1\}\}$ (the set of all TAs). This is a topological space with product topology, on which the basic open sets (called cylinders) are: $U \subseteq \{0, 1\}^{\mathcal{A}}$ is a cylinder, such that $p_n(U) = \{0, 1\}$ for all but finite many n , where p_n is the n -th projection. This means U is a cylinder if the truth values of its elements are at finitely many places fixed, and are arbitrary on everything else.

Note: These basic open sets are also closed. The open sets are unions of basic open sets. The idea is to use Tychonoff's Theorem which tells us that $\{0, 1\}^{\mathcal{A}}$ is compact. i.e. the intersection of a family of closed subsets w/ the finite intersection property (FIP) is non-empty. Finite intersection property means the intersection of finitely many sets is non-empty.

For $\alpha \in \Sigma$ let $T_\alpha \subseteq \{0, 1\}^{\mathcal{A}}$ be the set of TA that satisfy α . This T_α is a finite union of cylinders, hence T_α is closed. For the family $\{T_\alpha : \alpha \in \Sigma\}$ of closed sets we have (FIP). Tychonoff tells us, that $\bigcup_{\alpha \in \Sigma} T_\alpha \neq \emptyset$ so there is a TA satisfying Σ . \square

For a list of tautologies: useful might be book p. 26-27

1.2 A PARSING ALGORITHM

To prove 1.1 We essentially need to show that we have enough parenthesis to make the reading of a prop. sent. unique. That is given a TA v there is at most one truth value we can assign to a prop. sent.

Lemma 2.1. Every prop. sent. has the same number of left and right parenthesis.

Proof. Let $M =$ set of prop. sent. w/ $\#$ left parenthesis = $\#$ right parenthesis and $PS =$ set of all prop. sent. We have $M \subseteq PS$. Since atoms have no parenthesis, they are in M . we just need to show that M is closed under the 5 construction operations.

$\mathcal{E}_\neg = (\neg\alpha) \dots$ \square

Lemma 2.2. *No proper initial segment of a prop. sent. is itself a prop. sent.*

Proof. Let $\alpha = \alpha_1\alpha_2\ldots\alpha_n$ be a prop. sent. By proper initial segment we understand $\beta = \alpha_1\ldots\alpha_i$ for $1 \leq i < n$. We will prove that every proper initial segment has an excess of left parenthesis, then we use the previous lemma. Let PS = set of all prop. sent. and PF = set of prop. sent. s.t. no proper initial segment has # left parenthesis = # right parenthesis, we will prove that these sets are the same.

Let $\alpha \in PF$. By induction on the fla. building operations

- Atoms: since the empty sequence is not a prop. sent. they have no proper initial segment.
- If the above is true for α, β then the proper initial segments of $(\neg\alpha)$ are of the form

$(\neg\alpha$
 $(\neg\alpha'$ where α' is a proper initial segment of α
 $($ or
 $(\neg$

Therefore \mathcal{E}_{\neg} preserves this property and under $\mathcal{E}_{\wedge}, \mathcal{E}_{\vee}, \mathcal{E}_{\rightarrow}, \mathcal{E}_{\leftrightarrow}$ this is also the case. □

Parsing algorithm

We now give a parsing algorithm procedure. For input we take some expression τ and the algorithm will determine if τ is a prop. sent. If so, it will generate a unique construction tree (in form of a rooted tree) for τ . (i.e. the construction tree gives us unique readability) That there is a unique way to perform the algorithm is implied by 2.2

0. create the root and label it τ
1. HALT if all leaves are labeled w/ prop. atom and return: “ τ is a prop. sent.”
2. select a leaf of the graph which is not labeled w/ prop. atom
3. if the first symbol of label under consideration is not a left parenthesis, then halt and return: “ τ is not a prop. sent.”
4. if the second symbol of the label is “ \neg ” then GOTO 6.
5. scan the expression from left to right
if we reach a proper initial segment of the form “ $(\beta$ ” where $\#lp(\beta) = \#rp(\beta)$ and β is followed by one of the five sentential connectives $\wedge, \vee, \rightarrow, \leftrightarrow$ and the remainder of the expression is of the form β' , where $\#lp(\beta') = \#rp(\beta')$

Then: create two child nodes (left, right) to the selected element and label them (left := β , right := β') GOTO 1.


Else: HALT and return “ τ is not a prop. sent.”

6. if the expression is of the form $(\neg\beta)$ where $\#lp(\beta) = \#rp(\beta)$

Then: construct one childnode and label it β and GOTO 1.

Else: HALT and return: “ τ is not a prop. sent.”

Example 1.2. The parsing algorithm applied to $((\neg(A_1 \rightarrow A_2)) \vee A_3)$ returns the following construction tree.



Correctness of the parsing algorithm

- The algorithm always halts, because a child's label is shorter than the label of a parent.
- If the algorithm halts with the conclusion that τ is a prop. sent. then we can prove inductively (starting from the leaves) that each label is a prop. sent
- Unique way to make choices in the algorithm: in particular β, β' in step 5. If there was a shorter choice for β it would be a proper initial segment of β but such prop. sent. cannot exist. (This also works under the assumption that a longer choice exists).
- rejections are made correctly

Back to proving the existence and uniqueness of $\bar{\nu}$ in 1.1. Let α be a prop. sent. of \bar{S} . We apply the parsing algorithm to α to get a unique construction tree. For the leaves, use ν to get the truth values then work our way up using the conditions (1-6) in 1.5.

1.3 INDUCTION AND RECURSION

Generalization of induction principle:

Let U be a set and $B \subseteq U$ our initial set. $\mathcal{F} = \{f, g\}$ a class of functions containing just f and g , where

$$f : U \times U \rightarrow U, \quad g : U \rightarrow U$$

We want to construct the smallest subset $C \subseteq U$ such that $B \subseteq C$ and C is closed under all elements of \mathcal{F} .

Definition 1.8. Closedness, Inductiveness: We say $S \subseteq U$ is

- **closed** under f and g iff for all $x, y \in S$ it holds $f(x, y) \in S$ and $g(x) \in S$
- **inductive** if $B \subseteq S$ and S is closed under \mathcal{F}

closed

inductive

One way is from the top down

$$C^* := \bigcap_{\substack{B \subseteq S \\ \text{inductive}}} S$$

Another is from bottom up: We call $C_1 := B$,

$$C_i := C_{i-1} \cup \{f(x, y) : x, y \in C_{i-1}\} \cup \{g(x) : x \in C_{i-1}\}$$

and $C_* := \bigcup_{n \geq 1} C_n$. Exercise: show that $C^* = C_* =: C$.

Example 1.3. 1. Let U be the set of all expressions, B the set of atoms and $\mathcal{F} = \{\mathcal{E}_{\Box} : \Box \in \{\neg, \wedge, \vee, \rightarrow, \leftrightarrow\}\}$. Then C would be the set of all propositional formulas.

2. Let U be \mathbb{R} , B the set containing 0 and $\mathcal{F} = \{S\}$, $S(x) = x + 1$. Then C would be the set of the natural numbers.

Induction principle

C generated from B by use of elements of \mathcal{F} if $S \subseteq C$ such that $B \subseteq S$ and S is closed under all elements of \mathcal{F} , then $S = C$

Proof. $S \subseteq C$ is clear. S is inductive, so $C \subseteq S$. □

Question: under what conditions do we get “generalized unique readability?” The goal would be to define a function on C recursively i.e. to have rules for computing $\bar{h}(x)$ for $x \in B$ with some rules of computing $\bar{h}(f(x, y))$ and $\bar{h}(g(x))$ from $\bar{h}(x)$ and $\bar{h}(y)$.

Example 1.4. Suppose that G is some additive group, generated from B (the set of generators), $h = B \rightarrow H$ where $(H, \cdot, ^{-1}, 1)$ a group. When is there an extension \bar{h} of h s.th. $\bar{h} : G \rightarrow H$ is a grouphomomorphism.

- $\bar{h}(0) = 1$
- $\bar{h}(a + b) = \bar{h}(a) \cdot \bar{h}(b)$
- $\bar{h}(-a) = \bar{h}(a)^{-1}$

This is not always possible. **Note:** that it is possible if G is generated freely by the elements of B and the set of atoms is independent (one element of B cannot be generated in finitely many steps by other elements of B).

Definition 1.9. Freely generated set: C is freely generated from B by f, g if

freely generated

- C is generated from B by f, g
- $f|_{C^2}$ and $g|_C$ are such that
 1. $f|_{C^2}$ and $g|_C$ are one-to-one (injective)
 2. $\text{ran}(f|_{C^2})$ and $\text{ran}(g|_C)$ and B are p.w. disjoint

Theorem 3.1. Recursion Theorem: $C \subseteq U$ freely generated from B by f, g and V a set and $h : B \rightarrow V, F : V^2 \rightarrow V, G : V \rightarrow V$ Then $\exists! \bar{h} : C \rightarrow V$ s.th.

- for all a in B it holds $\bar{h}(a) = h(a)$
- for all x, y in C it holds
 1. $\bar{h}(f(x, y)) = F(\bar{h}(x), \bar{h}(y))$
 2. $\bar{h}(g(x)) = G(\bar{h}(x))$

Note : if given conditions are satisfied then h extends uniquely to a homomorphism

$$(C, f, g) \rightarrow (V, F, G)$$

Before we proof the recursion theorem, we will show how unique readability easily follows from it.

Note : Recursion Theorem implies unique readability for propositional formulas. What we need to check is that the Assumptions of recursion theorem are satisfied.

Claim: The formula building operations are one-to-one.

proof of claim. \mathcal{F}_\vee is one to one, suppose $(\alpha \vee \beta) = (\delta \vee \gamma)$ then $\alpha \vee \beta = \delta \vee \gamma$ And α, δ are prop. formulas, so they equal to each other (else one is an initial segment of the other, hence not a prop. fla.) By the same argument we get β is equal to γ . \square

Claim: Disjointment of ranges

proof of claim. • if $(\alpha \vee \beta) = A$ then A starts with (which can not be the case

- if $(\alpha \vee \beta) = (\gamma \rightarrow \delta)$ then by the same argument α is γ but \vee and \rightarrow are different
 - if $(\alpha \vee \beta) = (\neg \gamma)$, then $\alpha \vee \beta = \neg \gamma$, so α would start with a \neg , -no
- For all other connectives the proof is similar. \square

Proof of the Rec Thm.

$v : C \rightarrow V$ is called acceptable if $\forall x, y \in C$

acceptable

1. if $x \in B \cap \text{dom}(v)$ then $v(x) = h(x)$
2. if $f(x, y) \in \text{dom}(v)$ then $x, y \in \text{dom}(v)$ and similarly for g
 - $v(f(x, y)) = F(v(x), v(y))$
 - $v(g(x)) = G(v(x))$

And when $U = \{\Gamma_v : v \text{ acceptable}\}$, we define $\bar{h} := \text{function w/ graph } \bigcup \Gamma_v$

Claim 1: \bar{h} is a function.

proof of claim.

$$S := \{x \in C : \exists \text{ at most one } y \text{ with } (x, y) \in \bigcup \Gamma_v\}$$

We want $S = C$, we have $S \subseteq C$, it is enough to show that S is inductive.

- $x \in B \cap \text{dom}(v)$ for some v acceptable.
then $v(x) = h(x)$ by 1. also $x \notin \text{ran}(f|_{C^2})$ and $x \notin \text{ran}(g|_C)$
- $x, y \in S$ We want $f(x, y), g(x) \in S$
there are v_1, v_2 acceptable s.t. $f(x, y) \in \text{dom}(v_1) \cap \text{dom}(v_2)$

⊗

Claim 2: \bar{h} is acceptable.

proof of claim. $\bar{h} : C \rightarrow V$ by definition. if $x \in B \cap \text{dom } \bar{h}$ then there is a v acceptable, s.t. $x \in \text{dom}(v)$ then $\bar{h}(x) = v(x) = h(x)$ if $f(x, y) \in \text{dom } \bar{h}$ then $f(x, y) \in \text{dom}(v)$ form some v acceptable. Hence $x, y \in \text{dom}(v)$ and therefore $x, y \in \text{dom}(\bar{h})$ and we have

$$\bar{h}(f(x, y)) = v(f(x, y)) = F(v(x), v(y)) = F(\bar{h}(x), \bar{h}(y))$$

⊗

Claim 3: The domain of \bar{h} equals C .

proof of claim. it is enough to show that the domain of \bar{h} is inductive. $B \subseteq \text{dom}(\bar{h})$ bc. $B \subseteq \text{dom}(h)$ where h is acceptable. Now we need to show closure under f, g . suppose $x', y' \in \text{dom}(\bar{h})$ then $x' \in \text{dom}(v_1)$ for some acceptable v_i lets assume $f(x', y') \notin \text{dom}(\bar{h})$ then we extend \bar{h} to a function with the same graph as \bar{h} . Then $\Gamma \cup \{(f(x', y'), F(x', y'))\}$ is the graph of an acceptable function.

⊗

Claim 4: \bar{h} is uniquely constructed

proof of claim. Suppose both $\bar{h}, \bar{\bar{h}}$ work, we show that $S = \{x \in C : \bar{h}(x) = \bar{\bar{h}}(x)\}$ is the whole set C . it is enough to show that S is inductive. Let $x \in B$ then $\bar{h}(x) = h(x) = \bar{\bar{h}}(x)$. Then for $x, y \in S$

$$\bar{h}(f(x, y)) = F(\bar{h}(x), \bar{h}(y)) = F(\bar{\bar{h}}(x), \bar{\bar{h}}(y)) = \bar{\bar{h}}(f(x, y))$$

$$\bar{h}(g(x)) = G(\bar{h}(x)) = G(\bar{\bar{h}}(x)) = \bar{\bar{h}}(g(x))$$

and $f(x, y), g(x) \in S$, therefore S is inductive.

⊗

1.4 SENTENTIAL CONNECTIVES

Definition 1.10. Tautological equivalence relation: For α, β prop. sent. we define $\alpha \sim \beta$ iff $\alpha \models \beta$ (alternative notation: \models). This defines an equivalent relation.

Example 1.5. $A \rightarrow B \models \neg A \vee B$

Note : A k -place boolean function is a function of the form $f : \{0, 1\}^k \rightarrow \{0, 1\}$ and we define $0, 1$ as the 0-place boolean functions.

If α is a prop. sent. then it determines a k -place boolean function, where k is the number of atoms, α is built up from. If α is $(A_1 \vee \neg A_2)$ then $B_\alpha : \{0, 1\}^2 \rightarrow \{0, 1\}$ and assign its values corresponding a truth value of α . That is for any TA $v : \{A_1, A_2\} \rightarrow \{0, 1\}$ we define $B_\alpha(v(A_1), v(A_2)) = \bar{v}(\alpha)$

Theorem 4.1. If α, β are prop. sent. with at most n prop. Atoms (combined), then

1. $\alpha \models \beta$ iff $\forall x \in \{0, 1\}^n$ it holds $B_\alpha(x) \leq B_\beta(x)$
2. $\alpha \models \beta$ iff $\forall x \in \{0, 1\}^n$ it holds $B_\alpha(x) = B_\beta(x)$
3. $\models \alpha$ iff $\forall x \in \{0, 1\}^n$ it holds $B_\alpha(x) = 1$

tautological
equivalence

\sim

\models

\models

Theorem 4.2. (Post): ¹ Let G be an n -ary boolean function for $n \geq 1$. Then there is a prop. sent. α such that $B_\alpha = G$. We say α realizes G .

n -ary boolean function

Proof. 1. if G is constantly equal to 0 then set α to $A_1 \wedge \neg A_1$.

2. Otherwise the set of inputs $\{\vec{x}_1, \vec{x}_2, \dots, \vec{x}_k\}$ for which $G(\vec{x}_i) = 1$ holds is not empty. We denote $\vec{x}_i = (x_{i1}, x_{i2}, \dots, x_{in})$ and define a matrix $(x_{ij})_{k \times n}$. We further set

$$\beta_{ij} = \begin{cases} A_j & \text{iff } x_{ij} = 1 \\ \neg A_j & \text{iff } x_{ij} = 0 \end{cases}$$

Example:

$$(x_{ij}) = \begin{pmatrix} 0 & 1 & 0 \\ 1 & 1 & 0 \end{pmatrix} \rightsquigarrow \begin{pmatrix} \neg A_1 & A_2 & \neg A_3 \\ A_1 & A_2 & \neg A_3 \end{pmatrix} = (\beta_{ij})$$

We define γ_i as $\beta_{i1} \wedge \beta_{i2} \wedge \dots \wedge \beta_{in}$ for $1 \leq i \leq k$ and α as $\gamma_1 \vee \gamma_2 \vee \dots \vee \gamma_k = \bigvee_{i=1}^k \gamma_i$. Then $B_\alpha = G$ is fulfilled.

□

Note : α as constructed in the proof is in the so-called Disjunctive normal form (DNF).

Corollary 4.2-A. Every prop. sent. is tautologically equivalent to a sentence in DNF

DNF
Disjunctive normal form

Corollary 4.2-B. $\{\neg, \wedge, \vee\}$ is a complete set of logical connectives, i.e. every prop. sent. is tautologically equivalent to a sentence built up from atoms and \neg, \wedge, \vee .

completeness
(connectives)

Theorem 4.3. Both $\{\neg, \wedge\}$ and $\{\neg, \vee\}$ are complete.

Proof. Its sufficient to show that every k -place boolean function is realisable by a prop. sent. built up using only \neg and \wedge . This is, because $\alpha \wedge \beta \models \neg(\neg\alpha \vee \neg\beta)$. We prove this by induction on the number of disjunctions of a prop. sent. α in DNF. Suppose the statement is true for $k \leq n$. For $n+1$ and $\alpha = \bigvee_{j=1}^{n+1} \gamma_j$ there exists an $\alpha' \models \bigvee_{j=1}^n \gamma_j$ and

$$\alpha = \bigvee_{j=1}^{n+1} \gamma_j \models \alpha' \vee \gamma_{n+1} \models \neg(\neg\alpha' \wedge \neg\gamma_{n+1})$$

□

Note : We used the observation that, if $\alpha \models \beta$ and we replace a subsequence of α by a so called tautological equivalence then the result is also tautologically equivalent to β .

Example 1.6. $\{\rightarrow, \wedge\}$ is not complete.: Let $\alpha \in PS$ built up from only \rightarrow, \wedge from the atoms A_1, \dots, A_n then we claim

$$A_1 \wedge A_2 \wedge \dots \wedge A_n \models \alpha$$

We can also say $\{\rightarrow, \wedge\}$ is not complete bc. $\neg A$ is not tautological equivalent to a sent. built up from \rightarrow, \wedge

Proof. Let $C := \{\alpha \in PS \text{ built up from } \rightarrow, \wedge \text{ and } A_1, \dots, A_n \text{ for which } \bigwedge_{i=1}^n A_i \models \alpha\}$ we want to show that $C = \{\alpha \in PS \text{ built up from } \rightarrow, \wedge \text{ and } A_1, \dots, A_n\}$

- We have $\{A_1, A_2, \dots, A_n\} \subseteq C$
- for $\alpha, \beta \in C$ it holds

- (1) $A_1 \wedge \dots \wedge A_n \models \alpha \rightarrow \beta$
- (2) $A_1 \wedge \dots \wedge A_n \models \alpha \wedge \beta$

Therefore C is closed under the fla. building operations and we have proven our claim.

□

Note : $\{\wedge, \vee, \rightarrow, \leftrightarrow\}$ is still not complete.

Note : The number of n -ary boolean functions existing is 2^{2^n} . We define a notation for $n = 0$: \perp (for TV = 0) and \top (for TV = 1). We can conclude that $\{\neg, \rightarrow\}$ and $\{\rightarrow, \perp\}$ are both complete, it holds $\neg A \models A \rightarrow \perp$.

Definition 1.11. Satisfiability: A set of prop. sent. Σ is called **satisfiable** if there exists a TA that satisfies every member of Σ .

satisfiable

¹Emil Post

1.5 COMPACTNESS THEOREM

Theorem 5.1. Compactness Theorem:

Σ is satisfiable iff every finite subset $\Sigma_0 \subseteq \Sigma$ is satisfiable. (i.e. Σ is finitely satisfiable)

finitely satisfiable

Proof. Let Σ be a finitely satisfiable set of prop. sent. Outline of the proof:

1. extend Σ to a maximal finitely satisfiable set Δ of prop. sent.
2. construct a truth assignment using Δ
1. Let $\alpha_1, \alpha_2, \dots$ be an enumeration of all prop. sent. and define Δ_n inductively by $\Delta_0 := \Sigma$

$$\Delta_{n+1} := \begin{cases} \Delta_n \cup \{\alpha_{n+1}\} & \text{if finitely satisfiable} \\ \Delta_n \cup \{\neg\alpha_{n+1}\} & \text{otherwise} \end{cases}$$

Claim: Δ_n is finitely satisfiable for each n

proof of claim. By regular induction over n . Δ_0 is finitely satisfiable. Let us assume Δ_n is finitely satisfiable. If $\Delta_{n+1} = \Delta_n \cup \{\alpha_{n+1}\}$ then we are finished. Otherwise let $\Delta' \subseteq \Delta_n$ be a finite subset that $\Delta' \cup \{\alpha_{n+1}\}$ is not satisfiable. It holds $\Delta' \models \neg\alpha_{n+1}$. Let us assume that $\Delta_n \cup \{\neg\alpha_{n+1}\}$ is not finitely satisfiable. Then there exists a finite subset $\Delta'' \subseteq \Delta_n$ such that $\Delta'' \cup \{\neg\alpha_{n+1}\}$ is not satisfiable. It therefore holds $\Delta'' \models \alpha_{n+1}$. But $\Delta' \cup \Delta''$ is a finite subset of Δ_n and by above observations $\Delta' \cup \Delta'' \models \alpha_{n+1}$ and $\Delta' \cup \Delta'' \models \neg\alpha_{n+1}$. A contradiction to the assumption that Δ_n is finitely satisfiable. \square

We set $\Delta := \bigcup_{i \in \mathbb{N}} \Delta_i$ and get

- (a) $\Sigma \subseteq \Delta$
- (b) (Maximality): for every prop. sent. α it holds $\alpha \in \Delta$ or $\neg\alpha \in \Delta$
- (c) (Satisfiability): Δ is finitely satisfiable. (For every finite subset there exists a Δ_n which is a superset.)
2. Let ν be a TA for the prop. atoms A_1, A_2, \dots such that $\nu(A) = 1$ iff $A \in \Delta$

Claim: For every prop. sent. φ it holds $\bar{\nu}(\varphi) = 1$ iff $\varphi \in \Delta$.

proof of claim. Let $S = \{\varphi \in PS \text{ s.t. } \bar{\nu}(\varphi) = 1 \text{ iff } \varphi \in \Delta\}$.

- $PS \supseteq S$ is clear.
- $PS \subseteq S$
 - (a) $\{A_1, A_2, \dots\} \subseteq S$ by definition of ν
 - (b) closure under \neg : Let $\varphi \in S$ then we get by maximality and satisfiability of Δ :

$$\begin{aligned} \bar{\nu}(\neg\varphi) &= 1 \\ \text{iff } \bar{\nu}(\varphi) &= 0 \\ \text{iff } \varphi &\notin \Delta \\ \text{iff } (\neg\varphi) &\in \Delta \end{aligned}$$

closure under \rightarrow : Let $\varphi_1, \varphi_2 \in S$ similarly

$$\begin{aligned} \bar{\nu}(\varphi_1 \rightarrow \varphi_2) &= 0 \\ \text{iff } \bar{\nu}(\varphi_1) &= 1 \text{ and } \bar{\nu}(\varphi_2) = 0 \\ \text{iff } \varphi_1 &\in \Delta \text{ and } \varphi_2 \notin \Delta \\ \text{iff } (\varphi_1 \rightarrow \varphi_2) &\notin \Delta \end{aligned}$$

The closure under the other fla. building operations are similar. \square

By this claim $\bar{\nu}$ satisfies Σ . \square

Corollary 5.1-A. If $\Sigma \models \tau$ then there exists a finite subset $\Sigma' \subseteq \Sigma$ s.t. $\Sigma' \models \tau$

Proof. Recall: $\Sigma \models \tau$ iff $\Sigma \cup \{\neg\tau\}$ is not satisfiable. Suppose $\Sigma \models \tau$ but no finite subset does. Then $\forall \Sigma' \subseteq \Sigma$ finite $\Sigma' \cup \{\neg\tau\}$ is satisfiable. By the compactness theorem $\Sigma \cup \{\neg\tau\}$ is satisfiable which is a contradiction to $\Sigma \models \tau$. \square

Note : 5.1 and 5.1-A are equivalent.

CHAPTER 2

Predicate - / first order logic

The definitions, lemmata, propositions and theorems as well as the notes in this chapter are sourced from [EE01, chapter 2].

Definition 2.1. A First order Language: consists of infinitely many distinct symbols such that no symbol is a proper initial segment of another symbol. The symbols are divided into 2 groups:

1. logical symbols

(These elements have a fixed meaning and the equivalence symbol $=$ is optional)

$(,), \neg, \rightarrow, v_1, v_2, \dots, =$

logical symbols

2. parameters

(a) quantifier symbol: \forall (the range is subject of interpretation)

(b) predicate symbols: for every $n > 0$ we have a set of n -ary predicates P

(c) constant symbols: Some set of constants

(d) function symbols: for every $n > 0$ we have a set of n -ary function symbols

parameters

Note :

- The sets in Group 2, (b)-(d) can also be the empty set
- We could drop constants and instead introduce 0-ary function symbols
- to specify language we need to specify the parameters and say if $=$ is included
- In the book [EE01] they assume that some n -place predicate symbol is present for some n .

Example 2.1. • $\mathcal{L}_{\text{set}} = \{\in\}$, $=$ included and the binary predicate symbol \in "element in"

- $\mathcal{L}_{\text{arith}} = \{<, 0, S, E, +, \cdot\}$

$=$ included

$<$ is a binary rel. symbol

0 is a constant

S is a unary function symbol

E is a unary function symbol (exponentiation)

$+, \cdot$ are binary function symbols

- $\mathcal{L}_{\text{ring}} = \{=, +, \cdot, -, 0, 1\}$

$=$ included

$0, 1$ are constants

$-$ is a unary function symbol (additive inverse)

$+, \cdot$ binary function symbols

2.1 FORMULAS

Definition 2.2. Expression: An **expression** is any finite sequence of symbols. There exist two kinds of expressions that make sense “grammatically” The intuition should be:

expression

Terms: – points to an object

– they are built up from variables and constants using function symbols

Formulas: – They express assertions about objects,

– they are built up from atomic formulas

– atomic formulas these are built up from terms using predicate symbols and $=$, if included

Definition 2.3. Term Building Operations: For every $n > 0$ and for every n -place function symbol f , let \mathcal{F}_f be an n -place term building operation, that is $\mathcal{F}_f(t_1, \dots, t_n) := ft_1 \dots t_n$ (polish notation for $f(t_1, \dots, t_n)$). **Term**, the set of terms is defined as the set of expressions that are built up from variables and constants by applying the term building operations finitely many times.

Term

Example 2.2. Let $\mathcal{L} = \mathcal{L}_{arith}$, the set of terms will contain $0, v_{42}, S0, SSS0, Sv_1, +SOv_1$

Definition 2.4. Atomic formula: Any expression of the form

$= t_1 t_2$ or $P t_1, \dots, t_n$, where t_1, \dots, t_n are terms and P is an n -ary predicate symbol

is called **atomic formula**.

atomic formula

Note : Atomic formulas are not defined inductively.

Example 2.3. *cont.* $= v_1 v_{42}$ and $< S0SS0$ are atomic formulas, but $\neg = v_1 v_{42}$ is not.

Definition 2.5. Formulas: We define $\varepsilon_{\neg}, \varepsilon_{\rightarrow}, Q_i$ to be the fla. building operations, defined as follows $\varepsilon_{\neg}(\alpha) := (\neg\alpha)$, $\varepsilon_{\rightarrow} := (\alpha \rightarrow \beta)$ and $Q_i(\gamma) := \forall v_i \gamma$. The set of formulas is the set of expressions built up from atomic formulas by applying the fla. building operations finitely many times.

formula

Example 2.4. *cont.* $\forall v_1 (= Sv_1 0)$ is a formula we get by applying Q_1 on the atomic formula $= Sv_1 0$.

Free variables

Example 2.5. We introduce the \exists **quantifier** as an abbreviation: $\exists y \alpha$ means $\neg \forall y \neg \alpha$. “Every non-zero natural number is a successor” $\forall x (x \neq 0 \rightarrow \exists y S(y) = x)$ is different then “if a number is not 0, then it is a successor” $x \neq 0 \rightarrow \exists y S(y) = x$. We say, x occurs **bounded** in the first formula, for the latter x occurs free in the formula. If you have an expression without free variables, it is either true or false, on the other hand if a variable occurs free in a formula, the truth value of it depends on the variable itself.

 \exists quantifier

bounded variable

Definition 2.6. Free variables: Let x be a variable. We say “ x occurs **free** in φ ”, if

1. If φ is an atomic fla. then x occurs **free** in φ iff x occurs in φ
2. If $\varphi \equiv (\neg\alpha)$ then x occurs free in φ iff x occurs free in α
3. If $\varphi \equiv (\alpha \rightarrow \beta)$ then x occurs free in φ iff x occurs free in α or β
4. If $\varphi \equiv \forall v_i \alpha$ then x occurs free in φ iff x occurs free in α and $x \neq v_i$

A formula α is called a **sentence**, if no variable occurs free in α

sentence

Note : The above definition is well defined thanks to the recursion theorem.

To prove this, define the function h on the set of atoms: $h(\alpha) = \{v \in V : v \text{ occurs free in } \alpha\}$, which is the set of all variables v_i that occur free in α . we now want to extend h to \bar{h} on the set of all formulas. We observe

- $\bar{h}(\neg\alpha) = \bar{h}(\alpha)$
- $\bar{h}(\alpha \rightarrow \beta) = \bar{h}(\alpha) \cup \bar{h}(\beta)$
- $\bar{h}(Q_i(\alpha)) = \bar{h}(\alpha) \setminus \{v_i\}$

Then x occurs free in α iff $x \in \bar{h}(\alpha)$.

Note : From now on, we will now use $\neq, \wedge, \vee, \leftrightarrow, \exists v_i$ as abbreviations in our formulas (all can be expressed in terms of $\neg, \rightarrow, \forall_i$), so there is no ambiguity on the in the sense of the meaning of the formulas. Note: We will sometimes drop the $(,)$ and not always be using polish notation.

2.2 SEMANTICS OF FIRST ORDER LOGIC

The equivalent scheme to our TA in predicate logic. The meaning of formulas is given by **structures**, which also determine the scope of the quantifier \forall , the meaning of all parameters.

Definition 2.7. structure: A **structure** \mathcal{A} for a first order language \mathcal{L} is a non-empty set set A called **universe** or **underlying set** of \mathcal{A} together with an interpretation of each parameters of \mathcal{L} i.e.

- \forall ranges over the universe A
- for an n -ary pred. symbol $P \in \mathcal{L}$ its interpretation $P^{\mathcal{A}}$ is a subset of A^n
- for a constant $c \in \mathcal{L}$ its interpretation $c^{\mathcal{A}}$ is an element of A
- for an n -ary function symbol $f \in \mathcal{L}$ its interpretation $f^{\mathcal{A}}$ is a total function

$$f^{\mathcal{A}} : A^n \rightarrow A$$

Note : $A \neq \emptyset$, and all functions $f^{\mathcal{A}}$ are total.

Example 2.6. Let $\mathcal{L} = \{\in\}$ where \in is a binary relation " An example of an \mathcal{L} structure is $(\mathbb{N}, \in^{\mathbb{N}})$ where $\in^{\mathbb{N}} = \{(x, y) \in \mathbb{N}^2 : x < y\}$

Definition 2.8. Assignment: Let φ be a \mathcal{L} -fla. and \mathcal{A} a \mathcal{L} -structure. Let V be the set of all variables in \mathcal{L} . Any function $s : V \rightarrow A$ is called assignment. We define the extention \bar{s} of s to the set of all \mathcal{L} -terms by

- if $x \in V$ then $\bar{s}(x) := s(x)$
- if $c \in \mathcal{L}$ is a constant symbol, then $\bar{s}(c) := c^{\mathcal{A}}$
- for t_1, \dots, t_n \mathcal{L} -terms and $f \in \mathcal{L}$ an n -ary function symbol,

$$\bar{s}(ft_1 \dots t_n) := f^{\mathcal{A}}(\bar{s}(t_1), \dots, \bar{s}(t_n))$$

Note : in the previous definition point 3. for $n = 1$ yields a commutative diagram.

Theorem 2.1. For any given assignment s there exists a unique extention \bar{s} as in the previous definition.

Proof. will follow from recursion theorem and unique decomposition of terms. \square

structures

structure

interpretation

assignment

Satisfaction and Models

Definition 2.9. Satisfy: We define ' \mathcal{A} satisfies φ with s ' and write $\mathcal{A} \models \varphi [s]$ or $\models_{\mathcal{A}} \varphi [s]$, if (inductively over the complexity of the formula φ)

 $\models_{\mathcal{A}}$

1. if φ is atomic:
 - $\mathcal{A} \models = t_1, t_2 [s]$ iff $\bar{s}(t_1) = \bar{s}(t_2)$
 - $\mathcal{A} \models Pt_1, \dots, t_n [s]$ iff $(\bar{s}(t_1), \dots, \bar{s}(t_n)) \in P^{\mathcal{A}}$
2. suppose $\mathcal{A} \models \varphi [s]$ and $\mathcal{A} \models \psi [s]$ are defined, then
 - $\mathcal{A} \models \neg \varphi [s]$ iff $\mathcal{A} \not\models \varphi [s]$
 - $\mathcal{A} \models \varphi \rightarrow \psi [s]$ iff $\mathcal{A} \models \varphi [s]$ or $\mathcal{A} \not\models \psi [s]$
 - $\mathcal{A} \models \forall x \varphi [s]$ iff for all $a \in A$ $\mathcal{A} \models \varphi [s(x|a)]$ where

$$s(x|a)(v) := \begin{cases} s(v) & \text{if } v \neq x \\ a & \text{if } v = x \end{cases}$$

Example 2.7. $\mathcal{L} = \{\forall, \leq, S, 0\}$ an example of an \mathcal{L} -structure $\mathcal{N} = (\mathbb{N}, \leq^{\mathcal{N}}, S^{\mathcal{N}}, 0^{\mathcal{N}})$. Together with a specified assignment $s : v_n \mapsto n - 1$

- $s(v_1) = 0$
- $\bar{s}(0) = 0^{\mathcal{N}}$ (a constant is always mapped to its realisation, the interpretation of constant 0 in the structure \mathcal{N})
- $\bar{s}(Sv_1) = S^{\mathcal{N}}(\bar{s}(v_1)) = S^{\mathcal{N}}(0) = 1$
- $\mathcal{N} \models \forall v_1 (S(v_1) \neq v_1) [s]$
 iff for all $a \in \mathbb{N}$ we have that $\mathcal{N} \models (S(v_1) \neq v_1)[s(v_1|a)]$
 iff ...
 iff for all $a \in \mathbb{N}$ we have $S^{\mathcal{A}}(a) \neq a$, which is true in our structure of the natural numbers.
- Is it true in \mathcal{N} that $\mathcal{N} \models S(0) \leq S(v_1) [s]$? Yes because

$$\begin{aligned} \mathcal{N} \models S(0) \leq S(v_1) [s] \\ \text{iff } 1 \leq 1 \end{aligned}$$

To know wheter $\mathcal{A} \models \varphi [s]$ it suffices to know where s maps the variables that occur free in φ . This results from the Coincidence Lemma 2.2

Theorem 2.2. Coincidence Lemma: Suppose $s_1, s_2 : V \rightarrow A$ agree on all variables that occur free in φ then

$$\mathcal{A} \models \varphi [s_1] \text{ iff } \mathcal{A} \models \varphi [s_2]$$

Proof. By complexity of φ

1. if φ is Pt_1, \dots, t_n note: any var that occur in φ occur free in φ , so s_1, s_2 agree on all variables that occur in the terms t_1, \dots, t_n .
 So we Claim: for t a term, s_1, s_2 assignments that agree on all variables of t then $\bar{s}_1(t) = \bar{s}_2(t)$

proof of claim. By complexity of t

- $t = v_m$ then $\bar{s}_1(t) = s_1(v_m) = s_2(v_m) = \bar{s}_2(t)$
- $t = c$ then $\bar{s}_1(t) = c^{\mathcal{A}} = \bar{s}_2(t)$
- $t = ft_1 \dots t_n$ inductively, assume $\bar{s}_1(t_i) = \bar{s}_2(t_i)$ for all $1 \leq i \leq n$ then TODO

⊠

2. if φ is $= t_1, t_2$ is similar
3. if φ is $\neg \alpha$ then $\mathcal{A} \models \neg \alpha [s_1]$ iff $\mathcal{A} \not\models \alpha [s_1]$ iff $\mathcal{A} \not\models \alpha [s_2]$ iff $\mathcal{A} \models \neg \alpha [s_2]$

4. if φ is $\alpha \rightarrow \beta$ then $\mathcal{A} \models \alpha \rightarrow \beta [s_1]$ iff .. or .. iff for s_2 iff ... or ..
5. if φ is $\forall x \alpha$ then the assumption is that s_1, s_2 .. the free variables of α are the free variables of φ except for x . but because $s_1(x|a) = s_2(x|a)$ they both agree on all free variables of α .

$$\begin{aligned} \mathcal{A} \models \forall x \varphi [s_1] & \text{ iff for all } a \in A \mathcal{A} \models \varphi [s_1(x|a)] \\ & \text{ iff for all } a \in A \mathcal{A} \models \varphi [s_2(x|a)] \\ & \text{ iff } \mathcal{A} \models \forall x \varphi [s_2] \end{aligned}$$

□

Notation: if all free variables of φ are among v_1, \dots, v_n and the assignment $s(v_i) = a_i$ for all $1 \leq i \leq n$ we write

$$\mathcal{A} \models \varphi[a_1, a_2, \dots, a_n] := \mathcal{A} \models \varphi [s]$$

Note : If σ is a sentence then we can not have $\mathcal{A} \models \sigma$ and $\mathcal{A} \not\models \sigma$ because $A \neq \emptyset$.

Corollary 2.2-A. If σ is a sentence then either $\mathcal{A} \models \sigma [s]$ for all $s : V \rightarrow A$ or $\mathcal{A} \not\models \sigma [s]$ for all $s : V \rightarrow A$.

Notation: $\mathcal{A} \models \sigma$ and we say “ σ is true in \mathcal{A} ” or “ \mathcal{A} is a model of σ ” or “ σ holds in \mathcal{A} ”.

Definition 2.10. Model: \mathcal{A} is said to be “a model of a set of sentences Σ ” $\mathcal{A} \models \Sigma$, if for every sentence $\sigma \in \Sigma$ it holds $\mathcal{A} \models \sigma$

Example 2.8. $\mathcal{L} = \{0, 1, +, -, \cdot\}$ A Structure could be $\mathcal{R} = (\mathbb{R}, 0, 1, +, -, \cdot)$ or $\mathcal{C} = (\mathbb{C}, 0, 1, +, -, \cdot)$ then the sentence $\sigma : \exists x(x \cdot x = -1)$ then $\mathcal{R} \not\models \sigma$ but $\mathcal{C} \models \sigma$

Note : $\wedge, \vee, \leftrightarrow, \exists$ work as expected. That is $\mathcal{A} \models (\alpha \wedge \beta) [s]$ iff $\mathcal{A} \models \alpha [s]$ and $\mathcal{A} \models \beta [s]$
 $\mathcal{A} \models (\alpha \vee \beta) [s]$ iff $\mathcal{A} \models \alpha [s]$ or $\mathcal{A} \models \beta [s]$ $\mathcal{A} \models \exists x \alpha [s]$ iff $\mathcal{A} \models \neg \forall x \neg \alpha [s]$
 iff $\mathcal{A} \not\models \forall x \neg \alpha [s]$
 iff it is not true that for all $a \in A \mathcal{A} \models \neg \alpha [s(x|a)]$
 iff there is $a \in A$ such that $\mathcal{A} \models \alpha [s(x|a)]$

Logical implication

Let Γ be a set of \mathcal{L} -formulas, φ a \mathcal{L} -formula.

Definition 2.11. Logical implication: We say “ Γ logically implies φ ” $\Gamma \models \varphi$, if for every \mathcal{L} -structure \mathcal{A} and for every assignment $s : V \rightarrow A$

$$\text{if } \mathcal{A} \models \gamma [s] \text{ for every } \gamma \in \Gamma, \text{ then } \mathcal{A} \models \varphi [s]$$

Definition 2.12. Logical equivalence: Two formulas φ, ψ are called logically equivalent, if $\varphi \models \psi$ and $\psi \models \varphi$.

Definition 2.13. Valid: A formula φ is called valid, if $\models \varphi$ i.e. $\emptyset \models \varphi$ i.e. for every \mathcal{L} -structure \mathcal{A} and every $s : V \rightarrow A$ it is $\mathcal{A} \models \varphi [s]$

Example 2.9. 1. $\forall x_1 P x_1 \models P x_2$
 Suppose $\mathcal{A} \models \forall x_1 P x_1 [s]$. then for all $a \in A$ it is $\mathcal{A} \models P x_1 [s(x_1|a)]$ in particular, $a \in P^{\mathcal{A}}$ for $a = s(x_2)$

2. $\forall P x_2 \not\models \forall x_1 P x_1$
 We need a counterexample to $\forall P x_2 \models \forall x_1 P x_1$. Let $A = \{a_1, a_2\}$ $s(x_2) = a_1$ and $P^{\mathcal{A}} = \{a_1\}$ then $\mathcal{A} \models P x_2 [s]$.

3. Is the following valid? $\models \exists x (P x \rightarrow \forall y P y)$ yes

4. $\Gamma, \alpha \models \varphi$ iff $\Gamma \models \alpha \rightarrow \varphi$. (on next problem set, quite important)

2.3 DEFINABILITY IN A STRUCTURE

By choosing a language \mathcal{L} , we specify which constant, function and relation symbols we can use when constructing formulas. If we then fix a structure \mathcal{A} , some n -ary relations that may are not in our structure, can still be expressed by a formula φ . For example given the structure $\mathcal{N} = (\mathbb{N}; 0, +)$ with equality and the usual addition on \mathbb{N} , we can view the set

$$P_{<} := \{(n, m) \in \mathbb{N} : \varphi(n, m)\}, \quad \varphi(n, m) \equiv \exists x (\neg(x = 0) \wedge n + x = m)$$

That gives us the usual interpretation of $<$ in the natural numbers, namely $n < m$ iff $\varphi(n, m)$ iff $(n, m) \in P_{<}$.

Definition 2.14. definability in a structure: We say that a general n -ary relation P on A (we will just call it P , it does not have to be in the language) is definable in \mathcal{A} , if there is a \mathcal{L} -formula φ with free variables among $\{v_1, \dots, v_n\}$ such that

$$P = \{(a_1, \dots, a_n) \in A^n : \mathcal{A} \models \varphi[a_1, \dots, a_n]\}$$

We also say that φ defines P in the structure \mathcal{A} .

TODO definability by points: We say that a general n -ary relation P on A is definable by points in \mathcal{A} , if there is a \mathcal{L} -formula φ with free variables among $\{v_1, \dots, v_{n+m}\}$ and $b_1, \dots, b_m \in A$ such that

$$P = \{(a_1, \dots, a_n) \in A^n : \mathcal{A} \models \varphi[a_1, \dots, a_n, b_1, \dots, b_m]\}$$

definable by points

Example 2.10. 1. $x = x$ would define the entire universe.

2. $\neg x = x$ would define the empty set.

Example 2.11. 1. TODO

2. $\mathcal{R} = (\mathbb{R}, 0, 1, +, -, \cdot)$ Q: is $[0, \infty)$ definable in \mathcal{R} Yes because $\exists y (y \cdot y = x)$ Indeed we can even define the \leq relation on \mathbb{R}^2 by $x \leq z \Leftrightarrow \exists y (x + y \cdot y = z)$

Definition 2.15. definability of classes of structures: Let Σ be a set of sentences. τ a sentence. The class of models of Σ is the class $\text{Mod } \Sigma = \{\mathcal{A} : \mathcal{A} \models \Sigma\}$. Let K be a class of structures. We are going to call K an elementary class (EC) if there is a single sentence τ such that $K = \text{Mod } \tau$. K is called an elementary class in the wider sense (EC_Δ) if there is a set of sentences Σ such that $K = \text{Mod } \Sigma$

Example 2.12. Let $\mathcal{L} = \{0, 1, +, \cdot\}$ and τ be a sentence that expresses the field axioms (the unary inverse functions are not in our language but are definable.) $\text{Mod } \tau$ is the class of all the fields, which is EC. The class of all fields of characteristic 0: Let $\sigma_p : \neg(1 + \dots + 1 = 0)$ and $\Sigma = \{\tau\} \cup \{\sigma_p : p \in \mathbb{P}\}$ yields that $\text{Mod } \Sigma$ is the class of fields with characteristic 0, therefore it is EC_Δ , we will later see that it is not EC.

Example 2.13. Let E be a binary relation, $\mathcal{L} = \{E\}$ then a graph is a realisation $\mathcal{G} = (V, E^\mathcal{G})$ such that $V \neq \emptyset$, $E^\mathcal{G}$ is irreflexive and symmetric. By definition the universe is not empty, we still have to check irreflexive and symmetric.

- irreflexive: $\forall x (\neg xEx)$
- symmetric: $\forall x \forall y (xEy \rightarrow yEx)$

We take τ to be $\forall x \forall y ((\neg xEx) \wedge (xEy \rightarrow yEx))$ Then $\text{Mod } \tau$ is the class of all graphs and is therefore EC Note: the class of all finite graphs is neither EC nor EC_Δ . proof later.

Until now, we have no notion that tells us when two graphs are the same or at least similar.

2.4 HOMOMORPHISMS OF STRUCTURES

Definition 2.16. Homomorphism: Suppose that \mathcal{A}, \mathcal{B} are two \mathcal{L} -structures. then a Homomorphism of \mathcal{A} into \mathcal{B} is a map $h : A \rightarrow B$ satisfying the below conditions

- for every n -ary predicate $P \in \mathcal{L}$ it is

$$(a_1, \dots, a_n) \in P^{\mathcal{A}} \text{ iff } (h(a_1), \dots, h(a_n)) \in P^{\mathcal{B}}$$

(this def. a strong Homomorphism, other textbooks maybe only require “ \rightarrow ” direction)

- for every n -ary function $f \in \mathcal{L}$ and for all $\underline{a} = (a_1, \dots, a_n) \in A^n$ it holds

$$h(f^{\mathcal{A}}(\underline{a})) = f^{\mathcal{B}}(h(a_1), \dots, h(a_n))$$

- for every constant symbol $c \in \mathcal{L}$ it is $h(c^{\mathcal{A}}) = c^{\mathcal{B}}$
(could also skip this if we consider constants as 0-ary functions)

Note : Intuativly a Homomorphism of \mathcal{A} into \mathcal{B} is a map $A \rightarrow B$ that preserve all function and relation symbols in some sense, (in general however not the definable relations)

Definition 2.17. Isomorphism:

- $h : A \rightarrow B$ is called isomorphism of \mathcal{A} into \mathcal{B} if h is a homomorphism and injective (in other textbooks: an isomorphic embedding of \mathcal{A} into \mathcal{B})
- $h : A \rightarrow B$ is called isomorphism of \mathcal{A} onto \mathcal{B} , if h is a bijective homomorphism $A \rightarrow B$
- \mathcal{A} and \mathcal{B} are called isomorphic if there is an isomorphism of \mathcal{A} onto \mathcal{B}

Note :

Example 2.14. $\mathcal{L} = \{+, \cdot\}$ $\mathcal{N} = (\mathbb{N}, +^{\mathbb{N}}, \cdot^{\mathbb{N}})$ and $\mathcal{B} = (B, +^{\mathcal{B}}, \cdot^{\mathcal{B}})$ where $B = \{0, 1\}$ and

$+^{\mathcal{B}}$	e	0
e	e	0
0	e	e

let $h : \mathbb{N} \rightarrow B$ a Homomorphism? $h(n) = \begin{cases} e & \text{if } n \text{ is even} \\ 0 & \text{else} \end{cases}$ need at first that

$h(m+n) = h(m) +^{\mathcal{B}} h(n)$ and $h(m \cdot n) = h(m) \cdot^{\mathcal{B}} h(n)$. it is indeed a Homomorphism.

Definition 2.18. Substructure: Suppose we have two \mathcal{L} structures with $A \subseteq B$. We call \mathcal{A} a substructure of \mathcal{B} (notation: $\mathcal{A} \subseteq \mathcal{B}$ or we might say \mathcal{B} is an extension of \mathcal{A}), if

- for every n -ary relation $P^{\mathcal{A}} = P^{\mathcal{B}}|_A$
- for every n -ary function $f^{\mathcal{A}} = f^{\mathcal{B}}|_A$
- for every constant symbol c in \mathcal{L} it is $c^{\mathcal{A}} = c^{\mathcal{B}}$

Example 2.15. $\mathcal{L} = \{\leq\}$ then $\mathcal{N} = (\mathbb{N}, \leq)$ and $\mathcal{P} = (\mathbb{N}^+, \leq^{\mathcal{P}})$ where $\leq^{\mathcal{P}}$ is the restriction of \leq to the positive natural numbers. $\mathcal{P} \subseteq \mathcal{N}$ and there exists a isomorphic embedding $id : \mathbb{N}^+ \rightarrow \mathbb{N}$ from \mathcal{P} into \mathcal{N} They are even isomorphic ($h : \mathbb{N} \rightarrow \mathbb{N}^+, h(n) = n+1$) so in fact $\mathcal{P} \cong \mathcal{N}$.

Example 2.16. $(\mathbb{Q}, +) \subseteq (\mathbb{C}, +)$

Note : If $\mathcal{A} \subseteq \mathcal{B}$ then in particular \mathcal{A} is closed under all constant and functions in \mathcal{B} . So suppose that \mathcal{B} is a substructure and $A \subseteq B$ and $A \neq \emptyset$ and A is closed under $f^{\mathcal{B}}$, $c^{\mathcal{B}}$. Can then A be made into a substructure \mathcal{A} of \mathcal{B} . $f^{\mathcal{A}}$ would be the restriction of $f^{\mathcal{B}}$ to A^n , constants $c^{\mathcal{A}} = c^{\mathcal{B}}$ and if $P \in \mathcal{L}$ is an n -ary predicate then $P^{\mathcal{A}}$ should be $P^{\mathcal{B}} \cap A^n$. If \mathcal{L} has no const. or function symbols then any subset can be made into a substructure of a structure on \mathcal{L} .

Our next question will be: what is the relation of the above notions with truth and satisfiability. The answer will be given by the so called Homomorphism theorem.

Theorem 4.1. Homomorphism Theorem: Let $h : A \rightarrow B$ be a homomorphism of \mathcal{A} into \mathcal{B} and $s : V \rightarrow A$ an assignment then $h \circ s : V \rightarrow B$ is an assignment and it holds

1. for all terms t it is $h(\overline{s(t)}) = \overline{(h \circ s)(t)}$
2. for all quantifier free formulas φ that do not contain $=$ we have

$$\mathcal{A} \models \varphi[s] \text{ iff } \mathcal{B} \models \varphi[h \circ s]$$

3. if h is additionally injective then we can drop the requirement “no $=$ ” in (2)
4. if h is homomorphism of \mathcal{A} onto \mathcal{B} then we can drop the requirement “q.f.” in (2)

Proof. 1. problem set

2.
 - case $\varphi \equiv Pt$ then $\mathcal{A} \models Pt[s]$ iff $\overline{s(t)} \in P^{\mathcal{A}}$ iff $h(\overline{s(t)}) \in P^{\mathcal{B}}$ iff $\overline{(h \circ s)(t)} \in P^{\mathcal{B}}$ iff $\mathcal{B} \models Pt[h \circ s]$
 - case $\varphi \equiv \neg\psi$ $\mathcal{A} \models \neg\psi[s]$ iff $\mathcal{A} \not\models \psi[s]$ iff $\mathcal{A} \not\models \psi[s]$ iff
 - case $\varphi \equiv \psi \rightarrow \alpha$
3. $\mathcal{A} \models t_1 t_2[s]$ iff $\overline{s(t_1)} = \overline{s(t_2)}$ iff $h(\overline{s(t_1)}) = h(\overline{s(t_2)})$ iff (by (a)) $\overline{(h \circ s)(t_1)} = \overline{(h \circ s)(t_2)}$ iff $\mathcal{B} \models t_1 t_2[h \circ s]$
4. $\varphi \forall s : V \rightarrow A \mathcal{A} \models \varphi[s]$ iff $\mathcal{B} \models \varphi[h \circ s]$, want $\mathcal{A} \models \forall x \varphi[s]$ iff $\mathcal{B} \models \forall x \varphi[h \circ s]$ 1. $\mathcal{B} \models \forall x \varphi[h \circ s]$ iff for all $s : V \rightarrow A$, $a \in A$ (req. surjectivity) it is $\mathcal{B} \models \varphi[(h \circ s)(x|h(a))]$ iff $\mathcal{B} \models \varphi[h \circ (s(x|a))]$ iff (inductive assumption) $\mathcal{A} \models \varphi[s(x|a)]$ because a was arbitrary it is $\mathcal{A} \models \forall x \varphi[s]$ 2. Suppose $\mathcal{B} \not\models \forall x \varphi[h \circ s]$ then there exists a $b \in B$ such that $\mathcal{B} \models \neg \varphi[(h \circ s)(x|b)]$ by surjectivity we can find $a \in A$ such that $h(a) = b$ and it is $\mathcal{B} \models \neg \varphi[(h \circ s)(x|h(a))]$ By the inductive assumption $\mathcal{A} \models \neg \varphi[s(x|a)]$ and $\mathcal{A} \not\models \forall x \varphi[s]$

□

Definition 2.19. elementarily equivalent: \mathcal{A} and \mathcal{B} are called elementarily equivalent (notation: $\mathcal{A} \equiv \mathcal{B}$) if \mathcal{A} and \mathcal{B} satisfy the same sentences.

Note : $\mathcal{A} \cong \mathcal{B}$ implies $\mathcal{A} \equiv \mathcal{B}$ The converse is not true. For instance DLO (dense linear order) w/o endpoints is complete, so two structures on DLO are equivalent $(\mathbb{Q}, <) \equiv (\mathbb{R}, <)$ but they are not isomorphic because the universes have different cardinality.

Example 2.17. $\mathcal{N} = (\mathbb{N}, \leq)$ and $\mathcal{P} = (\mathbb{N}^{>0}, \leq)$ $h : n \mapsto n - 1 : \mathcal{P} \rightarrow \mathcal{N}$ isomorphism. so i.p. $\mathcal{N} \equiv \mathcal{P}$. but $id : \mathcal{P} \rightarrow \mathcal{N}$ is only isomorphic embedding, so for example $\mathcal{P} \models \forall y(x \neq y \rightarrow x \leq y) \llbracket 1 \rrbracket$ and $\mathcal{N} \not\models \forall y(x \neq y \rightarrow x \leq y) \llbracket 1 \rrbracket$, but $\mathcal{N} \models \forall y(x \neq y \rightarrow x \leq y) \llbracket h(1) \rrbracket$

Definition 2.20. Automorphism: An automorphism is an isomorphism of the form $h : A \rightarrow A$ from \mathcal{A} onto \mathcal{A}

Note : Every structure has a trivial automorphism $id : A \rightarrow A$

Definition 2.21. Rigid: \mathcal{A} is called rigid, if the only automorphism on \mathcal{A} is the trivial automorphism.

Example 2.18. If every element is definable then the structure is rigid. For example $(\mathbb{N}, 0, S)$ and $(\mathbb{N}, <)$ every element is definable, therefore the structures are rigid.

Corollary 4.1-A. Let h be an automorphism on \mathcal{A} , $R \subseteq A^n$ definable in \mathcal{A} then for all $a \in A^n$ $a \in R$ iff $(h(a_1), \dots, h(a_n)) \in R$. Suppose φ defines R in \mathcal{A} , then

$$\mathcal{A} \models \varphi[a_1, \dots, a_n] \quad \text{iff} \quad \mathcal{A} \models \varphi[h(a_1), \dots, h(a_n)]$$

Note : Corollary 4.1-A can be used to show that some $R \subseteq A^n$ is not definable in \mathcal{A}

Example 2.19. $\mathcal{R} = (\mathbb{R}, <)$ then \mathbb{N} is not definable in \mathcal{R} . What do automorphisms of \mathcal{R} look right? $h : \mathbb{R} \rightarrow \mathbb{R}$ is a bijection and $x < y$ iff $h(x) < h(y)$ so h is strictly increasing. for example $x \mapsto x + \frac{1}{2}$ or $x \mapsto x^3$.

2.5 UNIQUE READABILITY FOR TERMS

Definition 2.22. : We define K on symbols from which terms are built up (variables, constants, function symbols). $K(s) = 1 - n$ where s is a symbol and n is the number of terms that need to follow s in order to obtain a term. $K(x) = 1 = K(c)$ and $K(f) = 1 - n$ where f is an n -ary function symbol. We now extend K to the set of all expressions which are built up from above symbols (variables, constants, function symbols): $K(s_1, \dots, s_n) = K(s_1) + \dots + K(s_n)$ (unique because no symbol is a finite sequence of other symbols)

Lemma 5.1. *Let t be a term. Then $K(t) = 1$*

Proof. $K(x) = 1 = K(c)$ and $K(ft_1, \dots, t_n) = 1 - n + n = 1$ □

Definition 2.23. terminal segment: A terminal segment of string of symbols (s_1, \dots, s_n) is $(s_k, s_{k+1}, \dots, s_n)$ for some $1 \leq k \leq n$.

Lemma 5.2. *Any terminal segment of terms is a concatenation of one or more terms.*

Proof. True for variables and constants. $ft_1 \dots t_n$ the only non trivial case is $t'_k t_{k+1} \dots t_m$ where t_k is $t''_k t'_k$ □

Corollary 5.2-A. *If t_1 is a proper initial segment of a term t then $K(t_1) < 1$.*

Proof. let t be $t_1 t_2$ where t_1 is a proper initial segment then $K(t) = 1$ and $K(t_2) \geq 1$ therefore $K(t_1) \leq 0$ □

Proposition 5.3. Unique readability for terms: *The set of terms is freely generated from the set of variables (Var) and the set of constant symbols ($Const$) by the term building operations \mathcal{F}_f (for each function symbol f).*

Proof. By Definition 1.9:

- disjointment of ranges: Let f and g be two distinct function symbols then

$$\text{ran } \mathcal{F}_f \cap \text{ran } \mathcal{F}_g = \emptyset$$

$$\text{ran } \mathcal{F}_f \cap Var = \emptyset$$

$$\text{ran } \mathcal{F}_f \cap Const = \emptyset$$

- $\mathcal{F}_f|_{\text{terms}}$ are one-to-one (injective): assume $ft_1 \dots t_n = ft'_1 \dots t'_n$ and assume $t_1 \neq t'_1$ then one is a proper initial segment of the other. By Corollary 5.2-A its K -value has to be less than 1 so it is not a term. By induction we get $t_1 = t'_1, \dots, t_n = t'_n$.

□

Definition 2.24. : Extend K to the set of all expressions by (i) $K(()) = -1$ $K(()) = 1$ (ii) $K(\forall) = 1$ (iii) $K(\neg) = 0$ (iv) $K(\rightarrow) = -1$ (v) $K(P) = 1 - n$ for an n -ary rel. symb. P . (vi) $K(=) = -1$ (vii) $K(s_1 \dots s_n) = K(s_1) + K(s_n)$ for $s_i \in Terms \cup \{(), \forall, \neg, \rightarrow, P, \dots, =\}$

The idea is that K tells us the number of symbols that at least need to follow to obtain a formula.

Lemma 5.4. *for every formula φ it is $K(\varphi) = 1$*

Proof. induction on φ □

Lemma 5.5. *for every proper initial segment α' of a formula α we have $K(\alpha') < 1$*

Corollary 5.5-A. *No proper initial segment of a formula is a formula. The set of flas. is freely generated from the set of atomic flas. by operations $\mathcal{E}_{\neg}, \mathcal{E}_{\rightarrow}, Q_i$*

Proof. • \mathcal{E}_{\neg}, Q_i are one to one

- $\mathcal{E}_{\rightarrow}|_{\text{Flas.}}$ then itemwise and use of prev. lemmas
- p.w. disjointness of ranges

□

2.6 A PARSING ALGORITHM FOR FIRST ORDER LOGIC

2.7 DEDUCTIONS (FORMAL PROOFS)

Definition 2.25. Modus Ponens: We will use one rule of inference, Modus Ponens(MP). Our notation will be:

$$\frac{\alpha, \alpha \rightarrow \beta}{\beta}$$

And it reads as follows: "If α and $\alpha \rightarrow \beta$ then β ." This rule is the formalisation of the rather informal statement: "If we know a statement α is true, and this statement implies another statement β , then β must also be true."

Definition 2.26. Deduction: A formal proof (deduction) of a formula φ from a set of formulas Σ is a finite sequence of formulas $(\alpha_0, \alpha_1, \dots, \alpha_n)$ such that $\alpha_n = \varphi$ and for every $i < n$ α_i is either a logical axiom or $\alpha_i \in \Sigma$ or α_i is obtained from α_k and α_l where $0 \leq k, l < i$ by the use of MP, in particular $\alpha_k = \beta \rightarrow \alpha_i$ and $\alpha_l = \beta$. If a deduction of φ from Σ exists, we say " φ is deducible from Σ " or " φ is a theorem of Σ ".

Note : Deductions are not unique. However we do have an induction principle: If a set of formulas contains all logical axioms and all of Σ and is closed under MP, then it contains all theorems of Σ .

Logical axioms

Definition 2.27. Generalization: ψ is a generalization of φ if $\psi = \forall x_{i_1} \dots \forall x_{i_k} \varphi$

Definition 2.28. Logical axioms: Let x, y be variables and α, β formulas. then the logical axioms are generalizations of the following formulas:

1. tautologies
2. $\forall x \alpha \rightarrow \alpha_t^x$ where t is substitutable for x in α
3. $\forall x (\alpha \rightarrow \beta) \rightarrow (\forall x \alpha \rightarrow \forall x \beta)$
4. $\alpha \rightarrow \forall x \alpha$ where x does not occur free in α

if our language contains = then

1. $x = x$
2. $x = y \rightarrow (\alpha \rightarrow \alpha')$ where α' is obtained from α by replacing some of the occurrences of x with y .

Ad axiom group (2), Substitution:

Definition 2.29. Substitution: Let α, β be formulas, x a variable and t a term then α_t^x is expression obtained from α by substituting t for x . We define substitution inductive as follows:

1. if α is atomic then $\alpha_t^x = \alpha$ expression obtained from α by replacing all x 's by t 's
2. $(\neg \alpha)_t^x = \neg(\alpha_t^x)$
3. $(\alpha \rightarrow \beta)_t^x = (\alpha_t^x \rightarrow \beta_t^x)$
4. $(\forall y \alpha)_t^x = \begin{cases} \forall y (\alpha_t^x) & \text{iff } x \neq y \\ \forall x \alpha & \text{iff } x = y \end{cases}$

Example 2.20. • $\alpha_x^x = \alpha$

MP

- Let $\alpha = \neg\forall yx = y$ what is $\forall x\alpha \rightarrow \alpha_z^x$?

$$\forall x\neg\forall yx = y \rightsquigarrow \neg\forall yz = y$$

What is $\forall x\alpha \rightarrow \alpha_y^x$ $\forall x\neg\forall yx = y$ is true in all structures with a universe A with $|A| \geq 2$.

$$\forall x\neg\forall yx = y \rightsquigarrow \neg\forall yy = y$$

and $\neg\forall yy = y$ is an antitautology (it is always false).

•

So we have to define substitutable

Definition 2.30. substitutable: Let x be a variable, t a term. Then t is substitutable for x in α if

1. α atomic then t is SUB for x in α
2. then t is SUB for x in $\neg\alpha$ iff then t is SUB for x in α
3. then t is SUB for x in $\alpha \rightarrow \beta$ iff then t is SUB for x in α and β
4. then t is SUB for x in $\forall y\alpha$ iff either
 - x does not occur free in $\forall y\alpha$ or
 - y does not occur in t and t is SUB for x in α

Example 2.21. For instance the following is a logical axiom.

$$\forall x_3(\forall x_1(Ax_1 \rightarrow \forall x_2Ax_2) \rightarrow (Ax_2 \rightarrow \forall x_2Ax_2))$$

It is a generalization of $\forall x_1(Ax_1 \rightarrow \forall x_2Ax_2) \rightarrow (Ax_2 \rightarrow \forall x_2Ax_2)$ which is by point two a substitution with $\alpha = Ax_1 \rightarrow \forall x_2Ax_2$. Then $\alpha_{x_2}^{x_1} = Ax_2 \rightarrow \forall x_2Ax_2$ And x_2 is indeed substitutable for x_1 in α because it does not get bounded.

$$\forall x_1(\forall x_2Bx_1x_2 \rightarrow \forall x_2Bx_2x_2)$$

is a generalization of point (2), but x_2 is not substitutable for x_1 in α , therefore it is not a logical axiom.

Ad (1): tautologies

Definition 2.31. Tautologies of first order language: Tautologies are the formulas obtained from tautologies of propositional logic by replacing all propositional atoms by formulas of first order logic.

An alternative definition is: Divide all formulas of first order logic into two groups:

1. atomic formulas and generalizations of first order formulas (these are called prime formulas)
2. all other formulas i.e. of the form $\neg\alpha$ and $\alpha \rightarrow \beta$ (non-prime formulas)

So any first order formula is built up from the prime formulas using finitely many times the formula building operations. \mathcal{E}_\neg \mathcal{E}_\rightarrow We have unique readability because the set of formulas is freely generated.

Example 2.22.

$$\neg(\forall y(Px \rightarrow Py)) \rightarrow (Px \rightarrow \forall y\neg Py)$$

is built up from $\neg(\forall y(Px \rightarrow Py))$ and $Px \rightarrow \forall y\neg Py$. which itself $\forall y(Px \rightarrow Py)$ and Px and $\forall y\neg Py$ where they are all prime formulas.

Example 2.23. Is the following a tautology?

$$(\forall y(\neg Py) \rightarrow \neg Px) \rightarrow (Px \rightarrow \neg\forall y\neg Py)$$

We construct the construction tree into prime formulas and then assign truth values to them and evaluate the truth value of the whole formula. It is indeed a tautology.

Note :

- $\forall x(Px \rightarrow Px)$ is a prime formula which corresponds to a propositional atom, and therefore not a tautology. But it is a generalization of a tautology and therefore by (1) a logical axiom.
- $\forall xPx \rightarrow Px$ is not a tautology but is a logical axiom by group (2).

Note : $\Gamma \models_{\text{taut}} \varphi$ from propositional logic can be translated to first order logic.

Lemma 7.1. If $\Gamma \models_{\text{taut}} \varphi$ then $\Gamma \models \varphi$

Proof. Problem set. □

Note : The converse fails. For instance $\forall xPx \models Pc$. However Pc is a different propositional atom then $\forall xPx$ they have no connection between them when viewed in propositional logic.

We will prove $\Gamma \models \varphi$ iff $\Gamma \vdash \varphi$ (the first direction is completeness and the converse soundness.)

Theorem 7.2. $\Gamma \vdash \varphi$ iff $\Gamma \cup \Lambda \models_{\text{taut}} \varphi$

Proof. • Let $\Gamma \vdash \varphi$ and v be a truth assignment that satisfies every element in $\Gamma \cup \Lambda$. Induction on deduction of φ from Γ .

- if $\varphi \in \Gamma \cup \Lambda$ then we are done
- if φ is obtained from $\alpha, \alpha \rightarrow \varphi$ by MP then v satisfies α and $\alpha \rightarrow \varphi$
 $\{\alpha, \alpha \rightarrow \varphi\} \models_{\text{taut}} \varphi$

- Assume $\Gamma \cup \Lambda \models_{\text{taut}} \varphi$. Then by the compactness theorem for propositional logic there are $\gamma_1, \dots, \gamma_n \in \Gamma$ and $\lambda_1, \dots, \lambda_m \in \Lambda$ such that

$$\gamma_1 \rightarrow \gamma_2 \rightarrow \dots \rightarrow \gamma_n \rightarrow \lambda_1 \rightarrow \dots \rightarrow \lambda_m$$

is a tautology (always grouped to the left) because $\Gamma \cup \{\alpha\} \models_{\text{taut}} \beta$ iff $\Gamma \models_{\text{taut}} (\alpha \rightarrow \beta)$ □

2.8 GENERALIZATION AND DEDUCTION THEOREM

Note : Intuitively if Γ does not assume anything about x and Γ proves φ then Γ proves $\forall x\varphi$

Theorem 8.1. Generalization theorem: If $\Gamma \vdash \varphi$ and x does not occur free in Γ , then $\Gamma \vdash \forall x\varphi$

Proof. We use axiom group 4, $\alpha \rightarrow \forall x\alpha$ if x is not occurring free in α . Since x does not occur free in $\sigma \in \Gamma$, if $\varphi \in \text{Thm } \Gamma$ then $\forall x\varphi \in \text{Thm } \Gamma$. Induction principle: S the set of formulas. If $\Lambda \cup \Gamma \subseteq S$ and S is closed under MP then S contains $\text{Thm}(\Gamma)$. It is enough to show that $\{\varphi : \Gamma \vdash \forall x\varphi\}$ contains $\Gamma \cup \Lambda$. and is closed under MP.

1. if φ is a logical axiom then $\forall x\varphi$ is a generalization and therefore also a logical axiom, so $\Gamma \vdash \forall x\varphi$
2. Lets assume $\varphi \in \Gamma$. then x does not occur free in any element of Γ , then $\varphi \rightarrow \forall x\varphi$ is a logical axiom and $\Gamma \vdash \forall x\varphi$ by MP.
3. Closedness under MP. suppose φ is obtained from $\psi, \psi \rightarrow \varphi$ by MP. Then by induction hypothesis $\Gamma \vdash \forall x\psi$ and $\Gamma \vdash \forall x(\psi \rightarrow \varphi)$ Then $\forall x(\psi \rightarrow \varphi) \rightarrow (\forall x\psi \rightarrow \forall x\varphi)$ is a logical axiom in group 3. Then by MP $\Gamma \vdash \forall x\psi \rightarrow \forall x\varphi$
By MP again $\Gamma \vdash \forall x\varphi$

□

Note : Suppose x has free occurrence in Γ for example $Px \not\models \forall xPx$ so we can not have $Px \vdash \forall xPx$ (want \models iff \vdash)

Note : Proof of Generalization theorem can be used to obtain a deduction of $\forall x\varphi$ from Γ from a deduction of φ from Γ .

Lemma 8.2. Rule T: If $\Gamma \vdash \alpha_1, \Gamma \vdash \alpha_2, \dots, \Gamma \vdash \alpha_n$ and $\{\alpha_1, \alpha_2, \dots, \alpha_n\} \models_{\text{taut}} \beta$ then $\Gamma \vdash \beta$.

Proof. $\alpha_1 \rightarrow \alpha_2 \rightarrow \dots \rightarrow \alpha_n \rightarrow \beta$ is a logical axiom because it is a tautology. Apply MP n -times. □

Theorem 8.3. Deduction theorem: If $\Gamma \cup \{\gamma\} \vdash \varphi$ then $\Gamma \vdash (\gamma \rightarrow \varphi)$

Proof. Assume $\Gamma \cup \{\gamma\} \vdash \varphi$. $\Gamma \cup \{\gamma\} \vdash \varphi$ iff $\Gamma \cup \{\gamma\} \cup \Lambda \models_{\text{taut}} \varphi$
 iff $\Gamma \cup \Lambda \models_{\text{taut}} \gamma \rightarrow \varphi$ (exercise sheet 1, ex 7)
 iff $\Gamma \vdash (\gamma \rightarrow \varphi)$ □

Note : Deduction theorem is an equivalence. $\Gamma \vdash \gamma \rightarrow \varphi$ then $\Gamma \cup \{\gamma\} \vdash \varphi$. the statement follows by MP.

Corollary 8.3-A. (Contraposition): If $\Gamma \cup \{\varphi\} \vdash \neg\psi$ then $\Gamma \cup \{\psi\} \vdash \neg\varphi$

Proof. Suppose $\Gamma \cup \{\varphi\} \vdash \neg\psi$ then by deduction theorem $\Gamma \vdash \varphi \rightarrow \neg\psi$. We observe that $\{\varphi \rightarrow \neg\psi\} \models_{\text{taut}} \psi \rightarrow \neg\varphi$.

By rule T: $\Gamma \vdash \psi \rightarrow \neg\varphi$ and by the converse of the deduction theorem, by MP we have $\Gamma \cup \{\psi\} \vdash \neg\varphi$ □

Definition 2.32. Inconsistence: A set of fls. Γ is called inconsistent, if for some (equivalent to all) fla. β it is $\beta, \neg\beta \in \text{Thm } \Gamma$.

Note : If Γ is inconsistent, then for $\alpha \in \text{Thm } \Gamma$. Then $(\beta \rightarrow (\neg\beta \rightarrow \alpha))$ is a tautology. Use β from definition of inconsistency and use MP twice.

Corollary 8.3-B. (Reductio ad absurdum): If $\Gamma; \varphi$ inconsistent, then $\Gamma \vdash \neg\varphi$.

Proof. Suppose that $\Gamma; \varphi$ is inconsistent. then for any β $\Gamma; \varphi \vdash \beta$ and $\Gamma; \varphi \vdash \neg\beta$. By the deduction theorem $\Gamma \vdash \varphi \rightarrow \beta$ and $\Gamma \vdash \varphi \rightarrow \neg\beta$, therefore $\{\varphi \rightarrow \beta, \varphi \rightarrow \neg\beta\} \models_{\text{taut}} \neg\varphi$. By Rule T: $\Gamma \vdash \neg\varphi$. □

Note : strategies for finding deductions can be found in the textbook [EE01].

Theorem 8.4. Generalization on constants: Suppose $\Gamma \vdash \varphi$ and c is a constant symbol that does not occur in Γ . Then there is a variable y (y does not occur in φ) s.th. $\Gamma \vdash \forall y(\varphi)_y^c$. and moreover also there is a deduction of $\forall y(\varphi)_y^c$ in which c does not occur.

Proof. We will take a deduction $\langle \alpha_1, \dots, \alpha_n \rangle$ of φ from Γ . Pick the variable y as the first variable in any α_i for each i . **Claim:** $\langle (\alpha_1)_y^c, \dots, (\alpha_n)_y^c \rangle$ is a deduction of $(\varphi)_y^c$ from Γ .
proof of claim. We need to verify that every member $(\alpha)_y^c$ is actually provable from Γ .

- if $\alpha_k \in \Gamma$ then c does not occur in α_k then $(\alpha)_y^c = \alpha_k$
- if $\alpha_k \in \Lambda$ then $(\alpha_k)_y^c$ is also a logical axiom.
- lets say α_k was obtained by $\alpha_i, \alpha_i \rightarrow \alpha_k$ $i < k$ by MP. Now take $(\alpha_i \rightarrow \alpha_k)_y^c = (\alpha_i)_y^c \rightarrow (\alpha_k)_y^c$. (induction hypothesis) $(\alpha_k)_y^c$ is obtained from $(\alpha_i)_y^c$ nad $(\alpha_i \rightarrow \alpha_k)_y^c$. by MP.

⊠

Because formal proofs are finite, there is a $\Gamma_0 \subseteq \Gamma$ finite such that Γ_0 consists of the elements of Γ used in our deduction $\langle (\alpha_1)_y^c, \dots, (\alpha_n)_y^c \rangle$ (is therefore deduction of $(\varphi)_y^c$ from Γ_0). And because we assumed that y does not occur in Γ_0 , so we can use the generalization theorem on $\Gamma_0 \vdash (\varphi)_y^c$ and yield $\Gamma_0 \vdash \forall y(\varphi)_y^c$ □

Alphabetic Variants

We will formalize and proof the statement "You can always rename your bound variables". Why is that important? Suppose we want to proof that it is provable that $\forall x \forall y P(x, y) \rightarrow \forall y P(y, y)$. If we want to use a logical axiom of group 2, we would need to check if y is actually SUB for x . We obviously do not have that because y would get bounded. $\vdash \forall x \forall y P(x, y) \rightarrow \forall x \forall z P(x, z)$
 $\vdash \forall x \forall z P(x, z) \rightarrow \forall y P(y, y)$

Theorem 8.5. Existence of alphabetic variants: Let φ be a fla., x a variable, t a term. Then there exists a fla. φ' such that φ differs from φ' only in the choice of names of the bound variables. And

1. $\varphi' \vdash \varphi$ as well as $\varphi \vdash \varphi'$
2. t is SUB for x in φ'

Proof. Define φ' inductively on complexity of φ .

- if φ is atomic, then $\varphi' = \varphi$
- $(\neg\varphi)' = \neg\varphi'$
 1. $\varphi' \vdash \varphi$ and $\varphi \vdash \varphi'$, we want: $\neg\varphi' \vdash \neg\varphi$ as well as $\neg\varphi \vdash \neg\varphi'$ Ok by Contraposition.
 2. ok by definition of SUB
- $(\varphi \rightarrow \psi)' = \varphi' \rightarrow \psi'$
 1. By assumption: We want $(\varphi \rightarrow \psi) \vdash (\varphi \rightarrow \psi)'$, it is enough to show $\varphi \rightarrow \psi; \varphi' \vdash \psi'$ We have
$$\varphi \rightarrow \psi; \varphi' \vdash \varphi$$

$$\varphi \rightarrow \psi; \varphi' \vdash \psi$$
 2. ok by definition of SUB
- $(\forall y\varphi)'$

Case 1: No occurrence of y in t . or $x = y$ (that is, t is substitutable for x in φ). We define $(\forall y\varphi)' = \forall y\varphi'$. All we need to check is part (a). We have that $\forall y\varphi \vdash \varphi$ because $\forall y\varphi \rightarrow \varphi$ is an axiom group 2. So $\forall y\varphi \vdash \varphi'$ and therefore by the generalization theorem $\forall x\varphi \vdash \forall y\varphi'$

Case 2: If y does occur in t and $x \neq y$. let z be the variable that is the first variable that does not occur in φ', x, t then set $(\forall y\varphi)' = \forall z(\varphi')^y_z$

2. want t SUB for x in $(\forall y\varphi)'$
 z does not occur in t (choice of z) t is SUB for x in φ' . (ind assumption) Then t is SUB für x in $\forall z(\varphi')^y_z$ iff t is SUB for x in $(\varphi')^y_z$ because $x \neq z$.
1. $\varphi \vdash \varphi'$ (by ind. assumption) Then $\forall y\varphi \vdash \forall y\varphi'$, because

$$\vdash \forall y(\varphi \rightarrow \varphi') \rightarrow (\forall y\varphi \rightarrow \forall y\varphi') \text{ (axiom of group 3)}$$

then

$$\forall y(\varphi \rightarrow \varphi') \text{ gen thm}$$

and by MP:

$$\forall y\varphi \rightarrow \forall y\varphi'$$

We have $\forall y\varphi' \vdash (\varphi')^y_z$ (axiom of group 2, z does not occur in φ') By Gen Thm. $\forall y\varphi' \vdash \forall z(\varphi')^y_z$ Then
 Want $\forall z(\varphi')^y_z \vdash \forall y\varphi$
 $\forall z(\varphi')^y_z \vdash ((\varphi')^y_z)^z_y$ (ax of group 2.), y is SUB for z in $(\varphi')^y_z$ bc. φ' does not contain z so all occurrences of z in $(\varphi')^y_z$ are free. (we substituted z for free occ of y .) (Replacement lemma $((\varphi')^y_z)^z_y = \varphi'$, see problem set.) So we have $\forall z(\varphi')^y_z \vdash \varphi$ We also know that $\varphi' \vdash \varphi$ by the inductive hypothesis. So $\forall z(\varphi')^y_z \vdash \varphi$ So $\forall z(\varphi')^y_z \vdash \forall y\varphi$ (Gen Thm.)

□

Note : φ' constructed in proof is also called an alphabetic variant of φ
 if our language contains equality:

1. $\vdash \forall x x = x$ (ax 5.)
2. $\vdash \forall x \forall y (x = y \rightarrow y = x)$ p.122
3. $\vdash \forall x \forall y \forall z (x = y \rightarrow (y = z \rightarrow x = z))$ (Exercise 11. in [EE01])
4. $\vdash \forall x_1 \forall x_2 \forall y_1 \forall y_2 (x_1 = y_1 \rightarrow (x_2 = y_2 \rightarrow (Px_1x_2 \rightarrow Py_1y_2)))$, similarly for any n -ary predicate. p.128
5. $\vdash \forall x_1 \forall x_2 \forall y_1 \forall y_2 (x_1 = y_1 \rightarrow (x_2 = y_2 \rightarrow (fx_1x_2 = fy_1y_2)))$, similarly for n -ary formula symbol, p.122

2.9 SOUNDNESS AND COMPLETENESS

In first order logic it holds:

- soundness: If $\Gamma \vdash \varphi$ then $\Gamma \models \varphi$
- completeness: If $\Gamma \models \varphi$ then $\Gamma \vdash \varphi$

For the proof of soundness we will have to show that all our axioms are valid. For this we will need the following two lemmas.

Lemma 9.1. pre-substitution lemma: *Let be a map TODO*

Lemma 9.2. Substitution lemma: *If t SUB x in φ then $\mathcal{A} \models \varphi_t^x[s]$ iff $\mathcal{A} \models \varphi[s(x|\bar{s}(t))]$*

Proof. 1. φ atomic: use pre-substitution lemma.

2. φ is of the form $\neg\psi$ or $\psi \rightarrow \theta$ - use induction

3. φ is of the form $\forall y\psi$ and x does not occur free in φ
 $\varphi_t^x = \varphi$ wts. $\mathcal{A} \models \varphi_t^x[s]$ iff $\mathcal{A} \models \varphi[s(x|\bar{s}(t))]$
 By 2.2, this is indeed the case, so the lemma holds.

4. φ is $\forall y\psi$ where x occurs free in φ and t is SUB for x in φ . Then it must be: y does not occur in t and t is SUB for x in ψ .
 then $\bar{s}(t) = \bar{s}(y|a)(t)$ for every $a \in A$. Moreover we also have, that $\varphi_t^x = \forall y\psi_t^x$ bc. $x \neq y$
 Then $\mathcal{A} \models \varphi_t^x[s]$ iff $\mathcal{A} \models \forall y\psi_t^x[s]$
 iff $\mathcal{A} \models \psi_t^x[s(y|a)]$ and for all $a \in A$.
 iff $\mathcal{A} \models \psi[s(y|a)(x|\bar{s}(y|a)(t))]$ (inductive assumption) and for all $a \in A$
 By above: iff $\mathcal{A} \models \psi[s(y|a)(x|\bar{s}(t))]$ for all $a \in A$
 iff $\mathcal{A} \models \forall y\psi[s(x|\bar{s}(t))]$

□

Theorem 9.3. *If $\Gamma \vdash \varphi$ then $\Gamma \models \varphi$*

Proof. Proof by induction on φ . We have to show:

- (i) that every logical axiom is valid
- (ii) logical implication is preserved by MP

On (ii) Assume 1. we have to show that if $\Gamma \vdash \varphi$ then $\Gamma \models \varphi$

- $\varphi \in \Lambda$ by 1.
- $\varphi \in \Gamma$ then $\Gamma \models \varphi$
- φ follows by MP from $\psi, \psi \rightarrow \varphi$ then by assumption $\Gamma \models \psi$ and $\Gamma \models \psi \rightarrow \varphi$ Therefore $\Gamma \models \varphi$

On (i) [EE01, exercise 6, section 2.2] consists in showing that if a logical axiom is valid, then also its generalization. So generalizations of valid formulas are valid, we therefore may only consider logical axioms that are not generalizations of another logical axiom.

Ax of 1. can be found in [EE01, exercise 3, section 2.3]

Ax of 2. wts. $\forall x\varphi \rightarrow \varphi_t^x$ is valid, where t is SUB for x in φ .
 simple case: $\forall xPx \rightarrow Pt$ is valid. Let $\mathcal{A} \models \forall xPx[s]$ then $\mathcal{A} \models \forall xPx[s(x|a)]$ for every $a \in A$. so i.p. for $a = \bar{s}(t)$ this means $\bar{s}(t) \in P^{\mathcal{A}}$ that is $\mathcal{A} \models Pt$. In more generality we will need the substitution lemma: We have $\mathcal{A} \models \forall x\varphi[s]$
 this is equivalent to $\forall a \in A$ we have $\mathcal{A} \models \varphi[s(x|a)]$ then in particular $\mathcal{A} \models \varphi[s(x|\bar{s}(t))]$
 and by the substitution lemma we have the equivalence to $\mathcal{A} \models \varphi_t^x[s]$

Ax of 3. can be found in [EE01, exercise 3, section 2.2]

Ax of 4. can be found in [EE01, exercise 4, section 2.2]

Ax of 5. $x = x$: $\mathcal{A} \models x = x[s]$ because $s(x) = s(x)$

Ax of 6. $x = y \rightarrow (\alpha \rightarrow \alpha')$ where α is atomic fla, and α' is obtained from α by replacing some occurrences of x 's with y 's. By the deduction theorem, is enough to show that the set of formulas $\{x = y, \alpha\} \models \alpha'$. Let \mathcal{A} be a structure, s an assignment such that $\mathcal{A} \models x = y[s]$
Claim: for every term t if t' is obtained from t by replacing some x 's by y 's, then $\bar{s}(t) = \bar{s}(t')$.
proof of claim. Induction on terms. ☒

- α of the form $t_1 = t_2$ then α' is $t'_1 = t'_2$, use prev. claim.
- α of the form $Pt_1 \dots t_n$ similar

□

Corollary 9.3-A. $\vdash \varphi \leftrightarrow \psi$ then φ, ψ are logically equivalent.

Corollary 9.3-B. φ' an alphabetic variant of φ then φ, φ' are logically equivalent.

Definition 2.33. : A set of formulas Γ is called satisfiable, whenever there is a structure \mathcal{A} with an assignment into A that for all $\sigma \in \Gamma$ $\mathcal{A} \models \sigma[s]$

Corollary 9.3-C. If Γ is satisfiable then Γ is consistent

Note : Corollary 9.3-C is equivalent to Soundness (see Exercises)

completeness

Theorem 9.4. Completeness Theorem: $\Gamma \models \varphi \implies \Gamma \vdash \varphi$

Theorem 9.5. Completeness Theorem': Every consistent set of formulas is satisfiable.

Note :

- The completeness Theorem is equivalent to completeness theorem'
- The completeness Theorem holds for language of any cardinality.
- We will assume for simplicity that the Language is countable.

Proof. Let Γ be a consistent set of formulas in some language \mathcal{L} The idea of the proof:

- 1.-3. build a new set of formulas Δ
 - $\Gamma \subseteq \Delta$
 - Δ consistent and maximal
 - For every fla φ and every variable x there is constant c $\neg \forall x \varphi \rightarrow \neg \varphi_c^x \in \Delta$
- 4. Build \mathcal{A} by A is the set of terms (in expanded language) such that Every fla in Δ w/o. equality (=) is satisfiable in \mathcal{A}
- accomodate =

1. Add a countable infinite set of new constant symbols to the language \mathcal{L} and call it \mathcal{L}'

Claim: Γ is a consistent set of formulas in \mathcal{L}' .

proof of claim. Why? If not, then $\Gamma \vdash \beta \wedge \neg \beta$ where deduction is in \mathcal{L}' and there occurs finitely many new constant symbols in this deduction. By generalization on constants the new constants in the proof can be replaced by new variables. We get a deduction in the old language \mathcal{L} and that contradicts the assumption that Γ is consistent. ☒

2. Want to add for every formula φ and every variable x $\neg \forall x \varphi \rightarrow \varphi_c^x$ and need to stay consistent. Fix enumeration of pairs (φ, x) where φ is a \mathcal{L}' -fla., x variable.

$$\theta_1 := \forall x_1 \varphi_1 \rightarrow \neg \varphi_{1c_1}^{x_1}$$

where c_1 is the first new constant that does not occur in φ_1 :

$$\theta_n := \forall x_n \varphi_n \rightarrow \neg \varphi_{nc_n}^{x_n}$$

where c_n is the first new constant that does not occur in φ_n and does not occur in θ_k for $k < n$.

$$\Theta = \{\theta_1, \dots\}$$

Claim: $\Gamma \cup \Theta$ is consistent.

proof of claim. Suppose it is not. Then let m be minimal such that $\Gamma \cup \{\theta_1 \dots \theta_{m+1}\} \vdash \beta \wedge \neg \beta$. Then by (Raa) $\Gamma \cup \{\theta_1 \dots \theta_m\} \vdash \neg \theta_{m+1}$ θ_{m+1} is of the form

$$\forall x_m \varphi_m \rightarrow \neg \varphi_{n_{c_m}^{x_m}}$$

then by (Rule T)

$$\Gamma \cup \{\theta_1 \dots \theta_m\} \vdash \neg \forall x \varphi$$

and

$$\Gamma \cup \{\theta_1 \dots \theta_m\} \vdash \varphi_c^x$$

(star..TODO)

star: $\Gamma \cup \{\theta_1, \dots, \theta_m\} \vdash \forall x \varphi$ By generalization on constants: $\Gamma \cup \{\theta_1, \dots, \theta_m\} \vdash \forall x (\varphi_c^x)_x^c$ since c does not occur on the left. also $(\varphi_c^x)_x^c = \varphi$ bc c does not occur in φ . Now we have

$$\Gamma \cup \{\theta_1 \dots \theta_m\} \vdash \neg \forall x \varphi$$

and

$$\Gamma \cup \{\theta_1, \dots, \theta_m\} \vdash \forall x (\varphi_c^x)_x^c$$

which is a contradiction to minimality of m or the consistentness of Γ . \square

3. Extend $\Gamma \cup \Theta$ to maximal consistent set. Λ is the set of logical axioms in \mathcal{L}' we know that $\Gamma \cup \Theta$ is consistent. so we know that there is no β

$$\Gamma \cup \Theta \cup \Lambda \models_{\text{taut}} \beta \wedge \neg \beta$$

So we find v a truth assignment on prime flas. that satisfies $\Gamma \cup \Theta \cup \Lambda$ and we are going to use this truth assignment to find the maximal set

$$\Delta := \{\varphi : \bar{v}(\varphi) = 1\}$$

Then for every φ either $\varphi \in \Delta$ or $\neg \varphi \in \Delta$ so we have maximality and we also have consistency bc. $\Delta \vdash \varphi$ then $\Delta \models_{\text{taut}} \varphi$ because $\Lambda \subseteq \Delta$ and that means $\bar{v}(\varphi) = 1$ so $\varphi \in \Delta$. So we have that Δ is consistent. and we say that Δ is deductively closed i.e. $\Delta \vdash \varphi$ then $\varphi \in \Delta$.

4. Construction of an \mathcal{L}' structure \mathcal{A} from Δ . We will firstly replace $=$ with E bin. predicate symbol. $A =$ set of all \mathcal{L}' -terms

E^A def. by $u E^A t$ iff $u = t \in \Delta$

f^A def by $f^A(t_1, \dots, t_n) = f t_1 \dots t_n$

$c^A := c$

P^A then $P^A t_1, \dots, t_n$ iff $P t_1 \dots t_n \in \Delta$ We take the assignment $s : \text{Var} \rightarrow A$ by $s(x) = x$

Claim 1: $\bar{s}(t) = t$ for every term t **Claim 2:** for every φ let φ^* be obtained from φ by replacing each $=$ with E then $\mathcal{A} \models \varphi^* [s]$ iff $\varphi \in \Delta$

proof of claim. • φ atomic then φ is Pt

$$\mathcal{A} \models \varphi^* [s] \text{ iff } \mathcal{A} \models Pt [s] \text{ iff } \bar{s}(t) \in P^A \text{ iff } t \in P^A$$

φ is uEt then

$$\mathcal{A} \models \varphi^* [s] \text{ iff } \mathcal{A} \models uEt [s] \text{ iff } \bar{s}(u) E \bar{s}(t) \text{ iff } u = t \in \Delta$$

- $\neg \varphi$

$$\mathcal{A} \models \neg \varphi^* [s] \text{ iff } \mathcal{A} \not\models \varphi [s] \text{ iff } \varphi \notin \Delta \text{ iff } \neg \varphi \in \Delta$$

- $\varphi \rightarrow \psi$

$$\mathcal{A} \models \varphi^* \rightarrow \psi^* [s] \text{ iff } \mathcal{A} \not\models \varphi^* [s] \text{ or } \mathcal{A} \models \psi^* [s] \text{ iff } \mathcal{A} \models \neg \varphi^* [s] \text{ or } \mathcal{A} \models \psi^* [s] \text{ iff } \neg \varphi \in \Delta \text{ or } \psi \in \Delta \text{ iff } (\varphi \rightarrow \psi) \in \Delta$$

- $\forall x \varphi$ wts. $\mathcal{A} \models \forall x \varphi^* [s] \text{ iff } \forall x \varphi \in \Delta$ Suppose $\mathcal{A} \models \forall x \varphi^* [s]$ then $\mathcal{A} \models \varphi^* [s(x|c)]$ where c is such that $\neg \forall x \varphi \rightarrow \neg \varphi_c^x \in \Delta$ Provided that we have substitutability we have by substitution lemma we know $\mathcal{A} \models (\varphi_c^x)^* [s]$ By the inductive hyphothesis $\varphi_c^x \in \Delta$ and $\neg \varphi_c^x \notin \Delta$ so we do not have $\neg \forall x \varphi \notin \Delta$ and by maximality of Δ we have $\forall x \varphi \in \Delta$.

Suppose $\mathcal{A} \not\models \forall x \varphi^* [s]$ then $\mathcal{A} \not\models \varphi^* [s(x|t)]$ for some t . By the substitution lemma (providet that t is SUB for x in φ) we can replace x by t in the formula.

$\mathcal{A} \not\models (\varphi_t^x)^* [s]$ by the inductive hyphothesis $\varphi_t^x \notin \Delta$ then $\forall x \varphi \notin \Delta$ because Δ is deductively closed. If t is not SUB for x in φ , we know that there exists a logically equivalent alphabetic variant φ' of φ such that t is SUB for x in φ' .

\square

So at this point we have: If \mathcal{L} does not contain $=$ then take \mathcal{A} reduction to \mathcal{L} and \mathcal{A} w/s satisfies Δ .

5. Define \mathcal{A}/E and assignment

Claim: $E^{\mathcal{A}}$ is a congruence on the structure \mathcal{A} compatible with the predicates and formulas.

- $E^{\mathcal{A}}$ is equivalence relation
- $P^{\mathcal{A}}$ compatible w/ $E^{\mathcal{A}}$ i.e. $P^{\mathcal{A}}t_1, \dots, t_n$ iff $P^{\mathcal{A}}s_1, \dots, s_n$ whenever $t_i E^{\mathcal{A}} s_i$ for all $1 \leq i \leq n$.
- $f^{\mathcal{A}}$ compatible w/ $E^{\mathcal{A}}$ i.e. $f^{\mathcal{A}}(\underline{t}) E^{\mathcal{A}} f^{\mathcal{A}}(\underline{s})$ iff $\underline{t} E^{\mathcal{A}} \underline{s}$

Definition 2.34. : \mathcal{A}/E is the structure w/ universe A/E and $([t_1], \dots, [t_n]) \in P^{\mathcal{A}/E}$ iff $(t_1, \dots, t_n) \in P^{\mathcal{A}}$ $f^{\mathcal{A}/E}([t_1], \dots, [t_n]) = [f^{\mathcal{A}}(t_1, \dots, t_n)]$ Let $h : A \rightarrow A/E : t \mapsto [t]$ quotient map. note h is surjective. $E^{\mathcal{A}/E}$ realized by equality on A/E : $[t] E^{\mathcal{A}/E} [s]$ iff $t E^{\mathcal{A}} s$ iff $[t] = [s]$

Claim: \mathcal{A}/E satisfies Δ w/ $h \circ s$.

proof of claim. Let $\varphi \in \Delta$, $\mathcal{A} \models \varphi^*[s]$ by (4) Want to show $\mathcal{A} \models \varphi^*[s]$ iff $\mathcal{A}/E \models \varphi^*[h \circ s]$ by Homomorphism Thm. (φ^* has no occurrence of $=$, surjectivity) realisation of E in \mathcal{A}/E is the equality in A/E . Take the reduct of \mathcal{A}/E to \mathcal{L} . \square

\square

Exam exam

Corollary 9.5-A. *compactness statements*

1. $\Gamma \models \varphi$ then there is a finite subset $\Gamma_0 \subseteq \Gamma$ s.th. $\Gamma_0 \models \varphi$
2. every finitly satisfiable set of formulas is satisfiable

Proof. 1. $\Gamma \models \varphi$ then by completeness $\Gamma \vdash \varphi$ where the deduction uses only formulas from some $\Gamma_0 \subseteq \Gamma$ finite. By soundness, $\Gamma_0 \models \varphi$

2. Γ finitly satisfiable. Suppose Γ is not satisfiable then by completeness Γ is not consistent. So there has to be some $\Gamma_0 \subseteq \Gamma$ such that $\Gamma_0 \vdash \beta \wedge (\neg\beta)$ so Γ is not finitly satisfiable (by soundness). \square

2.9.1 Sizes of models

Let Γ be a consistent set of formulas. Is it possible to

- Example 2.24.** 1. For each $n \in \mathbb{N}$ there is Γ such that all models of Γ that have size n .
2. DLO (Dense liniear order) w/o endpoints: no finite models

Lemma 9.6. Γ such that all models are finite. Then there has to be $m \in \mathbb{N}$ such that $|\mathcal{A}| \leq m$ for every model $\mathcal{A} \in \text{Mod } \Gamma$

Proof. \square

Suppose Γ has models of arbitrarily large finite size.

Idea: Expand language by new constant symbols c_0, c_1, \dots

$$\theta_1 := c_0 \neq c_1$$

$$\theta_2 := c_0 \neq c_1 \wedge c_1 \neq c_2 \wedge c_0 \neq c_2$$

\vdots

$$\theta_n := \bigwedge_{i,j=0, i \neq j}^n c_i \neq c_j$$

$$\Theta := \{\theta_1, \theta_2, \dots\}$$

$\Gamma \cup \theta$ finitely satisfiable. By the compactness theorem there exists a $\Theta_0 \subseteq \Theta$ finite TODO. then there is a maximal element θ_k By the compactness theorem reduct to language of Γ is an infinite models of Γ which is a contradiction to all models of Γ are finite.

Note : There is no sentence in the language of groups / rings / ... that would be satisfied in all finite groups/rings/... and not satisfied in all infinite groups/rings/...

Lemma 9.7. 1. *FIN, the class of all finite \mathcal{L} -structures is not EC_Δ*

2. *INF, the class of infinite \mathcal{L} -structures is EC_Δ but not EC .*

Proof. 1. Suppose FIN is EC_Δ . By definition there is a set of formulas Γ such that

$$\text{Mod } \Gamma = \{ \text{the collection of all finite } \mathcal{L}\text{-structures} \} = \text{FIN}$$

But then Γ has only finite models, but of arbitrarily large size. That contradicts our previous lemma.

2.

$$\begin{aligned} \varphi_1 & \quad \exists x x = x \\ \varphi_2 & \quad \exists x_1 \exists x_2 x_1 \neq x_2 \\ & \quad \vdots \\ \varphi_n & \quad \exists x_1 \dots \exists x_n \bigwedge_{i,j=0, i \neq j}^n x_i \neq x_j \end{aligned}$$

and $\Gamma := \{\varphi_1, \dots\}$ and INF is indeed EC_Δ . Suppose $\text{INF} = \text{Mod}(\tau)$ then $\text{Mod}(\neg\tau)$ would be FIN, a contradiction to (a). □

Recall the proof of completeness theorem. \mathcal{L} , $|\mathcal{L}| = \aleph_0$ Γ consistent set of \mathcal{L} -formulas. \mathcal{A}/E countable.

2.9.2 Completeness for uncountable languages

Use (AC) in the form of Zorn's Lemma and Zermelo's Theorem

Theorem 9.8. Zorn's Lemma: *P partially ordered set such that every chain has an upperbound in P then P contains a maximal element.*

Theorem 9.9. Zermelos's Theorem: *Every set can be well-ordered. That is linearly ordered such that every non-empty set has a smallest element.*

ω is the first infinite ordinal. then it is also a cardinal and is called \aleph_0

$$A_0 = \{(\varphi_\alpha, x_\alpha) : \alpha < \lambda\}$$

\vdots

$$|\mathcal{A}/E| \leq \lambda$$

CHAPTER 3

Model Theory

The sections 3.1 to 3.4 are sourced from [EE01, chapter 2] and the theory of o-minimality (from 3.5 onwards) can be found in [Van98].

3.1 LÖWENHEIM-SKOLEM-THEOREM

Theorem 1.1. LST: Let Γ be a set of \mathcal{L} -formulas. $|\mathcal{L}| = \lambda$ and let's assume Γ is satisfiable in some infinite structure.

Then for every cardinal $\kappa \geq \lambda$, Γ is satisfiable in a structure of cardinality κ .

Proof. add κ many new constants to the language \mathcal{L} .

$$\mathcal{L}' = \mathcal{L} \cup \{c_\alpha : \alpha < \kappa\}$$

$$\Sigma = \{c_\alpha \neq c_\beta : \alpha \leq \beta, \alpha, \beta \leq \kappa\}$$

Then $\Gamma \cup \Sigma$ is finitely satisfiable in \mathcal{L}' . This is because Γ is satisfiable in some infinite structure. By compactness $\Gamma \cup \Sigma$ is satisfiable. We have $\mathcal{A} \models \Gamma \cup \Sigma$ then $|\mathcal{A}| \geq \kappa$.

By the proof of completeness theorem, $\Gamma \cup \Sigma$ has a model of size $\leq \kappa$. Hence it is exactly of size κ . Take reduct TODO \square

Example 3.1. Language of ZFC $\mathcal{L} = \{\in\}$ is countable. Löwenheim-Skolem guaranties that ZFC has a countable model. called skolems paradox. ZFC knows that there are uncountable sets. explanation: some bijections are missing

Example 3.2. 1. $\overline{\mathbb{R}}$ real field. $\text{Thm}(\overline{\mathbb{R}})$ has a countable model. \mathbb{R}_{alg}

2. $\mathcal{N} = (\mathbb{N}, 0, S, +, \cdot)$

Claim: there exists a countable structure \mathcal{M} such that $\mathcal{N} \equiv \mathcal{M}$ but $\mathcal{N} \not\cong \mathcal{M}$ One way is to add new constant c to language $\Sigma = \{0 < c, S0 < c, \dots\}$ is fin satisfiable. So $\Sigma \cup \text{Th}(\mathcal{N})$ is fin satisfiable by compactness it is satisfiable

Take the reduct to original language. \mathcal{M} . and \mathcal{M} not isomorphic to \mathcal{N} , bc A bijection of $M \rightarrow \mathbb{N}$ would have to map c somewhere but for every $S^k 0 < c$ for every k wont be preserved by any map.

3.2 THEORIES AND COMPLETENESS

Definition 3.1. Theory: A theory T is a set of sentences that is closed under logical implication.

$$T \models \sigma \implies \sigma \in T$$

Note : If \mathcal{L} is a language. Then

- there is a smallest \mathcal{L} -theory. The set of all valid \mathcal{L} -sentences.
- and also a largest \mathcal{L} -theory. The set of all \mathcal{L} -sentences.

Definition 3.2. Theory of structures: Let \mathcal{K} some class of \mathcal{L} - structures. Then

$$\text{Th}(\mathcal{K}) = \{\sigma : \sigma \text{ } \mathcal{L}\text{-sentence and for every } K \in \mathcal{K} \sigma \in \text{Th}(K)\}$$

Note : $\text{Th}(\mathcal{K})$ is a theory.

$$\text{if } \text{Th}(\mathcal{K}) \models \sigma \text{ then } \sigma \in \text{Th}(\mathcal{K})$$

Example 3.3. • $\mathcal{L} = \{0, 1, +, \cdot, -\}$ \mathcal{F} the class of fields then $\text{Th}(\mathcal{F})$ is the set of sentences true in every field.

Recall that $\text{Mod}(\Sigma)$ is the class of all models of Σ . $\text{Th}(\text{Mod } \Sigma)$ might not be the set Σ but it is the set of all sentences true in all models of Σ . Which is the set of all sentences that are logically implied by Σ

Or in other words: The set of all consequences of Σ

Definition 3.3. C_n : $C_n(\Sigma) := \text{Th}(\text{Mod } \Sigma)$

Note : Σ is a theory iff $C_n(\Sigma) = \Sigma$

Definition 3.4. : We say that a theory T is complete, if for every sentence σ either $\sigma \in T$ or $\neg\sigma \in T$.

Example 3.4. \mathcal{A} a \mathcal{L} -structure, then $\text{Th}(\mathcal{A})$ is complete.

Note : $\text{Th}(\mathcal{K})$ is complete, iff any $K_1, K_2 \in \mathcal{K}$ are elementarily equivalent.

A theory T is complete iff any two models are elementarily equivalent.

Example 3.5. • The theory of fields is not complete.

- The theory of algebraically closed fields of characteristic 0 is complete (That is non-trivial)

Definition 3.5. axiomatizability:

- A theory T is finitely axiomatizable if there is a sentence σ such that $C_n(\sigma) = T$.
- A theory T is axiomatizable, if there is a decidable set Σ such that $C_n(\Sigma) = T$.

Example 3.6. • The theory of fields (common theory of all fields) is finitely axiomatizable.

- The theory of fields of characteristic 0 is axiomatizable. $\Psi \cup \{1 + 1 \neq 0, 1 + 1 + 1 \neq 0, \dots\}$ It is however not finitely axiomatizable. If $\Psi_0 \subseteq \Psi \cup \{1 + 1 \neq 0, 1 + 1 + 1 \neq 0, \dots\}$ finite, then Ψ_0 has a model of characteristic p for some sufficiently large p .

Theorem 2.1. If $C_n(\Sigma)$ is finitely axiomatizable then there exists a finite subset $\Sigma_0 \subseteq \Sigma$ such that $C_n(\Sigma_0) = C_n(\Sigma)$

Proof. Suppose $C_n(\Sigma)$ is finitely axiomatizable. So $C_n(\sigma) = C_n(\Sigma)$. Then there is a finite subset $\Sigma_0 \subseteq \Sigma$ such that $\Sigma_0 \models \sigma$. And we get $C_n(\Sigma_0) = C_n(\Sigma)$ \square

Definition 3.6. : A theory T is \aleph_0 -categorical, if any two infinite countable models of T are isomorphic. Furthermore for some infinite cardinal κ a theory T is called κ -categorical, if every two models of cardinality κ are isomorphic.

Theorem 2.2. Los-Vaught test: For a theory T in a countable language with only infinite models it holds

If T is κ -categorical for some infinite cardinality κ then T is complete.

Proof. Let T be κ -categorical. Want: If $\mathcal{A}, \mathcal{B} \models T$ then $\mathcal{A} \equiv \mathcal{B}$. Note: both \mathcal{A} and \mathcal{B} are infinite. By LST there exists structures \mathcal{A}' and \mathcal{B}' with $\mathcal{A} \equiv \mathcal{A}'$ and $\mathcal{B} \equiv \mathcal{B}'$ and $|\mathcal{A}'|, |\mathcal{B}'| = \kappa$. By κ -categorical we have $\mathcal{A}' \cong \mathcal{B}'$ so $\mathcal{A} \equiv \mathcal{B}$ \square

Note : completeness does not imply categorical.

- The theory of natural numbers is not \aleph_0 -categorical. See Example 3.2
- RCF not κ -categorical for all infinite cardinalities κ Not \aleph_0 categorical real clo of $\mathbb{Q}(\pi)$, real closure of \mathbb{Q} not uncountable categorical $\mathbb{R}, \mathbb{R}(\varepsilon), 0 < \varepsilon < \frac{1}{n}$ for every $n \in \mathbb{N}$.

3.3 THEORY OF ALGEBRAIC CLOSED FIELDS

Theorem 3.1. *The theory of algebraic closed fields of characteristic p ACF_p , where p is either 0 or prime is complete. [EE01, Theorem 26J, p.158]*

Proof. Note that we have a

- countable language
- with no finite models

Let $\mathcal{K}_1, \mathcal{K}_2 \models ACF_p$ such that $|K_1| = |K_2| = \kappa$ uncountable. F_1 prime field of \mathcal{K}_1 , F_2 prime field of \mathcal{K}_2 .

Note F_1, F_2 are determined by p if $p = 0$ then $F_1 = F_2 = \mathbb{Q}$ and if p prime then $F_1 = F_2 = \mathbb{F}_p$
 Define $F := F_1 = F_2$. B_1 transcendence base of \mathcal{K}_1 over F B_2 transcendence base of \mathcal{K}_2 over F

- B is transcendence base of K over F if B is a \subseteq -maximal subset of K which is algebraically closed then
- $B \subseteq K$ is algebraically TODO

$F(B_1), F(B_2)$ subfields of $\mathcal{K}_1, \mathcal{K}_2$

- $\text{alg cl } F(B_1) = \mathcal{K}_1$
- $\text{alg cl } F(B_2) = \mathcal{K}_2$

Fact: Let F subfield of K . if F is countable and K uncountable, then any transe basis B of K oer F is of cardinality $|K|$, hence uncountable.

Steinitz: Two ACF are isomorphic iff they have the same characteristic and there transcendence spaces have the same cardinality. \square

Lefschetz Principle

Proposition 3.2. Lefschetz Principle: *Let $\mathcal{C} = (\mathbb{C}, 0, 1, +, \cdot, -)$ For a sentence in the language of \mathcal{C} Then the following are equivalent:*

- $\mathcal{C} \models \sigma$
- $\mathcal{A} \models \sigma$ for every $\mathcal{A} \models ACF_0$
- $ACF_0 \models \sigma$
- for all sufficiently large primes p $ACF_p \models \sigma$
- For infinitely many primes p ACF_p models σ

Proof. Sketch:

(a), (b), (c) are equivalent by completeness of ACF_0

(c) \implies (d) $ACF_0 \models \sigma$ so there is $T_0 \subseteq ACF_0$ such that $T_0 \models \sigma$ therefore there exists a sufficiently large prime p such that $ACF_p \models \sigma$.

(d) \implies (e) TODO

(e) \implies (c) If $ACF_0 \models \sigma$ than $ACF_0 \models \sigma$

\square

Example of the Lefschetz Principle:

Proposition 3.3. Ax{Grothendieck: ¹ *Let $f : \mathbb{C}^n \rightarrow \mathbb{C}^n$ be a polynomial map. If f is injective, then f is surjective.*

¹Alexander Grothendieck

Proof. Our language is $\mathcal{L} = \{0, 1, +, -, \cdot\}$. Note that there is an \mathcal{L} -sentence Φ_d such that a Field F

$F \models \Phi_d$ iff for every polynomial map $f : F^n \rightarrow F^n$ whose TODO coord. function is of degree at most d , if f is injective then f is surjective.

By Lefschetz principle it is enough to show for sufficiently large primes p , $\text{ACF}_p \models \Phi_d$ for all $d \in \mathbb{N}$. Since ACF_p is complete, it is enough to show that every injective polynomial map $f : K^n \rightarrow K^n$ is surjective, where $K = \text{TODO}$. Let $f : K^n \rightarrow K^n$ be a polynomial map.

Then there is a finite subfield K_0 of K such that all coefficients of f come from K_0 . Let $y \in K^n$. Then there is a finite subfield K_1 of K such that $y \in K_1$ and $K_0 \subseteq K_1 \subseteq K$. Since $f : K_1^n \rightarrow K_1^n$ is injective and K_1 finite, $f|_{K_1}$ is surjective onto K_1 . So there is $x \in K_1^n$ such that $f(x) = y$. \square

Note : Later, a purely geometric proof was found by Borel.

Another use of Łoś-Vaught

Proposition 3.4.

$$(\mathbb{Q}, <_{\mathbb{Q}}) \equiv (\mathbb{R}, <_{\mathbb{R}})$$

Proof. $\mathcal{L} = \{<\}$ and note that both $(\mathbb{Q}, <_{\mathbb{Q}}), (\mathbb{R}, <_{\mathbb{R}})$ are DLO without endpoints, i.e. they satisfy the following axioms

1. $\forall x \forall y (x < y \vee x = y \vee y < x)$
2. $\forall x \forall y (x < y \rightarrow \neg(y < x))$
3. $\forall x \forall y \forall z ((x < y \wedge y < z) \rightarrow x < z)$
4. $\forall x \forall y (x < y \rightarrow \exists z (x < z \wedge z < y))$
5. $\forall x \exists y \exists z (y < x \wedge x < z)$

TODO \square

3.4 NONSTANDARD ANALYSIS

1. Language \mathcal{L} : $=, \forall$ ranging over \mathbb{R} ,

- P_R TODO

2. standard structure for \mathcal{L} : \mathcal{R} with universe \mathbb{R} , $c_r^{\mathcal{R}} = r$, $P_R^{\mathcal{R}} = R$, $f_F^{\mathcal{R}} = F$.

3. Nonstandard structure for \mathcal{L} : \mathcal{R}^* , which is constructed using the compactness theorem

$$\Gamma := \text{Th}(\mathcal{R}) \cup \{c_r P_{<} v_1 : r \in \mathbb{R}\}$$

Compactness theorem \implies there exists a \mathcal{L} -structure \mathcal{R}^* with $\mathcal{R}^* \models \Gamma[(v_1|a)]$ for some $a \in R^*$. We have $\mathcal{R} \equiv \mathcal{R}^*$. Moreover, $h : \mathbb{R} \rightarrow R^*$ defined by $r \mapsto c_r^*$ is an isomorphism of \mathcal{R} into \mathcal{R}^*

- h is injective:
- TODO

Note : WMA \mathcal{R} substructure of \mathcal{R}^* (se PS)

Notation: We will write *B instead of $P_B^{\mathcal{R}^*}$.

Example 3.7. what is ${}^*\mathbb{R}$? We have that $\mathcal{R} \models \forall x P_{\mathbb{R}}$, hence $\mathcal{R}^* \models \forall x {}^*\mathbb{R}$, so ${}^*\mathbb{R} = R^* =$ universe of \mathcal{R}^* . **Note :** Let F be an n -ary operator on \mathbb{R} . Then F is the restriction of *F to \mathbb{R} . ${}^*c_r = r$.

Idea: If we want to show that *R or *F has certain property, then we show

- R or F have that property.
- property can be expressed in \mathcal{L}

TODO

$$\mathcal{R}^* \supseteq \mathcal{R} \text{ such that } \mathcal{R}^* \equiv \mathcal{R}. \mathcal{F} = \{x \in \mathcal{R}^* : \exists r \in \mathbb{R} {}^*|x|^* \leq r\} \mathcal{I} = \{x \in \mathcal{R}^* : \forall r \in \mathbb{R} {}^*|x|^* < r\}$$

Proposition 4.1. 1. \mathcal{F} is a subring of \mathcal{R}^*

2. \mathcal{I} is an ideal in \mathcal{R}^*

Proof. 1. Let $x, y \in \mathcal{F}$ then there exists $a, b \in \mathbb{R}^{>0}$ such that $^*|x|^* \leq a$ and $^*|y|^* \leq b$. then

$$\begin{aligned} ^*|x \pm y|^* &\leq ^*|x|^* + ^*|y|^* \leq a + b \in \mathbb{R}^{>0} \\ ^*|x \cdot y|^* &= ^*|x|^* \cdot ^*|y|^* \leq a \cdot b \in \mathbb{R}^{>0} \end{aligned}$$

2. $x, y \in \mathcal{I}$ then $\forall a \in \mathbb{R}^{>0}$ we have $|x| < \frac{a}{2}$ Then

$$|x \pm y| \leq \frac{a}{2} + \frac{a}{2} = a$$

Let z be finite then $|z| < b \in \mathbb{R}^{>0}$ Let $a \in \mathbb{R}^{>0}$ then $|x| < \frac{a}{b}$ so

$$|xz| < \frac{a}{b}b = a$$

□

Definition 3.7. infinitely close: x, y are called to be infinitely close ($x \simeq y$), if $y - x \in \mathcal{I}$

\simeq

Proposition 4.2. 1. \simeq is an equivalence relation

2. \simeq is congruent with $^*+, ^*\cdot, ^*-$

Lemma 4.3. Suppose $\neg x \simeq y$ and at least one of x, y is finite then there exists $q \in \mathbb{R}$ such that q is between x and y

Proof. $y - x \notin \mathcal{I}$, wlog. $x < y$ then there exists $b \in \mathbb{R}$ such that $0 < b < y - x$ and by the archimedian property there is $m \in \mathbb{N}^{>0}$ such that $x < mb$. Let m be the smallest such. i.e. $(m-1)b \leq x < mb$. And $mb < y$. □

Proposition 4.4. For every $x \in \mathcal{F}$ there is exactly one $r \in \mathbb{R}$ such that $x \sim r$

Proof. Let $S := \{r \in \mathbb{R} : r < x\}$. S is bounded in \mathbb{R} because $|x| < r_0$ for some $r_0 \in \mathbb{R}^{>0}$. Then $r := \sup S$. Claim: $r \simeq x$. Lets assume by contradiction that this is not the case. By the previous lemma, there is $q \in \mathbb{R}$ such that $r < q < x$ or $x < q < r$. but neither of this things can happen.

- $r < q < x$ is contradiction to r not being an upper bound.
- $x < q < r$ is contradiction to r is not the least upper bound.

□

A consequence of that:

Corollary 4.4-A. for each $x \in \mathcal{F}$ there is a unique way of writing of x in the form $r + i$ where $r \in \mathbb{R}$ and $i \in \mathcal{I}$

Note : If $x = r + i$ then we also write $\text{st}(x) = r$.

Proposition 4.5. • $\text{st} : \mathcal{F} \rightarrow \mathbb{R}$

- $\text{st}(x) = 0$ iff $x \in \mathcal{I}$
- $\text{st}(x^* + y) = \text{st}(x) + \text{st}(y)$
- $\text{st}(x^* \cdot y) = \text{st}(x) \cdot \text{st}(y)$

Note : this says that st is a homomorphism of \mathcal{F} onto field $\overline{\mathbb{R}}$ with $\ker(\text{st}) = \mathcal{I}$ and $\mathcal{F}/\mathcal{I} \cong \overline{\mathbb{R}}$

Definition 3.8. Convergence (non-standard definition): F converges at a to b if whenever $x \simeq a$ and $x \neq a$ then $^*F(x) \simeq b$.

Note : This definition is equivalent to $\varepsilon - \delta$ definition of convergence in Analysis.

- Suppose F converges to b at a in $\varepsilon - \delta$ -sense

$$\mathcal{R} \models \forall \varepsilon > 0 \exists \delta > 0 \forall z (|z - a| < \delta \implies |F(z) - b| < \varepsilon)$$

$$\mathcal{R}^* \models \forall \varepsilon > 0 \exists \delta > 0 \forall z (|z - a| < \delta \implies |F(z) - b| < \varepsilon)$$

Let $\varepsilon > 0$ and $\delta > 0$ corresponding to ε . Let $x \simeq a$ then $|x - a| < r$ for all positive $r \in \mathbb{R}^{>0}$ so in particular $|x - a| < \delta$, therefore $|F(x) - b| < \varepsilon$ but ε was arbitrarily, so $\text{st}(F(x)) = b$.

- Suppose F converges to b at a in the non-standard-sense. Then $\forall \varepsilon \in \mathbb{R}^{>0}$

$$\mathcal{R}^* \models \exists \delta > 0 \forall x (|x - a| < \delta \rightarrow |F(x) - b| < \varepsilon)$$

Because $\delta \in \mathcal{I}$ works. But then

$$\mathcal{R} \models \exists \delta > 0 \forall x (|x - a| < \delta \rightarrow |F(x) - b| < \varepsilon)$$

Note : If F converges to b at a then b is unique such that for every $i \in \mathcal{I}$ the standard part $\text{st}(F(a + i)) = b$. And we use the general notation $\lim_{x \rightarrow a} F(x) = b$

Corollary 4.5-A. F continuous at a then $x \simeq a \implies {}^*F(x) \simeq {}^*F(a)$

Derivatives From Analysis: If $F : \mathbb{R} \rightarrow \mathbb{R}$ then $F'(a) = \lim_{h \rightarrow 0} \frac{F(a+h) - F(a)}{h}$

Definition 3.9. : We will say that $F'(a) = b$ iff $\forall dx \in \mathcal{I}, dx \neq 0$ then

$$\text{st}\left(\frac{F(a + dx) - F(a)}{dx}\right) = b$$

$dF := {}^*F(a + dx) - F(a)$ then $F'(a) = b$ iff $\forall dx \in \mathcal{I}, dx \neq 0$ we have $\text{st}\left(\frac{dF}{dx}\right) = b$ $\frac{dF}{dx}$ is an actual division.

Example 3.8. $F(x) = x^2$

$$\frac{dF}{dx} = \frac{(a + dx)^2 - a^2}{dx} = \frac{2dxa + (dx)^2}{dx} = 2a + dx$$

and $\text{st} \frac{dF}{dx} = 2a$

Proposition 4.6. (standard) If $F'(a)$ exists, then F is continuous at a .

Proof. Assume $F'(a)$ (in the standard sense) exist, then $F'(a)$ is a finite number and $F'(a) \simeq \frac{F(a+dx) - F(a)}{dx}$. Therefore $F(a + dx) - F(a)$ has to be infinitesimal ($\in \mathcal{I}$). Which means $F(a + dx) \simeq F(a)$. \square

Proposition 4.7. Chain rule: Suppose $G'(a)$ and $F'(G(a))$ exist then $(F \circ G)'(a) = F'(G(a)) \cdot G'(a)$

Proof. Note: ${}^*(F \circ G) = {}^*F \circ {}^*G$ because $\mathcal{R} \models \forall x F_{f \circ g}(x) = (F_f \circ F_g)(x)$

$$dG := {}^*G(a + dx) - {}^*G(a)$$

$$\begin{aligned} dF &:= {}^*(F \circ G)(a + dx) - {}^*(F \circ G)(a) \\ &= {}^*F({}^*G(a + dx)) - {}^*F({}^*G(a)) \\ &= {}^*F({}^*G(a) + dG) - {}^*F({}^*G(a)) \end{aligned}$$

We know that $G(a)$ exists so G is continuous at a and therefore $dG \simeq 0$

- case $dG \neq 0$ then $\frac{dF}{dG} \simeq F'(G(a))$. We can re-write

$$\frac{dF}{dx} = \frac{dF}{dG} \frac{dG}{dx} = F'(G(a)) \cdot G'(a)$$

- case $dG = 0$ then $dF = 0$ and $G'(a) = \frac{dG}{dx} = 0$ and therefore $(dx \neq 0)$

$$\frac{dF}{dx} = 0 = F'(G(a)) \frac{dG}{dx}$$

\square

3.5 O-MINIMALITY

Example 3.9. $\overline{\mathbb{R}} = (\mathbb{R}, +, -, \cdot, 0, 1, \leq)$ $\overline{\mathbb{R}} = (\mathbb{R}, \leq)$ TODO or at least in a very similar language, then by quantifier elimination (QE, Tarski) all the definable sets of \mathbb{R} are finite unions of points and intervals.

Definition 3.10. o-minimality: Let $\mathcal{L} = \{\leq, \dots\}$, \mathcal{M} is an \mathcal{L} -structure such that $\mathcal{M} \models \text{DLO}$ and the only definable subsets of M are finite union of points and intervalls. Then \mathcal{M} or equivalent $\text{Th}(\mathcal{M})$ is called o-minimal.

o-minimal is not a first order property so to say that a theory is o-minimal is non trivial. **Note** : Cell decomposition means Suppose X is definable in an o-minimal structure \mathcal{M} , $X \subseteq M^n$ then X is a finite union of cells (in dimension 1 these are points or intervalls) in M^2 it is either the graph of a continuous function or everything inbetween two graphs of continuous functions. (its an inductive definition)

Note : Have Dedekind complete for definable subsets of M : For $X \subseteq M$ definable then $\inf X, \sup X$ exist in $M_{\pm\infty}$.

Note : If M contains infinitely small elements, for example $M = {}^*\mathbb{R}$ then $(0, 1) \subseteq M$ is not connected. $O_1 = \{x : \forall n \in \mathbb{N}^* 0 < x < \frac{1}{n}\}$ is open and so is its complement in $(0, 1)$. We have

Note that O_1 is however not definable in M . If O_1 would be definable it would be a finite union of points and intervals. It is convex, and not a point. But it is also not an intervall, because then it would have by Dedekind completeness that $\sup O_1$ exists in $M_{\pm\infty}$, a contradiction. TODO:

Definition 3.11. definably connectedness: $X \subseteq M^m$ is said to be definably connected, if X is definable and X is not the disjoint union of two definable, non-empty open sets.

- Lemma 5.1.**
1. The definably connected subsets of M are the intervalls (including singletons) and \emptyset .
 2. The image of a definable connected subset $X \subseteq M^n$ under a definable continuous map $f : X \rightarrow M^n$ is definably connected. (f is called to be definable, if its graph $\Gamma f \subseteq M^{mn}$ is).
 3. (IVP) If $f : [a, b] \rightarrow M$ definable and continuous, then f assumes all values between $f(a)$ and $f(b)$.

Proof. Exercise □

3.6 O-MINIMAL ORDERED GROUPS AND RINGS

Definition 3.12. ordered group: A ordered group is a group with a linear order such that

$$\forall x \forall y \forall z (x < y \rightarrow (zx < zy \wedge xz < yz))$$

Example 3.10. • $(\mathbb{R}, <, +)$

- $(\mathbb{R}^{>0}, <, \cdot)$
- non-example: $(\mathbb{R}^*, \cdot, <)$

Recall:

- (G, \cdot) is divisible, if $\forall n \forall g \exists x (xg = x^n)$, equivalent to $\forall n G^n = G$.
- (G, \cdot) is torsion-free, if no element has finite order except for 1.

Proposition 6.1. $(M, <, \cdot, \dots)$ o-minimal such that $(M, <, \cdot)$ ordered group, then $(M, <, \cdot)$ abelian, divisible and torsion-free.

Lemma 6.2. If G is a definable subgroup of M then G is convex.

Proof. Suppose G is not convex, then there exists $1 < a < g$ for some $g \in G$ and $a \in M \setminus G$. Then

$$1 < a < g < ag < g^2 < ag^2 < g^3 < \dots$$

but elements alternate being in G and outside of G so G is not definable (finite union). \square

Lemma 6.3. *The only definable subsets of M that are subgroups are $\{1\}$ and M .*

Proof. Suppose $G \neq \{1\}$ wts. $G = M$. From the previous lemma we know that G is convex. The idea is $s := \sup G$ then $1 < s$ and $(1, s) \subseteq G$. If $s = +\infty$ then $G = M$. Suppose $s \neq +\infty$ then Take $1 < g < s$ then $g^{-1}s \in (1, s)$ So $s = gg^{-1}s \in G$ and $s < sg$ thats a contradiction with $s = \sup G$. \square

Proof. of Proposition.

- (M, \cdot) abelian: For any $a \in M$ we can look at $C_a = \{x \in M : xa = ax\}$ it is a definable subgroup and contains a it therefore is non-trivial and we have $C_a = M$ for every $a \in M$, so abelian.
- For any $n \in \mathbb{N}^{>0}$ look at $\{x^n : x \in M\}$ non trivial definable subgroup of M , hence $= M$.
- Every ordered group is torsion-free.

\square

Definition 3.13. Ordered ring: A ring (assumed to always be associative, with 1) equipped with a linear order such that

1. $0 < 1$
2. $<$ translation invariant
3. $<$ invariant under multiplication by positive elements

Note :

- The additive group $(R, <, +)$ of an ordered ring is a ordered group.
- Ordered rings have no zero-divisors $\forall x \forall y xy = 0 \rightarrow (x = 0 \vee y = 0)$
- $x^2 \geq 0$
- $k \mapsto k \cdot 1 : \mathbb{Z} \rightarrow \text{ring}$ is a strictly increasing embedding with respect to the usual ordering on \mathbb{Z} that means our characteristic is 0.

Note :

- A division ring is a field without commutativity of multiplication, so

$$\forall x x \neq 0 \rightarrow \exists y xy = 1$$

- Suppose ordered ring is also a division ring. Then such y are unique and $yx = 1$. Further, $x > 0 \rightarrow y > 0$.

Also the additive group is divisible, the underlying set is DLO w/o endpoints and $(x, y) \rightarrow x \cdot y$
 $x \rightarrow x^{-1}$ are continuous with respect to intervall topology.

Definition 3.14. ordered field: An ordered field is an ordered division ring with commutative multiplication.

Definition 3.15. real closed field: ordered field R such that if $f(X) \in R[X]$ and $a < b$ are such that $f(a) < 0 < f(b)$ then there exists a $c \in (a, b)$ such that $f(c) = 0$

RCF

Example 3.11. • $(\mathbb{R}, +, \cdot, <)$ is RCF

- $(\mathbb{Q}, +, \cdot, <)$ is not a RCF

Proposition 6.4. $(M, <, +, \cdot, \dots)$ \mathcal{o} -minimal such that $(M, <, +, \cdot)$ is ordered ring then $(M, <, +, \cdot)$ is RCF.

Proof. • wts. $(M; <, +, \cdot)$ is ordered division ring. For all $a \in M$ aM is additive subgroup of $(M, +)$ hence $aM = M$ if $a \neq 0$.

- wts. commutativity of \cdot . $\text{Pos}(M) := \{a \in M : a > 0\}$ is a subgroup of the multiplicative group of M . Let $a \in M$ then bc $M = aM$ we have $b \in M$ such that $1 = a \cdot b$ and by note $0 < a$ then $0 < a^{-1}$. so multiplication is commutative on M
- IVP property for polynomials: The ring operations are continuous, see note and use lemma about IVP (c).

□

Cell decomposition

Base step:

Proposition 6.5. Monotonicity Theorem: Suppose $f : (a, b) \rightarrow M$ definable, then there are $a < c_1 < \dots < c_k < b$ such that for (a, c_1) , (c_i, c_{i+1}) , (c_k, b) subsets of (a, b) we have: f is either constant or strictly monotonic and continuous.

Lemma 6.6. \exists subinterval on which f is const or injective.

Lemma 6.7. If f injective, then strictly monotone on a subinterval.

Lemma 6.8. If f strictly monotone, then f continuous on a subinterval.

Proof. Proof of Monotonicity theorem: Consider

$$X := \left\{ x \in (a, b) : \begin{array}{l} \text{on some subinterval containing } x, \\ f \text{ is either constant or strictly monotone and continuous} \end{array} \right\}$$

remark: X is a definable set. Look at $(a, b) - X$ is finite. If not, it would contain subinterval use lemma to get contradiction. WMA: $X = (a, b)$ in particular we may assume f continuous. By subdividin (a, b) further WMA that we are in one of the following cases

Case 1: $\forall x \in (a, b)$ f constant on some neighborhood of x

Case 2: $\forall x \in (a, b)$ f is strictly monotone increasing on some neighborhood of x

Case 3: $\forall x \in (a, b)$ f is strictly monotone decreasing on some neighborhood of x

Case 1: $x_0 \in (a, b)$ then $s := \{x : x_0 < x < b \wedge f \text{ cont. on } [x_0, x]\}$ wts $s = b$ suppose $s < b$ then f constant on neighborhood of s contradiction with definition of s so f continuous on $[x_0, b)$. f constant on $(a, x_0]$ similar.

Case 2: $x_0 \in (a, b)$ then $s := \{x : x_0 < x < b \wedge f \text{ strictly incr. on } [x_0, x]\}$. wts: $s = b$ assume $s < b$ then f is strictly increasing on some neighborhood of s so f strictly increasing on $[x_0, s + \delta)$ for some $\delta > 0$, a contradiction to definition of s .

Case 3: similar to Case 2.

□

Proof of Lemma 1:

Proof. Statement: “There exists a subinterval on which f is constant or injective.”

- If $y \in R$ so that f^{-1} is infinite (it has to be a finite union of points and intervals), then $f^{-1}(y)$ contains an interval and $f(x) = y$ on that interval.
- Suppose $f^{-1}(y)$ is finite for every $y \in R$. $f(I)$ is infinite and is definable because f is definable, so it contains an interval J . We can define an inverse to f on J $g : J \rightarrow I$, $g(y)$ is the first $x \in I$ such that $f(x) = y < ($ this is definable). g is necessarily injective. $g(J)$ infinite, so contains a subinterval on which f is injective.

□

Proof of Lemma 2:

Proof. Statement: “If f injective, then strictly monotone on a subinterval.”

Suppose f is injective. $f : I = (a, b) \rightarrow R$ pick $x \in (a, b)$ then $(a, x) = \{y \in (a, x) : f(y) < f(x)\} \uplus \{y \in (a, x) : f(x) < f(y)\}$ is definable disjoint union of definable sets, so one of the subsets has to contain an interval (c, x) with $a \leq c$, similarly for (x, d) . So for all $x \in I$ we have x satisfies one of the following:

- $\Phi_{++}(x)$ iff $\exists c_1, c_2 (c_1 < x < c_2 \wedge \forall c \in (c_1, x) f(c) > f(x) \wedge \forall c \in (x, c_2) f(c) > f(x)$
- $\Phi_{--}(x)$
- $\Phi_{+-}(x)$
- $\Phi_{-+}(x)$

The set of all x that satisfy each Φ is definable, it therefore is a finite union of points and intervals. After passing to subinterval (a, b) of I WMA that each $x \in I$ satisfies the same $\Phi_{\pm\pm}$.

- $\Phi_{-+}(x)$, on the left everybody is smaller, on the right everybody is bigger.

$$\forall x \in I s(x) := \sup\{s \in (x, b) : f(x) < f(s)\}$$

If $s(x) < b$ then $\Phi_{-+}(s(x))$, and therefore there is an element $s' > s(x)$ such that $f(x) \leq f(s(x)) < f(s')$ so $s(x) \geq s'$ which is a contradiction to definition to $s(x)$, therefore $s(x) = b$ for every $x \in (a, b)$. Then f has to be strictly increasing on (a, b) .

- $\Phi_{+-}(x)$ similar (monotonic decreasing)
- $\Phi_{++}(x) \forall x \in I$.

$$B := \{x \in I : \forall y \in I (x < y \rightarrow f(y) > f(x))\}$$

B is definable, if B is infinite, it has to contain a subinterval on which f is strictly increasing. WMA B is finite. We restrict ourselves to subinterval and may assume $B = \emptyset$. So by injectivity:

$$\textcircled{*} \quad \forall x \in I \exists y \in I x < y \wedge f(x) > f(y)$$

Let $c \in I$. Claim: for every large enough $y \in I$ we have $f(y) < f(c)$.

proof of claim. By contradiction. suppose we can not find a neighborhood of b such that for all elements in this neighborhood $f(y) < f(c)$ otherwise $f(y) > f(c)$ for all large enough y . Let $d < b$ be minimal such that

$$\forall y \in (d, b) f(y) > f(c)$$

- case $f(d) > f(c)$: $\Phi_{++}(d)$, contradiction with minimality of d .
- case $f(d) < f(c)$: By $\textcircled{*}$ there has to be an e with $d < e < b$ and $f(e) < f(d)$. So $f(e) < f(c)$ which is a contradiction to $\Phi_{++}(d)$

⊠

Define $y(c)$ to be the least element of $[c, b)$ for which

$$\forall y y(c) < y < b f(c) > f(y)$$

c satisfies Φ_{++} , therefore $c < y(c)$ and $f(y(c)) < f(c)$ if $y(c) < y < b$. The minimality of $y(c)$ implies that $y(c)$ satisfies Ψ_{+-} , where

$$\Psi_{+-}(v) \text{ iff } \exists v_1, v_2 \in I (v_1 < v < v_2 \wedge \forall z_1, z_2 (v_1 < z_1 < v \wedge v < z_2 < v_2) \rightarrow f(z_1) > f(z_2))$$

But c was arbitrarily so $\forall x \in I \exists v \in I (x < v \wedge \Psi_{+-}(v))$ On subinterval Ψ_{+-} we have a contradiction with Φ_{++} , similarly on subinterval for Ψ_{-+} .

- Φ_{--} similar to above

□

Proof of Lemma 3:

Proof. Statement: “If f strictly monotone, then f continuous on a subinterval.”

WMA: $f : (a, b) \rightarrow R$ strictly monotone increasing. $f(I)$ infinite and definable, so $f(I)$ contains an interval J . Let $r, s \in J$ $r < s$ and $d, e \in I$ with $d < e$ and $f(d) = r$ and $f(e) = s$. restrict f to (d, e) and we get an increasing bijection $(d, e) \rightarrow (r, s)$. Our topology is the order topology, so f is continuous on (d, e) \square

So we have proved the monotonicity Theorem. **Note :** If $f : (a, b) \rightarrow R$ is definable, then $\lim_{x \rightarrow c^-} f(x)$ exists in R_∞ for $c \in (a, b]$. And further $\lim_{x \rightarrow c^+} f(x)$ exists in R_∞ for $c \in [a, b)$. If furthermore $f : [a, b] \rightarrow R$ is continuous and definable, then f assumes a minimum and maximum on $[a, b]$

One of the important tools in o-minimality theory is the cell decomposition theorem:

Definition 3.16. Cell: Let (i_1, \dots, i_n) a sequence in $\{0, 1\}$. An (i_1, \dots, i_n) -cell is defined inductively:

- (0)-cell: $\{r\} \subseteq R$,
- (1)-cell: $(a, b) \subseteq R$, $a < b$, $a, b \in R$.
- $(i_1, \dots, i_k, 0)$ -cell: $\Gamma f \subseteq R^{k+1}$, where f is definable and continuous function $f : X \rightarrow R$, where X is a (i_1, \dots, i_k) -cell
- $(i_1, \dots, i_k, 1)$ -cell: is a the set

$$(f, g) = \{(\underline{x}, x_{k+1}) \in R^{k+1} : f(\underline{x}) < x_{k+1} < g(\underline{x})\}$$

$f : X \rightarrow R, g : X \rightarrow R$, $f < g$, f, g are definable and continuous on X , which is a (i_1, \dots, i_k) -cell. f may be constantly $-\infty$ and g may be constantly ∞ .

Note Cells have nice topological properties:

- every $(1, \dots, 1)$ -cell are precisely the cells that are open in their ambient space. continuity of the functions is important.
- The union of finitely many non-open cells has empty interior.
- Each cell is locally closed i.e. open in its closure.
- Each cell is homeomorphic to an open cell under a coordinate projection Example $(1, 0, 0, 1)$ -cell or $(1, 0)$ -cell with $(x_1, x_2) \mapsto x_2$
- If $A \subseteq R^{n+1}$, then $\pi A \subseteq R^n$ cell $\pi(x_1, \dots, x_{n+1}) \mapsto (x_1, \dots, x_n)$
- Every cell is definably connected. You can proof this by induction on the cell. $\{r\}$ and open intervals are definably connected. If the projection of a cell is definably connected, then the fibre above it is either an open interval or a single point. It is even definably path connected. If there would exist an open disjoint cover there exists an open disjoint cover of the fibre, which is not possible.

Definition 3.17. decomposition: A decomposition of R^m is a finite partition of R^m into cells defined inductively:

- decomposition of $R^1 = R$:

$$\{(-\infty, a_1), \{a_1\}, (a_1, a_2) \dots (a_k, \infty)\}$$

- A decomposition of R^{n+1} is a finite partition of R^{n+1} into cells C such that the collection of πC is a decomposition of R^n .

Theorem 6.9. Cell decomposition:

- (I_m) Let $A_1, \dots, A_k \subseteq R^m$ definable sets. Then there is a decomposition of R^m partitioning each A_i .
- (II_m) Given a definable function $f : A \rightarrow R$, $A \subseteq R^m$ there is a decomposition \mathcal{D} of R^m partitioning A such that for every $B \in \mathcal{D}$ $f|_B : B \rightarrow R$ is continuous.

Proof. By induction on m . Base step:

- (I_1) o-minimality
- (II_1) monotonicity theorem.

Proof idea: Suppose we have

$$\left\{ \begin{array}{l} (I_1) \dots (I_m) \\ (II_1) \dots (II_m) \end{array} \right\} \implies (I_{m+1}), (II_{m+1})$$

□

Definition 3.18. : A definably connected component of a non-empty definable Subset $X \subseteq R^m$ is a definably-connected subset of X which is maximal wrt being definably connected

Example 3.12. $X \subseteq R^m$ definable. Then it is definably connected iff X definably path connected i.e.

$$\forall x, y \in X \exists f : [0, 1] \rightarrow X \text{ definable and continuous with } f(0) = x \wedge f(1) = y$$

Proposition 6.10. Suppose $X \subseteq R^m$ is definable and non-empty, then X has only finitely many definably connected components. The components are both open and closed in X and they form a finite partition of X .

Proof. Let $\{C_1, \dots, C_k\}$ be a partition of X into cells. $I \subseteq \{1, \dots, k\}$ then $C_I := \bigcup_{i \in I} C_i$. Let C' be the maximal among the C_I that is definably connected. Claim: For $Y \subseteq X$ definable connected such that $Y \cap C' \neq \emptyset$ then $Y \subseteq C'$.

proof of claim. $C_Y := \bigcup \{C_i : C_i \cap Y \neq \emptyset\}$ Then $Y \subseteq C_Y$. and C_Y is definably (finite union) connected union of definably connected set Y and finitely many cells that have non-empty intersection with Y . Then $C_Y \cap C'$ contains $Y \cap C' \neq \emptyset$. So if we take $C_Y \cup C'$ has to be again definably connected. By maximality $C_Y \cup C' = C'$ and $Y \subseteq C_Y \subseteq C'$. \square

Hence

- C' definably connected component of X
- The sets C' form a finite partition of X
- C' are the only definable connected components of X

The closure in X of a definably connected subset of X is definably connected. (see topology) So the C' are closed in X . They are also open because the complement in X is a finite union of closed subsets. \square

Note : The above Proposition is not true if we drop the requirement “definable”

Definition 3.19. Definable families: Let $S \subseteq R^{m+n}$ definable. For $a \in R^m$ we put

$$S_a = \{\underline{x} \in R^n : (a, \underline{x}) \in S\} \subseteq R^n$$

S describes the family of sets $(S_a)_{a \in R^m}$. And the sets S_a are called the fibers of S .

Example 3.13. $\mathcal{R} = (\mathbb{R}, <, +, \cdot)$

$$ax^2 + bxy + cy^2 + dx + ey + f = 0$$

defines $S \subseteq R^6 \times R^2$. The fibers are:

- points, circles, ellipse, hyperbola, parabola
- and the limiting cases: \emptyset , 2 lines intersecting each other, 2 parallel lines, one line, RR^2

Note : In o-minimal structures there are only finitely many homomorphism types in a definably family. (If there are infinitely many fibres, then only finitely many are not homeomorphic to each other).

Proposition 6.11. 1. C cell in R^{m+n} , $a \in \pi_m^{m+n} C$ (where $\pi_m^{m+n}(x_1, \dots, x_{m+n}) = (x_1, \dots, x_m)$) Then C_a is a cell in R^n

2. \mathcal{D} decomposition of R^{m+n} , and $a \in R^m$ then

$$\mathcal{D}_a = \{C_a : C \in \mathcal{D} \wedge a \in \pi_m^{m+n}(C)\}$$

is a decomposition of R^m .

Proof. 1. induction on n . If $n = 1$, $a \in \pi_m^{m+1}C$ Then C_a is one of the below

- If C is a $(i_1, \dots, i_m, 0)$ -cell then $C = \Gamma f$, $f : \pi_m^{m+1}C \rightarrow R$ definably continuous. $a \in \pi_m^{m+1}C$ then $C_a = \{f(a)\} \subseteq R$
- If C is a $(i_1, \dots, i_m, 1)$ -cell then $C = (f, g)$, $C_a = (f(a), g(a))$

Suppose the statement holds for some n then let $C \subseteq R^{m+n+1}$ be a cell. Consider the two projections $\pi_{m+n}^{m+n+1}, \pi_m^{m+n}$ and

$$\pi_m^{m+n} \circ \pi_{m+n}^{m+n+1} : R^{m+n+1} \rightarrow R^m$$

Two options: Either $C = \Gamma f$, then

$$C_a = \Gamma f_a \text{ where } f_a : (\pi_{m+n}^{m+n+1}C)_a \rightarrow R$$

and $f_a(x) = f(a, x)$

Or $C = (f, g)_D$ i.e. $f, g : D \rightarrow R$, $D \subseteq R^{m+n}$ cell, $D = \pi_{m+n}^{m+n+1}$ Then $C_a = (f_a, g_a)_E$, $E = D_a$. in both cases, C_a is a cell.

2. Exercise. □

Corollary 6.11-A. Let $S \subseteq R^m \times R^n$ a definable family then there exists $M_S \in \mathbb{N}$ such that for all $a \in R^m$ $S_a \subseteq R^n$ has a partition into M_S many cells.

Proof. $S \subseteq R^m \times R^n$ \mathcal{D} decomposition of $R^m \times R^n$ that partitions S . Then S is a finite union of cells from \mathcal{D} , each fiber S_a is a finite union of $C_a, C \in \mathcal{D}$ but C_a is a cell by Proposition. A bound: $|\mathcal{D}|$. □

Note : There is a uniform bound on $\#$ of definable connected components of sets in definable family.

Theorem 6.12. $\mathcal{R} = (R; <, \dots)$ o -minimal \mathcal{L} -structure, $\mathcal{R}' = (R'; <, \dots)$ \mathcal{L} -structure. If $R \equiv R'$ then R' is o -minimal.

Proof. $S \subseteq R$ definable, $S = \{r \in R : \mathcal{R} \models \varphi(x)[r]\}$ might use parameters from R . If φ is a \mathcal{L} -formula over \emptyset . Then

$$\begin{aligned} \mathcal{R} \models \exists x_1, x_2, x_3 (x_1 \neq x_2 \wedge x_1 \neq x_3 \wedge x_2 \neq x_3 \\ \wedge \forall c ((x_1 < c < x_2 \rightarrow \varphi(c)) \wedge (c = x_3 \rightarrow \varphi(c)) \\ \wedge \neg(x_1 < c < x_2 \vee c = x_3) \rightarrow \neg\varphi(c))) \end{aligned}$$

But if φ uses parameters, we TODO For all $S \subseteq R^{m+1}$ need formula $\forall \underline{a} \in R^m$ “ $S_{\underline{a}}$ is finite union of points and intervals” Idea: Subset of R definable by formula w/ param is just a fiber in a definable family that is parameter-free definable. $S_a \subseteq R$ definable. By the note, there is some number M_S that only depends on the TODO Such that for each $\underline{a} \in R^m$ S_a is a finite union of at most M_S cells.

Then

$$\begin{aligned} \mathcal{R} \models \forall \underline{z} \exists x_1 \dots \exists x_{M_S+1} (\forall y (y < x_1 \rightarrow \varphi_{\underline{z}}(y)) \vee \forall y (y < x_1 \rightarrow \neg\varphi_{\underline{z}}(y))) \\ \wedge (\forall y (x_1 < y < x_2 \rightarrow \varphi_{\underline{z}}(y)) \vee \forall y (x_1 < y < x_2 \rightarrow \neg\varphi_{\underline{z}}(y))) \\ \dots \end{aligned}$$

By elementary equivalence: $\mathcal{R}' \models \dots$ □

CHAPTER 4

Boolean Algebra

From [Kri98]? Our language in this chapter will be $\mathcal{L} = \{0, 1, +, \cdot, \neg\}$, where $+$, \cdot are binary operations and \neg is a unary operation.

The axioms for boolean algebras are

1. (Associativity of $+$, \cdot) $\forall x, y, z (x + (y + z) = (x + y) + z \wedge x \cdot (y \cdot z) = (x \cdot y) \cdot z)$
2. (Commutativity of $+$, \cdot) $\forall x, y (x + y = y + x \wedge x \cdot y = y \cdot x)$
3. (Idempotence) $\forall x (x + x = x \wedge x \cdot x = x)$
4. (Distributivity) $+, \cdot: \forall x, y, z (x \cdot (y + z) = x \cdot y + x \cdot z \wedge x + (y \cdot z) = (x + y) \cdot (x + z))$ =TODO
5. (Absorbtion) $\forall x, y (x \cdot (x + y) = x \wedge x + (x \cdot y) = x)$
6. (De Morgan's Laws) $\forall x, y (\overline{x + y} = \overline{x} \cdot \overline{y} \wedge \overline{x \cdot y} = \overline{x} + \overline{y})$
7. (Laws of 0, 1 and \neg)

$$\forall x (\quad x + 0 = x \quad \wedge \quad x \cdot 0 = 0 \quad \wedge \quad x + 1 = 1 \quad \wedge \quad x \cdot 1 = x \\ \wedge \quad x + \overline{x} = 1 \quad \wedge \quad x \cdot \overline{x} = 0 \quad \wedge \quad \overline{\overline{x}} = x \quad)$$

Definition 4.1. Boolean Algebra: The theory of boolean algebras is the deductive closure of (1)-(7) above.

Note : Every boolean algebra \mathcal{B} can be partially ordered by

$$x \leq y \quad \text{iff} \quad x + y = y$$

It is easy to see that \leq is reflexive, antisymmetric and transitive. In this ordering the smallest set is 0 and the largest one is 1. In this notion the supremum and infimum of two elements are equal to

$$\sup\{x, y\} = x + y, \quad \inf\{x, y\} = x \cdot y \quad (\text{Exercise})$$

Definition 4.2. Alternative Def: Boolean Algebra: A boolean algebra is a set B with

- distinguished elements 0, 1 (called zero and unit of B)
- a unary operation $'$ on B (called **complementation**)
- two binary operations \vee called **join** and \wedge called **meet** s.t. for all $x, y, z \in B$

1. $x \vee 0 = x \quad x \wedge 1 = x$
2. $x \vee x' = 1 \quad x \wedge x' = 0$
3. $x \vee y = y \vee x \quad x \wedge y = y \wedge x$
4. $(x \vee y) \vee z = x \vee (y \vee z) \quad (x \wedge y) \wedge z = x \wedge (y \wedge z)$
5. $x \vee (y \wedge z) = (x \vee y) \wedge (x \vee z) \quad x \wedge (y \vee z) = (x \wedge y) \vee (x \wedge z)$

Example 4.1. Let $X \neq \emptyset$ be a set, $B := \mathcal{P}(X)$ the power set of X , $0 := \emptyset$ and $1 := S$,

$$' : \mathcal{P}(S) \rightarrow \mathcal{P}(S), x' := S \setminus x \quad x \vee y := x \cup y, \quad x \wedge y := x \cap y \text{ for } x, y \in \mathcal{P}(S)$$

Example 4.2. $X \neq \emptyset$ and $S \subseteq \mathcal{P}(X)$ such that

- $\emptyset \in S$
- $X \in S$
- S is closed under (finite) intersections and unions and complements.

Then $(S; \emptyset, X, \cup, \cap, -)$ is called a boolean algebra of sets and \leq corresponds to \subseteq .

Note : Conversely, every boolean algebra is isom. to a boolean algebra of sets. (it can be embedded in)

4.1 STONE REPRESENTATION THEOREM

Definition 4.3. (Ultra-)Filter: Suppose $\mathcal{B} \models BA$. A non-empty subset $F \subseteq B$ is called a filter of \mathcal{B} , if

- $0 \notin F$
- $\forall a \forall b (a \in F \wedge b \in F \rightarrow a \cdot b \in F)$
- $\forall a \forall b (a \in F \wedge a \leq b \rightarrow b \in F)$

An **Ultrafilter** of \mathcal{B} is a filter \mathcal{F} such that $\forall a \in B (a \in \mathcal{F} \vee \bar{a} \in \mathcal{F})$.

Ultrafilter

We define $S(B)$, the stonospace of \mathcal{B} to be $S(B) := \{\text{the set of all ultrafilters of } \mathcal{B}\}$

$S(B)$

Note : $\mathcal{B} \models BA$ then for a filter F .

1. $1 \in F$
2. $F \subseteq B$ satisfying TODO
3. $\langle a \rangle = \{x \in S(B) : a \in x\}$ where a runs through B forms a basis for a topology on the stonospace of \mathcal{B}

Recall: $(X, \tau), \tau \subseteq \mathcal{P}(X)$ is called a topological space, if

T1 $\emptyset, X \in \tau$

T2 $\forall I (\forall (\sigma_i)_i \in \tau^I : \bigcup_{i \in I} (\sigma_i) \in \tau$

T3 $\forall n \in \mathbb{N} (\forall (\sigma_i)_i \in \tau^{\{1, \dots, n\}} : \bigcup_{1 \leq i \leq n} (\sigma_i) \in \tau$

And $\tau' \subseteq \mathcal{P}(X)$ is a base for the topology τ on X , if every open set in τ is the union of sets in τ' .
Back on $\langle a \rangle = \{x \in S(B) : a \in x\}$:

- $\emptyset = \langle 0 \rangle, S(B) = \langle 1 \rangle, \langle a \rangle \cap \langle b \rangle = \langle a \cdot b \rangle$

Every filter on B can be extended to an ultrafilter on \mathcal{B} (Zorn's Lemma).

In fact, suppose $\mathcal{F} \subseteq B$ has FIP. i.e. for any $n \in \mathbb{N} \forall f_1, \dots, f_n \in \mathcal{F}$ it is $f_1 \cdot \dots \cdot f_n \neq 0$. (Exercise)

Definition 4.4. Stone space: A stone space is a non-empty topological space which

1. has a basis of clopen sets
2. is compact (every open cover contains a finite subcover)
3. and hausdorff (every two distinct points can be separated by open sets)

$$\forall x \forall y (x \neq y \rightarrow \exists \sigma_x \exists \sigma_y (x \in \sigma_x \wedge y \in \sigma_y \wedge \sigma_x \cap \sigma_y = \emptyset))$$

Theorem 1.1. Stone Representation Theorem:

1. $\mathcal{B} \models BA$ then $S(B)$ is a Stone-space
2. If S is a Stone space then the clopen subsets of S form a boolean algebra denoted by $\mathcal{B}(S)$.
3. Every boolean algebra \mathcal{B} is isomorphic to the boolean algebra $\mathcal{B}(S(B))$ with $a \mapsto \langle a \rangle$. Hence \mathcal{B} is isomorphic to a subalgebra of boolean algebra $\mathcal{P}(S(B))$

4. Every stonospace S is homeomorphic to the stonospace $S(\mathcal{B}(S))$

$$x \mapsto \{a \in S(B) : x \in a\}$$

Proof. 1. We have the base for a topology $\langle a \rangle = \{x \in S(B) : a \in x\}$. $\langle a \rangle$ is clopen : It is clearly open.

$$\langle a \rangle^c = \overline{\langle a \rangle} = \{x \in S(B) : a \notin x\} = \{x \in S(B) : \bar{a} \in x\} = \langle \bar{a} \rangle$$

hausdorff: Let $x, y \in S(B)$ such that $x \neq y$. then $\exists a \in B$ $a \in x \wedge \bar{a} \in y$

then $x \in \langle a \rangle, y \in \langle \bar{a} \rangle$.

compact: Fact X is topological space then T_{FAE}

Every open cover of X contains a finite subcover if any family of closed sets which has FIP, has non-empty intersection.

Supposed $(F_i)_{i \in I}$ a family of closed subsets of $S(B)$ such that $I \neq \emptyset$ and $\bigcap_{i \in I} F_i = \emptyset$. we want to show that there is a finite intersection $\exists i_1, \dots, i_k \in I$ $\bigcap_{1 \leq m \leq k} F_{i_m}$

WMA that $F_i = \langle a_i \rangle$ for some $a_i \in B$. Assume $\bigcap_{i \in I} \langle a_i \rangle = \emptyset$.

If $a_i \cdot \dots \cdot i_k \neq 0$ For all $\{i_1, \dots, i_k\} \subseteq I$ Then $\{a_i : i \in I\} \subseteq B$ has FIP, so it extends to an ultrafilter on \mathcal{B} (using Zorns lemma).

X set, a filter / ultrafilter on X is some \mathcal{B} boolean algebra, then a filter / ultrafilter of $\mathcal{F} \subseteq \mathcal{P}(X)$ s.th. If \mathcal{F} has FIP then \mathcal{F} extends to UF \mathcal{U} of \mathcal{B} i.e. $\forall i \in I$ $a_i \in \mathcal{U}$ so $\mathcal{U} \in \bigcap_{i \in I} F_i$ but we assumed $\bigcap_{i \in I} \langle a_i \rangle = \emptyset$

So there exists some $i_1 \dots i_k$ such that $\bigcap_{1 \leq j \leq k} \langle a_{i_j} \rangle = \emptyset$

□

Definition 4.5. Atomic, Atomless: An atom is an element of a boolean algebra such that $a \neq 0$ and there is no element in the boolean algebra that is strictly inbetween 0 and a .

$$\forall y(0 \leq y \leq a \rightarrow (y = 0 \vee y = a))$$

A boolean algebra \mathcal{B} is called atomic, if

$$\forall a(a \neq 0 \rightarrow \exists y(y \leq a \wedge y \text{ is atomic}))$$

A boolean algebra is atomless if it contains no atoms

Note : There exists boolean algebras that are neither atomic nor atomless.

Note : Axioms for atomic boolean algebras: add

$$\forall a(a \neq 0 \rightarrow \exists y(y \leq x \wedge y \neq 0 \wedge \forall z(0 \leq z \leq y \rightarrow (z = 0 \vee z = y))))$$

Axioms for atomless: add

$$\forall y y \neq 0 \rightarrow \exists z(0 < z < y)$$

4.2 LINDENBAUM-TARSKI ALGEBRAS

Let \mathcal{L} be a first order Language, \mathcal{L}_0 the set of all \mathcal{L} -sentences and \sim the logical equivalence relation. On the quotient set \mathcal{L}_0/\sim we can define \wedge, \vee, \neg py passing to representatives. This is well defined and does not depend on the choice of representatives.

Definition 4.6. Lindenbaum-Tarski algebra: With the above notation

$$B_L = (\mathcal{L}_0/\sim; \perp/\sim, \top/\sim, \vee, \wedge, \neg)$$

forms then a boolean algebra. (it is called Lindenbaum-Tarski algebra for \mathcal{L})

Note that \perp is logically equivalent to $\exists x x \neq x$ and \top is logically equivalent to $\forall x x = x$

The construction can be extended to equivalence modulo some \mathcal{L} -theory T (or $T \subseteq \mathcal{L}_0$) Let $n \in \mathbb{N}$ and $\underline{x} = (x_1, \dots, x_n)$ For $\varphi, \psi \in \mathcal{L}_{\underline{x}}$, where $\mathcal{L}_{\underline{x}}$ are the \mathcal{L} -formulas with free variables among $\{x_1, \dots, x_n\}$

Define $\varphi \leq_T \psi$ iff $T \models \forall x(\varphi \rightarrow \psi)$

We can define T -equivalence: $\varphi \sim_T \psi$ iff $\varphi \leq_T \psi$ and $\psi \leq_T \varphi$

$$\mathcal{B}_n = (\mathcal{L}_x / \sim_T; \perp / \sim_T, \top / \sim_T, \wedge, \vee, \neg)$$

Is then again a boolean algebra, whose isomorphism type depends only on T and it is called the n -th Lindenbaum-Tarski-algebra of T . In the case we take the 0-th L-T algebra of \emptyset $B_L = B_0(\emptyset^{\models 0})$

Definition 4.7. Recap:

- The deductive closure of a set of sentences Σ is $\{\varphi : \sigma \models \varphi\}$
- A contradiction is any sentence of the form $\varphi \wedge \neg\varphi$
- A set of sentences is consistent, if its deductive closure does not contain a contradiction.
- A \mathcal{L} -theory is a set of sentences that is consistent and deductively closed.

The question we know ask ourselves is: what is the stone space of a Lindenbaum-Tarski algebra?

Note :

- \mathcal{L} -theories are indeed exactly the filters of \mathcal{B}_L
- complete \mathcal{L} -theories are exactly the ultrafilters of \mathcal{B}_L

Let S_L equal the set of all complete \mathcal{L} -theories then our compactness theorem

$$\Gamma \models \varphi \implies \exists \Gamma' \subseteq \Gamma \text{ finite } \Gamma' \models \varphi$$

is equivalent to

Theorem 2.1. Compactness Theorem *: S_L with stone topology is compact.

Two things we would like to show:

- Compactness theorem \implies Compactness theorem *
- By showing that $S_L = S(\mathcal{B}_L)$

proof of claim. \subseteq Let T be complete \mathcal{L} -Theory. by consistency and abuse of notation $0 \notin T$. and T is closed under conjunction. So for all $\varphi, \psi \in T$ we have $\varphi \wedge \psi \in T$. $\varphi \in T$ and $\varphi \leq \psi$ then $\models \varphi \rightarrow \psi$ so $\varphi \models \psi$ and $\psi \in T$ bc. T is deductively closed.

\supseteq Let $x \in S(\mathcal{B}_L)$ completeness: By maximality of x , $\forall \varphi$ either $\varphi / \sim \in x$ or $\neg\varphi / \sim \in x$.
deductively closedness: $x \models \gamma$ then by compactness Theorem $\exists x' \subseteq x, x' \models \gamma$ and $x' \in x$, so by if $x' \in x$ and $x' \leq \gamma$, then $\gamma \in x$ hence deductive closure
consistency: $0 \notin x$ and x is deductively closed.

⊠

- Compactness theorem * \implies compactness theorem $\Gamma = \{\gamma_i : i \in I\}$ set of \mathcal{L} -sentences. We want: $\Gamma \models \varphi$ then $\exists \Gamma' \subseteq \Gamma$ finite $\Gamma' \models \varphi$

proof of claim. Suppose, by contradiction that it is not the case.

$\forall I' \stackrel{\text{fin}}{\subseteq} I \{ \gamma_i : i \in I' \} \cup \{ \neg\varphi \}$ is consistent.

$$\implies \forall I' \stackrel{\text{fin}}{\subseteq} I \bigcap_{i \in I'} \langle \gamma_i \rangle \cap \langle \neg\varphi \rangle \neq \emptyset$$

$$\{ \langle \gamma_i \rangle : i \in I \} \cup \{ \langle \neg\varphi \rangle \}$$

is a collection of closed sets with FIP. By using compactness of S_L with stone topology,

$$\bigcap_{i \in I} \langle \gamma_i \rangle \cap \langle \neg\varphi \rangle \neq \emptyset$$

hence $\Gamma \not\models \varphi$.

⊠

From here on are the lecture notes of last year

As they are in a different notation than this year I will rephrase them after we have discussed them in the lecture.

Lemma 2.2. *Let $(B, ', \vee, \wedge, 0, 1)$ be a boolean algebra. Then it holds*

- a) $0' = 1, 1' = 0$
- b) $x \vee x = x, x \wedge x = x$
- c) $(x')' = x$
- d) $(x \vee y)' = x' \wedge y', (x \wedge y)' = x' \vee y'$
- e) $x \vee y = y$ iff $x \wedge y = x$

Lemma 2.3. a) $x \leq y \Leftrightarrow x \vee y = y$ defines a partial ordering on B (inclusion) and it holds

- b) $x \vee y$ is the least upper bound of $\{x, y\}$ in B
 $x \wedge y$ is the greatest lower bound of $\{x, y\}$ in B
- c) $0 \leq x \leq 1$ for all $x \in B$

Note : A boolean algebra is a complemented distributive lattice.

Definition 4.8. Opposite of boolean algebra: Let $(B, ', \vee, \wedge, 0, 1)$ be a boolean algebra. The boolean algebra B^{op} is defined by

$$B^{\text{op}} := B, \quad 0^{\text{op}} := 1, \quad 1^{\text{op}} := 0, \quad ' \text{ stays the same as for } B, \quad \vee^{\text{op}} := \wedge, \quad \wedge^{\text{op}} := \vee$$

Note: $(B^{\text{op}})^{\text{op}} = B$

Definition 4.9. Subalgebra: A subalgebra of B is a subset $A \subseteq B$ s.t. $0, 1 \in A$ and A is closed under $', \wedge, \vee$. The subalgebra generated by $P \subseteq B$ is defined to be the smallest subalgebra containing P . Equivalently it is the intersection of all Subalgebras of B that contain P .

Example 4.3. Power set algebra: Let S be a set then $\mathcal{P}(S)$ defines a boolean algebra on S . $B := \{x \in \mathcal{P}(S) : x \text{ is finite or cofinite}\}$ is a subalgebra of $\mathcal{P}(S)$ w/ set of generators $\{\{s\} : s \in S\}$

Note : We will prove the Tarski-Stone Theorem: every boolean algebra is isomorphic to an algebra on a set.

Example 4.4. Lindenbaum Algebra of Σ : Let A be a set of prop. atoms, $\text{Prop}(A)$ the set of prop. generated by A . Further let $\Sigma \subseteq \text{Prop}(A)$ and p, q, r range over $\text{Prop}(A)$.

We say p is Σ -equivalent to q iff $\Sigma \models_{\text{taut}} p \leftrightarrow q$. Σ -Equivalence is an equivalent relation on $\text{Prop}(A)$ and $\text{Prop}(A)/\Sigma$ is a boolean algebra with

$$0 := \perp/\Sigma, \quad 1 := \top/\Sigma, \quad (p/\Sigma)' := (\neg p)/\Sigma, \quad (p/\Sigma \vee q/\Sigma) := (p \vee q)/\Sigma, \quad (p/\Sigma \wedge q/\Sigma) := (p \wedge q)/\Sigma$$

a set of generators is $\{a/\Sigma : a \in A\}$

Definition 4.10. Homomorphisms of boolean algebras: Let B, C be boolean algebras. A map $\phi : B \rightarrow C$ is a (homo)morphism of boolean algebras iff $\forall x, y \in B$ it holds

- $\phi(0_B) = 0_C$
- $\phi(x') = \phi(x)'$
- $\phi(x \vee y) = \phi(x) \vee \phi(y)$
- $\phi(x \wedge y) = \phi(x) \wedge \phi(y)$

If $\phi : B \rightarrow C$ is bijective too, we call ϕ an isomorphism and $\phi^{-1} : C \rightarrow B$ is also a morphism of boolean algebras.

Note : $\phi(B)$ is subalgebra of C

Example 4.5. Let S, T be sets then a function $f : S \rightarrow T$ induces a morphism of boolean algebras $\mathcal{P}(T) \rightarrow \mathcal{P}(S) : y \mapsto f^{-1}(y)$. If $S \subseteq T$ and f the inclusion map $S \hookrightarrow T$ then we get a boolean algebra morphism $\mathcal{P}(T) \rightarrow \mathcal{P}(S)$.

• $id_B : B \rightarrow B$ • $x \mapsto x' : B \rightarrow B^{\text{op}}$ are both isomorphism

Note : A boolean algebra morphism $\phi : B \rightarrow C$ is injective iff $\ker \phi = 0_B$

Lemma 2.4. Let $X_1, \dots, X_m \subseteq S$ and \mathcal{A} a boolean algebra on S generated by $\{X_1, \dots, X_m\}$. Then \mathcal{A} is finite and isomorphic to $\mathcal{P}(\{1, 2, \dots, n\})$ for some $n \leq 2^m$.

Proof. TODO □

Definition 4.11. Trivial algebras:

- B is trivial if $|B| = 1$ (equivalently $0 = 1 \in B$) according to 2.4 B is isomorphic to $\mathcal{P}(\emptyset)$
- If $|S| = 1$ then $|\mathcal{P}(S)| = 2$ TODO

Definition 4.12. Ideal: An ideal of B is a subset of $I \subseteq B$ s.t.

(I1) $0 \in I$

(I2) $\forall a, b \in B$ it holds $a \leq b$ and $b \in I \implies a \in I$ and $a, b \in I \implies a \vee b \in I$

Example 4.6. $F_{\text{in}} = \{F \subseteq S : F \text{ finite}\}$ is ideal in $\mathcal{P}(S)$.

Note : If I is an ideal of B then $I \vee b := \{x \in B : x = a \vee b \text{ for some } a \in I\}$ is the smallest ideal w/ respect of \subseteq of B that contains $I \cup \{b\}$.

Example 4.7. • For a boolean algebra morphism $\phi : B \rightarrow C$ the kernel $\ker(\phi)$ is an ideal in B .

- If I is an ideal in B then $a =_I b \iff a \vee x = b \vee x$ for some $x \in I$ defines an equivalent relation and $B/_I$ is a boolean algebra w/

$$0 := 0/_I \quad 1 := 1/_I \quad (a/_I)' := a'/_I \quad a/_I \vee b/_I := (a \vee b)/_I \quad a/_I \wedge b/_I := (a \wedge b)/_I$$

Then $\phi : B \rightarrow B/_I : b \mapsto b/_I$ is a boolean algebra morphism w/ $\ker(\phi) = I$

CHAPTER 5

Set Theory of ZFC

The contents on this chapter are at least partially sourced on [Kri98].

Example 5.1. Russel's paradox: Let $A = \{a : a \notin a\}$. If any collection of elements is a set, then A would be a set. Question: is $A \in A$? if yes, then $A \notin A$, if not then $A \in A$

Trying to resolve this, we will introduce the ZFC (Zermelo-Frankel axioms w/ choice) System. Let $\mathcal{L} = \{\in\}$ be a Language of first order, where $\in \dots$ binary relation "being element of" For (\mathcal{U}, \in) If $\mathcal{A} = (\mathcal{U}, \in^{\mathcal{A}}) \models \text{ZFC}$, then the elements of the universe \mathcal{U} are called sets. We will show roughly that some definably sets are not sets (in the sense of ZFC), others are not. The latter will be called classes.

5.1 FIRST AXIOMS OF ZFC

Definition 5.1. Axiom of extensionality:

$$\forall x \forall y (x = y \leftrightarrow \forall u (u \in x \leftrightarrow u \in y))$$

In other words, two sets are the same if they have the same elements. This will give us later uniqueness in construction of other sets.

Definition 5.2. Pairing Axiom: for any two sets a, b one can form a set whose elements are precisely a, b

$$\forall x \forall y \exists z \forall u (u \in z \leftrightarrow (u = x \vee u = y))$$

Our notation will be $z = \{x, y\}$

In words: For any two sets there exists a set whose members are those two sets.

Note :

- $\{x, y\}$ is unique by 5.1
- $\{x\}$ is a set. from 5.2, take $x = y$

Lemma 1.1. Let x, y be sets. We define the ordered pair $(x, y) := \{\{x\}, \{x, y\}\}$. Then it holds $(x, y) = (a, b)$ iff $x = a$ and $y = b$

Proof. By cases

- if $x = y$, then $(x, y) = \{\{x\}\}$ therefore $a = b$ and by 5.1 it holds $x = a$.
- if $x \neq y$, then $\{\{x\}, \{x, y\}\} = \{\{a\}, \{a, b\}\}$ iff $\{x\} = \{a\}$ and $\{x, y\} = \{a, b\}$. That is, iff $x = a$ and $y = b$.

□

Note : The set (x, y) exists, because its obtained by repeatedly using 5.2

Lemma 1.2. Let x, y, a, b be sets. Then $(x, y) = (a, b)$ iff $x = a$ and $y = b$

Proof. • case $x = y$, then $(x, y) = \{\{x\}\}$ is a singleton then (a, b) is a singleton, wlog $\{a\} = \{a, b\}$ then $a = b = x$.

- case $x \neq y$ and $\{\{x\}, \{x, y\}\} = \{\{a\}, \{a, b\}\}$ then $\{x\} = \{a\}$ and $\{x, y\} = \{a, b\}$ because by 5.1 a singleton can not be equal to a set of size 2.

□

Definition 5.3. n-tuples: Define (x_1, \dots, x_n) inductively:

- (x_1, x_2) already defined
- $(x_1, \dots, x_n) := (x_1(x_2, \dots, x_n))$

Lemma 1.3. For all $n > 1$ $(x_1, \dots, x_n) = (y_1, \dots, y_m)$ iff $n = m$ and $\forall i \leq n \ x_i = y_i$

Proof. Exercise □

Definition 5.4. Union Axiom: For every set x there is a set z consisting of all elements of the elements of x .

$$\forall x \exists z \forall y (y \in z \leftrightarrow \exists u (u \in x \wedge y \in u))$$

We call z the union of x , notation: $\cup_x := z$

Definition 5.5. :

$$x \cup y := \bigcup_{\{x, y\}}$$

Example 5.2. 1. $\bigcup_{(x, y)} = \{x, y\}$.

2. $(x_1, x_2, \dots, x_n) = \bigcup_{\{x_1\}, x_2, \dots, x_n}$

Note :

- $\forall x_1, \dots, x_n$ then there is exactly one set with elements x_1, \dots, x_n
- $x \cup (y \cup z) = (x \cup y) \cup z$

Definition 5.6. Power set Axiom: Let $x \subseteq y$ be the abbreviation for $\forall z (z \in x \rightarrow z \in y)$ The Powerset Axiom states, that for every set x there exists a set z consisting of all subsets $y \subseteq x$ that are themselves sets.

$$\forall x \exists z \forall y (y \in z \leftrightarrow y \subseteq x)$$

Notation: $\mathcal{P}(x) := z$.

Or in words: “For every set x there is a set z consisting of all subcollections of x that are themselves sets.” class relations

5.1.1 Classes and functions

Definition 5.7. Classes: All the unary \mathcal{L} -definable relations (w/ parameters) are called classes.

Example 5.3. • $\varphi(x) : x = x$ defines a class that is not a set

- $\varphi(x) : \exists u (u \in x \wedge \forall v (v \in u \rightarrow v \in x))$

Definition 5.8. Class functions: Suppose we have a formula $\phi(x_1, \dots, x_n, y)$. Then we say ϕ defines a class function R_ϕ iff

$$\forall x_1 \dots \forall x_n \forall y \forall y' ((\phi(x, y) \wedge \phi(x, y')) \rightarrow y = y')$$

We can then define the domain and image of the class function.

$$\text{dom } R_\phi : \{\underline{x} : \exists y \phi(\underline{x}, y)\}$$

$$\text{im } R_\phi : \{y : \exists \underline{x} \phi(\underline{x}, y)\}$$

Note that $R_\phi(\underline{x}) = y$ iff $\phi(\underline{x}, y)$

Definition 5.9. Axiom of replacement / substitution: Let $\varphi(x, y, \underline{a})$ a \mathcal{L} -f.a., w/ free variables among x, y and set-parameters \underline{a} . Suppose φ defines a class function on \mathcal{U} , then the following is an axiom:

$$\forall u \exists z \forall y (y \in z \leftrightarrow \exists x (x \in u \wedge \varphi(x, y, \underline{a})))$$

i.e. the image of a set under a class function is a set.

Definition 5.10. Axiom scheme of comprehension: Let $\psi(x, \underline{a})$ be an \mathcal{L} -formula. Then the following is an axiom:

$$\forall u \exists z \forall v (v \in z \leftrightarrow (v \in u \wedge \forall \psi(v, \underline{a})))$$

i.e. all elements of a set that satisfy a given \mathcal{L} -formula form a set.

Note : 5.10 follows from 5.9

Definition 5.11. Set existence:

$$\exists x x = x$$

i.e. $U \neq \emptyset$. - this is clear when we view it as a universe of a structure.

Note on the existence of the empty set: Let u be any set (there exists one by 5.11), $\psi(x) \equiv x \neq x$ then by 5.10 $\emptyset := \{x \in u : \psi(x)\}$ is a set.

Note : We can derive pairing from replacement 5.9, extensionality, powerset and set existence. From set existence: \emptyset is a set By powerset, replacement(comprehension): $\{\emptyset\}$ is a set.

$$\mathcal{P}(\{\emptyset\}) = \{\emptyset, \{\emptyset\}\}$$

is a set. Then by defining a class function $R_\phi(x) = y$,

$$\phi(x, y) \equiv (x = \emptyset \wedge y = a) \vee (x = \{\emptyset\} \wedge y = b)$$

Note : If the domain of a class function is a set then the graph of the class function is a set. R_ϕ defined by $\phi(x, y, \underline{a})$ then the domain $u := \text{dom } R_\phi \in \mathcal{U}$ Image $v := \text{im } R_\phi \in \mathcal{U}$ by replacement

$$\{(x, y) : x \in u \wedge y \in v \wedge \phi(x, y, \underline{a})\}$$

is the graph of R_ϕ The above would be a set if $u \times v$ which can be shown by using comprehension (exercise).

Definition 5.12. Function: A function $f : a \rightarrow b$ where a, b are sets is a subset of $a \times b$ that satisfies the following

- $\forall x (x \in a \rightarrow \exists y \in b (x, y) \in f)$
- $\forall x \forall y \forall y' (((x, y) \in f \wedge (x, y') \in f) \rightarrow y = y')$

Families of sets and cartesian products

Suppose we have a function $a : I \rightarrow X$. set a_i to the unique $x \in X$ s.th. $(i, x) \in a$

Definition 5.13. :

$$\bigcup_{i \in I} a_i = \{z \in \bigcup X : \exists i \in I z \in a_i\}$$

$$\bigcap_{i \in I} a_i = \{z \in \bigcup X : \forall i \in I z \in a_i\}$$

$$\prod_{i \in I} a_i = \{f : I \rightarrow \bigcup X : \forall i \in I f(i) \in a_i\}$$

Note : If $I = \emptyset$ then $\bigcap_{i \in I} a_i = \bigcup X$

Class relations and well ordering

Types of well ordered sets

Definition 5.14. : Let R class relation, C class Then R defines a strict ordering on C , if

- (i) $\forall x \forall y \forall z (R(x, y) \rightarrow (C(x) \wedge C(y)))$
- (ii) $\forall x \forall y \forall z \neg (R(x, y) \wedge R(y, x))$
- (iii) $\forall x \forall y \forall z (R(x, y) \wedge R(y, z)) \rightarrow R(x, z)$

The ordering is linear, if additionally

- (iv) $\forall x \forall y (C(x) \wedge C(y)) \rightarrow (x = y \vee R(x, y) \vee R(y, x))$

Definition 5.15. Well ordering: Let R be a strict ordering on C x is a set such that $\forall y \in x \ C(y)$ Then x is called well-ordered by R , if

$$\forall \emptyset \neq y \subseteq x \text{ has a smallest element}$$

i.e.

$$\forall y \left((\emptyset \neq y \wedge y \subseteq x) \rightarrow \exists y' (y' \in y \wedge \forall z (z \in y \rightarrow (R(y', z) \vee y' = z))) \right)$$

Definition 5.16. Initial segment: Let x be a set, well ordered by R . Then $y \subseteq x$ is called an initial segment of x , if

$$\forall s \forall t (s \in x \wedge t \in x) \rightarrow ((t \in y \wedge R(s, t)) \rightarrow s \in y)$$

notation: If $y \in x$, x is well-ordered Then $\delta_y(x) = \{z \in x : z < y\} < \text{strict}, \implies y \notin \delta_y(x)$

Note : If x is well-ordered and $y \subseteq x$ then

y is an initial segment of x iff $y = x$ or $y = \delta_z(x)$ for some $z \in x$

Let $y \subseteq x$ an initial segment. Suppose $x \neq y$ that means $x \setminus y \neq \emptyset$ Let z be the smallest element of $x \setminus y$ (exists by well-ordering of x)

Definition 5.17. Proper class: A class C is a proper class if it is not a set. i.e. if C is given by $\phi(x, \underline{a})$ then there is not $z \in \mathcal{U}$ such that $\forall x x \in z$ iff $\phi(x, \underline{a})$

Example 5.4. \mathcal{U} is a proper class If \mathcal{U} was a set then $\{x : x \notin x\}$ would be a set.
the class of all ordinals is a proper class.

Definition 5.18. well-ordering (class): A class relation R defining a strict ordering on a class C is a well-ordering, if
For every $x \in C$ the class initial segment $\delta_x(C) = \{y : R(y, x)\}$ is a set that is well ordered by R

5.2 ORDINALS

Definition 5.19. Transitivity of sets: A set x is called transitive if

$$\forall y (y \in x \rightarrow y \subseteq x)$$

Note : It corresponds to Transitivity of the belonging relation \in . $z \in y \in x \rightarrow z \in x$

Definition 5.20. Ordinal: An ordinal is a transitive set which is well ordered by \in .

Note : The collection of all ordinals form a class relation notation: Ord, On
Proof: Write down formula

Example 5.5. • $\emptyset, \{\emptyset\}, \{\emptyset, \{\emptyset\}\}$ are ordinals

Lemma 2.1. Characterization of ordinals: α ordinal

- the initial segments of α are α itself and the elements of α
- if $\beta \in \alpha$ then β is an ordinal.
- $\alpha \notin \alpha$

Proof. Problem set □

Lemma 2.2. Let $\alpha, \beta \in \text{On}$ TODO then either $\alpha = \beta$, or $\alpha \in \beta$ or $\beta \in \alpha$.

Proof. Let $\gamma := \alpha \cap \beta$ Claim: γ is initial segment of both α and β
proof of claim. $x \in y \in \gamma$ then $x \in y \in \alpha$ and $x \in y \in \beta$. but α, β are ordinals, so $x \in \alpha$ and $x \in \beta$ and $x \in \gamma$ ☒

Then by previous lemma, either

- $\gamma = \alpha$ and $\gamma = \beta$: done

- $\gamma = \alpha$ and $\gamma \in \beta$: $\alpha \in \beta$
- $\gamma \in \alpha$ and $\gamma = \beta$: $\beta \in \alpha$.
- $\gamma \in \alpha$ and $\gamma \in \beta$ we have $\gamma \in \alpha \cap \beta = \gamma$ which is impossible

□

Proposition 2.3. *On is well-ordered by \in*

Proof. We need to show that if $\alpha \in \text{On}$ then $\delta_\alpha(\text{On})$ is a set that is well ordered by \in .

$\delta_\alpha(\text{On}) = \{\beta \in \text{On} : \beta \in \alpha\} = \alpha$ Because α is a well-ordered set, and by the lemma TODO On is linearly ordered by \in □

Lemma 2.4. *On, the class of all ordinals is a proper class, well-ordered by \in*

Proof. Suppose On would be a set z .

On is well ordered by \in On is transitive: $y \in x \in \text{On}$ then $y \in \text{On}$ by lemma so On would be an ordinal and $\text{On} \in \text{On}$ which is not possible by Lemma TODO. □

Note :

- If $\alpha \in \text{On}$ then the initial segments of α are α and the elements of α .
- If $\alpha \in \text{On}$ and $\beta \in \alpha$ then $\beta \in \text{On}$
- $\alpha, \beta \in \text{On}$ then $\alpha \subseteq \beta$ iff $\alpha \in \beta$ or $\alpha = \beta \rightarrow$: If $\alpha \neq \beta$. Then by linear ordering of \in
- $\alpha \subseteq \beta$ iff $\alpha = \beta$ or $\alpha \in \beta$.

Lemma 2.5. : *If $\alpha \in \text{On}$ then $\alpha \cup \{\alpha\} \in \text{On}$ and $\alpha \cup \{\alpha\}$ is the successor of α in the ordering \in*

Proof. • $\alpha \cup \{\alpha\}$ transitive:

$x \in y \in \alpha \cup \{\alpha\}$ if $y \in \alpha$ then $x \in \alpha$ hence $x \in \alpha \cup \{\alpha\}$. else $y = \alpha$ then $x \in \alpha \cup \{\alpha\}$

- $\alpha \cup \{\alpha\}$ well-ordered:
 $\emptyset \neq x \subseteq \alpha \cup \{\alpha\}$ if $x \cap \alpha \neq \emptyset$ then there is $x_0 \in x \cap \alpha$ smallest. $x_0 \in \alpha$ and is smallest element in $\alpha \cup \{\alpha\}$ otherwise $\emptyset \neq x \subseteq \{\alpha\}$ then α is the smallest element.
- $\alpha \cup \{\alpha\}$ is successor of α
 $\alpha \in \alpha \cup \{\alpha\}$ assume $\alpha \in \beta \in \alpha \cup \{\alpha\}$ if $\beta \in \alpha$ then $\alpha \in \beta \in \alpha$, so by transitivity, $\alpha \in \alpha$ which is not possible.
else $\beta = \alpha$

□

Note : If α, β are ordinals then we will use $\alpha \in \beta$, $\alpha < \beta$, $\alpha \subsetneq \beta$ interchangeable.

$\gamma \in \text{On}$ then $\gamma = \{\alpha \in \text{On} : \alpha \in \gamma\}$

Lemma 2.6. *X a set of ordinals, then $\sup X = \bigcup X$ is an ordinal and $\forall \alpha \in X \alpha \subseteq \bigcup X$ and $\bigcup X$ is smallest with this property.*

Proof. • $\bigcup X$ transitive: $x \in y \in \bigcup X$. then $\exists \alpha \in X$ such that $y \in \alpha$. then $x \in \alpha$ hence $x \in \bigcup X$

- $\bigcup X$ well-ordered by \in : $\bigcup X$ contained in On and is a set, but On is well-ordered, so $\bigcup X$ is well-ordered.
- $\alpha \in X$ then $\alpha \subseteq \bigcup X$.
- $\alpha \in X$ then $\alpha \subseteq \bigcup X$. Let $\beta \in \bigcup X$ then there exists $\alpha \in X$ such that $\beta \in \alpha$ so β is not an upper bound for X .

□

Lemma 2.7. *Suppose that $\alpha, \beta \in \text{On}$ and $f : \alpha \rightarrow \beta$ that is strictly increasing i.e. $\forall \gamma, \delta \in \alpha \gamma < \delta \rightarrow f(\gamma) < f(\delta)$ Then $\alpha \subseteq \beta$ and $\forall \gamma \gamma \leq f(\gamma)$*

Proof. By contradiction, Let $\gamma \in \alpha$ be the smallest element with $f(\gamma) < \gamma$ then by minimality of γ , $f(\gamma) \leq f(f(\gamma))$. Because f is strictly increasing $f(f(\gamma)) < f(\gamma)$. So we get $f(\gamma) \leq f(f(\gamma)) < f(\gamma)$ but $f(\gamma) \notin f(\gamma)$ because $f(\gamma) \in \beta$.

Suppose $\beta \in \alpha$ then $f(\beta) < f(\alpha) < \beta$ so $f(\beta) < \beta$, a contradiction. \square

Theorem 2.8. $f : \alpha \rightarrow \beta$ isomorphism between (α, \in) , (β, \in) and $\alpha, \beta \in \text{On}$ then $\alpha = \beta$ and f is unique such isomorphism, hence $f = \text{id}_\alpha$.

Proof. $\alpha = \beta$:

Apply previous lemma to f, f^{-1} hence $\alpha \subseteq \beta$ and $\beta \subseteq \alpha$ uniqueness:

$\gamma \in \alpha$ then $\gamma \leq f(\gamma)$ and $\gamma \leq f^{-1}(\gamma)$ by prev lemma we get $\gamma \leq f(\gamma) \leq \gamma$ so $f(\gamma) = \gamma$ \square

Note : $y \mapsto \beta_y$ for $y \in Y$ is function defined on Y and maps to $Z = \{\beta(x) : x \in Y\}$ (is a set by replacement)

Theorem 2.9. Let $(X, <_X)$ be well-ordered, then there is a unique isomorphism onto an ordinal (α, \in)

Proof. uniqueness:

Suppose we have $f : (X, <_X) \rightarrow (\alpha, \in)$, $g : (X, <_X) \rightarrow (\beta, \in)$ isomorphisms then $f \circ g^{-1}$ and by prev thm: $\alpha = \beta$ and $f \circ g^{-1} = \text{id}_\alpha$ so $f = g$.

Existence:

define $y = \{x \in X : (\delta_x, <_X) \text{ is isomorphic to an ordinal}\}$ where $\delta_x := \delta_x(X)$.

For each $y \in Y$ there is a unique ordinal $\beta_y \in \text{On}$ such that $(\delta_y, <_X)$ and $(\beta(y), \in)$ are isomorphic.

Claim: Y is initial segment of X

proof of claim. If $x <_X y \in Y$, $f : \delta_y \rightarrow \beta(y)$ isomorphism, then f maps $\delta_x \subseteq \delta_y$ to initial segment of β , hence to an ordinal. \boxtimes

$y \mapsto \beta_y$ for $y \in Y$ is function defined on Y and maps to $Z = \{\beta(x) : x \in Y\}$ (is a set by replacement)

Claim: $Z = \{\beta(x) : x \in Y\}$ is an initial segment in On

proof of claim. if $\gamma \in \beta(x)$, $x \in Y$ have isomorphism between $(\delta_x, <_X)$ and (β_x, \in) so its preimage y and $(\delta_y, <_X)$ is mapped to the initial segment determined by gamma, so to (γ, \in) hence $(\delta_y, <_X)$ isom to γ \boxtimes

So Z is initial segment of On and Z is a set. So $\alpha := Z$ is itself an ordinal and $y \mapsto \beta_y$ isomorphism between Y and α .

Assuming $Y \subsetneq X$, then there is a minimal $x_0 \in X \setminus Y$ $\delta_{x_0} = Y$ (Y is initial segment) $Y \cong \alpha$ hence $x_0 \in Y$, a contradiction. \square

5.2.1 Transfinite induction

Suppose $\phi(x)$ (possibly with parameters) to prove

$$\forall \alpha \in \text{On} \phi(\alpha) \text{ iff } \forall \alpha \forall \beta ((\beta < \alpha \rightarrow \phi(\beta)) \rightarrow \phi(\alpha)) \quad (5.1)$$

5.1 $\implies \forall \alpha \in \text{On} \phi(\alpha)$

Suppose there is $\alpha \in \text{On}$ such that $\neg \phi(\alpha)$ then let α be smallest with the property.

proof by induction on ordinals (proof by transfinite induction) is a proof of $\forall \alpha \in \text{On} \phi(\alpha)$ by proving 5.1

Let F be a class function in one variable and a a set contained in $\text{dom}(F)$ Then $F|_a = \{(x, y) \in (a, b) : F(x) = y\}$ where $b = \{F(x) : x \in a\}$ (which is a set by replacement).

Let H be any class function in one variable.

Definition 5.21. H-inductive: A function f is called H-inductive, if 1. $\alpha := \text{dom}(f) \in \text{On}$ and 2. $\forall \beta \in \alpha f|_\beta \in \text{dom}(H)$ and 3. $f(\beta) = H(f|_\beta)$

$f : \alpha \rightarrow X$ then H gives you a way to extend the f . H extends f to a function on $\alpha \cup \{\alpha\}$

$$f(\alpha) = H(f)$$

Lemma 2.10. For every class function H and ordinal α there is at most one H-inductive function on α with domain α .

Proof. Suppose not. $f, g : \alpha \rightarrow X$ different H-inductive functions. Let x_0 the smallest element of α such that $f(x_0) \neq g(x_0)$. By x_0 smallest, $f|_{x_0} = g|_{x_0}$ By H-inductiveness

$$f(x_0) = H(f|_{x_0}) = H(g|_{x_0}) = g(x_0)$$

A contradiction. □

Lemma 2.11. *Let H be a class function, $\alpha \in \text{On}$ such that any function $f : \beta \rightarrow X$ where $\beta \in \text{On}$ belongs to $\text{dom}(H)$ then there is an H -inductive function $f : \alpha \rightarrow X$.*

Proof. $\tau = \{\beta < \alpha : \text{there is H-inductive } f_\beta : \alpha \rightarrow X\}$ τ is a set and initial segment of α hence $\tau \in \text{On}$ and $\tau \subseteq \alpha$

$\beta \mapsto f_\beta$ for $\beta \in \tau$ is well-defined function by uniqueness of f_β .

Moreover for $\gamma < \beta < \tau$ we have $f|_\beta|_\gamma = f|_\gamma$ (H-ind, uniqueness)

$f := \bigcup_{\beta < \tau} f_\beta$ is H-inductive function (graphs agree on intersection, each of the f_β are H-ind).

The domain

$$\text{dom}(f) = \sup_{\beta < \tau}(\beta) = \bigcup_{\beta < \tau}(\beta) = \sigma \in \text{On}$$

If $\sigma = \alpha$ we are finished, otherwise we can define

\tilde{f} such that $\tilde{f}|_\sigma = f$ and $\tilde{f}(\sigma) = H(f)$ \tilde{f} is now H-inductive, and $\text{dom}(\tilde{f}) = \sigma \cup \{\sigma\}$ a contradiction σ , the domain of f . □

Theorem 2.12. Transfinite Recursion: *Let A be a class, M be a class of all functions $f : \alpha \rightarrow X$ for $\alpha \in \text{On}$ arb. X a subset of A and H a class function in one variable defined on all of M with values in A . Then there exists a unique class function F defined on On such that $\forall \alpha F(\alpha) = H(F|_\alpha)$*

Proof. define F by

$$F(\alpha) = y \text{ iff there is an H-inductive function } f : \alpha \rightarrow X, X \text{ subset of } A \text{ and } y = H(f)$$

It is well defined by the prev two lemmas TODO □

5.2.2 Axiom of Choice

Definition 5.22. Axiom of Choice (AC): For every set X and $A \subseteq \mathcal{P}(X)$ that consists of pairwise disjoint, non-empty subsets of X there is a set $T \subseteq X$ such that $\forall a \in A \# a \cap T = 1$. T as above is called a transversal.

Definition 5.23. (AC'): (Existence of choice function.) For every set X there exists a function $\pi : \mathcal{P} \setminus \{\emptyset\} \rightarrow X$ such that for every non-empty subset $a \subseteq X$ $\pi(a) \in a$

Definition 5.24. (AC''): Let $(X_i)_{i \in I}$ be an indexed family of non-empty sets then $\prod_{i \in I} X_i \neq \emptyset$. ($(X_i)_{i \in I}$ could also be thought of as a function $i \mapsto X_i : I \rightarrow X$)

In the next Problemset: $AC \leftrightarrow AC' \leftrightarrow AC''$

Theorem 2.13. Zermelo: “Well-ordering theorem”: Every set can be well-ordered.

Proof. By contradiction. Suppose X is a set which can not be well-ordered. Choose a choice function $\pi : \mathcal{P}(X) \setminus \{\emptyset\} \rightarrow X$ define a class function H by $H(f) = y$ iff f is a function with (i) $\text{dom } f = \alpha \in \text{On}$ (ii) $\text{im } f \subsetneq X$ (iii) $y = \pi(X \setminus \text{im } f)$

note:

- H is defined on class of all H-inductive functions, whose image is a proper subset of X . $\text{im } f \subsetneq X$.
- each H-inductive function is injective

If f is a H -inductive function that is also surjective, then we are done because f induces well-ordering on X .

By our assumption, every H -inductive function has to be not surjective. So H is defined on all H -inductive functions, and we can use Transfinite recursion theorem and get an H -inductive class function $F : \text{On} \rightarrow X$ which is injective

Suppose $\alpha < \beta < \gamma$ and $F(\alpha) = F(\beta)$. $F(\alpha) = \pi(X \setminus \text{im } F|_\alpha)$ $F(\beta) = \pi(X \setminus \text{im } F|_\beta)$

Then F is an injection from a proper class into $\text{im } F$, a set, which is impossible. \square

Theorem 2.14. Zorn's Lemma: *Let (X, \leq) be a partially ordered set (poset) such that all linearly ordered subsets (called chains) have an upperbound. Then (X, \leq) has a maximal element. i.e. $\exists y \in X \forall x \in X y \not< x$*

Proof. Let $A := \{Y \subseteq X : \exists x \in X \forall y \in Y y < x\}$ Take $\pi : \mathcal{P}(X) \setminus \{\emptyset\} \rightarrow X$ a choice function.

Define

$$p : A \rightarrow X, \quad p(Y) := \pi(\{x \in X : \forall y \in Y y < x\})$$

Define a class function H by $H(f) = y$ iff (i) f is a function with $\text{dom } f \in \text{On}$ (ii) $\text{im } f \in A$ (iii) $y = \pi(\text{im } f)$

We get

- any H -inductive $f : \alpha \rightarrow X$ is strictly increasing. $(\star) f : \alpha \rightarrow X, f(\alpha \cup \{\alpha\}) = H(f) > \text{im } f$
- The image of any H -inductive function $f : \alpha \rightarrow X$ is linearly ordered, so by assumption has an upperbound.
i.e. $\exists x_f \in X \forall \beta < \alpha f(\beta) < x_f$

The idea now is similar to above theorem. Suppose $f : \alpha \rightarrow X$ is H -inductive but H is not defined on f , then the image of f has no strict majorant, so there is $\beta < \alpha$ such that $x_f = f(\beta)$. Then x_f has to be maximal for X .

If H is actually defined on all H -inductive functions, then there is an H -inductive class function $F : \text{On} \rightarrow X$. F is strictly increasing by \star . So have injective of proper class into set. a contradiction. \square

5.2.3 Ordinal arithmetic

Or how to think of the natural numbers to be contained in On

Definition 5.25. :

- $0 := \emptyset$ (smallest ordinal)
- $\beta \in \text{On}$ then its successor ordinal is $\beta + 1 := \beta \cup \{\beta\}$
- (β is a successor ordinal if there is $\alpha \in \text{On}$ such that $\beta = \alpha + 1$)
- $\beta \in \text{On}$ is a limit ordinal if $\beta \neq 0$ and β is not a successor ordinal
- $\beta \in \text{On}$ is a natural number / finite ordinal, if $\beta = 0$ or for every $\alpha \leq \beta$ α is a successor ordinal or $\alpha = 0$

Example 5.6. $0 = \emptyset, 1 = \emptyset \cup \{\emptyset\} = \{\emptyset\}, \dots$

Note : $(X, <_X), (Y, <_Y)$ well-ordered then we can also order their disjoint union $(X \times \{0\}) \cup (Y \times \{1\})$, $X \times Y$ can be well ordered as follows:

$$(a, i) < (b, j) \text{ iff } \begin{cases} i = j = 0 & a <_X b \\ \vee i = j = 1 & a <_Y b \\ \vee i < j \end{cases}$$

and we will call this the ordering on the sum of X and Y .

$(x_0, x_1) < (x_1, y_1)$ iff $y_0 <_Y y_1 \vee (y_0 = y_1 \wedge x_0 <_X x_1)$ (reverse lexicographical ordering)

Definition 5.26. : Let $\alpha, \beta \in \text{On}$ then

- (i) $\alpha + \beta$ is the unique $\gamma \in \text{On}$ such that γ is order-isomorphic to the sum of α and β

(ii) $\alpha \cdot \beta$ is the unique $\gamma \in \text{On}$ such that γ is order-isomorphic to the product of α and β

Properties of $+$ and \cdot :

Lemma 2.15. 1. $+$ is associative, 0 is a 2-sided add. identity

2. \cdot is associative

3. $\alpha \cdot 0 = 0$, $\alpha \cdot 1 = \alpha = 1 \cdot \alpha$

4. $\alpha \cdot (\beta + \gamma) = \alpha \cdot \beta + \alpha \cdot \gamma$

5. λ limit ordinal then $\alpha \cdot \lambda = \sup_{\beta < \lambda} \alpha \cdot \beta$

Proof. Problem set □

Note : Right now it would be consistent to assume \cdot is commutative, but no longer after the next axiom

Definition 5.27. Axiom of infinity: There exists an infinite ordinal.
i.e. there exists an ordinal that is not a natural number.

Note :

- The natural numbers form an initial segment in On .
- Let ω be the smallest infinite ordinal, i.p. ω is a limit ordinal

Example 5.7. 1. $\omega \cdot 2 \stackrel{?}{=} 2 \cdot \omega$, observations:

- $\omega \cdot 2 = \omega + \omega$
- $2 \cdot \omega = \omega$
- $\omega + \omega \neq \omega$

2. $\omega + 2 \neq 2 + \omega$, observations:

- $2 + \omega = \omega$
- $\omega + 2$ has maximal element

Exponentiation

Definition 5.28. : α^β is defined recursively on β :

1. $\alpha^0 = 1$
2. $\alpha^{\beta+1} = \alpha^\beta \cdot \alpha$
3. $\alpha^\lambda = \sup_{\delta < \lambda} \alpha^\delta$ for a limit ordinal λ

Note : can also define exponentiation as given by the class function $EXP : \text{On} \times \text{On} \rightarrow \text{On}$ defined by

$$Exp(\alpha, \beta) = \gamma \text{ iff}$$

There exists a function $f : \beta + 1 \rightarrow \text{On}$ s.th. $\forall \xi < \beta$

- $f(\beta) = \gamma$
- if $\xi = 0$ then $f(\xi) = 1$
- if $\xi = \delta + 1$ then $f(\xi) = f(\delta) \cdot \alpha$
- if ξ is a limit ordinal then $f(\xi) = \sup_{\delta < \xi} f(\delta)$

TODO uniqueness follows from recursion theorem.

Cardinals

Definition 5.29. : Given a set X , the cardinality of X (denoted by $|X|$, $\text{card}(X)$) is the smallest ordinal for which there is a bijection with X .

Note : Every set has a cardinality. This is bc every set can be well-ordered (by Zermelo's Theorem, which is equivalent to AC) and then we get an order-preserving bijection. In fact the statement " $\text{card}(X)$ is defined for each set X " is equivalent to AC.

Definition 5.30. : We call two sets X, Y equinumerous, if there is a bijection between them.

X, Y equinumerous if $|X| = |Y|$

Theorem 2.16. Let X, Y be non-empty sets Then the following is equivalent.

- (i) there is an injection of X into Y
- (ii) there is a surjection of Y into X
- (iii) $|X| \leq |Y|$

Proof. " $i \implies ii$ " don't need AC " $ii \implies i$ " need AC

□

Theorem 2.17. Cantor-Schröder-Bernstein: $|X| = |Y|$ iff there is an injection $X \rightarrow Y$ and an injection $Y \rightarrow X$

Proof.

□

Proof is clear with AC, but can proof it without AC need to do some kind of back and fourth argument

Theorem 2.18. Cantor: For every set X we have $|\mathcal{P}(X)| > |X|$

Proof. Suppose its not, then there exists a set X with $|X| \geq |\mathcal{P}(X)|$.

We can find a surjection $\pi : X \rightarrow \mathcal{P}(X)$ $Y = \{x \in X : x \notin \pi(x)\}$ Let $y \in X$ be such that $\pi(y) = Y$ Either $y \in Y$ or $y \notin Y$ If $y \in Y$ then by definition of Y , $y \notin Y$ If $y \notin Y$ then by definition $y \in Y$ □

Note this says: there is no largest set.

5.2.4 Cardinals

Definition 5.31. Cardinal: $\kappa \in \text{On}$ is called a cardinal, if $\kappa = |\kappa|$

Note : The class Card of all cardinals is a proper class.

proof of claim. Suppose Card is a set. then $\sup \text{Card} = \gamma \in \text{On}$ Then $|\mathcal{P}(\gamma)| > |\gamma| \geq |\lambda|$ for every cardinals $\lambda \in \text{Card}$ that is a contradiction □

Definition 5.32. finite sets: A set X is called finite, if $|X|$ is a finite ordinal, otherwise X is called infinite.

Note : In particular a set is finfinite iff ω injects into it.

Proposition 2.19. Galileo: A set a is infinite iff it properly injects into itself.

Proof. \implies Suppose a is infinite, then by note use ω injects into it show ω injects properly in it self

\Leftarrow want: set finite then it does not inject properly into itself show: every finite ordinal is a cardinal inductively, then $\kappa = |\kappa|$

□

\aleph -function

Note : Card is cofinal in On For every $\alpha \in \text{On}$ there is $\gamma \in \text{Card}$ with $\alpha < \gamma$

And Card is a proper subclass of On , well ordered by the same ordering \in as in On

There is a unique function (class function) from the class of all ordinals to the infinite cardinals that preserves $\in \dots$ this function is called \aleph

Definition 5.33. : Instead of $\aleph(0)$ we will write subscript \aleph_0

- $\aleph_0 = |\omega|$
- $\aleph_{\alpha+1}$ is the smallest cardinal larger than \aleph_α

Note : For every cardinal κ there is a smallest cardinal κ^+ such that $\kappa < \kappa^+$ this is not the successor in the sense of the ordinals. It is

- $n^+ = n + 1$
- $\aleph_\alpha^+ = \aleph_{\alpha+1}$

Note : For every ordinal γ we have $|\gamma| \leq \gamma < |\gamma^+|$

Similarly as with ordinals we will call cardinals of the form κ^+ successor cardinals. and non-zero, non-successor cardinals will be called limit cardinals.

successor cardinals
limit cardinals

Proposition 2.20. *The function \aleph is continuous with respect to the interval topology induced by \in . i.e. If λ is a limit ordinal then $\aleph_\lambda = \sup_{\delta < \lambda} \aleph_\delta$*

Proof. let $\gamma := \sup_{\delta < \lambda} \aleph_\delta$ have $|\gamma| \leq \gamma < |\gamma^+|$ Assume $\gamma < \aleph_\lambda$ then there exist a $\xi_0 < \lambda$ such that $|\gamma| = \aleph_{\xi_0}$

Then we would have $\gamma < |\gamma^+| = \aleph_{\xi_0+1} \leq \sup_{\delta < \lambda} \aleph_\delta$ which is a contradiction.

we have shown that $\aleph_\lambda \leq \sup_{\delta < \lambda} \aleph_\delta$ (TODO: other direction?) \square

Definition 5.34. countable sets: A set X is called to be countable iff $|X| \leq \aleph_0$ and we say X is uncountable otherwise.

Continuum hypothesis (CH)

Hypothesis 2.21. (CH) :

$$|\mathcal{P}(\omega)| = \aleph_1$$

i.e. there is no cardinality between $|\mathbb{R}|$ and $|\mathbb{N}|$ or $|\mathbb{R}|$ is the first uncountable cardinality.

It can be shown that CH is independent of ZFC. (method to show this is called forcing, very popular method)

Cardinal arithmetic

Definition 5.35. : Let κ, λ be cardinals then $\kappa \otimes \lambda := |\kappa \times \lambda|$ $\kappa \oplus \lambda := |(\kappa \times \{0\}) \cup (\lambda \times \{1\})|$

Note : \oplus, \otimes are commutative and associative.

Theorem 2.22. *If κ is infinite, then $\kappa \otimes \kappa = \kappa$*

Proof. By induction on κ . What is ment by that is induction on α where $\aleph_\alpha = \kappa$.

Base case: We should check it for ω $\aleph_0 \otimes \aleph_0 = \aleph_0$ is like finding bijection of ω^2 onto ω . (diagonal)

Now suppose: $\forall \beta < \kappa |\beta| \otimes |\beta| = |\beta|$

On The cartesian product define order such that we get order isomorphism, then this is also a bijection.

Define ordering on $\kappa \times \kappa$

$$(\alpha, \beta) \prec (\alpha', \beta') \iff \begin{cases} \max\{\alpha, \beta\} < \max\{\alpha', \beta'\} \\ \max\{\alpha, \beta\} = \max\{\alpha', \beta'\} \text{ and } \alpha < \alpha' \\ \max\{\alpha, \beta\} = \max\{\alpha', \beta'\} \text{ and } \alpha = \alpha' \text{ and } \beta < \beta' \end{cases}$$

Claim: \prec is a well-ordering (Exercise)

Claim: $(\kappa \times \kappa, \prec)$ is order-isomorphic to (κ, \in) From the claim we immediately get $|\kappa \times \kappa| = |\kappa|$
proof of claim. $\kappa \times \kappa = \bigcup_{\alpha < \kappa} \alpha \times \alpha$ increasing union. then $\{\xi \times \xi\}$ is an initial segment of $\kappa \times \kappa$ (with respect to \prec) Consider $(\xi \times \xi, \prec)$ then by our inductive assumption this will have to be isomorphic to some ordinal $\gamma \in \text{On}$. We know that the cardinality of that ordinal γ $|\gamma| = |\xi \times \xi| \stackrel{\text{ind. Hyp.}}{=} |\xi| < |\kappa|$
So also $|\gamma| < |\kappa|$. so then $(\kappa \times \kappa, \prec)$ is order-isomorphic to (κ, \in) \boxtimes

\square

Corollary 2.22-A. *For every infinite cardinal κ we have $\kappa \oplus \kappa = \kappa$*

Proof. Proof of corollary: $\kappa \oplus \kappa = |\kappa \times 2| \leq |\kappa \times \kappa| = |\kappa|$ □

Definition 5.36. : Define for κ, λ cardinals $\kappa^\lambda := |\{f : f \text{ is a function from } \lambda \text{ to } \kappa\}|$

Note $2^\kappa = |\mathcal{P}(X)|$

Lemma 2.23. If $\lambda \geq \omega$ and $2 \leq \kappa \leq \lambda$ then $\kappa^\lambda = 2^\lambda$.

Proof.

$$2^\lambda = 2^{\lambda \otimes \lambda} = 2^{\lambda^\lambda} \geq \lambda^\lambda \geq \kappa^\lambda$$

$2^{\lambda^\lambda} \geq \lambda^\lambda$ think of function that constantly maps to 0 is part TODO image 5 The other implication is obvious. □

Definition 5.37. : A function $f : \alpha \rightarrow \beta$, where α, β are ordinals is said to be cofinal, if $\text{im } f$ is unbounded in β i.e.

$$\forall \gamma \in \beta \exists \xi \in \alpha \gamma \leq f(\xi)$$

The cofinality of $\beta \in \text{On}$ (denoted by $\text{cof}(\beta)$) is the smallest ordinal α such that there exists a function $f : \alpha \rightarrow \beta$ that is cofinal. Note: $\text{cof}(\beta) \leq \beta$,

Example 5.8. $\text{cof}1 = \text{cof}(\{\emptyset\}) = 1$ and in more generality $\text{cof}(\alpha + 1) = \text{cof}(\alpha \cup \{\alpha\}) = 1$

Note : $\text{cof}(\beta)$ is always a cardinal. Let $\beta \in \text{On}$ and let $f : \text{cof}(\beta) \rightarrow \beta$ be cofinal. Then $|\text{cof}(\beta)| \leq \text{cof}(\beta)$ and there exists a bijection $h : |\text{cof}(\beta)| \rightarrow \text{cof}(\beta)$, so $f \circ h$ yields cofinal map and by minimality of $\text{cof}(\beta)$ have $|\text{cof}(\beta)| = \text{cof}(\beta)$

If β limit ordinal then there is a strict increasing cofinal map $h : \text{cof}(\beta) \rightarrow \beta$ Let $f : \text{cof}(\beta) \rightarrow \beta$ be cofinal define $h : \text{cof}(\beta) \rightarrow \beta$ by

$$h(\xi) := \max\{f(\xi), \sup_{\gamma < \xi} (h(\gamma) + 1)\}$$

Proposition 2.24. Suppose α, β are limit ordinals, $f : \alpha \rightarrow \beta$ strictly increasing and cofinal. Then $\text{cof}(\alpha) = \text{cof}(\beta)$

Proof. One side is obvious

- $\text{cof}(\alpha) \geq \text{cof}(\beta)$ is clear
- $\text{cof}(\alpha) \leq \text{cof}(\beta)$ Let $g : \text{cof}(\beta) \rightarrow \beta$ be cofinal define $h : \text{cof}(\beta) \rightarrow \alpha$ by

$$h(\xi) = \min\{\gamma < \alpha : f(\gamma) > g(\xi)\}$$

cofinal in α

□

Corollary 2.24-A. For a limit ordinal α , $\text{cof}(\aleph_\alpha) = \text{cof}(\alpha)$

Proof. Use 2.20 and 2.24 on $f = \aleph$ □

Corollary 2.24-B. For every ordinal β , $\text{cof}(\beta) = \text{cof}(\text{cof}(\beta))$

Proof. By cases

- $\beta, \text{cof}(\beta)$ are limit ordinals: there exists a strictly increasing map $\text{cof}(\beta) \rightarrow \beta$ result follows from 2.24
- $\beta, \text{cof}(\beta)$ are not limits then $\beta = \alpha \cup \{\alpha\}$, $\text{cof}(\beta) = 1$ and $\text{cof}(\text{cof}(\beta)) = \text{cof}(1) = 1$

□

Definition 5.38. regular ordinals: An ordinal β is called regular, if $\text{cof}(\beta) = \beta$ (it is fixed point of cofinality map)

Note : regular ordinals are cardinals and the first regular, infinite cardinal is ω

Lemma 2.25. κ^+ is regular for $\kappa \geq \omega$

Proof. If $f : \alpha \rightarrow \kappa^+$ is cofinal, then $\kappa^+ = \bigcup_{\gamma < \alpha} f(\gamma) = \sup_{\gamma < \alpha} f(\gamma)$. Each $f(\gamma)$ is an ordinal of cardinality less than κ^+ hence is less or equal than κ

$$\kappa^+ = \left| \bigcup_{\gamma < \alpha} f(\gamma) \right| \leq |\alpha \times \kappa| \leq \max\{|\alpha|, |\kappa|\}$$

because $\kappa < \kappa^+$ we have $\kappa^+ \leq |\alpha| = \alpha$ TODO check □

Note : α limit ordinal then $\text{cof}(\aleph_\alpha) = \alpha$

If \aleph_α is regular then $\text{cof} \aleph_\alpha = \text{cof} \alpha \leq \alpha \leq \aleph_\alpha$ So $\alpha = \aleph_\alpha$

Definition 5.39. : Let κ be a cardinal.

- κ is called weakly inaccessible, if α is a regular limit cardinal strictly greater than ω
- κ is called (strongly) inaccessible, if $\kappa > \omega$, κ regular and for every $\lambda < \kappa$ we have $2^\lambda < \kappa$

We will see that κ inaccessible implies that κ is not a union of fewer than κ sets each of cardinality less than κ **Note :** Existence of inaccessible cardinals does not follow from ZFC.

Why care then?

Axiom of existence of inaccessible cardinals:

$$(IC) \text{inaccessible cardinals exist} \quad (5.2)$$

(AC) has less desirable consequences e.g. Banach-Tarski-Paradox

If we drop AC we could not prove that the lebesgue measure is countably additive, so we may want to replace AC by something weaker (e.g. DC, dependent choice). Various nice results that depend on AC still hold.

LM the axiom that states: every set of reals is lebesgue measurable

Theorem 2.26. *Con... consistency. $\text{Con}(ZF+DC+LM)$ is equal to $\text{Con}(ZF + IC)$*

Note: We do now that ZF can not prove IC, but we don't know yet if ZF can prove the negation of IC.

Lemma 2.27. *If κ is an infinite cardinal and $\lambda \geq \text{cof}(\kappa)$ then $\kappa^\lambda > \kappa$*

Proof. Fix a cofinal map $f : \lambda \rightarrow \kappa$. Consider any function $G : \kappa \rightarrow \kappa^\lambda$. It suffices to show that G can not be surjective.

κ^λ is technically the set of functions $\lambda \rightarrow \kappa$ define $h : \lambda \rightarrow \kappa$ by

$$h(\xi) = \min\{\alpha \in \kappa : G(\alpha)(\xi) \leq f(\xi)\}$$

If $h \in \text{im } G$ then there is $\alpha \in \kappa$ s.t. $G(\alpha) = h$ pick $\xi < \lambda$ s.t. $f(\xi) \geq \alpha$ Then $h(\xi) = G(\alpha)(\xi)$ but by construction of h , $G(\alpha)(\xi) \leq f(\xi)$ so contradiction. □

Corollary 2.27-A. *If $\lambda \geq \omega$ then $\text{cof}(2^\lambda) > \lambda$*

Proof. We know that $(2^\lambda)^\lambda = 2^{(\lambda \otimes \lambda)} = 2^\lambda$ So if $\text{cof}(2^\lambda) \leq \lambda$ then by lemma $(2^\lambda)^\lambda > 2^\lambda$, a contradiction. □

Theorem 2.28. (König): *Let I be a set, $(A_i)_{i \in I}$ and $(B_i)_{i \in I}$ indexed families of sets.*

If $\forall i \in I |A_i| < |B_i|$ then $|\bigcup_{i \in I} A_i| < |\prod_{i \in I} B_i|$

Proof. Exercise with hints. □

Note: above thm is equivalent to AC

One can use König theorem to show that: “ κ inaccessible implies that κ is not a union of fewer than κ sets each of cardinality less than κ ” TODO: check if actually true

Next time : ZF can not proof ZFC

Today we are going to state our last axiom, the axiom of foundation. We are also going to show a relative consistency theorem. We are going to finish prob next course thursday

Definition 5.40. Axiom of foundation (AF): Being an element of, does not admit an infinite decreasing chain.

$$\forall x (x \neq \emptyset \rightarrow \exists y (y \in x \wedge \forall z (z \in y \rightarrow z \notin x)))$$

Note : Suppose we have a sequence of sets $(u_n)_{n \in \omega}$ s.t. $\forall n u_{n+1} \in u_n$ then $\{u_n : n \in \mathbb{N}\}$ would contradict AF. and have $\forall x (x \notin x)$ too.

Definition 5.41. : Class function $V : \text{On} \rightarrow \mathcal{U}$ by transfinite induction on ordinals and set $V_\beta := \bigcup_{\alpha < \beta} \mathcal{P}(V_\alpha)$ $V_0 = \emptyset$ If $\alpha \leq \beta$ then $V_\alpha \subseteq V_\beta$ (an example of increasing sequence, in contrast to note on axiom AF) $V_{\beta+1} = \mathcal{P}(V_\beta)$

If λ is a limit ordinal then $V_\lambda = \bigcup_{\alpha < \lambda} V_\alpha$

V is also a class given by $V(x) \equiv \exists \alpha \in \text{On } x \in V_\alpha$

V also gives us a way to associate a rank to each set.

Definition 5.42. : For every x such that $V(x)$ we define $\text{rk}(x) := \min\{\alpha : x \in V_\alpha\}$

Note : The rank $\text{rk}(x)$ is always a successor ordinal.

Lemma 2.29. $V(x)$ iff $\forall y y \in x \rightarrow V(y)$ And also if $V(x)$ then $\forall y (y \in x \rightarrow \text{rk}(y) < \text{rk}(x))$

Proof. “ \implies ” direction: $V(x)$ and $\text{rk}(x) = \beta + 1$ then $x \in V_{\beta+1}$ so $x \subseteq V_\beta$ hence $\forall y y \in x \rightarrow y \in V_\beta$ Also $\text{rk}(y) \leq \beta < \beta + 1 = \text{rk}(x)$

“ \impliedby ”-direction: Suppose $\forall y y \in x \rightarrow V(y)$ Note that $\text{rk} : V \rightarrow \text{On}$ is bounded on x . Else, $\{\text{rk } y : y \in x\}$ is unbounded in On and it is a set (image of function rk of a set) Take $\bigcup \{\text{rk } y : y \in x\} \in \text{On}$ and it would be an ordinal bigger than any other set, TODO

Suppose $\{\text{rk } y : y \in x\}$ is bounded by β then $\forall y y \in x \rightarrow y \in V_\beta$ so $x \in V_{\beta+1}$. \square

Lemma 2.30. For every ordinal $\alpha \in \text{On}$ we have $V(\alpha)$ and $\text{rk}(\alpha) = \alpha + 1$.

Proof. Exercise \square

Definition 5.43. inductive closure: For any set x we define the function with domain ω $f(0) := x$ $f(n+1) := \bigcup_{y \in f(n)} y = \bigcup f(n)$ and we define the closure of x to be the union $\text{cl } x := \bigcup_{n < \omega} f(n)$

Note :

- $x \subseteq \text{cl } x$
- cl is transitive
- If z is transitive set that contains x then $\text{cl } x \subseteq z$ ($\text{cl } x$ is the unique transitive closure of x)

Theorem 2.31. (AF) holds iff $\forall x V(x)$

Proof. “ \impliedby ”-direction: Suppose $\forall x V(x)$. We need to show that any set contains an element Let $a \neq \emptyset$. Let $y \in a$ be of minimal rank. Then for every $c \in y$ we know $\text{rk } c < \text{rk } y$ so $c \notin a$ by minimality of $\text{rk } y$. So $y \in a$ s.t. $y \cap a = \emptyset$

“ \implies ” direction: Lets assume the axiom of foundation and by contradiction that x is a set for which $\neg V(x)$ Have $x \subseteq \text{cl } x$ **Claim:** $Y = \{y \in \text{cl } x : \neg V(y)\} \neq \emptyset$

proof of claim. If $Y = \emptyset$ then $\forall y y \in Y \rightarrow V(y)$ and rk bounded on Y . \boxtimes

Let $y \in Y$. Then $\neg V(y)$ so i.p. $y \not\subseteq V$, so for some $z \in y$ have $\neg V(z)$ but because $\text{cl } x$ is transitive, $z \in \text{cl } x$ hence $z \in Y$ hence $\forall y y \in Y \rightarrow Y \cap Y \neq \emptyset$, a contradiction with (AF). \square

By Gödel's 2-nd incompleteness theorem ZFC can not prove its own consistency. There is a way to express consistency in the formal level, by coding and using peano arithmetic, and the above statement says that ZFC can not prove this sentence. All we can hope for are relative consistency results, and therefore relate two theories with each other. For example some theories are less debated about and it shows a way to prove independence of certain axioms from others.

- ZFC The axioms 1. extensionality 2. union 3. power-set 4. ax scheme of replacement 5. set ax 6. axiom of infinity 7. AC 8. AF (recall pairing and comprehension follow from the other)
- ZFC^- is ZFC without (AF)
- ZF is ZFC without (AC)
- ZF^- is ZF without (AF)

We will use a tool that is in set theory called Relativization

Let C be a class, $\phi(\underline{x}, \underline{a})$, $\underline{a} \in C$

$\phi^C(\underline{x}, \underline{a})$ defined by induction on compl of ϕ

- If ϕ is atomic then $\phi^C = \phi$
- $(\neg\phi)^C = \neg(\phi^C)$, $(\phi \vee \psi)^C = (\phi^C) \vee (\psi^C)$
- $(\exists y\phi)^C = \exists y(C(y) \wedge \phi^C)$
- $(\forall y\phi)^C = \forall y(C(y) \rightarrow \phi^C)$

Theorem 2.32. Suppose $(\mathcal{U}, \in) \models ZF^-$ then V constructed in \mathcal{U} is such that $(V, \in) \models ZF$ i.e. assume ZF^- is consistent (has a model) and we get a model of ZF

Proof. ZF^- Take 1. extensionality 2. union 3. power-set 4. ax scheme of replacement 5. set ax 6. axiom of infinity (recall pairing and comprehension follow from the other) Need to check if $(\mathcal{U}, \in) \models ZF^-$ and V class defined by $V = \bigcup_{\alpha \in \text{On}} V_\alpha$, $V_\beta := \bigcup_{\alpha < \beta} \mathcal{P}(V_\alpha)$ then $(V, \in) \models ZF^-$ (AF will follow from prev. result)

1. Let $x, y \in V$ wts $(\forall z \in V z \in x \leftrightarrow z \in y) \leftrightarrow x = y$ $x, y \in V$ so $x, y \subseteq V$ and can use Axiom of extensionality in \mathcal{U} $(\forall z \in \mathcal{U} z \in x \leftrightarrow z \in y) \leftrightarrow x = y$
2. Let $x \in V$ $U_x = \{z : \exists y \in x z \in y\} \subseteq V$ so $\bigcup_x \in V$.
3. Let $x \in V$ then $x \subseteq V$. so every subset of x is a subset of V hence is an element of V . Powerset $\mathcal{P}(x) \subseteq V$ hence $\mathcal{P}(x) \in V$
4. $\varphi(x, y)$ with parameters from V and assume that $\varphi(x, y)$ defines a class function in V i.e.

$$\left(\forall x \exists^{\leq 1} y \varphi(x, y) \right)^V$$

i.e.

$$\forall x (V(x) \rightarrow (\exists^{\leq 1} y \varphi(x, y)))$$

Then $\psi(x, y) : \Leftrightarrow V(x) \wedge V(y) \wedge \varphi^V(x, y)$ defines a class function in \mathcal{U} , use replacement in \mathcal{U} That yields

$$\forall a \exists b y \in b \text{ iff } (\exists x \in A \varphi(x, y)) \text{ iff } \exists x \in a \varphi^V(x, y) \wedge V(y)$$

$b \subseteq V$ hence $b \in V$ and b is the image of a under the class function given by φ .

5. Since we proved that every ordinal is in V , $\emptyset \in V$.
6. enough to show that $\omega \in V$. in fact we know that any ordinal is in V .

□

An ordinal α is called regular, if it is equal to its own cardinality $\text{cof } \alpha = \alpha$ A cardinal κ is called inaccessible if $\kappa > \omega$ and κ is regular.

It suffices to say $\forall \lambda < \omega 2^\lambda < \kappa$ Suppose now that $(\mathcal{U}, \in) \models ZFC$

Lemma 2.33. If κ is an inaccessible cardinal, then $|V_\kappa| = \kappa$. Moreover for every $a \subseteq V_\kappa$ we have $a \in V_\kappa$ iff $|a| < \kappa$

Proof. Recall, if $\alpha \in \text{On}$ then $\alpha \subseteq V_\alpha$. i.p. $\kappa \subseteq V_\kappa$ so $|\kappa| \leq |V_\kappa|$

The other direction: For this we are going to show inductively that for all $\xi < \kappa$ that $|V_\xi| < \kappa$ Then $\kappa \geq |V_\kappa|$ follows. If $|V_\xi| < \kappa$ then $|V_{\xi+1}| = |\mathcal{P}(V_\xi)| \leq |2^{|V_\xi|}| < \kappa$ by κ inaccessible. Suppose $|V_\xi| < \kappa$ for all $\xi < \lambda < \kappa$ where λ is a limit ordinal.

$$|V_\lambda| = \left| \bigcup_{\xi < \lambda} V_\xi \right| \leq \sup \xi < \lambda |V_\xi| < \kappa$$

here we use regularity of κ .

So have $|V_\kappa| = \kappa$. Left to show: for every $a \subseteq V_\kappa$ have $a \in V_\kappa$ iff $|a| < \kappa$

- “ \Leftarrow ”-direction: Assume $a \subseteq V_\kappa$ and $|a| < \kappa$ $\text{rk} : a \rightarrow \text{On}$ is not cofinal in κ because $\text{cof } \kappa = \kappa > |a|$ So for some ordinal β have $a \subseteq V_\beta$ and then $a \in V_{\beta+1} \subseteq V_\kappa$

- “ \implies ”-direction: it suffices $a \in V_\kappa$ then $|a| < \kappa$ Exercise. □

Lemma 2.34. *Let $(\mathcal{U}, \in) \models ZFC$. If κ is inaccessible, then $V_\kappa \models ZFC$.*

Proof. Will check (AC) and replacement, the remaining axioms are an exercise.

(AC) : Suppose $a \in V_\kappa$ and a is a family of pairwise disjoint, non-empty sets. By (AC) in \mathcal{U} there is a transversal T in \mathcal{U} for the set a . What is left to show is $T \in V_\kappa$. Have $a \subseteq V_\kappa$ then every subset of a is a subset of V_κ so in particular $T \subseteq V_\kappa$. We do know that $a \in V_\kappa$ so $|a| < \kappa$ and $T \subseteq a$ we have $|T| \leq |a|$ so by the previous lemma $T \in V_\kappa$.

(RE) : $\varphi(x, y)$ a formula with parameters in V_κ that defines a class function in V_κ . i.e.

$$\forall x \in V_\kappa \exists^{<\kappa} y \in V_\kappa \varphi^{V_\kappa}(x, y)$$

let $a \in V_\kappa$ then $\psi(x, y) \equiv x \in V_\kappa \wedge y \in V_\kappa \wedge \varphi(x, y)$ does define a class function on \mathcal{U} with domain contained in V_κ . So $f[a] \subseteq V_\kappa$ and $|f[a]| < \kappa$ so $f[a] \in V_\kappa$. □

Now we are ready to state our meta-theorem, it is not a statement in first order language of set theory.

Theorem 2.35. *If ZFC is consistent then $ZFC + \text{“There are no strongly inaccessible cardinals”}$ is also consistent*

Proof. Assume we have a model of ZFC, $(\mathcal{U}, \in) \models ZFC$.

- If \mathcal{U} does not contain any inaccessible cardinals then we are done.
- Assume that \mathcal{U} does contain inaccessible cardinals. Let κ be the smallest inaccessible cardinal.

What we want to show is that there are no inaccessible cardinals in V_κ . An ordinal is by definition a transitive set, well-ordered by \in . By (AF) α is transitive and α is linearly ordered by \in . i.e. α ordinal iff

$$\forall x, y \in \alpha (x \in y \vee y \in x \vee x = y) \wedge \forall x (x \in \alpha \rightarrow x \subseteq \alpha) \quad (5.3)$$

Claim 1: The ordinals in V_κ are the ordinals below κ i.e. $Ord^{V_\kappa} = \kappa$.

proof of claim. If $\alpha < \kappa$ then $\alpha \subseteq V_\kappa$ ($\text{rk } \alpha = \alpha + 1$ and $V_{\alpha+1} \subseteq V_\kappa$ so $\alpha \in V_\kappa$ ($\alpha + 1 < \kappa$)) α is an ordinal so 5.3 holds. and $\alpha \in \text{On}^{V_\kappa}$. If $\alpha \in \text{On}^{V_\kappa}$ then $\alpha \in V_\kappa$ and α is transitive and totally ordered by \in . Hence α is an ordinal. Left to show α is below κ . Have $|\alpha| < \kappa$, by the Lemma 2.33 $\alpha < \kappa$. □

Claim 2: The cardinals in V_κ are the cardinals that are below κ .

proof of claim. Suppose λ is a cardinal in V_κ . in particular λ is an ordinal in V_κ and therefore an ordinal in \mathcal{U} and by the lemma 2.33 $\lambda = |\lambda| < \kappa$. left to show: λ is an actual cardinal in \mathcal{U} . Suppose there is a bijection $f : \lambda \rightarrow \alpha$ to some smaller ordinal α . Then by Claim 1, $\alpha \in \text{On}^{V_\kappa}$ and $f \subseteq V_\kappa$ (the graph $f \subseteq \lambda \times \alpha \subseteq \lambda \times \lambda$ so $f \subseteq \mathcal{P}\mathcal{P}\mathcal{P}(V_{\lambda+1}) =: V_\beta$, $\text{rk } \lambda = \lambda + 1$ and have $\beta < \kappa$ by inaccessibility). $f \subseteq V_\kappa$ and $|f| < \kappa$ so by lemma 2.33 $f \in V_\kappa$. f is a bijection in V_κ between λ and a smaller ordinal, a contradiction.

Suppose $\lambda < \kappa$ and λ cardinal, then by lemma 2.33 $\lambda \in V_\kappa$ and $\lambda \in \text{On}^{V_\kappa}$. If there would be a bijection in V_κ between λ and a strictly smaller ordinal α then α would be an actual ordinal. By an argument as before would get a bijection in \mathcal{U} between λ and smaller ordinal. □

If $\lambda \in V_\kappa$ is a cardinal in V_κ then $\lambda < \kappa$ and by claim 2 it is a cardinal in \mathcal{U} , λ can not be inaccessible. by choice of κ .

Note that only \mathcal{U} knows that λ is not inaccessible. We therefore need to check that also V_κ knows that λ is inaccessible. Reasons for λ to not be inaccessible:

- $\lambda \leq \omega$ (in \mathcal{U}) hence $(\lambda \leq \omega)^{V_\kappa}$ bc. $\omega \in V_\kappa$
- $\lambda \leq 2^\xi$ for some $\xi < \lambda < \kappa$ hence $\xi, 2^{\xi i} \in V_\kappa$ and have $(\xi < \lambda < 2^\xi)^{V_\kappa}$
- λ is not regular, if we have a function $f : \alpha \rightarrow \lambda$ cofinal and α an ordinal $\alpha < \lambda$, then $\alpha \in V_\kappa$ hence $f \in V_\kappa$ then λ is not regular in V_κ . □

Note: $Ord^{V_\kappa} = \kappa$. but Ord^{V_κ} is no set in V_κ .

CHAPTER 6

Outernotes, Abbreviations, Overview

List of Outernotes

	freely generated	7	sentence	12
	acceptable	7	structures	13
	tautological equivalence	8	structure	13
	\sim	8	interpretation	13
	\models	8	assignment	13
	\models	8	$\models_{\mathcal{A}}$	14
	n -ary boolean function	9	definable by points . .	16
	DNF		onto	17
	Disjunctive nor-		isomorphic	17
	mal form	9	MP	20
	completeness (con-		Chapter 3	30
	nectives)	9	\simeq	34
	satisfiable	9	RCF	37
	finitely satisfiable . .	10	Chapter 4	43
	Chapter 2	11	Ultrafilter	44
	logical symbols	11	$S(B)$	44
	parameters	11	Chapter 5	49
	expression	12	Ord	52
	Term	12	On	52
	atomic formula	12	successor cardinals . .	59
	formula	12	limit cardinals	59
	\exists quantifier	12	Chapter 6	65
	bounded variable . . .	12	Chapter 6	69
Chapter 1	2			
Language	2			
expression	2			
prop. fla.	2			
prop. sentence	2			
construction sequence	3			
closure	3			
Truth assigment . . .	3			
TA	3			
satisfy	4			
satisfiable	4			
taut. implication \models .	4			
closed	6			
inductive	6			

List of Abbreviations

prop.	-	propositional	2	fla.	-	formula	3
exp.	-	expression(s)	2	TV	-	truth value	3
sent.	-	sentence(s)	3	taut.	-	tautological	4
seq.	-	sequence	3	w/	-	with	4
TA	-	truth assignment	3	lp / rp	-	left / right parenthesis	5
				i.e.	-	id est (that is)	9
				MP	-	Modus Ponens	20
				SUB	-	substitutable	21
				WMA	-	We may assume	38

List of definitions

Propositional logic

2

Def. 1.1	Language of PL	2
Def. 1.2	Expression / prop. sentence	2
Def. 1.3	Construction sequence	3
Def. 1.4	Closedness of a set	3
Def. 1.5	Truth assignment	3
Def. 1.6	Satisfaction	4
Def. 1.7	Tautological implication	4
Def. 1.8	Closedness, Inductiveness	6
Def. 1.9	Freely generated set	7
Def. 1.10	Tautological equivalence relation	8
Def. 1.11	Satisfiability	9

Predicate - / first order logic

11

Def. 2.1	A First order Language	11
Def. 2.2	Expression	12
Def. 2.3	Term Building Operations	12
Def. 2.4	Atomic formula	12
Def. 2.5	Formulas	12
Def. 2.6	Free variables	12
Def. 2.7	structure	13
Def. 2.8	Assignment	13
Def. 2.9	Satisfy	14
Def. 2.10	Model	15
Def. 2.11	Logical implication	15
Def. 2.12	Logical equivalence	15
Def. 2.13	Valid	15
Def. 2.14	definability in a structure	16
Def. 2.15	definability of classes of structures	16
Def. 2.16	Homomorphism	17
Def. 2.17	Isomorphism	17
Def. 2.18	Substructure	17
Def. 2.19	elementarily equivalent	18
Def. 2.20	Automorphism	18
Def. 2.21	Rigid	18
Def. 2.22		19
Def. 2.23	terminal segment	19
Def. 2.24		19
Def. 2.25	Modus Ponens	20
Def. 2.26	Deduction	20
Def. 2.27	Generalization	20
Def. 2.28	Logical axioms	20
Def. 2.29	Substitution	20
Def. 2.30	substitutable	21
Def. 2.31	Tautologies of first order language	21
Def. 2.32	Inconsistence	23
Def. 2.33		26
Def. 2.34		28

Model Theory

30

Def. 3.1	Theory	30
Def. 3.2	Theory of structures	30
Def. 3.3	C_n	31
Def. 3.4		31
Def. 3.5	axiomatizability	31
Def. 3.6		31
Def. 3.7	infinitely close	34
Def. 3.8	Convergence (non-standard definition)	34
Def. 3.9		35
Def. 3.10	o-minimality	36
Def. 3.11	definably connectedness	36
Def. 3.12	ordered group	36
Def. 3.13	Ordered ring	37
Def. 3.14	ordered field	37
Def. 3.15	real closed field	37
Def. 3.16	Cell	40
Def. 3.17	decomposition	40
Def. 3.18		41
Def. 3.19	Definable families	41

Boolean Algebra

43

Def. 4.1	Boolean Algebra	43
Def. 4.2	Alternative Def: Boolean Algebra	43
Def. 4.3	(Ultra-)Filter	44
Def. 4.4	Stone space	44
Def. 4.5	Atomic, Atomless	45
Def. 4.6	Lindenbaum-Tarski algebra	45
Def. 4.7	Recap	46
Def. 4.8	Opposite of boolean algebra	47
Def. 4.9	Subalgebra	47
Def. 4.10	Homomorphisms of boolean algebras	47
Def. 4.11	Trivial algebras	48
Def. 4.12	Ideal	48

Set Theory of ZFC

49

Def. 5.1	Axiom of extensionality	49
Def. 5.2	Pairing Axiom	49
Def. 5.3	n-tuples	50
Def. 5.4	Union Axiom	50
Def. 5.5		50
Def. 5.6	Power set Axiom	50
Def. 5.7	Classes	50
Def. 5.8	Class functions	50
Def. 5.9	Axiom of replacement / substitution	50
Def. 5.10	Axiom scheme of comprehension	51
Def. 5.11	Set existence	51
Def. 5.12	Function	51
Def. 5.13		51
Def. 5.14		51
Def. 5.15	Well ordering	52
Def. 5.16	Initial segment	52
Def. 5.17	Propper class	52
Def. 5.18	well-ordering (class)	52
Def. 5.19	Transitivity of sets	52
Def. 5.20	Ordinal	52
Def. 5.21	H-inductive	54
Def. 5.22	Axiom of Choice (AC)	55

Def. 5.23	(AC')	55
Def. 5.24	(AC'')	55
Def. 5.25		56
Def. 5.26		56
Def. 5.27	Axiom of infinity	57
Def. 5.28		57
Def. 5.29		58
Def. 5.30		58
Def. 5.31	Cardinal	58
Def. 5.32	finite sets	58
Def. 5.33		58
Def. 5.34	countable sets	59
Def. 5.35		59
Def. 5.36		60
Def. 5.37		60
Def. 5.38	regular ordinals	60
Def. 5.39		61
Def. 5.40	Axiom of foundation (AF)	61
Def. 5.41		62
Def. 5.42		62
Def. 5.43	inductive closure	62
Outernotes, Abbreviations, Overview		65
Bibliography		69

List of Theorems

Propositional logic		2
Truth assignments		3
Thm. 1.1	Unique readability	3
Thm. 1.2	Compactness theorem	4
A parsing algorithm		4
Lemma 2.1		4
Lemma 2.2		5
Induction and recursion		6
Thm. 3.1	Recursion Theorem	7
Sentential connectives		8
Thm. 4.1		8
Thm. 4.2	(Post)	9
Thm. 4.3		9
Compactness Theorem		10
Thm. 5.1	Compactness Theorem	10
Predicate - / first order logic		11
Formulas		12
Semantics of first order logic		13
Thm. 2.1		13
Thm. 2.2	Coincidence Lemma	15
definability in a structure		16
Homomorphisms of structures		17
Thm. 4.1	Homomorphism Theorem	18

Unique readability for terms	19
Lemma 5.1	19
Lemma 5.2	19
Prop. 5.3 Unique readability for terms . .	19
Lemma 5.4	19
Lemma 5.5	19
A parsing algorithm for first order logic	20
Deductions (formal proofs)	20
Lemma 7.1	22
Thm. 7.2	22
Generalization and deduction theorem	22
Thm. 8.1 Generalization theorem	22
Lemma 8.2 Rule T	23
Thm. 8.3 Deduction theorem	23
Thm. 8.4 Generalization on constants . .	23
Thm. 8.5 Existence of alphabetic variants	24
Soundness and completeness	25
Lemma 9.1 pre-substitution lemma	25
Lemma 9.2 Substitution lemma	25
Thm. 9.3	26
Thm. 9.4 Completeness Theorem	26
Thm. 9.5 Completeness Theorem'	26
Lemma 9.6	28
Lemma 9.7	29
Thm. 9.8 Zorn's Lemma	29
Thm. 9.9 Zermelos's Theorem	29

Model Theory		30
Löwenheim-Skolem-Theorem		30
Thm. 1.1	LST	30
Theories and completeness		30
Thm. 2.1		31
Thm. 2.2	Los-Vaught test	31
Theory of algebraic closed fields		32
Thm. 3.1		32
Prop. 3.2	Lefschetz Principle	32
Prop. 3.3	Ax-Grothendieck	33
Prop. 3.4		33
Nonstandard Analysis		33
Prop. 4.1		34
Prop. 4.2		34
Lemma 4.3		34
Prop. 4.4		34
Prop. 4.5		34
Prop. 4.6		35
Prop. 4.7	Chain rule	35
o-minimality		36
Lemma 5.1		36
o-minimal ordered groups and rings		36
Prop. 6.1		36
Lemma 6.2		37
Lemma 6.3		37
Prop. 6.4		38
Prop. 6.5	Monotonicity Theorem	38
Lemma 6.6		38
Lemma 6.7		38
Lemma 6.8		38
Thm. 6.9	Cell decomposition	41

Prop. 6.10	41	Lemma 2.11	55
Prop. 6.11	42	Thm. 2.12 Transfinite Recursion	55
Thm. 6.12	42	Thm. 2.13 Zermelo	56
Boolean Algebra	43	Thm. 2.14 Zorn's Lemma	56
Stone Representation Theorem	44	Lemma 2.15	57
Thm. 1.1 Stone Representation Theorem	45	Thm. 2.16	58
Lindenbaum-Tarski Algebras	45	Thm. 2.17 Cantor-Schröder-Bernstein	58
Thm. 2.1 Compactness Theorem *	46	Thm. 2.18 Cantor	58
Lemma 2.2	47	Prop. 2.19 Galileo	58
Lemma 2.3	47	Prop. 2.20	59
Lemma 2.4	48	Hyp. 2.21 (CH)	59
Set Theory of ZFC	49	Thm. 2.22	59
First axioms of ZFC	49	Lemma 2.23	60
Lemma 1.1	49	Prop. 2.24	60
Lemma 1.2	49	Lemma 2.25	61
Lemma 1.3	50	Thm. 2.26	61
Ordinals	52	Lemma 2.27	61
Lemma 2.1 Characterization of ordinals	52	Thm. 2.28 (König)	61
Lemma 2.2	53	Lemma 2.29	62
Prop. 2.3	53	Lemma 2.30	62
Lemma 2.4	53	Thm. 2.31	62
Lemma 2.5	53	Thm. 2.32	63
Lemma 2.6	53	Lemma 2.33	64
Lemma 2.7	54	Lemma 2.34	64
Thm. 2.8	54	Thm. 2.35	64
Thm. 2.9	54	Outernotes, Abbreviations, Overview	65
Lemma 2.10	55	Bibliography	69

Bibliography

- [EE01] Herbert B Enderton and Herbert Enderton. *A Mathematical Introduction to Logic*. eng. United States: Elsevier Science and Technology, 2001. ISBN: 0122384520.
- [Kri98] J.L. Krivine. *Théorie des ensembles*. Nouvelle bibliothèque mathématique. Cassini, 1998. ISBN: 9782842250140. URL: <https://books.google.at/books?id=04A2AAAAAAJ>.
- [Van98] Lou Van den Dries. *Tame topology and o-minimal structures*. Vol. 248. London Mathematical Society lecture note series. Cambridge University Press, 1998. ISBN: 9780511525919.