



National Comprehensive
Cancer Network®

NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®)

Prostate Cancer

Version 5.2026 — January 23, 2026

**NCCN recognizes the importance of clinical trials and encourages participation when applicable and available.
Trials should be designed to maximize inclusiveness and broad representative enrollment.**

NCCN Guidelines for Patients® available at www.nccn.org/patients



NCCN Guidelines Version 5.2026

Prostate Cancer

***Daniel E. Spratt, MD/Chair §**

Case Comprehensive Cancer Center/University
Hospitals Seidman Cancer Center and Cleveland
Clinic Taussig Cancer Institute

***Sandy Srinivas, MD/Vice-Chair † ω**

Stanford Cancer Institute

Nabil Adra, MD, MSc †

Indiana University Melvin and Bren
Simon Comprehensive Cancer Center

Bilawal Ahmed, MD †

The University of Tennessee Health Science Center

Yi An, MD §

Yale Cancer Center/Smilow Cancer Hospital

Rhonda Bitting, MD †

Duke Cancer Institute

Brian Chapin, MD ω

The University of Texas
MD Anderson Cancer Center

Heather H. Cheng, MD, PhD †

Fred Hutchinson Cancer Center

Steve Y. Cho, MD φ

University of Wisconsin Carbone Cancer Center

Anthony Victor D'Amico, MD, PhD §

Dana-Farber/Brigham and Women's
Cancer Center | Mass
General Cancer Center

Neil Desai, MD, MHS §

UT Southwestern Simmons
Comprehensive Cancer Center

***Tanya Dorff, MD †**

City of Hope National Cancer Center

James A. Eastham, MD ω

Memorial Sloan Kettering Cancer Center

***Thomas A. Farrington, BSEE ¥**

Prostate Health Education Network (PHEN)

***Xin Gao, MD †**

Dana-Farber/Brigham and Women's
Cancer Center | Mass General Cancer Center

Shilpa Gupta, MD †

Case Comprehensive Cancer Center/University
Hospitals Seidman Cancer Center and Cleveland
Clinic Taussig Cancer Institute

Joseph E. Ippolito, MD, PhD ∩

Siteman Cancer Center at Barnes-
Jewish Hospital and Washington
University School of Medicine

R. Jeffrey Karnes, MD ω

Mayo Clinic Comprehensive Cancer Center

Amar U. Kishan, MD §

UCLA Jonsson Comprehensive Cancer Center

Michael R. Kuettel, MD, MBA, PhD §

Roswell Park Comprehensive Cancer Center

Joshua M. Lang, MD, MS †

University of Wisconsin Carbone Cancer Center

Daniel Lee, MD, MS ω

Abramson Cancer Center at
The University of Pennsylvania

Tamara Lotan, MD ≠

Johns Hopkins Kimmel Cancer Center

Andrew McDonald, MD, MS §

O'Neal Comprehensive Cancer Center at UAB

***Todd M. Morgan, MD ω**

University of Michigan Rogel Cancer Center

Rodrigo Pessoa, MD, PhD ω

Moffitt Cancer Center

Soroush Rais-Bahrami, MD, MBA ω

O'Neal Comprehensive Cancer Center at UAB

Mack Roach, III, MD §

UCSF Helen Diller Family
Comprehensive Cancer Center

Tyler Robin, MD, PhD §

University of Colorado Cancer Center

Stan Rosenfeld ¥

University of California San Francisco
Patient Services Committee Chair

Kristen R. Scarpato, MD, MPH ω

Vanderbilt-Ingram Cancer Center

Tyler Seibert, MD, PhD §

UC San Diego Moores Cancer Center

Ahmad Shabsigh, MD ω

The Ohio State University Comprehensive
Cancer Center - James Cancer Hospital
and Solove Research Institute

Russell Szmulewitz, MD ‡ †

The UChicago Medicine
Comprehensive Cancer Center

Benjamin A. Tepley, MD †

Fred & Pamela Buffett Cancer Center

Jonathan Tward, MD, PhD §

Huntsman Cancer Institute
at the University of Utah

Richard Valicenti, MD §

UC Davis Comprehensive Cancer Center

David VanderWeele, MD, PhD †

Robert H. Lurie Comprehensive Cancer
Center of Northwestern University

Jessica Karen Wong, MD §

Fox Chase Cancer Center

NCCN

Deborah Freedman-Cass, PhD

Emily Kovach

∩ Diagnostic/ Interventional radiology	§ Radiotherapy/Radiation oncology
‡ Hematology/Hematology oncology	ω Urology
† Medical oncology	¶ Surgery/Surgical oncology
φ Nuclear medicine	* Discussion Section Writing Committee
≠ Pathology	
¥ Patient advocate	

[NCCN Guidelines Panel Disclosures](#)



NCCN Guidelines Version 5.2026

Prostate Cancer

[NCCN Prostate Cancer Panel Members](#)

[Summary of Guidelines Updates](#)

[Initial Prostate Cancer Diagnosis \(PROS-1\)](#)

[Initial Risk Stratification and Staging Workup for Clinically Localized Disease \(PROS-2\)](#)

[Low-Risk Group \(PROS-3\)](#)

[Favorable Intermediate-Risk Group \(PROS-4\)](#)

[Unfavorable Intermediate-Risk Group \(PROS-5\)](#)

[High- or Very-High-Risk Group \(PROS-6\)](#)

[Regional Prostate Cancer \(PROS-7\)](#)

[Monitoring \(PROS-8\)](#)

[Radical Prostatectomy PSA Persistence/Recurrence \(PROS-9\)](#)

[Radiation Therapy Recurrence \(PROS-10\)](#)

[Workup for Second Biochemical Recurrence \(BCR2\) \(PROS-11\)](#)

[Treatment and Monitoring for Second Biochemical Recurrence \(M0\) \(PROS-12\)](#)

[Workup and Treatment of Metachronous Oligometastatic M1 CSPC \(PROS-13\)](#)

[Workup and Treatment of Low-Volume M1 \(Metachronous or Synchronous\) or Synchronous Oligometastatic CSPC \(PROS-14\)](#)

[Workup and Treatment of High-Volume M1 CSPC \(PROS-15\)](#)

[Workup and Treatment of M0 Castration-Resistant Prostate Cancer \(CRPC\) \(PROS-16\)](#)

[Workup and Treatment of M1 CRPC \(PROS-17\)](#)

[Systemic Therapy for M1 CRPC: Adenocarcinoma \(PROS-18\)](#)

[Principles of Life Expectancy Estimation \(PROS-A\)](#)

[Principles of Survivorship in Prostate Cancer \(PROS-B\)](#)

[Principles of Genetics and Molecular/Biomarker Analysis \(PROS-C\)](#)

[Principles of Quality of Life and Shared Decision-Making \(PROS-D\)](#)

[Principles of Imaging \(PROS-E\)](#)

[Principles of Active Surveillance and Observation \(PROS-F\)](#)

[Principles of Androgen Deprivation Therapy \(PROS-G\)](#)

[Principles of Risk Stratification and Biomarkers \(PROS-H\)](#)

[Principles of Focal/Subtotal Therapy or Whole Gland Ablative Therapy \(PROS-I\)](#)

[Principles of Radiation Therapy \(PROS-J\)](#)

[Principles of Surgery \(PROS-K\)](#)

[Principles of Local Secondary Therapy Post-Radiation \(PROS-L\)](#)

[Principles of Metastasis-Directed Therapy \(MDT\) \(PROS-M\)](#)

[Principles of Non-Hormonal Systemic Therapy \(PROS-N\)](#)

[Staging \(ST-1\)](#)

[Abbreviations \(ABBR-1\)](#)

Find an NCCN Member Institution: <https://www.nccn.org/home/member-institutions>.

NCCN Categories of Evidence and Consensus: All recommendations are category 2A unless otherwise indicated.

See [NCCN Categories of Evidence and Consensus](#).

NCCN Categories of Preference: All recommendations are considered appropriate.

See [NCCN Categories of Preference](#).

The NCCN Guidelines® are a statement of evidence and consensus of the authors regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult the NCCN Guidelines is expected to use independent medical judgment in the context of individual clinical circumstances to determine any patient's care or treatment. The National Comprehensive Cancer Network® (NCCN®) makes no representations or warranties of any kind regarding their content, use or application and disclaims any responsibility for their application or use in any way. The NCCN Guidelines are copyrighted by National Comprehensive Cancer Network®. All rights reserved. The NCCN Guidelines and the illustrations herein may not be reproduced in any form without the express written permission of NCCN. ©2026.



NCCN Guidelines Version 5.2026

Prostate Cancer

Updates in Version 5.2026 of the NCCN Guidelines for Prostate Cancer include:

[PROS-14](#)

- Treatment for Low-Volume M1 CSPC disease, regimen added to synchronous and metachronous settings
 - ▶ ADT with: Useful in certain circumstances, Niraparib/abiraterone (BRCA2m only) (category 2B)

[PROS-15](#)

- Treatment for High-Volume M1 CSPC disease, regimen added
 - ▶ ADT with: Useful in certain circumstances, Niraparib/abiraterone (BRCA2m only)

[PROS-15A](#)

- Footnote bbb added: Niraparib/abiraterone can be used following the completion of docetaxel in patients receiving triplet therapy or following EBRT to the primary tumor. The use of this combination in patients with low-volume disease is controversial because 78% of the patients in the trial had high-volume disease and there were very few patients with low-volume metachronous disease in particular. This regimen adds considerable toxicity, and patients with low-volume disease should be informed that the benefit in this setting is not clear.

[PROS-G \(4 of 6\)](#)

- ADT for patients with mCSPC, sub-header added: Additional option for patients with BRCA2m mCSPC
 - ▶ Bullet added: Orchiectomy, LHRH agonist, or LHRH antagonist plus niraparib and abiraterone
- Footnote g added: The fine-particle (category 2B; other recommended option) or standard formulation of abiraterone can be given with single-agent niraparib as a substitute for the combination niraparib/abiraterone tablet.

[PROS-N \(1 of 5\)](#)

- Non-hormonal systemic therapy for M1 castration-sensitive prostate cancer
 - ▶ Bullet 2 added: An additional option for patients with BRCA2-mutated mCSPC is the use of ADT with niraparib and abiraterone.
 - ◊ Sub-bullet 1 added: ADT with niraparib and abiraterone was compared to ADT plus abiraterone doublet therapy in a randomized phase 3 trial. Most patients had high-volume and/or synchronous disease. rPFS was improved in the BRCAm population; OS data are immature. Grade 3/4 adverse events were more common in the niraparib arm (75% vs. 59%) as were treatment-emergent adverse events leading to death (14 participants vs. 7 participants). The most common grade 3/4 adverse events were anemia (29.1% in the niraparib group versus 4.6% in the control group) and hypertension (26.5% versus 18.4%).
 - ◊ Sub-bullet 2 added: Niraparib/abiraterone can be used following the completion of docetaxel in patients receiving triplet therapy or following EBRT to the primary tumor.
 - ◊ Sub-bullet 3 added: The use of this combination in patients with low-volume disease is controversial because 78% of the patients in the trial had high-volume disease and there were very few patients with low-volume metachronous disease in particular. This regimen adds considerable toxicity, and patients with low-volume disease should be informed that the benefit in this setting is not clear.
 - ◊ Sub-bullet 4 added: The fine-particle (category 2B; other recommended option) or standard formulation of abiraterone can be given with single-agent niraparib as a substitute for the combination niraparib/abiraterone tablet.

[MS-1](#)

- The Discussion section was updated to reflect the changes in the algorithm.

Updates in Version 4.2026 of the NCCN Guidelines for Prostate Cancer include:

[PROS-J \(9 of 11\)](#)

- Table 1, sub-header modified: EBRT + Brachytherapy Boost (*combined with 45-50.4 Gy in 25-28 fx or 37.5 Gy in 15 fx*).

[MS-1](#)

- The remainder of the Discussion section was updated to reflect the changes in the algorithm.



NCCN Guidelines Version 5.2026

Prostate Cancer

Updates in Version 3.2026 of the NCCN Guidelines for Prostate Cancer include:

[PROS-G \(3 of 6\)](#)

- ADT for Patients with M0 CSPC BCR2
 - ▶ Sub-heading modified: *For High-Risk BCR2 Useful in Certain Circumstances*

[PROS-G \(3 of 6\)\(continued\)](#)

- ▶ Bullet 1 modified: Enzalutamide with or without Leuprolide ~~(for high-risk BCR2)~~
- ▶ Bullet 2 modified: Apalutamide with LHRH agonist or LHRH antagonist ~~(for high-risk BCR2)~~

[PROS-H](#)

- Section extensively revised.

[PROS-N \(3 of 4\)](#)

- Immunotherapy, bullet 2, sub-bullet 3 added: Pembrolizumab and berahyaluronidase alfa-pmph subcutaneous injection may be substituted for IV pembrolizumab. Pembrolizumab and berahyaluronidase alfa-pmph has different dosing and administration instructions compared to IV pembrolizumab.

[MS-1](#)

- The section of the Discussion regarding "Advanced Risk Stratification Tools" was updated to reflect the changes in the algorithm.

Updates in Version 2.2026 of the NCCN Guidelines for Prostate Cancer from Version 1.2026 include:

[MS-1](#)

- The sections of the Discussion regarding mCSPC and mCRPC were updated to reflect the changes in the algorithm.

Updates in Version 1.2026 of the NCCN Guidelines for Prostate Cancer from Version 2.2025 include:

[Global](#)

- Very-low-risk group has been removed from the Guidelines.
- Androgen receptor pathway inhibitors ("ARPIs") has replaced "other hormonal agents" throughout the Guidelines.
- References have been updated throughout the Guidelines.

[PROS-1](#)

- Workup
 - ▶ Top pathway
 - ◊ Bullet 3 modified: Perform and/or collect prostate-specific antigen (PSA) ~~and calculate PSA density.~~
 - ◊ Bullet 4 modified: ~~Obtain and~~ Review diagnostic prostate biopsies.
 - ▶ Bottom pathway
 - ◊ Bullet 3 modified: *Consider* ~~Perform~~ DRE to confirm clinical stage.
 - ◊ Last column, pathway added: See Workup and Treatment of High-Volume M1 CSPC (PROS-15).
- Footnote d added: Refer to the NCCN Distress Thermometer and Problem List, which includes social determinants of health. See NCCN Guidelines for Distress Management (DIS-A).
- Footnote h added: See "Number of Metastatic Sites" in the Principles of MDT (PROS-M). (Also for PROS-2A, 7A, -8A, -9A, -10A, -11, -12, -13, and -15A, and 18A)

[PROS-2](#)

- Additional Evaluation for Intermediate, High, and Very high, sub-bullets for soft-tissue imaging
 - ▶ Sub-bullet 1 modified: If regional ~~or distant~~ metastases are found, see PROS-7 ~~or~~
 - ▶ Sub-bullet added: If distant metastases are found, see PROS-14 for Low-Volume M1 (Metachronous or Synchronous) or Synchronous Oligometastatic CSPC or PROS-15 for High-Volume M1 CSPC.



NCCN Guidelines Version 5.2026

Prostate Cancer

Updates in Version 1.2026 of the NCCN Guidelines for Prostate Cancer from Version 2.2025 include:

[PROS-2A](#)

- Footnote removed: An ultrasound-, MRI-, or DRE-targeted lesion that is biopsied more than once and demonstrates cancer (regardless of percentage core involvement or number of cores involved) can be considered as a single positive core.

[PROS-3](#)

- Footnote n added: Principles of Focal/Subtotal Therapy or Whole Gland Ablative Therapy (PROS-I). (Also for PROS-4, -5, -6, and -7A)

[PROS-6](#)

- ≤5 y and asymptomatic
 - ▶ Initial therapy modified: Observation or RT ± ADT ~~or ADT ± RT~~.

[PROS-7A](#)

- Footnote x modified: Adverse laboratory/pathologic features include: positive margin(s), seminal vesicle invasion, *or* extracapsular extension, ~~or detectable PSA~~.
- Footnote removed: The Panel remains concerned about the problems of overtreatment related to the increased diagnosis of early prostate cancer from PSA testing. See NCCN Guidelines for Prostate Cancer Early Detection. Active surveillance is recommended for this subset of patients.
- Footnote removed: The fine-particle formulation of abiraterone can be used instead of the standard form (category 2B; other recommended option). (Also for PROS-9A, -10A, -15A)

[PROS-9](#)

- PSA persistence/recurrence
 - ▶ Life expectancy >5 y: treatment recommendations extensively revised based on No evidence of M1 and M1 disease.

[PROS-9A](#)

- Footnote q, sentence 1 modified: Observation involves monitoring the course of disease with the expectation to deliver *definitive or* palliative therapy for the development of symptoms or a change in examination or PSA that suggests symptoms are imminent.
- Footnote jj added: Treatment for a first biochemical recurrence has historically been referred to as 'salvage' therapy. In efforts to use language that is more sensitive to patients, the NCCN Guidelines for Prostate Cancer refer to treatment in this setting as 'secondary' therapy. (Also for PROS-10A)
- Footnote ll added: For patients with PSA progression who have not received treatment for a first BCR, continue monitoring or consider treatment for first PSA persistence/recurrence. See PROS-9 for RP PSA persistence/recurrent or PROS-10 for RT recurrence. (Also for PROS-10A)
- Footnote removed: If considering treatment, reinstate the PROS-10 algorithm.

[PROS-10](#)

- PSA recurrence or positive DRE
 - ▶ Life expectancy >5 y: treatment recommendations extensively revised based on No evidence of M1 and M1 disease.

[PROS-11](#)

- New page added: Second Biochemical Recurrence (BCR2).

[PROS-12](#)

- Page changed
 - ▶ New algorithm added: Treatment and Monitoring for Second Biochemical Recurrence (N0M0).
 - ▶ Algorithm removed: Treatment and Monitoring for Progressive M0 Castration-Sensitive Prostate Cancer (CSPC) After Maximal Pelvic Therapy.
- Footnote pp added: PSA level and PSADT should be considered when deciding whether to begin ADT for patients considered to have low risk disease. High-risk BCR has different definitions in different clinical trials that generally includes a PSADT ≤9 months and other adverse prognostic features.
- Footnote qq modified: ~~For patients with non-metastatic castration-sensitive disease, (by CT, MRI, or bone scan) who are not candidates for pelvic therapy, monitoring until diagnosis of metastatic disease is preferred. PSADT and Grade Group should be considered when deciding whether to begin ADT for patients with M0 disease. For ADT alone if favorable PSA response, an intermittent ADT approach should be considered. can be considered to reduce toxicity.~~

[PROS-13](#)

- New page added: Workup and Treatment of Metachronous Oligometastatic CSPC.



NCCN Guidelines Version 5.2026

Prostate Cancer

Updates in Version 1.2026 of the NCCN Guidelines for Prostate Cancer from Version 2.2025 include:

[PROS-14](#)

- Page header modified: Workup and Treatment of *Low-Volume M1 (Metachronous or Synchronous) or Synchronous Oligometastatic CSPC*.
- Synchronous low-volume metastases or Synchronous oligometastatic disease, second option modified: ADT with docetaxel and one of the following (*low-volume disease only*).
- Progression modified: Progression *on ADT ± ARPI*.

[PROS-15](#)

- Page header modified: Workup and Treatment of *High-Volume M1 CSPC*.

[PROS-15A](#)

- Footnote zz added: Concurrent MDT can be considered in select patients with oligometastatic disease. See Principles of MDT (PROS-M).
- Footnote removed: Stereotactic body RT (SBRT) to metastases can be considered in appropriate clinical situations. See Principles of Radiation Therapy (PROS-I).

[PROS-18](#)

- Systemic Therapy for M1 CRPC: Adenocarcinoma
 - ▶ Page extensively revised and reorganized by Pre-ARPI, Post-ARPI/Pre-Docetaxel, Post-ARPI/Post-Docetaxel, Additional Options Irrespective of Prior ARPI or Prior Docetaxel.

[PROS-18A](#)

- Footnote kkk modified: ~~Novel hormone ARPI~~ therapies include abiraterone, enzalutamide, darolutamide, or apalutamide. Abiraterone given as part of neoadjuvant/concomitant/adjuvant ADT with EBRT is not considered *post-ARPI prior novel hormone therapy*.
- Footnote mmm added: Principles of MDT (PROS-M).
- Footnote ooo, sentence 3 added: There are limited data to support the efficacy of Sipuleucel-T in the post-chemotherapy setting.
- Footnote removed: Principles of Radiation Therapy (PROS-I).
- Footnote removed: Visceral metastases refers to liver, lung, adrenal, peritoneal, and brain metastases. Soft tissue/lymph node sites are not considered visceral metastases.
- Footnote removed: Pan-cancer, tumor-agnostic treatments can be considered for patients with actionable mutations.
- Footnote removed: The fine-particle formulation of abiraterone can be used instead of the standard form (other recommended option).
- Footnote removed: The fine-particle (category 2B; other recommended option) or standard formulation of abiraterone can be given with single-agent niraparib as a substitute for the combination of niraparib/abiraterone tablet.

[PROS-B](#)

- Section renamed: Principles of Survivorship.

[PROS-B \(5 of 6\)](#)

- New page added: Cardiovascular Disease in Prostate Cancer.

[PROS-B \(6 of 6\)](#)

- New page added: Cardiovascular Disease in Prostate Cancer: Risk Assessment.

[PROS-D](#)

- Bullet 5 added: Consider geriatric assessment. See NCCN Guidelines for Older Adult Oncology for tools to aid optimal assessment and management of disease in older adults.
- Bullet 6 added: Refer to the NCCN Distress Thermometer and Problem List, which includes social determinants of health. See NCCN Guidelines for Distress Management (DIS-A).

[PROS-E \(2 of 5\)](#)

- Bone Imaging
 - ▶ Bullet 2 modified: Plain films, CT, MRI, or PET/CT or PET/MRI with F-18 piflufolastat prostate-specific membrane antigen (PSMA), Ga-68 PSMA-11, F-18 flutemetamol PSMA, *F-18 fluciclovine*, F-18 sodium fluoride, or C-11 choline can be considered for equivocal results on initial bone scan.



NCCN Guidelines Version 5.2026

Prostate Cancer

Updates in Version 1.2026 of the NCCN Guidelines for Prostate Cancer from Version 2.2025 include:

[PROS-E \(3 of 5\)](#)

- Full Body Imaging
 - ▶ Positron Emission Tomography
 - ◊ Bullet 1
 - Sub-bullet 1, last sentence modified: See Table 1 on PROS-E (5 of 5) ~~2 in the discussion section~~ for more details.
 - Sub-bullet 3 modified: PSMA-PET/CT or PET/MRI can be considered as an alternative to CT, MRI, and bone scans for initial staging in *unfavorable intermediate-, high-, and very-high-risk disease*, the detection...
 - Sub-bullet 4 added: PSMA-PET imaging should only be used in the setting of M1 CRPC to determine if a patient is a candidate for Lu-177-PSMA-617.
 - Sub-bullet 5 added: Changes in systemic therapy should not be made solely based on a positive PSMA-PET in patients with M0 CRPC.
 - Sub-bullet 6 added: Bone scans can be considered to confirm osseous uptake on PET scans.
 - Sub-bullet 8 modified: C-11 choline *or F-18 fluciclovine* PET/CT or PET/MRI...
 - Sub-bullet 9 modified: Studies suggest that PSMA-PET imaging has a higher sensitivity than C-11 choline *or F-18 fluciclovine* PET imaging, especially at very low PSA levels (*eg, <0.5 ng/mL*).

[PROS-E \(4 of 5\)](#)

- Sub-bullet modified (continued from PROS-E [3 of 5]) ...interpretation of scans. *The use of a standardized reporting system is encouraged. Several reporting systems have been proposed but will not have been validated or widely used.*
- Sub-bullet 3 modified: Table 1 on PROS-E (5 of 5) ~~in the discussion section~~ provides a summary of the main imaging tracers utilized for study in prostate cancer ~~both before definitive therapy and at recurrence~~.
- Imaging as Workup for Progression
 - ▶ Bullet 2 modified: Imaging for patients with progressive ~~m~~CRPC should included chest CT, bone imaging, and abdomen/pelvis CT with contrast or abdomen/pelvis MRI with and without contrast.
 - ▶ Bullet 3 modified: There is a lack of evidenced to support the use of PET imaging in ~~this the~~ CRPC setting *and it should only be used in the setting of M1 CRPC to determine if a patient is a candidate for Lu-177-PSMA-617. Changes in treatment should not be made solely based on a positive PSMA-PET in patients with M0 CRPC.*

[PROS-E \(5 of 5\)](#)

- New table added: FDA-Cleared PET Imaging Tracers Studied in Prostate Cancer.
- Footnote a added: Interpret with caution. Wherever possible, studies were included that used histopathologic confirmation, but not all studies used confirmatory histology as the gold standard. Values may vary depending upon the site of the lesion and phase of the disease process.
- Footnote b added: CLR: Correct localization rate. Patient-level positive predictive value + anatomic lesion co-localization. Preferred over sensitivity and specificity in analyses of patients with BCR.

[PROS-F](#)

- Principles of Active Surveillance: Section extensively revised.

[PROS-G \(1 of 6\)](#)

- Neoadjuvant, Concurrent, and/or Adjuvant ADT with RT
 - ▶ Bullet 4 modified: *First* M0 RP PSA persistence/recurrence ~~(without maximal pelvic therapy)~~.
 - ▶ Notes
 - ◊ Bullet 1 added: ADT is not recommended with RT for most patients with favorable intermediate-risk prostate cancer. If it is given (see Principles of Risk Stratification), the duration should be short term (4–6 months).
 - ◊ Bullet 4 modified: For high-risk ~~and very-high-risk~~ prostate cancer...
 - ◊ Bullet 6 modified: *First* M0 PSA persistence/recurrence:
 - Sub-bullet 3 added: For patients treated with secondary RT in the setting of first RP recurrence, if ADT is given, it should be for a duration of 6 to 24 months.



NCCN Guidelines Version 5.2026

Prostate Cancer

Updates in Version 1.2026 of the NCCN Guidelines for Prostate Cancer from Version 2.2025 include:

[PROS-G \(2 of 6\)](#)

- ADT for Patients with M0 CSPP, bullet 2 modified: *First* M0 RT recurrence ~~(without maximal pelvic therapy).~~

[PROS-G \(3 of 6\)](#)

- ADT for Patients with M0 CSPP *BCR2 After Maximal Pelvic Therapy*.
 - ▶ Useful in Certain Circumstances
 - ◊ Bullet 1 modified: Enzalutamide with or without leuprolide *(for high-risk BCR2)*.
 - ◊ Bullet 2 modified: Apalutamide with LHRH agonist or LHRH antagonist ~~(category 2B)~~ *(for high-risk BCR2)*.
 - ▶ Notes, bullet 1 modified: Monitoring until diagnosis of metastatic disease is preferred for patients with *M0 CSPP in the low-risk BCR2 setting non-metastatic castration-sensitive disease who are not candidates for pelvic therapy*.
 - ▶ Footnote c added: For patients receiving MDT with ADT, the ADT options are LHRH agonist or LHRH antagonist.

[PROS-G \(5 of 6\)](#)

- ADT Monotherapy
 - ▶ Notes
 - ◊ Bullet 1 modified: ADT monotherapy is appropriate for patients with life expectancy ≤5 years whose cancer progressed on observation of localized disease, who are symptomatic or who have N1M0 disease ~~and in select patients with high- or very-high-risk disease, where complications, such as hydronephrosis or metastasis, can be expected within 5 years.~~
 - ◊ Bullet 2 modified: ADT monotherapy is also used for asymptomatic patients with ~~high-risk, very-high-risk, and~~ regional disease and life expectancy ≤5 years whether or not RT is given.
 - Footnote h added: In the mCRPC setting, this option should only be used for select patients who are not candidates for other recommended mCRPC therapies.

[PROS-G \(6 of 6\)](#)

- Optimal ADT, bullet removed: Data are limited on long-term adherence to oral relugolix and the potential effects on optimal ADT. Ongoing monitoring for sustained suppression of testosterone (<50 ng/dL) can be considered, and relugolix may not be a preferred agent if adherence to the prescribed regimen is uncertain.

[PROS-I](#)

- New section added: Principles of Focal/Subtotal Therapy or Whole Gland Ablative Therapy.

[PROS-J](#)

- Principles of Radiation Therapy: Section extensively revised.

[PROS-K](#)

- Pelvic Lymph Node Dissection
 - ▶ Bullet 1, sub-bullet 5 modified: PLND can be performed using an open, laparoscopic, or robotic technique *and can provide staging and prognostic information*.
 - ▶ Bullet removed: While PLND at the time of RP has not been shown to improve oncologic outcomes, it can provide staging and prognostic information.

[PROS-L \(1 of 2\)](#)

- Bullet 1 modified: Patients with biopsy-proven recurrence in the prostate after prior RT and without distant metastatic disease can be considered for local therapy. *Monitoring is also an option (see PROS-10). ADT may also be included, but it should not be reflexively ordered.*
- Bullet 3 modified: Local therapy options for patients with recurrence in the prostate ~~only~~ include:...
 - ▶ Sub-bullet 2 modified: High-intensity focused ultrasound (HIFU) ~~(category 2B)~~.
 - ▶ Sub-bullet 3 added: Irreversible electroporation (IRE) (category 2B).
 - ▶ Sub-bullet removed: Non-surgical strategies.
- Bullet and sub-bullets removed: Local therapy options for patients with recurrence in the regional nodes with or without prostate recurrence include: ADT + pelvic lymph node radiation (if not previously done), ADT + pelvic lymph node reirradiation (category 2B), ADT + PLND (category 2B), Pelvic lymph node radiation, PLND.



NCCN Guidelines Version 5.2026

Prostate Cancer

Updates in Version 1.2026 of the NCCN Guidelines for Prostate Cancer from Version 2.2025 include:

[PROS-L \(1 of 2\)\(continued\)](#)

- Footnote a added: Reirradiation with LDR brachytherapy, HDR brachytherapy, and SBRT is supported by phase II trials with >36 months of median follow-up and should be strongly considered as an option in patients with >2 years interval from prior radiation who do not have ongoing moderate or higher grade radiation-associated toxicities. Phase II data with >36 months of median follow-up are available, albeit to a more limited scope, for secondary RP, cryotherapy, and HIFU.
- Footnote b added: Currently, all phase II and/or prospectively accrued registries reporting oncologic outcomes with >36 months of follow-up have pursued whole gland treatments. Focal therapy for prostate-only recurrences is the subject of active investigation. See Principles of Focal/Subtotal Therapy or Whole Gland Ablative Therapy (PROS-I).
- Footnote c added: There are no prospective data to guide the use and duration of ADT with reirradiation for prostate-only recurrences. The Panel recommends extrapolation from the definitive treatment setting to suggest that adding ADT or prolonging it in this setting likely offers an improvement in biochemical control. See Principles of Radiation Therapy (PROS-J).

[PROS-M](#)

- New section added: Principles of Metastasis-Directed Therapy.

[PROS-N \(1 of 4\)](#)

- Non-Hormonal Systemic Therapy for M1 Castration-Sensitive Prostate Cancer
 - ▶ Bullet 1 modified: ~~Patients with low-volume synchronous or high-volume castration-sensitive metastatic prostate cancer who are fit for chemotherapy should be considered for Triple therapy with ADT, certain ARPIs, and docetaxel is an option for certain patients with mCSPC who are fit for chemotherapy~~ based on phase 3 studies:

[PROS-N \(2 of 4\)](#)

- Non-Hormonal Systemic Therapy for M1 CRPC
 - ▶ Chemotherapy
 - ◊ Bullet 1
 - Sub-bullet 4 modified: Docetaxel retreatment can be attempted after progression on *an ARPI and docetaxel* ~~novel hormone therapy~~ in patients with mCRPC whose cancer has not yet demonstrated definitive evidence of progression on prior docetaxel therapy in *either* the castration-sensitive setting *or the mCRPC setting*.
 - Sub-bullet 5 added: Rare patients may have exposure to docetaxel without prior ARPI (ie, in the setting of EBRT to the primary tumor for low-volume synchronous mCSPC). Docetaxel rechallenge is not recommended for such patients in the pre-ARPI mCRPC setting.
 - ▶ Bullet 2, sub-bullet 5 added: Biweekly cabazitaxel at 16 mg/m² dose with G-CSF in patients ≥65 years in a Phase 3 trial. Clinical outcomes were comparable between the 2 groups.

[PROS-N \(3 of 4\)](#)

- Non-Hormonal Systemic Therapy for M1 CRPC
 - ▶ PARP Inhibitors With or Without ARPIs: Subsection extensively revised.
 - ▶ Immunotherapy
 - ◊ Bullet 2 modified: Pembrolizumab is an option for certain patients with *MSI-H/dMMR* mCRPC and *MSI-H, dMMR, or TMB ≥10 mut/mB* (PROS-18).
 - Sub-bullet 2 added: Limited data suggest that pembrolizumab may be associated with some benefit in patients with mCRPC and TMB ≥10 mut/mB. Lenis AT, et al. Clin Canc Res 2024;30:3894-3903.
 - ◊ Bullet removed: Patients with asymptomatic or minimally symptomatic mCRPC may consider immunotherapy.

[PROS-N \(4 of 4\)](#)

- Table added: Table 1. PARP Inhibitors with or without ARPIs.



NCCN Guidelines Version 5.2026

Prostate Cancer

INITIAL PROSTATE CANCER DIAGNOSIS^{a,b,c,d}

WORKUP

Clinically localized prostate cancer
(Any T, N0, M0 or Any T, NX, MX)

- Perform physical exam
- Perform digital rectal examination (DRE) to confirm clinical stage
- Perform and/or collect prostate-specific antigen (PSA)
- Review diagnostic prostate biopsies
- Estimate life expectancy ([Principles of Life Expectancy Estimation \[PROS-A\]](#))
- Inquire about known high-risk germline mutations and family history^e
 - ▶ Perform somatic and/or germline testing as appropriate^e
- Assess quality-of-life measures^f

Initial Risk
Stratification and
Staging Workup for
Clinically Localized
Disease ([PROS-2](#))

Regional prostate cancer
(Any T, N1, M0)

- Perform physical examination
- Perform bone and soft tissue imaging for staging^g
- Consider DRE to confirm clinical stage
- Perform and/or collect PSA and calculate PSA doubling time (PSADT)
- Estimate life expectancy ([Principles of Life Expectancy Estimation \[PROS-A\]](#))
- Inquire about known high-risk germline mutations and family history^e
 - ▶ Perform somatic and/or germline testing as appropriate^e
- Assess quality-of-life measures^f

Regional Prostate
Cancer ([PROS-7](#))

Workup and Treatment
of Low-Volume M1
(Metachronous
or Synchronous)
or Synchronous
Oligometastatic^h
Castration-Sensitive
Prostate Cancer (CSPC)
([PROS-14](#))

Metastatic prostate cancer
(Any T, Any N, M1)

Workup and
Treatment of
High-Volume M1
CSPC ([PROS-15](#))

^a See [NCCN Guidelines for Older Adult Oncology](#) for tools to aid optimal assessment and management of disease in older adults.

^b [NCCN Guidelines for Prostate Cancer Early Detection](#).

^c [Principles of Survivorship in Prostate Cancer \(PROS-B\)](#).

^d Refer to the NCCN Distress Thermometer and Problem List, which includes social determinants of health. See [NCCN Guidelines for Distress Management \(DIS-A\)](#).

^e [Principles of Genetics and Molecular/Biomarker Analysis \(PROS-C\)](#).

^f [Principles of Quality of Life and Shared Decision-Making \(PROS-D\)](#).

^g [Principles of Imaging \(PROS-E\)](#).

^h See Number of Metastatic Sites in the [Principles of MDT \(PROS-M\)](#).

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

INITIAL RISK STRATIFICATION AND STAGING WORKUP FOR CLINICALLY LOCALIZED DISEASEⁱ

Risk Group	Clinical/Pathologic Features (Staging, ST-1)		Additional Evaluation ^{g,i}	Initial Therapy
Low ^j	Has all of the following: • cT1–cT2a • Grade Group 1 • PSA <10 ng/mL		• Confirmatory testing can be used to assess the appropriateness of active surveillance (PROS-F 2 of 5)	PROS-3
Intermediate ^j	Has all of the following: • No high-risk group features • No very-high-risk group features • Has one or more intermediate risk factors (IRFs): ▶ cT2b–cT2c ▶ Grade Group 2 or 3 ▶ PSA 10–20 ng/mL	Favorable intermediate	Has all of the following: • 1 IRF • Grade Group 1 or 2 • <50% biopsy cores positive (eg, <6 of 12 cores) ^k	• Confirmatory testing can be used to assess the appropriateness of active surveillance (PROS-F 2 of 5) PROS-4
		Unfavorable intermediate	Has one or more of the following: • 2 or 3 IRFs • Grade Group 3 • ≥50% biopsy cores positive (eg, ≥6 of 12 cores) ^k	• Soft tissue imaging and consider bone imaging ^g ▶ If regional metastases are found, see PROS-7 ▶ If distant metastases are found, see PROS-14 for Low-Volume M1 (Metachronous or Synchronous) or Synchronous Oligometastatic ^h CSPC or PROS-15 for High-Volume M1 CSPC PROS-5
High	Has one or more high-risk features, but does not meet criteria for very high risk: • cT3–cT4 • Grade Group 4 or Grade Group 5 • PSA >20 ng/mL		Bone and soft tissue imaging ^g ▶ If regional metastases are found, see PROS-7 ▶ If distant metastases are found, see PROS-14 for Low-Volume M1 (Metachronous or Synchronous) or Synchronous Oligometastatic ^h CSPC or PROS-15 for High-Volume M1 CSPC	PROS-6
Very high	Has at least two of the following: • cT3–cT4 • Grade Group 4 or 5 • PSA >40 ng/mL		Bone and soft tissue imaging ^g ▶ If regional metastases are found, see PROS-7 ▶ If distant metastases are found, see PROS-14 for Low-Volume M1 (Metachronous or Synchronous) or Synchronous Oligometastatic ^h CSPC or PROS-15 for High-Volume M1 CSPC	PROS-6

Note: All recommendations are category 2A unless otherwise indicated.

[Footnotes on PROS-2A](#)



INITIAL RISK STRATIFICATION AND STAGING WORKUP FOR CLINICALLY LOCALIZED DISEASE

^g [Principles of Imaging \(PROS-E\)](#).

^h See Number of Metastatic Sites in the [Principles of MDT \(PROS-M\)](#).

ⁱ Tumor-based molecular assays and germline genetic testing are other tools that can assist with risk stratification. See CRIT-6 in the [NCCN Guidelines for Genetic/Familial High-Risk Assessment: Breast, Ovarian, Pancreatic, and Prostate](#) and HRS-3 in the [NCCN Guidelines for Genetic/Familial High-Risk Assessment: Colorectal, Endometrial, and Gastric](#) to determine if a patient is an appropriate candidate for germline genetic testing, and see [Principles of Risk Stratification and Biomarkers \(PROS-H\)](#) to determine if a patient is an appropriate candidate for tumor-based molecular assays.

^j For patients who are asymptomatic in low- and intermediate-risk groups with life expectancy ≤ 5 years, no imaging or treatment is indicated until the patient becomes symptomatic, at which time imaging can be performed [[Principles of Imaging \(PROS-E\)](#)] and androgen deprivation therapy (ADT) should be given [[Principles of Androgen Deprivation Therapy \(PROS-G\)](#)].

^k Percentage of positive cores in the intermediate-risk group is based on biopsies that include systematic biopsies with or without targeted MRI-guided biopsies. The Panel considers biopsies from a single region of interest (ROI) to count as a single sample.

^l Bone imaging should be performed for any patient with symptoms consistent with bone metastases.

Note: All recommendations are category 2A unless otherwise indicated.




NCCN Guidelines Version 5.2026

Prostate Cancer

LOW-RISK GROUP

EXPECTED
PATIENT
SURVIVAL^m

INITIAL THERAPYⁿ



Note: All recommendations are category 2A unless otherwise indicated.

[Footnotes on PROS-7A](#)




NCCN Guidelines Version 5.2026

Prostate Cancer

FAVORABLE INTERMEDIATE-RISK GROUP

EXPECTED
PATIENT
SURVIVAL^m

INITIAL THERAPYⁿ



Note: All recommendations are category 2A unless otherwise indicated.

[Footnotes on PROS-7A](#)




NCCN Guidelines Version 5.2026

Prostate Cancer

UNFAVORABLE INTERMEDIATE-RISK GROUP

EXPECTED
PATIENT
SURVIVAL^m

INITIAL THERAPYⁿ



Note: All recommendations are category 2A unless otherwise indicated.

[Footnotes on PROS-7A](#)




NCCN Guidelines Version 5.2026

Prostate Cancer

HIGH- OR VERY-HIGH-RISK GROUP

EXPECTED
PATIENT
SURVIVAL^m

INITIAL THERAPYⁿ



Note: All recommendations are category 2A unless otherwise indicated.


[Footnotes on PROS-7A](#)



NCCN Guidelines Version 5.2026

Prostate Cancer

REGIONAL PROSTATE CANCER (ANY T, N1, M0)



Note: All recommendations are category 2A unless otherwise indicated.

[Footnotes on PROS-7A](#)



FOOTNOTES

- ^c [Principles of Survivorship in Prostate Cancer \(PROS-B\)](#).
^h See Number of Metastatic Sites in the [Principles of MDT \(PROS-M\)](#).
^j For patients who are asymptomatic in low- and intermediate-risk groups with life expectancy ≤ 5 years, no imaging or treatment is indicated until the patient becomes symptomatic, at which time imaging can be performed [[Principles of Imaging \(PROS-E\)](#)] and ADT should be given [[Principles of Androgen Deprivation Therapy \(PROS-G\)](#)].
^m [Principles of Life Expectancy Estimation \(PROS-A\)](#).
ⁿ [Principles of Focal/Subtotal Therapy or Whole Gland Ablative Therapy \(PROS-I\)](#).
^o Active surveillance involves actively monitoring the course of disease with the expectation to intervene with potentially curative therapy if the cancer progresses. See [Principles of Active Surveillance and Observation \(PROS-F\)](#).
^p Confirmatory testing can be used to assess the appropriateness of active surveillance ([PROS-F 2 of 5](#)). If higher grade and/or higher T stage is found during confirmatory testing, see [PROS-2](#).
^q Observation involves monitoring the course of disease with the expectation to deliver definitive or palliative therapy for the development of symptoms or a change in examination or PSA that suggests symptoms are imminent. See [Principles of Active Surveillance and Observation \(PROS-F\)](#).
^r Criteria for progression are not well-defined and require physician judgment; however, a change in risk group strongly implies disease progression. See [Discussion](#).
^s The Panel recognizes that there is heterogeneity across the low-risk group, and that some factors may be associated with an increased probability of near-term grade reclassification, including high PSA density, a high number of positive cores (eg, ≥ 3), high genomic risk (from tissue-based molecular tumor analysis), and/or a known BRCA2 germline mutation. In some of these cases, upfront treatment with RP or prostate RT may be preferred based on shared decision-making with the patient. See [Principles of Active Surveillance and Observation \(PROS-F\)](#).
^t [Principles of Radiation Therapy \(PROS-J\)](#).
^u [Principles of Surgery \(PROS-K\)](#).

- ^v RTOG-ASTRO (Radiation Therapy Oncology Group - American Society for Radiation Oncology) Phoenix Consensus: 1) PSA increase by ≥ 2 ng/mL above the nadir PSA is the standard definition for PSA recurrence after external beam RT (EBRT) with or without hormone therapy; and 2) a recurrence evaluation should be considered when PSA has been confirmed to be increasing after radiation even if the increase above nadir is < 2 ng/mL, especially in candidates for secondary local therapy who are young and healthy. Retaining a strict version of the ASTRO definition allows comparison with a large existing body of literature. Rapid increase of PSA may warrant evaluation (prostate biopsy) prior to meeting the Phoenix definition, especially in patients who are younger or healthier.
^w PSA persistence/recurrence after RP is defined as when PSA does not fall to undetectable levels (PSA persistence) or undetectable PSA after RP with a subsequent detectable PSA that increases on ≥ 2 determinations (PSA recurrence) or increases to PSA > 0.1 ng/mL. Trials indicating noninferiority of early RT compared with adjuvant RT after RP have used a PSA threshold of 0.1 or 0.2 ng/mL to trigger treatment. Imaging and treatment at lower PSA levels may be appropriate in patients at high risk for progression based on pretreatment risk factors, pathologic parameters, timing of recurrence, and genomic classifier (GC) score, among other factors.
^x Adverse pathologic features include: positive margin(s), seminal vesicle invasion, or extracapsular extension.
^y Particular consideration to active surveillance may be appropriate for those patients in the favorable intermediate-risk group with a low percentage of Gleason pattern 4 cancer, low tumor volume, low PSA density, and/or low genomic risk (from tissue-based molecular tumor analysis). See [Principles of Active Surveillance and Observation \(PROS-F\)](#).
^z Active surveillance of unfavorable intermediate and high-risk clinically localized cancers is not recommended in patients with a life expectancy > 10 years (category 1).
^{aa} For details on the use of ADT and androgen receptor pathway inhibitors (ARPIs), see [Principles of Androgen Deprivation Therapy \(PROS-G\)](#) and [Discussion](#).
^{bb} RP + pelvic lymph node dissection (PLND) can be considered in patients who are younger and healthier without tumor fixation to the pelvic sidewall.
^{cc} Monitoring is not preferred for patients with multiple high-risk features.
^{dd} There is limited evidence that RP + PLND is beneficial in the setting of node-positive disease. Use of this approach should be limited to patients with > 10 -year life expectancy and resectable disease and should be used in the context of a clinical trial or planned multimodality approach.

Note: All recommendations are category 2A unless otherwise indicated.




NCCN Guidelines Version 5.2026

Prostate Cancer

MONITORING

See [NCCN Guidelines for Survivorship](#)

RECURRENCE



Note: All recommendations are category 2A unless otherwise indicated.

[Footnotes on PROS-8A](#)



MONITORING AND RECURRENCE FOOTNOTES

^g [Principles of Imaging \(PROS-E\)](#).

^h See Number of Metastatic Sites in the [Principles of MDT \(PROS-M\)](#).

^v RTOG-ASTRO Phoenix Consensus: 1) PSA increase by ≥ 2 ng/mL above the nadir PSA is the standard definition for PSA recurrence after EBRT with or without hormone therapy; and 2) a recurrence evaluation should be considered when PSA has been confirmed to be increasing after radiation even if the increase above nadir is < 2 ng/mL, especially in candidates for secondary local therapy who are young and healthy. Retaining a strict version of the ASTRO definition allows comparison with a large existing body of literature. Rapid increase of PSA may warrant evaluation (prostate biopsy) prior to meeting the Phoenix definition, especially in patients who are younger or healthier.

^w PSA persistence/recurrence after RP is defined as when PSA does not fall to undetectable levels (PSA persistence) or undetectable PSA after RP with a subsequent detectable PSA that increases on ≥ 2 determinations (PSA recurrence) or increases to PSA > 0.1 ng/mL. Trials indicating non-inferiority of early RT compared with adjuvant RT after RP have used a PSA threshold of 0.1 or 0.2 ng/mL to trigger treatment. Imaging and treatment at lower PSA levels may be appropriate in patients at high risk for progression based on pretreatment risk factors, pathologic parameters, timing of recurrence, and GC score, among other factors.

^{ee} PSA as frequently as every 3 mo may be necessary to clarify disease status, especially in patients at high risk of recurrence.

^{ff} Document castrate levels of testosterone if clinically indicated. Workup for progression should include bone and soft tissue evaluation. See [Principles of Imaging \(PROS-E\)](#).

⁹⁹ Treatment for patients with life expectancy ≤ 5 y whose cancer progressed on observation of localized disease is ADT. See [Principles of Androgen Deprivation Therapy \(PROS-G\)](#).

Note: All recommendations are category 2A unless otherwise indicated.




NCCN Guidelines Version 5.2026

Prostate Cancer

RADICAL PROSTATECTOMY PSA PERSISTENCE/RECURRENCE^{hh}

TREATMENT FOR FIRST PSA PERSISTENCE/RECURRENCE^{jj}



Note: All recommendations are category 2A unless otherwise indicated.

[Footnotes on PROS-9A](#)



RADICAL PROSTATECTOMY PSA PERSISTENCE/RECURRENCE FOOTNOTES

^c [Principles of Survivorship in Prostate Cancer \(PROS-B\)](#).

^g [Principles of Imaging \(PROS-E\)](#).

^h See Number of Metastatic Sites in the [Principles of MDT \(PROS-M\)](#).

^q Observation involves monitoring the course of disease with the expectation to deliver definitive or palliative therapy for the development of symptoms or a change in examination or PSA that suggests symptoms are imminent. See [Principles of Active Surveillance and Observation \(PROS-F\)](#).

^t [Principles of Radiation Therapy \(PROS-J\)](#).

^w PSA persistence/recurrence after RP is defined as when PSA does not fall to undetectable levels (PSA persistence) or undetectable PSA after RP with a subsequent detectable PSA that increases on ≥2 determinations (PSA recurrence) or increases to PSA >0.1 ng/mL. Trials indicating noninferiority of early RT compared with adjuvant RT after RP have used a PSA threshold of 0.1 or 0.2 ng/mL to trigger treatment. Imaging and treatment at lower PSA levels may be appropriate in patients at high risk for progression based on pretreatment risk factors, pathologic parameters, timing of recurrence, and GC score, among other factors.

^{aa} For details on the use of ADT and ARPIs, see [Principles of Androgen Deprivation Therapy \(PROS-G\)](#) and [Discussion](#).

^{ff} Document castrate levels of testosterone if clinically indicated. Workup for progression should include bone and soft tissue evaluation. See [Principles of Imaging \(PROS-E\)](#).

^{gg} Treatment for patients with life expectancy ≤5 y whose cancer progressed on observation of localized disease is ADT. See [Principles of Androgen Deprivation Therapy \(PROS-G\)](#).

^{hh} Recommendations for RP PSA persistence/recurrence may also apply to patients with undetectable PSA with multiple adverse features or lymph node metastases if treatment is being considered.

ⁱⁱ [Principles of Risk Stratification and Biomarkers \(PROS-H\)](#).

^{jj} Treatment for a first biochemical recurrence has historically been referred to as 'salvage' therapy. In efforts to use language that is more sensitive to patients, the NCCN Guidelines for Prostate Cancer refer to treatment in this setting as 'secondary' therapy.

^{kk} Monitoring should include physical exam, PSA every 3–6 months, and imaging for symptoms or increasing PSA.

^{ll} For patients with PSA progression who have not received treatment for a first biochemical recurrence (BCR), continue monitoring or consider treatment for first PSA persistence/recurrence. See [PROS-9](#) for RP PSA persistence/recurrence or [PROS-10](#) for RT recurrence.

Note: All recommendations are category 2A unless otherwise indicated.




NCCN Guidelines Version 5.2026

Prostate Cancer

RADIATION THERAPY RECURRENCE

TREATMENT FOR FIRST RECURRENCE^{jj}



[Footnotes on PROS-10A](#)

Note: All recommendations are category 2A unless otherwise indicated.



RADIATION THERAPY RECURRENCE FOOTNOTES

^c [Principles of Survivorship in Prostate Cancer \(PROS-B\)](#).

^g [Principles of Imaging \(PROS-E\)](#).

^h See Number of Metastatic Sites in the [Principles of MDT \(PROS-M\)](#).

^q Observation involves monitoring the course of disease with the expectation to deliver definitive or palliative therapy for the development of symptoms or a change in examination or PSA that suggests symptoms are imminent. See [Principles of Active Surveillance and Observation \(PROS-F\)](#).

^v RTOG-ASTRO Phoenix Consensus: 1) PSA increase by ≥ 2 ng/mL above the nadir PSA is the standard definition for PSA recurrence after EBRT with or without hormone therapy; and 2) A recurrence evaluation should be considered when PSA has been confirmed to be increasing after radiation even if the increase above nadir is < 2 ng/mL, especially in candidates for secondary local therapy who are young and healthy. Retaining a strict version of the ASTRO definition allows comparison with a large existing body of literature. Rapid increase of PSA may warrant evaluation (prostate biopsy) prior to meeting the Phoenix definition, especially in patients who are younger or healthier.

^{aa} For details on the use of ADT and ARPIs, see [Principles of Androgen Deprivation Therapy \(PROS-G\)](#) and [Discussion](#).

^{ff} Document castrate levels of testosterone if clinically indicated. Workup for progression should include bone and soft tissue evaluation. See [Principles of Imaging \(PROS-E\)](#).

^{gg} Treatment for patients with life expectancy ≤ 5 y whose cancer progressed on observation of localized disease is ADT. See [Principles of Androgen Deprivation Therapy \(PROS-G\)](#).

^{jj} Treatment for a first biochemical recurrence has historically been referred to as 'salvage' therapy. In efforts to use language that is more sensitive to patients, the NCCN Guidelines for Prostate Cancer refer to treatment in this setting as 'secondary' therapy.

^{kk} Monitoring should include physical exam, PSA every 3–6 mo, and imaging for symptoms or increasing PSA.

^{ll} For patients with PSA progression who have not received treatment for a first BCR, continue monitoring or consider treatment for first PSA persistence/recurrence. See [PROS-9](#) for RP PSA persistence/recurrence or [PROS-10](#) for RT recurrence.

^{mm} PSADT can be calculated to inform nomogram use and counseling.

ⁿⁿ PSADT and Grade Group should be considered when deciding whether to begin ADT. See [Principles of Androgen Deprivation Therapy \(PROS-G\)](#).

^{oo} [Principles of Local Secondary Post-Recurrence Therapy \(PROS-L\)](#).


Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

WORKUP FOR SECOND BIOCHEMICAL RECURRENCE (BCR2)




⁹ [Principles of Imaging \(PROS-E\)](#).

^h See Number of Metastatic Sites in the [Principles of MDT \(PROS-M\)](#).

Note: All recommendations are category 2A unless otherwise indicated.



TREATMENT AND MONITORING FOR SECOND BIOCHEMICAL RECURRENCE (N0M0)



^c [Principles of Survivorship in Prostate Cancer \(PROS-B\)](#).

^h See Number of Metastatic Sites in the [Principles of MDT \(PROS-M\)](#).

^{aa} For details on the use of ADT and ARPIs, see [Principles of Androgen Deprivation Therapy \(PROS-G\)](#) and [Discussion](#).

^{ff} Document castrate levels of testosterone if clinically indicated. Workup for progression should include bone and soft tissue evaluation. See [Principles of Imaging \(PROS-E\)](#).

^{kk} Monitoring should include physical exam, PSA every 3–6 months, and imaging for symptoms or increasing PSA.

^{pp} PSA level and PSADT should be considered when deciding whether to begin ADT for patients considered to have lower risk disease. High-risk BCR has different definitions in different clinical trials that generally include a PSADT ≤9 months and other adverse prognostic features.

^{qq} If favorable PSA response, an intermittent ADT approach should be considered.

^{rr} Enzalutamide with or without leuprolide is an option for patients who have the following high-risk criteria: M0 by CT, MRI, or bone scan; PSADT ≤9 months; PSA ≥2 ng/mL above nadir after RT or ≥1 ng/mL after RP with or without postoperative RT; and not considered a candidate for pelvic-directed therapy (Freedland SJ, et al. N Engl J Med 2023;389:1453-1465). See [Principles of Androgen Deprivation Therapy \(PROS-G\)](#).

^{ss} Apalutamide plus ADT is an option for patients with biochemical recurrence after RP who meet the following high-risk criteria: PSADT ≤9 months; PSA ≥0.5 ng/mL; and prior adjuvant or secondary RT or not considered a candidate for RT (Aggarwal R, et al. J Clin Oncol 2024;42:1114-1123.) See [Principles of Androgen Deprivation Therapy \(PROS-G\)](#).

Note: All recommendations are category 2A unless otherwise indicated.




NCCN Guidelines Version 5.2026

Prostate Cancer

WORKUP AND TREATMENT OF METACHRONOUS OLIGOMETASTATIC CSPC^{c,h,tt,uu,vv,ww}

WORKUP FOR METASTASES^{xx}



^c [Principles of Survivorship in Prostate Cancer \(PROS-B\)](#).

^e [Principles of Genetics and Molecular/Biomarker Analysis \(PROS-C\)](#).

^f [Principles of Quality of Life and Shared Decision-Making \(PROS-D\)](#).

^g [Principles of Imaging \(PROS-E\)](#).

^h See Number of Metastatic Sites in the [Principles of MDT \(PROS-M\)](#).

^{aa} For details on the use of ADT and ARPIs, see [Principles of Androgen Deprivation Therapy \(PROS-G\)](#) and [Discussion](#).

^{ff} Document castrate levels of testosterone if clinically indicated. Workup for progression should include bone and soft tissue evaluation. See [Principles of Imaging \(PROS-E\)](#).

^{tt} EBRT to sites of bone metastases can be considered if metastases are in weight-bearing bones or if the patient is symptomatic.

^{uu} Bone antiresorptive therapy is indicated for elevated fracture risk based upon FRAX in the castration-sensitive setting. See [PROS-B](#).

^{vv} The term "castration-sensitive" is used to define disease in patients who have not been treated with ADT and those who are not on ADT at the time of progression. The NCCN Prostate Cancer Panel uses the term "castration-sensitive" even when patients have had neoadjuvant, concurrent, or adjuvant ADT as part of RT provided they have recovered testicular function.

^{ww} ADT is strongly recommended in combination therapy for metastatic castration-sensitive disease. The use of ADT monotherapy in metastatic castration-sensitive disease is discouraged unless there are clear contraindications to combination therapy. If ADT monotherapy is given, intermittent ADT can be considered to reduce toxicity. See [Principles of Androgen Deprivation Therapy \(PROS-G\)](#).

^{xx} ADT alone ([PROS-G](#)) or observation are recommended for asymptomatic patients with metastatic disease or M0 CRPC and life expectancy ≤5 years.

^{yy} See [Principles of MDT \(PROS-M\)](#).

Note: All recommendations are category 2A unless otherwise indicated.




NCCN Guidelines Version 5.2026

Prostate Cancer

WORKUP AND TREATMENT OF LOW-VOLUME M1 (METACHRONOUS OR SYNCHRONOUS) OR SYNCHRONOUS OLIGOMETASTATIC CSPC^{c,h,tt,uu,vv,ww}

WORKUP FOR METASTASES^{xx}



Note: All recommendations are category 2A unless otherwise indicated.

[Footnotes on PROS-15A](#)




NCCN Guidelines Version 5.2026

Prostate Cancer

WORKUP AND TREATMENT OF HIGH-VOLUME M1 CSPC^{c,tt,uu,vv,ww,ccc}

WORKUP FOR METASTASES^{xx}



Note: All recommendations are category 2A unless otherwise indicated.

[Footnotes on PROS-15A](#)



FOOTNOTES

^c [Principles of Survivorship in Prostate Cancer \(PROS-B\)](#).

^e [Principles of Genetics and Molecular/Biomarker Analysis \(PROS-C\)](#).

^f [Principles of Quality of Life and Shared Decision-Making \(PROS-D\)](#).

^g [Principles of Imaging \(PROS-E\)](#).

^h See Number of Metastatic Sites in the [Principles of MDT \(PROS-M\)](#).

^t [Principles of Radiation Therapy \(PROS-J\)](#).

^{aa} For details on the use of ADT and ARPIs, see [Principles of Androgen Deprivation Therapy \(PROS-G\)](#) and [Discussion](#).

^{ff} Document castrate levels of testosterone if clinically indicated. Workup for progression should include bone and soft tissue evaluation. See [Principles of Imaging \(PROS-E\)](#).

^{tt} EBRT to sites of bone metastases can be considered if metastases are in weight-bearing bones or if the patient is symptomatic.

^{uu} Bone antiresorptive therapy is indicated for elevated fracture risk based upon FRAX in the castration-sensitive setting. See [PROS-B](#).

^{vv} The term "castration-sensitive" is used to define disease in patients who have not been treated with ADT and those who are not on ADT at the time of progression. The NCCN Prostate Cancer Panel uses the term "castration-sensitive" even when patients have had neoadjuvant, concurrent, or adjuvant ADT as part of RT provided they have recovered testicular function.

^{ww} ADT is strongly recommended in combination therapy for metastatic castration-sensitive disease. The use of ADT monotherapy in metastatic castration-sensitive disease is discouraged unless there are clear contraindications to combination therapy. If ADT monotherapy is given, intermittent ADT can be considered to reduce toxicity. See [Principles of Androgen Deprivation Therapy \(PROS-G\)](#).

^{xx} ADT alone ([PROS-G](#)) or observation are recommended for asymptomatic patients with metastatic disease or M0 CRPC and life expectancy ≤5 years.

^{zz} Concurrent MDT can be considered in select patients with oligometastatic disease. See [Principles of MDT \(PROS-M\)](#).

^{aaa} EBRT to the primary tumor is associated with an overall survival (OS) benefit in patients with low metastatic burden at the time of diagnosis of metastatic disease, which is defined by bone scan and CT or MRI as either non-regional, lymph-node-only disease OR <4 bone metastases and without visceral/other metastasis (Ali A, et al. JAMA Oncol 2021;7:555-563). See [Principles of Radiation Therapy \(PROS-J\)](#).

^{bbb} Niraparib/abiraterone can be used following the completion of docetaxel in patients receiving triplet therapy or following EBRT to the primary tumor. The use of this combination in patients with low-volume disease is controversial because 78% of the patients in the trial had high-volume disease and there were very few patients with low-volume metachronous disease in particular. This regimen adds considerable toxicity, and patients with low-volume disease should be informed that the benefit in this setting is not clear.

^{ccc} High-volume disease in this setting is defined based on CHARTED criteria (the presence of visceral metastasis or ≥4 bone lesions with ≥1 beyond the vertebral bodies and pelvis).


Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

WORKUP AND TREATMENT OF M0 CASTRATION-RESISTANT PROSTATE CANCER (CRPC)^{xx,ddd}



^c [Principles of Survivorship in Prostate Cancer \(PROS-B\)](#).

^g [Principles of Imaging \(PROS-E\)](#).

^{aa} For details on the use of ADT and ARPIs, see [Principles of Androgen Deprivation Therapy \(PROS-G\)](#) and [Discussion](#).

^{ff} Document castrate levels of testosterone if clinically indicated. Workup for progression should include bone and soft tissue evaluation. See [Principles of Imaging \(PROS-E\)](#).


^{kk} Monitoring should include physical exam, PSA every 3–6 mo, and imaging for symptoms or increasing PSA.

^{xx} ADT alone ([PROS-G](#)) or observation are recommended for asymptomatic patients with metastatic disease or M0 CRPC and life expectancy ≤5 years.

^{ddd} CRPC is prostate cancer that progresses clinically, radiographically, or biochemically despite castrate levels of serum testosterone (<50 ng/dL). Scher HI, et al. J Clin Oncol 2008;26:1148-1159.

Note: All recommendations are category 2A unless otherwise indicated.

WORKUP AND TREATMENT OF M1 CRPC^{xx,ddd}



^c [Principles of Survivorship in Prostate Cancer \(PROS-B\)](#).

^e [Principles of Genetics and Molecular/Biomarker Analysis \(PROS-C\)](#).

^t [Principles of Radiation Therapy \(PROS-J\)](#).

^{aa} For details on the use of ADT and ARPIs, see [Principles of Androgen Deprivation Therapy \(PROS-G\)](#) and [Discussion](#).

^{xx} ADT alone ([PROS-G](#)) or observation are recommended for asymptomatic patients with metastatic disease or M0 CRPC and life expectancy ≤5 years.

^{ddd} CRPC is prostate cancer that progresses clinically, radiographically, or biochemically despite castrate levels of serum testosterone (<50 ng/dL). Scher HI, et al. J Clin Oncol 2008;26:1148-1159.

^{eee} Histologic evidence of both adenocarcinoma and small cell carcinoma may be present, in which case treatment can follow either pathway. Treat as adenocarcinoma if biopsy is not feasible or not performed.

^{fff} Germline testing for HRR mutations is recommended if not performed previously. See [Principles of Genetics and Molecular/Biomarker Analysis \(PROS-C\)](#).

⁹⁹⁹ Document castrate levels of testosterone if progression occurs on ADT. See [Principles of Imaging \(PROS-E\)](#) and [Discussion](#).

^{hhh} For details on the efficacy and safety of these agents, see [Principles of Non-Hormonal Systemic Therapy \(PROS-N\)](#).

ⁱⁱⁱ Cabazitaxel 20 or 25 mg/m² plus carboplatin area under the curve [AUC] 4 mg/mL per min with growth factor support can be considered for fit patients with aggressive variant metastatic CRPC (mCRPC) (ie, visceral metastases, low PSA and bulky disease, high lactate dehydrogenase [LDH], high carcinoembryonic antigen [CEA], lytic bone metastases, NEPC histology) or unfavorable genomics (defects in at least 2 of PTEN, TP53, and RB1). Corn PG, et al. Lancet Oncol 2019;20:1432-1443.

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

SYSTEMIC THERAPY FOR M1 CRPC: ADENOCARCINOMA^{g,aa,jjj,kkk}

Pre-ARPI ^{aa,iii}	Post-ARPI ⁱⁱⁱ /Pre-Docetaxel ^{aa}	Post-ARPI ⁱⁱⁱ /Post-Docetaxel ^{aa}
Preferred: • Abiraterone (category 1) • Enzalutamide (category 1) Other Recommended: • Docetaxel^{hhh} (category 1) Useful in Certain Circumstances: • <u>Molecular Biomarker-Directed Therapy</u> ▶ <i>BRCA</i> mutation ◊ Niraparib/abiraterone^{mmm} (category 1) ◊ Olaparib/abiraterone^{mmm} (category 1) ◊ Talazoparib/enzalutamide^{mmm} (category 1) ▶ <i>HRRm</i> (other than <i>BRCA1/2</i>) ◊ Talazoparib/enzalutamide^{mmm} (category 1) • <u>Disease State-Specific Therapy</u> ▶ Bone metastases ◊ Radium-223^{ooo}/enzalutamide	Preferred: • Docetaxel^{hhh} (category 1) Useful in Certain Circumstances: • <u>Molecular Biomarker-Directed Therapy</u> ▶ <i>BRCA</i> mutation ◊ Olaparib^{mmm} (category 1, preferred) ◊ Rucaparib^{mmm} (category 1, preferred) ◊ Niraparib/abiraterone^{mmm} (category 2B) ◊ Talazoparib/enzalutamide^{mmm} (category 2B) ▶ <i>HRRm</i> (other than <i>BRCA1/2</i>) ◊ Olaparib^{mmm} ◊ Talazoparib/enzalutamide^{mmm} (category 2B) • <u>Disease State-Specific Therapy</u> ▶ PSMA-positive metastases ◊ Lutetium Lu 177 vipivotide tetraxetan (Lu-177-PSMA-617)^{qqq} ▶ Aggressive variantⁱⁱⁱ ◊ Cabazitaxel/Carboplatin^{hhh}	Preferred: • Cabazitaxel^{hhh} (category 1) • Docetaxel rechallenge^{hhh} Useful in Certain Circumstances: • <u>Molecular Biomarker-Directed Therapy</u> ▶ <i>BRCA</i> mutation ◊ Olaparib^{mmm} (category 1) ◊ Rucaparib^{mmm} ▶ <i>HRRm</i> (other than <i>BRCA1/2</i>) ◊ Olaparib^{mmm} ▶ Other FDA-approved agents for tissue agnostic indications^{hhh} • <u>Disease State-Specific Therapy</u> ▶ PSMA-positive metastases ◊ Lu-177-PSMA-617^{qqq} (category 1) ▶ Aggressive variantⁱⁱⁱ ◊ Cabazitaxel/carboplatin^{hhh} ▶ Palliation for symptomatic patients unable to tolerate other therapies ◊ Mitoxantrone^{hhh}

Additional Options Irrespective of Prior ARPI or Prior Docetaxel (Useful in Certain Circumstances)

- | | |
|--|---|
| <ul style="list-style-type: none">• <u>Disease State-Specific Therapy</u><ul style="list-style-type: none">▶ Asymptomatic without visceral metastases<ul style="list-style-type: none">◊ Sipuleucel-T^{hhh,ppp}▶ Oligometastatic^h/Oligoprogressive disease<ul style="list-style-type: none">◊ Metastasis-directed therapyⁿⁿⁿ with metastatic castration-resistant prostate cancer (mCRPC) systemic therapy▶ Symptomatic bone-predominant metastases<ul style="list-style-type: none">◊ Radium-223^{ooo} (category 1) | <ul style="list-style-type: none">• <u>Molecular Biomarker-Directed Therapy</u><ul style="list-style-type: none">▶ MSI-High (MSI-H)/dMMR<ul style="list-style-type: none">◊ Pembrolizumab^{hhh} (category 2B) |
|--|---|

Note: All recommendations are category 2A unless otherwise indicated.

[Footnotes on PROS-18A](#)



THERAPY FOR M1 CRPC: ADENOCARCINOMA

FOOTNOTES

^g [Principles of Imaging \(PROS-E\)](#).

^h See Number of Metastatic Sites in the [Principles of MDT \(PROS-M\)](#).

^{aa} For details on the use of ADT and ARPIs, see [Principles of Androgen Deprivation Therapy \(PROS-G\)](#) and [Discussion](#).

^{hhh} For details on the efficacy and safety of these agents, see [Principles of Non-Hormonal Systemic Therapy \(PROS-N\)](#).

ⁱⁱⁱ Cabazitaxel 20 or 25 mg/m² plus carboplatin AUC 4 mg/mL per min with growth factor support can be considered for fit patients with aggressive variant mCRPC (ie, visceral metastases, low PSA and bulky disease, high LDH, high CEA, lytic bone metastases, NEPC histology) or unfavorable genomics (defects in at least 2 of PTEN, TP53, and RB1). Corn PG, et al. Lancet Oncol 2019;20:1432-1443.

^{jjj} Document castrate levels of testosterone if progression occurs on ADT. Consider metastatic lesion biopsy. If small cell neuroendocrine is found, see [PROS-17](#).

^{kkk} Patients can continue through all treatment options listed. Best supportive care, which can include androgen-directed therapy or steroid, is always an appropriate option.

^{lll} ARPI therapies include abiraterone, enzalutamide, darolutamide, or apalutamide. Abiraterone given as part of neoadjuvant/concomitant/adjuvant ADT with EBRT is not considered post-ARPI.

^{mmm} PARP inhibitors with or without ARPI have different biomarker and previous treatment requirements. See [Principles of Non-Hormonal Systemic Therapy \(PROS-N\)](#).

ⁿⁿⁿ [Principles of MDT \(PROS-M\)](#).

^{ooo} Radium-223 should not be used in patients with visceral metastases. Concurrent use with systemic therapies other than enzalutamide should be pursued with caution. Concomitant use of denosumab or zoledronic acid is recommended. See [Principles of Radiation Therapy \(PROS-J\)](#).

^{ppp} Sipuleucel-T is recommended only for asymptomatic or minimally symptomatic, no liver metastases, life expectancy >6 months, and ECOG performance status 0–1. Benefit with sipuleucel-T has not been reported in patients with visceral metastases and is not recommended if visceral metastases are present. There are limited data to support the efficacy of Sipuleucel-T in the post-chemotherapy setting. Sipuleucel-T is not recommended for patients with small cell prostate cancer/NEPC.

^{qqq} Lu-177–PSMA-617 is a treatment option for patients with ≥1 PSMA-positive lesion and/or metastatic disease that is predominately PSMA-positive and with no dominant PSMA-negative metastatic lesions who have been treated previously with androgen receptor-directed therapy and a taxane-based chemotherapy or are considered appropriate to delay a taxane-based chemotherapy. Sartor O, et al. N Engl J Med 2021;385:1091-1103; Morris MJ, et al. Lancet 2024;404:1227-1239. See [Principles of Radiation Therapy \(PROS-J\)](#).

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF LIFE EXPECTANCY ESTIMATION

- Life expectancy estimation is critical to informed decision-making in prostate cancer early detection and treatment.
- Estimation of life expectancy is possible for groups of patients but challenging for individuals.
- Life expectancy can be estimated using:
 - ▶ The Social Security Administration tables (www.ssa.gov/OACT/STATS/table4c6.html)
 - ▶ The WHO's Life Tables by country (<http://apps.who.int/gho/data/view.main.60000?lang=en>)
 - ▶ The Memorial Sloan Kettering Male Life Expectancy tool (<https://www.mskcc.org/nomograms/prostate>)
 - ▶ University of California San Francisco (UCSF) Lee Schonberg Index (<https://eprognosis.ucsf.edu/leeschonberg.php>)
- If using a life expectancy table, life expectancy should be adjusted using the clinician's assessment of overall health as follows:
 - ▶ Best quartile of health - add 50%
 - ▶ Worst quartile of health - subtract 50%
 - ▶ Middle two quartiles of health - no adjustment
- Examples of upper, middle, and lower quartiles of life expectancy at selected ages are included in the [NCCN Guidelines for Older Adult Oncology](#) for life expectancy estimation.

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF SURVIVORSHIP BONE HEALTH IN PROSTATE CANCER

Treatment-Related Bone Loss

- ADT increases the risk of bone loss, and this risk is exacerbated with more potent androgen suppression, longer duration of therapy or delayed testosterone recovery, and concurrent prednisone use.
- The goal of osteoporosis screening is to identify patients at increased risk of sustaining a low-trauma fracture who would benefit from intervention to minimize the fracture risk. Risk assessment for treatment-related bone loss should take place for all patients initiating ADT of any duration. Fracture risk can be assessed using the Fracture Risk Assessment Tool (FRAX), the algorithm released by The University of Sheffield (<https://www.fraxplus.org/>). FRAX was developed to estimate the 10-year probability of hip fracture or major osteoporotic fractures combined (hip, spine, shoulder, or wrist) for an untreated individual using easily obtainable clinical risk factors for fracture with or without information on bone mineral density. When utilizing the FRAX algorithm select YES for secondary osteoporosis for individuals with hypogonadism. ADT should be considered “secondary osteoporosis” when using the FRAX algorithm. A previous major osteoporotic fracture (hip fracture or spine fracture) is considered clinical osteoporosis and warrants bone antiresorptive drug therapy independent of bone mineral density.
- A baseline dual-energy x-ray absorptiometry (DEXA) scan should be obtained before starting ADT in patients at increased risk for fracture based on FRAX screening and being considered for antiresorptive therapy (see Table 1). For patients at low risk of fracture based on the FRAX risk assessment, baseline DEXA scan can be omitted. The exact FRAX fracture risk threshold has not been defined in this population. One approach is to set the threshold at 10-year risk of major osteoporotic fracture (calculated without DEXA) greater than that of a 65-year old white woman with no additional risk factors (defined as 8.4% in the United States).
- Treatment for osteoporosis is advised according to guidelines for the general population from the Bone Health and Osteoporosis Foundation.¹ These guidelines (see Table 1) include recommendations for: 1) calcium (1000–1200 mg daily from food, with supplements if intake is insufficient); 2) vitamin D3 (serum levels of 30–50 ng/mL with supplements prescribed if needed); and 3) pharmacologic treatment for patients aged ≥50 years with low bone mass (T-score between -1.0 and -2.5, osteopenia) at the femoral neck or total hip by DEXA scan with a 10-year probability of hip fracture ≥3% or a 10-year probability of a major osteoporosis-related fracture ≥20% based on FRAX screening (see Table 2).
- Antiresorptive medications that increase bone mineral density and reduce disease-related skeletal complications during ADT for prostate cancer include denosumab (60 mg subcutaneously [SQ] every 6 months), zoledronic acid (5 mg IV annually), and alendronate (70 mg PO weekly) (see Table 2). Treatment with either denosumab, zoledronic acid, or alendronate sodium is recommended when the absolute fracture risk warrants drug therapy.
 - ▶ Choice of agent may depend on underlying comorbidities, whether the patient has been treated with zoledronic acid previously, logistics, and/or cost considerations.
 - ▶ Bisphosphonates (zoledronic acid or alendronate) can cause side effects of acute phase reaction, joint pain, hypocalcemia, osteonecrosis of the jaw, nephrotoxicity with need for dose modification for renal insufficiency, ocular toxicities, and atypical femoral fractures with prolonged use (>3 to 5 years).
 - ▶ Denosumab can cause side effects of hypocalcemia, osteonecrosis of the jaw, and atypical femoral fractures with prolonged use. The risk factors for denosumab-associated hypocalcemia include blastic bone metastases, renal impairment, vitamin D deficiency, the lack of prophylactic supplementation of calcium and/or vitamin D, preexisting hypoparathyroidism, hypomagnesemia, and gastric bypass. Although renal monitoring is not required, denosumab is not recommended in patients with a creatinine clearance <30 mL/min given the risk of severe hypocalcemia. Calcium, creatinine, and vitamin D levels should be checked prior to initiating therapy. Periodic monitoring of serum calcium levels is recommended with denosumab use. Stopping denosumab therapy can result in rebound bone loss and fractures; therefore it is recommended to administer at least one dose of a potent bisphosphonate (zoledronic acid 4 or 5 mg) to prevent rebound bone loss and presumably rebound fracture.²

Note: All recommendations are category 2A unless otherwise indicated.

[References on PROS-B 4 of 6](#)

PROS-B
1 OF 6



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF SURVIVORSHIP BONE HEALTH IN PROSTATE CANCER

Treatment-Related Bone Loss Continued

- ▶ The risk of osteonecrosis of the jaw is increased in patients who have tooth extractions, poor dental hygiene, or a dental appliance. To prevent osteonecrosis of the jaw, it is recommended that all patients have a comprehensive dental evaluation prior to initiating an osteoclast inhibitor.³ If invasive dental procedures are required, bone-targeted therapy should be withheld until the dentist indicates that the patient has healed completely from all dental procedure(s). Stopping denosumab represents a dilemma in this context, and the clinician must carefully weigh the risk of rebound spine fractures versus the risk of osteonecrosis of the jaw.
- Annual assessment of fracture risk using the FRAX risk assessment tool is recommended for all patients on ADT or those who remain hypogonadal after completion of ADT (see Table 1). Depending on the fracture risk and prior DEXA scan results, repeat DEXA scan in 1 to 2 years is recommended for those patients on ADT. For individuals initiated on antiresorptive therapy, a follow-up DEXA scan after 1 year of treatment is recommended by the International Society for Clinical Densitometry, although there is no consensus on the optimal approach to monitoring the effectiveness of bone treatment. Use of biochemical markers of bone turnover to monitor response to therapy is not recommended. There are currently no guidelines on how often to monitor vitamin D levels.
- For patients receiving antiresorptive therapy, there are currently no consensus guidelines on duration of treatment. Due to concerns of long-term risks of antiresorptive therapy, a “drug holiday” at 3 to 5 years can be considered based on agent utilized, stability of bone mineral density, prior fracture history, and future fracture risk. Bone mineral density should be monitored approximately every 1 to 2 years after suspending therapy, and therapy should generally be resumed if bone mineral density declines significantly or if the patient develops a new fragility fracture.

Table 1: Risk Assessment and Monitoring	
Clinical Scenario	Recommendation
Baseline at ADT initiation	DEXA recommended for most patients. In select individuals at low probability of fracture based on FRAX risk assessment tool, DEXA can be omitted
On ADT	DEXA every 1–2 years, dependent on FRAX risk assessment tool
On antiresorptive therapy	DEXA at 1 year

Note: All recommendations are category 2A unless otherwise indicated.

[References on PROS-B 4 of 6](#)



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF SURVIVORSHIP BONE HEALTH IN PROSTATE CANCER

Prevention of Symptomatic Skeletal-Related Events (SREs) in Patients with Bone-Metastatic CRPC

- In patients with CRPC who have bone metastases, denosumab and zoledronic acid have been shown to prevent disease-related skeletal complications, which include fracture, spinal cord compression, or the need for surgery or RT to bone.
- When compared to zoledronic acid, denosumab was shown to be superior in prevention of SREs in patients with metastatic CRPC (mCRPC), albeit with numerically higher hypocalcemia and osteonecrosis of the jaw risks. Initial studies investigated zoledronic acid and denosumab administered every 4 weeks. Subsequent studies demonstrated that every-12-week dosing of zoledronic acid compared to every-4-week dosing did not increase the risk of skeletal events.^{4,5} Every-12-week dosing of zoledronic acid is recommended for symptomatic SRE reduction when indicated. Every-12-week dosing of denosumab is under investigation and current data suggest noninferior symptomatic skeletal events compared to every-4-week dosing.⁵ Utilization of zoledronic acid and denosumab for symptomatic SRE reduction requires consideration of degree of benefit and risk associated with therapy to optimize use, dose, and schedule. It is important to recognize that testing of zoledronic acid and denosumab in bone-metastatic CRPC was conducted during an era when treatment options for mCRPC were largely limited to docetaxel chemotherapy. Subsequent studies investigating abiraterone, enzalutamide, cabazitaxel, radium-223, and Lu-177-PSMA-617 have demonstrated improvement of SREs with treatment. While radium-223 did improve symptomatic SREs in patients with bone mCRPC, the combination of radium-223 with abiraterone was associated with increased frequency of bone fractures, particularly in individuals not receiving an antiresorptive agent.⁶
- A phase 3 clinical trial that assessed the role for zoledronic acid in patients with castration-sensitive disease beginning ADT for bone metastases was negative.⁷ Therefore, use of osteoclast inhibitors for reduction of symptomatic SREs in metastatic castration-sensitive disease with bone metastases is not recommended. However, usage of these agents to prevent bone loss and fragility fractures at appropriate doses and dosing intervals should be utilized when clinically appropriate in this context (see Treatment-Related Bone Loss, [PROS-B 1 of 4](#)).

Table 2: Optimization of Bone Health in Patients with Prostate Cancer

Patient Population	Category	Intervention
All patients receiving ADT	Lifestyle modifications	• Weight-bearing exercises (30 minutes per day), balance training, safe movement strategies • Limit alcohol consumption • Smoking cessation
	Calcium and vitamin D supplementation	• Calcium 1000–1200 mg daily from food with supplements if needed • Maintain serum vitamin D3 levels of 30–50 ng/mL with supplements if needed
For treatment-related bone loss in patients receiving ADT	Antiresorptive agents	• Alendronate 70 mg PO weekly • Denosumab 60 mg SQ every 6 months • Zoledronic acid 5 mg IV annually
For prevention of symptomatic SREs in patients with bone-metastatic CRPC	Antiresorptive agents	• Denosumab 120 mg SQ every 4 weeks • Zoledronic acid 4 mg IV every 12 weeks

Note: All recommendations are category 2A unless otherwise indicated.

[References on PROS-B 4 of 6](#)



PRINCIPLES OF SURVIVORSHIP

BONE HEALTH IN PROSTATE CANCER: REFERENCES

- ¹ LeBoff MS, Greenspan SL, Insogna KL, et al. The clinician's guide to prevention and treatment of osteoporosis. *Osteoporos Int* 2022;33:2049-2102.
- ² Cummings SR, Ferrari S, Eastell R, et al. Vertebral fractures after discontinuation of denosumab: A post hoc analysis of the randomized placebo-controlled freedom trial and its extension. *J Bone Miner Res* 2018;33:190-198.
- ³ Yarom N, Shapiro CL, Peterson DE, et al. Medication-related osteonecrosis of the jaw: MASCC/ISOO/ASCO Clinical practice guideline. *J Clin Oncol* 2019;37:2270-2290.
- ⁴ Himelstein AL, Foster JC, Khatcheressian JL, et al. Effect of longer-interval vs standard dosing of zoledronic acid on skeletal events in patients with bone metastases: a randomized clinical trial. *JAMA* 2017;317:48-58.
- ⁵ Clemons M, Ong M, Stober C, et al. A randomised trial of 4- versus 12-weekly administration of bone-targeted agents in patients with bone metastases from breast or castration-resistant prostate cancer. *Eur J Cancer* 2021;142:132-140.
- ⁶ Smith M, Parker C, Saad F, et al. Addition of radium-223 to abiraterone acetate and prednisone or prednisolone in patients with castration-resistant prostate cancer and bone metastases (ERA 223): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet Oncol* 2019;20:408-419.
- ⁷ Smith MR, Halabi S, Ryan CJ, et al. Randomized controlled trial of early zoledronic acid in men with castration-sensitive prostate cancer and bone metastases: results of CALGB 90202 (alliance). *J Clin Oncol* 2014;32:1143-1150.

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF SURVIVORSHIP CARDIOVASCULAR DISEASE IN PROSTATE CANCER

- Individuals with prostate cancer are at increased risk for cardiovascular disease (CVD) and cardiovascular events, and CVD is a leading cause of death in this population.¹
 - ▶ The main cause for this increased risk is the cardiovascular effects of ADT and ARPIs.^{2,3}
 - ▶ Shared risk factors for prostate cancer and CVD (ie, smoking, poor health behaviors) also likely contribute to the increased risk of CVD in this population.⁴
- CVD develops over time because of well-defined risk factors such as hypertension, hyperlipidemia, use of tobacco products, obesity, diabetes, and the cardiotoxic effects of cancer treatment.
- CVD-related morbidity and mortality can occur in both the short- and long-term in prostate cancer survivors.
- Control of modifiable risk factors can decrease the risk of subsequent cardiovascular events.
- Therefore, prostate cancer survivors should be assessed for and counseled on:
 - ▶ Pre-existing and emerging CVD (eg, coronary artery disease, congestive heart failure, peripheral vascular disease, arrhythmias including atrial fibrillation)
 - ▶ Any increased risk of CVD based on prior treatment, comorbidity, or CVD risk factors (eg, hypertension, dyslipidemia, obesity, cigarette/tobacco use, diabetes mellitus)
 - ▶ Lifestyle modification and interventions for modifiable risk factors (see the ABCDEs of CVD Prevention, [Table 1](#))
- Aggressive risk factor management is recommended for those deemed to be at higher risk, and referral to a cardiologist or cardio-oncologist in patients at elevated CVD risk should be considered at any stage of the prostate cancer journey.
- Cooperation and shared care with primary care providers, and cardiovascular specialists as needed, is key to optimizing cardiac and vascular outcomes in prostate cancer survivors.
- For further information on cardiovascular risk assessment in cancer survivors, see the [NCCN Guidelines for Survivorship](#).

¹ Sturgeon KM, Deng L, Bluethmann SM, et al. A population-based study of cardiovascular disease mortality risk in US cancer patients. *Eur Heart J* 2019;40:3889-3897.
² Okwuosa TM, Morgans A, Rhee JW, et al. Impact of hormonal therapies for treatment of hormone-dependent cancers (breast and prostate) on the cardiovascular system: effects and modifications: a scientific statement from the American Heart Association. *Circ Genom Precis Med* 2021;14:e000082.
³ El-Taji O, Taktak S, Jones C, et al. Cardiovascular events and androgen receptor signaling inhibitors in advanced prostate cancer: A systematic review and meta-analysis. *JAMA Oncol* 2024;20:874-884.
⁴ Bergengren O, Pekala KR, Matsoukas K, et al. 2022 update on prostate cancer epidemiology and risk factors - a systematic review. *Eur Urol* 2023;84:191-206.

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF SURVIVORSHIP CARDIOVASCULAR DISEASE IN PROSTATE CANCER: RISK ASSESSMENT^a

Table 1: ABCDEs to Promote Cardiovascular Wellness in Cancer Survivors ^b	
A	• Awareness of risks and presentation of heart disease • Assessment of CVD and cardiovascular risk (consider use of a CVD risk assessment tool^c) • Aspirin use as appropriate (indicated for secondary prevention; clinician-survivor discussion required for primary prevention with careful weighing of benefits and risks)^d
B	• Blood pressure monitoring/management (with clinician-survivor discussion regarding the use of hypertension treatment and blood pressure goals)
C	• Cholesterol assessment/management (with clinician-survivor discussion regarding the use of statin therapy for primary prevention and lipid profile goals) • Cigarette/tobacco cessation (NCCN Guidelines for Smoking Cessation)
D	• Diet and weight management (see SNWM-1 in NCCN Guidelines for Survivorship) • Diabetes mellitus prevention/treatment
E	• Exercise (see SPA-1 in NCCN Guidelines for Survivorship)^e • Echocardiogram (ECHO) and/or electrocardiogram (ECG) based on individual risk (consider use of a CVD risk assessment tool^c)

^a Armenian SH, Lacchetti C, Barac A, et al. Prevention and monitoring of cardiac dysfunction in survivors of adult cancers: American Society of Clinical Oncology Clinical Practice Guideline. J Clin Oncol 2017;35:893-911.

^b Adapted with permission from Montazeri K, Unitt C, Foody JM, et al. ABCDE steps to prevent heart disease in breast cancer survivors. Circulation 2014;130:e157-e159.

^c The ASCVD Risk Estimator Plus from the American College of Cardiology is available at <https://tools.acc.org/ascvd-risk-estimator-plus/#!/calculate/estimate> and The American Heart Association's PREVENT Risk Calculator is available at <https://professional.heart.org/en/guidelinesand-statements/prevent-calculator>.

^d U.S. Preventive Services Task Force; Davidson KW, Barry MJ, Mangione CM, et al. Aspirin use to prevent cardiovascular disease: U.S. Preventive Services Task Force Recommendation Statement. JAMA 2022;327:1577-1584.

^e Gilchrist SC, et al. Circulation 2019;139:e997-e1012.

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF GENETICS AND MOLECULAR/BIOMARKER ANALYSIS

GERMLINE TESTING

For details regarding the nuances of genetic counseling and testing, see Principles of Cancer Risk Assessment and Counseling (EVAL-A) in the [NCCN Guidelines for Genetic/Familial High-Risk Assessment: Breast, Ovarian, Pancreatic and Prostate](#)

- **Pre-test Considerations**

- ▶ The Panel recommends inquiring about family and personal history of cancer, and known germline variants at time of initial diagnosis. Criteria for germline testing (see CRIT-6 in the [NCCN Guidelines for Genetic/Familial High-Risk Assessment: Breast, Ovarian, Pancreatic and Prostate](#) and HRS-3 in the [NCCN Guidelines for Genetic/Familial High-Risk Assessment: Colorectal, Endometrial, and Gastric](#)) should be reviewed at time of initial diagnosis and, if relevant, at recurrence.
- ▶ Germline testing is also recommended for patients with metastatic, regional (node positive), very-high-risk localized, or high-risk localized prostate cancer.
- ▶ Germline testing should be considered in appropriate individuals where it is likely to impact the prostate cancer treatment and clinical trial options, management of risk of other cancers, and/or potential risk of cancer in family members.

- **Testing**

- ▶ If criteria are met, multigene testing is recommended (see GENE-1 in the [NCCN Guidelines for Genetic/Familial High-Risk Assessment: Breast, Ovarian, Pancreatic, and Prostate](#)).

- **Post-test Considerations**

- ▶ Post-test genetic counseling is strongly recommended if a germline mutation (pathogenic/likely pathogenic variant) is identified. Cascade testing for relatives is critical to inform the risk for familial cancers in all relatives.
- ▶ Post-test genetic counseling is recommended if positive family history but no pathogenic variant OR if only germline variants of uncertain significance (VUS) are identified. This is to ensure accurate understanding of family implications and review indications for additional testing and/or follow-up (including clinical trials of reclassification).
- ▶ Resources are available to review the available data supporting pathogenic consequences of specific variants (eg, <https://www.ncbi.nlm.nih.gov/clinvar/>; <https://brcaexchange.org/about/app>).
- ▶ Individuals should be counseled to inform providers of any updates to family cancer history.

Note: All recommendations are category 2A unless otherwise indicated.



PRINCIPLES OF GENETICS AND MOLECULAR/BIOMARKER ANALYSIS

SOMATIC TUMOR TESTING

• Pre-test Considerations

- ▶ At present, tumor molecular and biomarker analysis is recommended for patients with metastatic disease for treatment decision-making, including understanding eligibility for biomarker-directed treatments, genetic counseling, and eligibility for clinical trials. Clinical trials may include established and/or candidate molecular biomarkers for eligibility.
- ▶ Tumor molecular profiles may change with subsequent treatments and re-evaluation may be considered at time of cancer progression for treatment decision-making.
- ▶ Patients should be informed that tumor molecular analysis by DNA sequencing has the potential to uncover germline findings. Confirmatory germline testing may be recommended [see Post-test Considerations (below) and Tumor Testing:s Potential Implications for Germline Testing in the Principles of Cancer Risk Assessment and Counseling (EVAL-A) in the [NCCN Guidelines for Genetic/Familial High-Risk Assessment: Breast, Ovarian, Pancreatic, and Prostate](#)].

• Testing

- ▶ Somatic testing for alterations in DNA damage response:
 - ◊ Multigene tumor testing for alterations in HRR genes, including but not limited to *BRCA1*, *BRCA2*, *ATM*, *PALB2*, *FANCA*, *RAD51D*, *CHEK2*, and *CDK12*, is recommended in patients with metastatic prostate cancer. This testing can be considered in patients with regional prostate cancer.
 - Loss of *BRCA1* and *BRCA2* may be especially associated with response to PARP inhibitor therapy compared to other HRR gene alterations.
 - ◊ Tumor testing for MSI-H or dMMR is recommended in patients with mCRPC and may be considered in patients with regional or castration-sensitive metastatic prostate cancer.
 - ◊ TMB testing is recommended in patients with mCRPC.

• Tumor Specimen and Assay Considerations

- ▶ The Panel strongly recommends a metastatic biopsy for histologic and molecular evaluation. This could include lymph node biopsy for patients with N1 disease.
 - ◊ When metastatic biopsy is unsafe or unfeasible, plasma circulating tumor DNA (ctDNA) assay is an option, preferably collected during biochemical (PSA) and/or radiographic progression in order to maximize diagnostic yield. When diagnostic yield is low, the risk of false negatives is higher, so ctDNA collection is not recommended when PSA is undetectable.
- ▶ Caution is needed when interpreting ctDNA-only evaluation due to potential interference from clonal hematopoiesis of indeterminate potential (CHIP), which can result in a false-positive biomarker signal.
- ▶ DNA analysis for MSI and immunohistochemistry for mismatch repair (MMR) are different assays measuring different biological effects caused by dMMR function. If MSI is used, testing using a next-generation sequencing assay validated for prostate cancer is preferred.

• Post-test Considerations

- ▶ Post-test genetic counseling is recommended if pathogenic/likely pathogenic variant (mutation) identified in any gene that has clinical implications if also identified in germline (eg, *BRCA1*, *BRCA2*, *ATM*, *PALB2*, *CHEK2*, *HOXB13*, *MLH1*, *MSH2*, *MSH6*, *PMS2*).
- ▶ Post-test genetic counseling to assess for the possibility of Lynch syndrome is recommended if MSI-H or dMMR is found.

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF QUALITY OF LIFE AND SHARED DECISION-MAKING

- Treatments for patients with localized prostate cancer have risks and side effects that must be considered in the context of the risk posed by the disease.¹⁻⁴
- Baseline urinary, sexual, and bowel function are strongly associated with functional outcomes among patients undergoing treatment.¹⁻⁴
- Thus, it is important to measure baseline disease-specific function (urinary, sexual, and bowel function), preferably using a standardized patient-reported outcomes instrument (eg, EPIC-26⁵).
- Shared decision-making in a multidisciplinary manner regarding initial management of localized prostate cancer should include an explanation of the potential benefits and potential harms of each option. The provider should explain the likelihood of cure, recurrence, disease progression, and disease-specific mortality with each management option, taking into account disease severity and competing risks. In addition to the primary intended effects of treatment, the clinician should discuss the side effects of each treatment and predicted impact on quality of life, including urinary, sexual, and bowel function. Patient preferences should be elicited and should be incorporated into the disease management decision.⁶
- Consider geriatric assessment. See [NCCN Guidelines for Older Adult Oncology](#) for tools to aid optimal assessment and management of disease in older adults.
- Refer to the NCCN Distress Thermometer and Problem List, which includes social determinants of health. See [NCCN Guidelines for Distress Management](#) (DIS-A).

¹ Sanda MG, Dunn RL, Michalski J, et al. Quality of life and satisfaction with outcome among prostate-cancer survivors. *N Engl J Med* 2008;358:1250-1261.

² Chen RC, Basak R, Meyer AM, et al. Association between choice of radical prostatectomy, external beam radiotherapy, brachytherapy, or active surveillance and patient-reported quality of life among men with localized prostate cancer. *JAMA* 2017;317:1141-1150.

³ Hoffman KE, Penson DF, Zhao Z, et al. Patient-reported outcomes through 5 years for active surveillance, surgery, brachytherapy, or external beam radiation with or without androgen deprivation therapy for localized prostate cancer. *JAMA* 2020;323:149-163.

⁴ Donovan JL, Hamdy FC, Lane JA, et al; ProtecT Study Group. Patient-reported outcomes after monitoring, surgery, or radiotherapy for prostate cancer. *N Engl J Med* 2016;375:1425-1437.

⁵ Szymanski KM, Wei JT, Dunn RL, Sanda MG. Development and validation of an abbreviated version of the expanded prostate cancer index composite instrument for measuring health-related quality of life among prostate cancer survivors. *Urology* 2010;76:1245-1250.

⁶ Makarov D, Fagerlin A, Finkelstein J et al. AUA White Paper on Implementation of Shared Decision Making into Urological Practice. American Urological Association 2022. Available at: <https://www.auanet.org/guidelines-and-quality/guidelines/best-practice-statements-and-whitepapers/shared-decision-making>.

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF IMAGING

Goals of Imaging

- Imaging is performed for the detection and characterization of disease to select treatment or guide change in disease management.
- Imaging techniques can evaluate anatomic or functional parameters.
 - ▶ Anatomic imaging techniques include ultrasound, CT, and MRI.
 - ▶ Functional imaging techniques include radionuclide bone scan, PET/CT, and advanced MRI techniques, such as spectroscopy and diffusion-weighted imaging (DWI).

Efficacy of Imaging

- The utility of imaging for patients with early PSA persistence/recurrence after RP depends on risk group prior to operation, pathologic Gleason grade and stage, PSA, and PSADT after recurrence. Low- and intermediate-risk groups with low serum PSAs postoperatively have a very low risk of positive bone scans or CT scans.
- Frequency of imaging should be based on individual risk, age, PSADT, Gleason score, and overall health.
- Bone scans are rarely positive in asymptomatic patients with PSA <10 ng/mL. The relative risk for bone metastasis or death increases as PSADT shortens. Bone imaging should be performed more frequently when PSADT is ≤8 months, where there appears to be an inflection point.

Ultrasound

- Ultrasound uses high-frequency sound waves to image small regions of the body.
 - ▶ Standard ultrasound imaging provides anatomic information.
 - ▶ Vascular flow can be assessed using Doppler ultrasound techniques.
- Endorectal ultrasound is used to guide transrectal biopsies of the prostate. Endorectal ultrasound can be considered for patients with suspected recurrence after RP to guide prostate bed biopsy.
- Advanced ultrasound techniques for imaging of the prostate and for differentiation between prostate cancer and prostatitis are under evaluation.

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF IMAGING

Bone Imaging

- The use of the term “bone scan” refers to the technetium-99m-MDP bone scan in which technetium is taken up by bone that is turning over and imaged with a gamma camera using planar imaging or 3D imaging with single-photon emission CT (SPECT).
 - ▶ Sites of increased uptake imply accelerated bone turnover and may indicate metastatic disease.
 - ▶ Osseous metastatic disease may be diagnosed based on the overall pattern of activity, or in conjunction with anatomic imaging.
- Plain films, CT, MRI, or PET/CT or PET/MRI with F-18 piflufolastat prostate-specific membrane antigen (PSMA), Ga-68 PSMA-11, F-18 flutemetamol PSMA, F-18 fluciclovine, F-18 sodium fluoride, or C-11 choline can be considered for equivocal results on initial bone scan.
- Ga-68 PSMA-11, F-18 piflufolastat PSMA, or F-18 flutemetamol PSMA-PET/CT or PET/MRI (full body imaging) can be considered as an alternative to bone scan.
- Bone imaging is indicated in the initial evaluation of patients at high risk for skeletal metastases.
- Bone imaging can be considered for the evaluation of the patient post-prostatectomy when PSA does not fall to undetectable levels, or when there is undetectable PSA after RP with a subsequent detectable PSA that increases on ≥2 subsequent determinations.
- Bone imaging can be considered for the evaluation of patients with an increasing PSA or positive DRE after RT if the patient is a candidate for additional local therapy or systemic therapy.
- Bone scans are helpful to monitor metastatic prostate cancer to determine the clinical benefit of systemic therapy. However, new lesions seen on an initial post-treatment bone scan, compared to the pretreatment baseline scan, may not indicate disease progression.
- New lesions in the setting of a falling PSA or soft tissue response and in the absence of pain progression at that site may indicate bone scan flare or an osteoblastic healing reaction. For this reason, a confirmatory bone scan 8 to 12 weeks later is warranted to determine true progression from flare reaction. Additional new lesions favor progression. Stable scans make continuation of treatment reasonable. Bone scan flare is common, particularly on initiation of new hormonal therapy, and may be observed in nearly half of patients treated with the newer agents, enzalutamide and abiraterone. Similar flare phenomena may exist with other imaging modalities, such as CT or PET/CT imaging.

- Bone scans and soft tissue imaging (CT or MRI) in patients with metastatic or non-metastatic prostate cancer may be obtained regularly during systemic therapy to assess clinical benefit.
- Bone scans should be performed for symptoms and as often as every 6 to 12 months to monitor ADT. The need for soft tissue images remains unclear. In CRPC, 8- to 12-week imaging intervals appear reasonable.
- **Plain Radiography**
 - ▶ Plain radiography can be used to evaluate symptomatic regions in the skeleton. However, plain films will not detect a bone lesion until nearly 50% of the mineral content of the bone is lost or gained.
 - ▶ CT or MRI may be more useful to assess fracture risk as these modalities permit more accurate assessment of cortical involvement than plain films where osteoblastic lesions may obscure cortical involvement.

Soft Tissue Imaging

- Soft tissue imaging of the pelvis, abdomen, and chest can include:
 - ▶ Chest CT and abdomen/pelvis CT or abdomen/pelvis MRI or
 - ▶ PSMA-PET/CT or PSMA-PET/MRI for bone and soft tissue (full body) imaging.
- **Computed Tomography**
 - ▶ CT provides a high level of anatomic detail, and may detect gross extracapsular disease, nodal metastatic disease, and/or visceral metastatic disease.
 - ▶ CT is generally not sufficient to evaluate the prostate gland.
 - ▶ CT may be performed with IV contrast, and CT technique should be optimized to maximize diagnostic utility while minimizing radiation dose.
 - ▶ CT can be used for examination of the pelvis and/or abdomen for initial evaluation ([PROS-2](#)) and as part of workup for recurrence or progression (see [PROS-9](#) through [PROS-18](#)).

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF IMAGING

• *Magnetic Resonance Imaging*

- ▶ The strengths of MRI include high soft tissue contrast and characterization, multiparametric image acquisition, multiplanar imaging capability, and the use of specific MRI sequences to assess function.
 - ◊ MRI can be performed with and without the administration of IV contrast material.
 - ◊ Resolution of MRI images in the pelvis can be augmented using a phased array/endorectal coil.
- ▶ Standard MRI techniques can be used for examination of the pelvis and/or abdomen for initial evaluation ([PROS-1](#)) and as part of workup for recurrence or progression (see [PROS-9](#) through [PROS-18](#)).
- ▶ MRI may be considered in patients after RP when PSA does not fall to undetectable levels or when an undetectable PSA becomes detectable and increases on ≥ 2 subsequent determinations, or after RT for increasing PSA or positive DRE if the patient is a candidate for additional local therapy. MRI-ultrasound fusion biopsy may improve the detection of higher grade (Grade Group ≥ 2) cancers.
- ▶ Multiparametric MRI (mpMRI) can be used in the staging and characterization of prostate cancer. mpMRI images are defined as images acquired with at least one more sequence in addition to the anatomical T2-weighted images, such as DWI or dynamic contrast-enhanced (DCE) images. mpMRI may be used to better risk stratify patients who are considering active surveillance. Additionally, mpMRI may detect large and poorly differentiated prostate cancer (Grade Group ≥ 2) and detect extracapsular extension (T staging) and is preferred over CT for abdomen/pelvis staging. mpMRI has been shown to be equivalent to CT scan for pelvic lymph node evaluation.

Full Body Imaging

• *Positron Emission Tomography*

- ▶ PSMA-PET refers to a growing body of radiopharmaceuticals that target PSMA on the surface of prostate cells. There are multiple PSMA radiopharmaceuticals at various stages of investigation. At this time, the NCCN Guidelines only recommend the currently FDA-approved PSMA agents: F-18 piflutolastat PSMA (also known as F-18 DCFPyL), F-18 flutolastat PSMA (also known as rh-

PSMA-7.3), and Ga-68 PSMA-11 (also known as PSMA HBED-CC). Throughout these Guidelines, “PSMA-PET” refers to any of these FDA-approved PSMA ligands. See Table 1 on [PROS-E \(5 of 5\)](#) for more details.

- ▶ F-18 flutolastat PSMA is a PET imaging agent that is part of a class of tracers referred to as radiohybrid (rh) ligands. These tracers have two binding sites for radionuclides for both imaging and treatment, but the significance of this remains to be determined.
- ▶ PSMA-PET/CT or PET/MRI can be considered as an alternative to CT, MRI, and bone scans for initial staging of unfavorable intermediate, high, and very-high-risk disease, the detection of biochemically recurrent disease, and as workup for progression.
- ▶ In the setting of M1 CRPC, PSMA-PET imaging should only be used to determine if a patient is a candidate for Lu-177-PSMA-617.
- ▶ Changes in systemic therapy should not be made solely based on a positive PSMA-PET in patients with M0 CRPC.
- ▶ Bone scans can be considered to confirm osseous uptake on PET scans.
- ▶ Synthesis of Ga-68 PSMA-11 requires that the PSMA-11 ligand is labeled with Ga-68 from a generator or cyclotron. Two commercial kits to perform this in nuclear pharmacies have been approved by the FDA.
- ▶ C-11 choline or F-18 flutolastat PET/CT or PET/MRI may be used to detect small-volume recurrent disease in soft tissues and in bone.
- ▶ Studies suggest that PSMA-PET imaging has a higher sensitivity than C-11 choline or F-18 flutolastat PET imaging, especially at very low PSA levels (eg, <0.5 ng/mL).
- ▶ Because of the increased sensitivity and specificity of PSMA-PET tracers for detecting micrometastatic disease compared to CT, MRI, and bone scan at both initial staging and BCR, PSMA-PET/CT or PSMA-PET/MRI can serve as a more effective frontline imaging tool for these patients.
- ▶ Histologic or radiographic confirmation of involvement detected by PET imaging is recommended whenever feasible due to the presence of false positives. Although false positives exist, literature suggests that these are outweighed by the increase in true positives detected by PET relative to CT, MRI, and bone scans. To reduce the false-positive rate, physicians should consider the

Note: All recommendations are category 2A unless otherwise indicated.

PRINCIPLES OF IMAGING

intensity of PSMA-PET uptake and correlative CT findings in the interpretation of scans. The use of a standardized reporting system is encouraged.

- ▶ PSMA imaging should be done before initiation of ADT because ADT may affect detection sensitivity.
- ▶ High variability among PET/CT or PET/MRI equipment, protocols, interpretation, and institutions provides challenges for application and interpretation of the utility of PET/CT or PET/MRI.
- ▶ Table 1 on [PROS-E \(5 of 5\)](#) provides a summary of FDA-approved PET imaging tracers utilized for study in prostate cancer.
- ▶ PET/CT or PET/MRI results may change treatment but may not change oncologic outcome.
- ▶ When patients with the worst prognosis move from one risk group to the higher risk group, the average outcome of both risk groups will improve even if treatment has no impact on disease. This phenomenon is known as the Will Rogers effect, in which the improved outcomes of both groups could be falsely attributed to improvement in treatment, but would be due only to improved risk group assignment. As an example, F-18 sodium fluoride PET/CT may categorize some patients as M1b who would have been categorized previously as M0 using a bone scan (stage migration). Absent any change in the effectiveness of therapy, the overall survival (OS) of both M1b and M0 groups would improve. The definition of M0 and M1 disease for randomized clinical trials that added docetaxel or abiraterone to ADT was based on CT and radionuclide bone scans. Results suggest that OS of M1 disease is improved, whereas progression-free but not OS of M0 disease is improved. Therefore, a subset of patients now diagnosed with M1 disease using F-18 sodium fluoride PET/CT might not benefit from the more intensive therapy used in these trials and could achieve equivalent OS from less intensive therapy aimed at M0 disease. Carefully designed clinical trials using proper staging will be necessary to prove therapeutic benefit, rather than making assumptions compromised by stage migration.

- ▶ Fluorodeoxyglucose (FDG)-PET/CT should not be used routinely for staging prostate cancer since data are limited in this setting.
- ▶ F-18 FDG-PET has been shown to be prognostic in patients with progressive CRPC.^{1,2}
- ▶ The increasing use of PSMA-PET has identified the potential for considerable biological diversity among disease foci within a given individual with prostate cancer, especially mCRPC, and that this heterogeneity can be detected with a combination of PSMA-PET and FDG-PET. Initial data suggest that metastases with PSMA-negative/FDG-positive mismatches may exist in patients with mCRPC undergoing Lu-PSMA radioligand therapy and that patients with these mismatches may have worse outcomes. Currently, no robust clinical trial data exist to support the incorporation of FDG-PET into routine clinical use alongside PSMA-PET. To overcome the limitations of PSMA-PET in PSMA-negative metastatic disease, the Panel currently recommends the use of contrast-enhanced CT or MRI in these patients, as the non-contrast CT component of PSMA-PET/CT is insufficient to detect visceral metastatic disease.

Imaging as Workup for Progression

- Workup for progression should include bone and soft tissue evaluation.
 - ▶ See Bone Imaging ([PROS-E \[2 of 5\]](#)).
 - ▶ See Soft Tissue Imaging ([PROS-E \[2 of 5\]](#)).
- Imaging for patients with progressive CRPC should include chest CT, bone imaging, and abdomen/pelvis CT with contrast or abdomen/pelvis MRI with and without contrast.
- There is a lack of evidence to support the use of PET imaging in the CRPC setting and it should only be used in the setting of M1 CRPC to determine if a patient is a candidate for Lu-177-PSMA-617. Changes in treatment should not be made solely based on a positive PSMA-PET in patients with M0 CRPC.

¹ Buteau JP, Martin AJ, Emmett L, et al. PSMA and FDG-PET as predictive and prognostic biomarkers in patients given [177Lu]Lu-PSMA-617 versus cabazitaxel for metastatic castration-resistant prostate cancer (TheraP): A biomarker analysis from a randomised, open-label, phase 2 trial. *Lancet Oncol* 2022;23:1389-1397.

² Pathmanandavel S, Crumbaker M, Nguyen A, et al. The prognostic value of posttreatment 68Ga-PSMA-11 PET/CT and 18F-FDG PET/CT in metastatic castration-resistant prostate cancer treated with 177Lu-PSMA-617 and NOX66 in a phase I/II trial (LuPIN). *J Nucl Med* 2023;64:69-74.



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF IMAGING

Table 1. FDA-Cleared PET Imaging Tracers Studied in Prostate Cancer

Tracer	Half-Life (min)	Excretion	Detection Rates ^a	Panel Recommendation
Ga-68 PSMA-11 (PSMA-HBED-CC)	68	Renal	• 40% sensitivity and 95% specificity to detect nodal involvement in primary staging of patients with intermediate-, high-, and very-high-risk disease • 92% patient-level PPV in BCR	• May be considered as an alternative to CT, MRI, and bone scans for initial staging of unfavorable intermediate-, high-, and very-high-risk disease, the detection of biochemically recurrent disease, and as workup for progression • May be considered for equivocal results on initial bone scan
F-18 piflufolastat-PSMA (18F-DCFPyL)	110	Renal	• 31%–42% sensitivity and 96%–99% specificity to detect nodal involvement in primary staging of patients with unfavorable intermediate-risk, high-risk, and very-high-risk disease • 85%–87% patient-level CLR^b in BCR	
F-18 flutolastat PSMA (18F-rh-PSMA-7.3)^{6,7}	110	Renal	• 23%–30% sensitivity and 93%–97% specificity to detect nodal involvement in primary staging of patients with unfavorable-intermediate-, high-, and very-high-risk disease • 82% patient-level PPV in BCR	
C-11 choline	20	Hepatic and Renal	• 53%–96% PPV in BCR	• May be used to detect small-volume recurrent disease in soft tissues and in bone • May be considered for equivocal results on initial bone scan
F-18 fluciclovine (FACBC)	110	Renal	• 87%–91% CLR^b in BCR	
F-18 sodium fluoride	110	Renal	• 77%–94% sensitivity, 92%–99% specificity, and 82%–97% PPV for bone metastases	• May be considered for equivocal results on initial bone scan

^a Interpret with caution. Wherever possible, studies were included that used histopathologic confirmation, but not all studies used confirmatory histology as the gold standard. Values may vary depending upon the site of the lesion and phase of the disease process.

^b CLR: Correct localization rate. Patient-level positive predictive value (PPV) + anatomic lesion co-localization. Preferred over sensitivity and specificity in analyses of patients with BCR.

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF ACTIVE SURVEILLANCE AND OBSERVATION

- The NCCN Prostate Cancer Panel and the NCCN Prostate Cancer Early Detection Panel ([NCCN Guidelines for Prostate Cancer Early Detection](#)) remain concerned about overdiagnosis and overtreatment of prostate cancer. The Prostate Cancer Panel recommends that patients and their physicians carefully consider active surveillance based on the patient's prostate cancer risk profile and estimated life expectancy. In settings where the patient's age and comorbidities suggest a shorter life expectancy, observation may be more appropriate. Shared decision-making, after appropriate counseling on the risks and benefits of the various options, is critical.
- Life expectancy is a key determinant for the choice between observation and active surveillance. Multiple tools exist to assist in life expectancy estimation. See [Principles of Life Expectancy Estimation \(PROS-A\)](#).
- There remains no high-level evidence showing clinical benefit of advanced risk stratification tools (eg, gene expression biomarkers, germline or somatic testing, or artificial intelligence (AI)-based digital pathology models), to guide the candidacy for active surveillance.

ACTIVE SURVEILLANCE¹

- Active surveillance involves actively monitoring the course of disease with the expectation to intervene with curative intent if the cancer progresses. It involves PSA monitoring and longitudinal serial pathologic assessments via a prostate biopsy, often guided by MRI.
- There are three phases of active surveillance: 1) candidacy, 2) confirmation of candidacy, and 3) active surveillance.
- **Candidacy for Active Surveillance:**
 - ▶ **NCCN Low-Risk:**
 - ◊ Active surveillance is the preferred management approach for patients with low-risk prostate cancer and a life expectancy ≥10 years.
 - ◊ The Panel recognizes that there is heterogeneity across this risk group.
 - Some factors may be associated with an increased probability of near-term grade reclassification including:
 - High PSA density (≥0.15)²
 - Increased number of positive cores³
 - Higher body mass index (BMI)³
 - Faster PSA kinetics³
 - Conversely a negative biopsy following diagnosis is associated with a decreased probability of grade reclassification.
 - The Canary PASS risk calculator can be used for more precise estimates of upgrading probability on active surveillance ([Canary Pass Risk Calculator](#))
 - Higher risk features alone do not necessitate immediate treatment, but should be used as part of a shared-decision making discussion and in tailoring the intensity of active surveillance.
 - ◊ For patients who are at high-risk of suboptimal adherence to an active surveillance protocol, a careful discussion should be had about the potential risks of active surveillance. For some of these patients, upfront treatment with definitive RP or RT may be preferred.

Note: All recommendations are category 2A unless otherwise indicated.

[References on PROS-F 5 of 5](#)

PROS-F
1 OF 5



PRINCIPLES OF ACTIVE SURVEILLANCE AND OBSERVATION

• **Candidacy for Active Surveillance (continued):**

▶ **NCCN Favorable Intermediate-Risk (FIR)**

- ◊ Patients with FIR prostate cancer and a life expectancy of >10 years may also consider active surveillance.
- ◊ The FIR group was not created for optimal selection of active surveillance candidacy, and there is considerable heterogeneity in this risk group. Not all patients with FIR disease are suitable candidates for active surveillance. For example:
 - A patient with Grade Group 1 in a single biopsy core, cT1c, PSA of 12 ng/mL, and PSA density of 0.15 ng/mL/g has FIR disease and a favorable prognostic outcome. Therefore, active surveillance is preferred.
 - A patient with GG2 in 5 of 12 cores, cribriform architecture, cT1c, PSA of 8 ng/mL, and PSA density 0.30 ng/mL/g has FIR disease, but active surveillance would not be the preferred strategy.
- ◊ In addition to prognostic factors listed for Low Risk, additional prognostic factors relevant for FIR include:
 - Increased percentage pattern 4.
 - Presence of unfavorable histology such as expansile/large cribriform histology and/or intraductal carcinoma.
- ◊ Active surveillance is not recommended for patients with FIR and unfavorable histology (eg, expansile/large cribriform histology, intraductal carcinoma). See [Discussion](#).
- ◊ Intensity of active surveillance should generally be greater for patients with FIR disease.
- ◊ Patients should be counseled that long-term data show that patients with GG2 disease managed with active monitoring or watchful waiting had an increase in the development of distant metastasis as compared to those receiving immediate radical treatment.⁴ Continued refinements in active surveillance protocols and active surveillance candidacy are expected to improve the safety of active surveillance for GG2 FIR disease, and patients should be counseled on the importance of longitudinal follow-up.

• **Confirmation of Candidacy:**

- ▶ Goals of confirmation testing are to help facilitate early identification of patients who may be at a higher risk of grade reclassification or cancer progression.
- ▶ An initial prostate biopsy may underestimate tumor grade or volume, thus confirmatory testing is strongly recommended for patients who are considering active surveillance.
- ▶ A repeat biopsy is a routine component of confirmatory testing.
 - ◊ The timing of repeat biopsy can be based on the patient's risk of progression. Patients with a higher risk should have a repeat biopsy earlier than those with a lower risk.
 - ◊ If the initial prostate biopsy was performed without mpMRI guidance, then the Panel recommends an earlier repeat biopsy (eg, within the first 6–12 months) and recommends mpMRI prior to the biopsy. A systematic biopsy should be included.
 - ◊ For most patients, the timing is generally within 6 to 24 months of diagnostic biopsy, but all patients on active surveillance should undergo a confirmatory prostate biopsy within 3 years of their diagnostic biopsy, irrespective of prior mpMRI findings.
- ▶ Confirmatory testing should also include mpMRI with calculation of PSA density.
- ▶ Advanced risk stratification tools lack high level evidence to obviate other testing at this time.
- ▶ Other forms of imaging are discouraged.

Note: All recommendations are category 2A unless otherwise indicated.

[References on PROS-F 5 of 5](#)

PROS-F
2 OF 5



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF ACTIVE SURVEILLANCE AND OBSERVATION

- **Active Surveillance Program**

- ▶ Patients who choose active surveillance should have regular follow-up.
- ▶ Key principles include:
 - ◊ PSA no more often than every 6 months, unless clinically indicated.
 - ◊ DRE no more often than every 12 months, unless clinically indicated.
 - ◊ Repeat prostate biopsy no more often than every 12 months, unless clinically indicated. While the intensity of surveillance may be tailored based on patient and tumor factors (eg, grade, tumor volume), most patients should have prostate biopsies every 1 to 3 years as part of their monitoring. Longer intervals between biopsies may be considered if maintained on active surveillance for >3 years without progression.
 - ◊ Consider repeat prostate MRI no more than every 12 months, unless clinically indicated.
 - ◊ In patients with a suspicious lesion on prostate MRI, MRI-targeted biopsy improves the detection of higher grade (GG ≥2) cancers.
 - ◊ Patients should be transitioned to observation when life expectancy is <10 years.
 - ◊ A metastatic staging evaluation (PSMA PET, bone scan, CT scan, or whole-body MRI) should not be performed unless patient develops unfavorable intermediate or higher-risk disease, or has concerning symptoms not explainable by other causes.

- **Considerations for Treatment of Patients on Active Surveillance**

- ▶ Low risk: Grade reclassification on repeat biopsy to GG2 or higher is the most common factor to transition off active surveillance to treatment. PSA alone or growth in size of the tumor on MRI (assuming organ confined) should not be used alone to stop active surveillance without histologic confirmation of progression.
- ▶ FIR: Grade reclassification, increase in percent pattern 4, increase in tumor volume on biopsy or imaging, or rise in PSA density are common factors to transition off active surveillance to treatment.
- ▶ Patient anxiety is commonly cited as a reason to transition off active surveillance. However, studies indicate patient anxiety is closely linked with provider education on the safety of active surveillance. Thus, if a patient is an excellent candidate for active surveillance and has anxiety, the provider should provide appropriate education on the safety of active surveillance.

- **Advantages of Active Surveillance**

- ▶ Approximately 50% of those eligible for active surveillance may safely avoid treatment for 10 years.⁵⁻⁷ This varies by baseline risk.
- ▶ Patients will avoid or delay possible side effects of treatment. Quality of life will be less affected while on active surveillance.
- ▶ Risk of unnecessary treatment of small, indolent cancers will be reduced.

- **Limitations of Active Surveillance**

- ▶ According to current data, the majority of patients on active surveillance receive treatment by 15 years post-diagnosis.
- ▶ Although the risk is very low for patients diagnosed with GG1 disease (<1% at 10 years and 5% at 15 years), it is possible for cancer to progress to a regional or metastatic stage.⁵⁻⁸
- ▶ The majority of high quality evidence with >10 years of follow-up primarily limited active surveillance to patients with low-risk disease, low-volume and low PSA density.
- ▶ In early series, GG2 disease had higher rates of progression to metastatic disease.⁸ However, there has been grade migration, stage migration with MRI, and refinement in active surveillance candidacy with assessment of percentage pattern 4 and unfavorable histologic variants. Long-term follow-up (10–15 years) of patients with GG2 FIR disease continues to develop.

Note: All recommendations are category 2A unless otherwise indicated.

[References on PROS-F 5 of 5](#)

PROS-F
3 OF 5



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF ACTIVE SURVEILLANCE AND OBSERVATION

OBSERVATION

- Observation involves monitoring with a history and physical examination no more often than every 12 months (without surveillance biopsies, imaging, or PSA monitoring) until symptoms develop or are thought to be imminent.
- Low risk
 - ▶ Observation is recommended for asymptomatic patients with life expectancy <10 years.
- FIR
 - ▶ Observation is recommended for asymptomatic patients with a life expectancy ≤5 years.
 - ▶ Observation is preferred for asymptomatic patients with a life expectancy between 5 to 10 years.
- Observation may be considered for asymptomatic patients with high-risk, very-high-risk, regional, and metastatic prostate cancer and a life expectancy ≤5 years.
- If patients under observation become symptomatic, an assessment of disease burden can be performed, and treatment per the appropriate disease setting or palliation can be considered.
- **Advantages of Observation**
 - ▶ Patients will avoid possible side effects of unnecessary confirmatory testing and definitive therapy with a low probability that this would compromise their survival.
- **Limitations of Observation**
 - ▶ There may be risk of local or systemic symptoms (eg, urinary retention, pathologic fracture) that could have been avoided with upfront definitive treatment.

Note: All recommendations are category 2A unless otherwise indicated.



PRINCIPLES OF ACTIVE SURVEILLANCE AND OBSERVATION

REFERENCES

- ¹ Ganz PA, Barry JM, Burke W, et al. NIH State-of-the-Science Conference Statement: Role of active surveillance in the management of men with localized prostate cancer. NIH Consens State Sci Statements 2011;28:1-27.
- ² Mcleod LC, Ellis WJ, Newcomb LF, et al. Timing of adverse prostate cancer reclassification on first surveillance biopsy: Results from the Canary prostate cancer active surveillance study. J Urol 2017;197:1026-1033.
- ³ Cooperberg MR, Zheng Y, Faino AV, et al. Tailoring intensity of active surveillance for low-risk prostate cancer based on individualized prediction of risk stability. JAMA Oncol 2020;6:e203187.
- ⁴ Wilt TJ, Vo TN, Langsetmo L, et al. Radical prostatectomy or observation for clinically localized prostate cancer: Extended follow-up of the prostate cancer intervention versus observation trial (PIVOT). Eur Urol 2020;77:713-724.
- ⁵ Tosoian JJ, Mamawala M, Epstein JI, et al. Intermediate and longer-term outcomes from a prospective active-surveillance program for favorable-risk prostate cancer. J Clin Oncol 2015;33:3379-3385.
- ⁶ Klotz L, Vesprini D, Sethukavalan P, et al. Long-term follow-up of a large active surveillance cohort of patients with prostate cancer. J Clin Oncol 2015;33:272-277.
- ⁷ Cooley LF, Emeka AA, Meyers TJ, et al. Factors associated with time to conversion from active surveillance to treatment for prostate cancer in a multi-institutional cohort. J Urol 2021;206:1147-1156.
- ⁸ Musunuru HB, Yamamoto T, Klotz L, et al. Active surveillance for intermediate risk prostate cancer: Survival outcomes in the sunnybrook experience. J Urol 2016;196:1651-1658.

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF ANDROGEN DEPRIVATION THERAPY

Neoadjuvant, Concurrent, and/or Adjuvant ADT with RT^a:

- Clinically localized (N0, M0) disease
- Regional (N1, M0) disease
- Positive lymph nodes (pN1 disease) and/or adverse features post-RP
- First M0 RP PSA persistence/recurrence

RT with one of the following ADT options:

- Luteinizing hormone-releasing hormone (LHRH) agonist monotherapy
 - Goserelin, leuprolide, or triptorelin
- LHRH agonist (as above) plus first-generation antiandrogen
 - Nilutamide, flutamide, or bicalutamide
- LHRH antagonist
 - Degarelix or relugolix

LHRH agonist or antagonist plus abiraterone^b (2 years; only for very-high-risk localized disease or positive lymph nodes)

Notes:

- ADT is not recommended with RT for most patients with favorable intermediate-risk prostate cancer. If it is given (see [Principles of Risk Stratification](#)), the duration should be short term (4–6 months).
- For unfavorable intermediate risk prostate cancer treated with RT, short-term ADT (ST-ADT) (4–6 months) is recommended. Concurrent/ adjuvant ADT is preferred over neoadjuvant ADT in this setting.
- For high- and very-high-risk prostate cancer treated with EBRT alone, long-term ADT (LT-ADT) (18–36 months) is recommended.
- For high-risk prostate cancer treated with combination EBRT + brachytherapy, a shortened duration of ADT (12 months) can be considered.
- For additional details on the use of RT with ADT by risk group, see [PROS-J](#).
- First M0 PSA persistence/recurrence:
 - The timing of secondary treatment for patients whose only evidence of cancer after definitive treatment is an increasing PSA is influenced by PSA velocity, patient anxiety, the short- and long-term side effects of ADT, and the underlying comorbidities of the patient.
 - Earlier treatment may be better than delayed treatment, although the definitions of early and late (what level of PSA) are controversial. Since the benefit of early treatment is not clear, treatment should be individualized until definitive studies are done. Patients with a shorter PSADT (or a rapid PSA velocity) and an otherwise long life expectancy should be encouraged to consider treatment earlier.
 - For patients treated with secondary RT in the setting of first RP recurrence, if ADT is given, it should be for a duration of 6 to 24 months.
 - Patients with prolonged PSADTs (>12 months) and who are older are candidates for observation or monitoring.
 - Patients who choose ADT monotherapy in the secondary treatment setting should consider intermittent ADT.

^a Specific recommendations, Categories of Evidence and Consensus, and Categories of Preference vary based on patient and disease characteristics (see [PROS-5](#) through [PROS-18](#)). This chart delineates the forms of ADT that can be used and provides some additional details.

^b The fine-particle formulation of abiraterone can be used instead of the standard form (category 2B; other recommended). Abiraterone should be given with concurrent steroid: prednisone 5 mg PO once daily (in the CSPC setting without docetaxel) or twice daily (in the CSPC setting with docetaxel and in the CRPC setting) with the standard formulation or methylprednisolone 4 mg PO twice daily with the fine-particle formulation. The standard formulation of abiraterone can be given at 250 mg/day following a low-fat breakfast in patients who will not take or cannot afford the standard dose of 1000 mg/day after an overnight fast.

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF ANDROGEN DEPRIVATION THERAPY

ADT for Patients with M0 CSPC^a:

- Regional (N1, M0) disease
- First M0 RT recurrence

ADT Options:

- Orchiectomy
- LHRH agonist monotherapy
 - Goserelin, leuprolide, or triptorelin
- LHRH agonist (as above) plus first-generation antiandrogen
 - Nilutamide, flutamide, or bicalutamide
- LHRH antagonist
 - Degarelix or relugolix
- LHRH agonist, LHRH antagonist, or orchiectomy plus abiraterone^b (only for positive lymph nodes)

Notes:

- M0 PSA persistence/recurrence:
 - The timing of secondary treatment for patients whose only evidence of cancer after definitive treatment is an increasing PSA is influenced by PSA velocity, patient anxiety, the short- and long-term side effects of ADT, and the underlying comorbidities of the patient.
 - Earlier treatment may be better than delayed treatment, although the definitions of early and late (what level of PSA) are controversial. Since the benefit of early treatment is not clear, treatment should be individualized until definitive studies are done. Patients with a shorter PSADT (or a rapid PSA velocity) and an otherwise long life expectancy should be encouraged to consider treatment earlier.
 - Patients with prolonged PSADTs (>12 months) and who are older are candidates for observation or monitoring.
 - Patients who choose ADT monotherapy in the secondary treatment setting should consider intermittent ADT.

^a Specific recommendations, Categories of Evidence and Consensus, and Categories of Preference vary based on patient and disease characteristics (see [PROS-5](#) through [PROS-18](#)). This chart delineates the forms of ADT that can be used and provides some additional details.

^b The fine-particle formulation of abiraterone can be used instead of the standard form (category 2B; other recommended). Abiraterone should be given with concurrent steroid: prednisone 5 mg PO once daily (in the CSPC setting without docetaxel) or twice daily (in the CSPC setting with docetaxel and in the CRPC setting) with the standard formulation or methylprednisolone 4 mg PO twice daily with the fine-particle formulation. The standard formulation of abiraterone can be given at 250 mg/day following a low-fat breakfast in patients who will not take or cannot afford the standard dose of 1000 mg/day after an overnight fast.

Note: All recommendations are category 2A unless otherwise indicated.



PRINCIPLES OF ANDROGEN DEPRIVATION THERAPY

ADT for Patients with M0 CSPC BCR2^a:

ADT Options^c:

- **Orchiectomy**
- **LHRH agonist monotherapy**
 - ▶ Goserelin, leuprolide, or triptorelin
- **LHRH agonist (as above) plus first-generation antiandrogen**
 - ▶ Nilutamide, flutamide, or bicalutamide
- **LHRH antagonist**
 - ▶ Degarelix or relugolix

For High-Risk BCR2:

- Enzalutamide with or without Leuprolide^d
- Apalutamide with LHRH agonist or LHRH antagonist^e

Notes:

- **Monitoring until diagnosis of metastatic disease is preferred for patients with M0 CSPC in the low-risk BCR2 setting.**
- **PSADT and Grade Group should be considered when deciding whether to begin ADT for patients with M0 disease.**
- **ADT monotherapy is an option for these patients, and intermittent ADT can be considered to reduce toxicity.**

^a Specific recommendations, Categories of Evidence and Consensus, and Categories of Preference vary based on patient and disease characteristics (see [PROS-5](#) through [PROS-18](#)). This chart delineates the forms of ADT that can be used and provides some additional details.

^c For patients receiving MDT with ADT, the ADT options are LHRH agonist or LHRH antagonist.

^d Enzalutamide with or without leuprolide is an option for patients who have the following high-risk criteria: M0 by CT, MRI, or bone scan; PSADT ≤9 months; PSA ≥2 ng/mL above nadir after RT or ≥1 ng/mL after RP with or without postoperative RT; and not considered a candidate for pelvic-directed therapy.

^e Apalutamide plus ADT is an option for patients with biochemical recurrence after RP who meet the following high-risk criteria: PSADT ≤ 9 months; PSA ≥0.5 ng/mL; and prior adjuvant or secondary RT or not considered a candidate for RT (Aggarwal R, et al. J Clin Oncol 2024;42:1114-1123).

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF ANDROGEN DEPRIVATION THERAPY

ADT for Patients with mCSPC^a:

ADT with treatment intensification (systemic therapy):

- Orchiectomy plus abiraterone,^b apalutamide, darolutamide, or enzalutamide
- LHRH agonist plus abiraterone,^b apalutamide, darolutamide, or enzalutamide
- LHRH antagonist plus abiraterone,^b apalutamide, darolutamide, or enzalutamide
- Orchiectomy plus docetaxel and either abiraterone,^b apalutamide, darolutamide, or enzalutamide
- LHRH agonist plus docetaxel and either abiraterone,^b apalutamide, darolutamide, or enzalutamide
- LHRH antagonist plus docetaxel and either abiraterone,^b apalutamide, darolutamide, or enzalutamide

ADT with treatment intensification (EBRT to the primary tumor):

EBRT with:

- Orchiectomy alone or with abiraterone,^b apalutamide, docetaxel, or enzalutamide
- LHRH agonist alone or with abiraterone,^b apalutamide, docetaxel,^f or enzalutamide
- LHRH antagonist alone or with abiraterone,^b apalutamide, docetaxel, or enzalutamide

Additional option for patients with BRCA2m mCSPC:

- Orchiectomy, LHRH agonist, or LHRH antagonist plus niraparib and abiraterone^g

ADT alone for select patients^h:

- **Orchiectomy**
- **LHRH agonist monotherapy^f**
 - ▶ Goserelin, leuprolide, or triptorelin
- **LHRH agonist (as above) plus first-generation antiandrogen**
 - ▶ Nilutamide, flutamide, or bicalutamide
- **LHRH antagonist**
 - ▶ Degarelix or relugolix

^a Specific recommendations, Categories of Evidence and Consensus, and Categories of Preference vary based on patient and disease characteristics (see [PROS-5](#) through [PROS-18](#)). This chart delineates the forms of ADT that can be used and provides some additional details.

^b The fine-particle formulation of abiraterone can be used instead of the standard form (category 2B; other recommended). Abiraterone should be given with concurrent steroid: prednisone 5 mg PO once daily (in the CSPC setting without docetaxel) or twice daily (in the CSPC setting with docetaxel and in the CRPC setting) with the standard formulation or methylprednisolone 4 mg PO twice daily with the fine-particle formulation. The standard formulation of abiraterone can be given at 250 mg/day following a low-fat breakfast in patients who will not take or cannot afford the standard dose of 1000 mg/day after an overnight fast.

Note: All recommendations are category 2A unless otherwise indicated.

^f A first-generation antiandrogen must be given with LHRH agonist for ≥7 days to prevent testosterone flare if metastases are present in weight-bearing bone or if there is a large prostate mass.

^g The fine-particle (category 2B; other recommended option) or standard formulation of abiraterone can be given with single-agent niraparib as a substitute for the combination niraparib/abiraterone tablet.

^h ADT is strongly recommended in combination therapy for metastatic castration-sensitive disease. The use of ADT monotherapy in metastatic castration-sensitive disease is discouraged unless there are clear contraindications to combination therapy. If ADT monotherapy is given, intermittent ADT can be considered to reduce toxicity. Close monitoring of PSA and testosterone levels and possibly imaging is required when using intermittent ADT, especially during off-treatment periods, and patients may need to switch to continuous ADT upon signs of disease progression.



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF ANDROGEN DEPRIVATION THERAPY

Secondary Hormone Therapy for Patients with M0 or M1 CRPC^a:

Castrate levels of testosterone (<50 ng/dL) should be maintained while additional therapies are applied:

- Orchiectomy, LHRH agonist, or LHRH antagonist with specific therapies noted on [PROS-16](#) [M0 CRPC], [PROS-17](#) [NEPC], and [PROS-18](#) [M1 CRPC]
- Orchiectomy, LHRH agonist, or LHRH antagonist with other secondary hormone options:
 - **Abiraterone^b or enzalutamide following progression on ARPIs (M1 only)**
 - **Abiraterone^b plus 0.5 mg/day dexamethasone following progression on either formulation of abiraterone (M1 only)**
 - **Antiandrogen withdrawalⁱ**
 - **Corticosteroids** (hydrocortisone, prednisone, or dexamethasone)ⁱ
 - **First-generation anti-androgen** (nilutamide, flutamide, or bicalutamide)ⁱ
 - **Ketoconazole plus hydrocortisoneⁱ**

Notes:

- Although the optimal sequence of therapies remains undefined, some data are emerging that can help with treatment selection in some cases. See [Discussion](#).

ADT Monotherapy^a:

ADT Options:

- **Orchiectomy**
- **LHRH agonist monotherapy^f**
 - Goserelin, leuprolide, or triptorelin
- **LHRH antagonist**
 - Degarelix or relugolix

Notes:

- **ADT monotherapy is appropriate for patients with life expectancy ≤5 years whose cancer progressed on observation of localized disease, who are symptomatic, or who have N1M0 disease.**
- **ADT monotherapy is also used for asymptomatic patients with regional disease and life expectancy ≤5 years whether or not RT is given.**

^a Specific recommendations, Categories of Evidence and Consensus, and Categories of Preference vary based on patient and disease characteristics (see [PROS-5](#) through [PROS-18](#)). This chart delineates the forms of ADT that can be used and provides some additional details.

^b The fine-particle formulation of abiraterone can be used instead of the standard form (category 2B; other recommended). Abiraterone should be given with concurrent steroid: prednisone 5 mg PO once daily (in the CSPC setting without docetaxel) or twice daily (in the CSPC setting with docetaxel and in the CRPC setting) with the standard formulation or methylprednisolone 4 mg PO twice daily with the fine-particle formulation. The standard formulation of abiraterone can be given at 250 mg/day following a low-fat breakfast in patients who will not take or cannot afford the standard dose of 1000 mg/day after an overnight fast.

^f A first-generation antiandrogen must be given with LHRH agonist for ≥7 days to prevent testosterone flare if metastases are present in weight-bearing bone or if there is a large prostate mass.

ⁱ In the mCRPC setting, this option should only be used for select patients who are not candidates for other recommended mCRPC therapies.

Note: All recommendations are category 2A unless otherwise indicated.



PRINCIPLES OF ANDROGEN DEPRIVATION THERAPY

Optimal ADT:

- Medical castration (ie, LHRH agonist or antagonist) and surgical castration (ie, bilateral orchiectomy) are equally effective.
- Patients who do not achieve adequate suppression of serum testosterone (<50 ng/dL) with medical or surgical castration can be considered for additional hormonal manipulations (with antiandrogens, LHRH antagonists, or steroids), although the clinical benefit remains uncertain. Consider monitoring testosterone levels 12 weeks after first dose of LHRH therapy, then upon increase in PSA. The optimal level of serum testosterone to affect “castration” has yet to be determined.

Monitor/Surveillance:

- ADT has a variety of adverse effects, including hot flashes, loss of libido, erectile dysfunction, shrinkage of penis and testicles, loss of muscle mass and strength, fatigue, anemia, breast enlargement and tenderness/soreness, depression and mood swings, hair loss, osteoporosis, greater incidence of clinical fractures, obesity, insulin resistance, alterations in lipids, and greater risk for diabetes and cardiovascular disease. The intensity and spectrum of these side effects vary greatly, and many are reversible or can be avoided or mitigated. See [NCCN Guidelines for Survivorship](#). Patients and their medical providers should be advised about these risks prior to treatment.

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF RISK STRATIFICATION AND BIOMARKERS

General Principles:

- Currently, the primary method for personalization of treatment from localized to advanced prostate cancer is based on prognostic risk stratification, rather than the use of predictive tools.
- NCCN uses multiple categories and subgroupings to capture prognostic risk to personalize treatment recommendations.
- The purpose of the NCCN categories and subgroupings are to provide a method for risk stratification to allow standardized treatment recommendations to be provided.
 - ▶ It is acknowledged that there are methods of risk stratification with superior prognostic performance to NCCN risk groups. However, they have not been routinely reported in clinical trials. This limits the ability to provide evidence-based guideline treatment recommendations using these methods. Thus, the NCCN Guidelines continue to use NCCN categories and subgroups of risk as a framework.
 - ▶ Clinical trials have established the benefit of various treatments in prostate cancer and have commonly enrolled patients across a spectrum of risk. Subgroup analyses, absolute benefit estimates, and expert opinion are used to provide treatment recommendations for each NCCN risk group or disease state.
 - ▶ There is intrinsic heterogeneity in prognosis within a given NCCN category and subgroup. Thus, treatment recommendations for adjacent subgroups or categories of risk may be appropriate when using additional risk stratification methods.
 - ▶ The Panel acknowledges the ability to personalize treatment decisions through additional tools and have created this section to assist.
- Tools that are prognostic or predictive in one disease state may not be in other disease states, or they may have other forms of clinical utility beyond prognostication and prediction of treatment benefit.
 - ▶ For example, germline homologous recombination deficiency (HRD) mutations do not have an established prognostic or predictive role in localized prostate cancer, but specific HRD mutations have been demonstrated to have a prognostic and predictive role in advanced disease. Additionally, the utility of germline testing extends to inform screening recommendations for other cancers and cascade germline testing for family members.
- Imaging is also a biomarker (ie, MRI, PSMA-PET/CT) and can aid in risk stratification. See [Principles of Imaging \(PROS-E\)](#).

Note: All recommendations are category 2A unless otherwise indicated.



PRINCIPLES OF RISK STRATIFICATION AND BIOMARKERS

Biomarker Categories:

- Biomarkers and risk stratification methods are tools that may assist in personalization of treatment. For clarity these tools are separated by type and category:
 - ▶ **Type:**
 - ◇ **Standard Tools:** These include clinical and/or pathologic variables routinely collected to assign a patient to an NCCN category and/or subgroup. Examples include TNM stage, Grade Group, PSA, and metastatic volume of disease.
 - ◇ **Clinical and Pathologic Tools:** These include clinical and/or pathologic tools that are generally derived from standard tools. Examples include multivariable models or nomograms, histologic variants, and PSA kinetics.
 - ◇ **Advanced Tools:** These involve an additional test above what is collected to assign an NCCN category or subgroup. These may include, but are not limited to, germline or somatic tests, gene expression tests, digital histopathology-based tests, additional imaging, and circulating markers.
 - ▶ **Category:**
 - ◇ **Prognostic:** Discriminates the risk of developing an oncologic endpoint (eg, distant metastasis). The relative benefit of a treatment (ie, the treatment effect or hazard ratio) is generally similar across a prognostic spectrum, although the absolute benefit of an intervention may vary by risk (ie, number needed to treat [NNT]).
 - Ideally, prognostic biomarkers independently discriminate and are associated with a clinically meaningful endpoint above and beyond standard tools relevant to that disease setting that ultimately helps guide a therapeutic decision.
 - ◇ **Predictive:** Discriminates a difference in the relative benefit of a specific treatment for an oncologic endpoint.
 - Ideally, predictive biomarkers have been demonstrated to measure differential treatment effects that ultimately help guide a therapeutic decision in the context of a randomized trial, specifically randomizing the treatment of interest.

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF RISK STRATIFICATION AND BIOMARKERS

Clinical and Pathologic Tools:

- An extensive number of prognostic clinical or pathologic tools have been reported based on highly variable evidence quality (retrospective or registry study vs. randomized trial), validation rigor, strength of endpoint (adverse pathology or BCR vs. distant metastasis), and univariable versus multivariable association with an outcome. Thus, while some of these tools may have value, these limitations hinder the ability to accurately provide guidance to specific treatment recommendations with confidence.
- A comprehensive list of these tools is outside the scope of this guideline.
 - ▶ Examples of such prognostic tools include multivariable models and nomograms (eg, CAPRA,¹ STAR-CAP,² MSKCC nomograms³), histopathology (ie, cribriform, intraductal carcinoma, absolute and percent Gleason pattern 4, total mm of cancer), and clinical variables (ie, PSA density, PSA velocity, PSA level, PSADT).

Advanced Tools:

- Advanced risk stratification tools generally offer either superior prognostic performance beyond clinical and pathologic tools AND/OR serve as predictive biomarkers for identifying patient groups that differentially benefit from a specific treatment. In general, these tools are only recommended when they have the potential ability to change management and should not be ordered reflexively.
 - ▶ Prognostic tools: Generally, the Panel recommends the use of prognostic tests that are validated in well-designed prospective studies with clinically meaningful endpoints based on disease settings that guide a specific treatment indication based on a specific score or result. These studies can be either prospective integral or integrated clinical trial(s) or post-hoc correlative analyses of prospective trials.
 - ▶ Predictive tools: Generally, the Panel recommends the use of predictive tests that are validated in a prospective, biomarker-directed, randomized clinical trial of the treatment of interest. Alternatively, if validation is post-hoc in a trial not designed to test the tool, it should be performed in more than one independent randomized trial of the treatment of interest.
- The Panel recognizes that there is an extensive number of advanced tools created with substantial variability in quality of reporting and model design, endpoint selection, and quality and caliber of validation. There are risks in using advanced tools to change management without robust validation, as they may drive patients or providers to inappropriate treatment options. If advanced tools are used, it is recommended to use tests that have robust validation, ideally with high-quality, long-term clinical trial data and across multiple clinical trials.
- Only advanced tools with high evidence quality are shown in Table 1. Other prognostic tools that are commonly used are described in the [Discussion](#).

Note: All recommendations are category 2A unless otherwise indicated.

[References](#)



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF RISK STRATIFICATION AND BIOMARKERS

Table 1. Advanced Prognostic Tools^a

Intermediate-Risk Prostate Cancer		
Tool	Category	Discussion
22-gene genomic classifier (GC) (Decipher) ⁴⁻⁶	Gene Expression	NRG/RTOG 0126 phase III randomized trial was profiled post-hoc with a prespecified analysis plan. ⁴ The study demonstrated the independent prognostic effect of GC on biochemical failure, secondary therapy, DM, PCSM, MFS, and OS. Patients receiving RT alone with low GC scores had 10-year DM rates of 4%, compared with 16% for GC high risk. These results suggest that the benefit of short-term ADT in NCCN intermediate-risk prostate cancer is likely to be smaller in patients with low GC scores (≤ 0.45) than in patients with high GC scores (≥ 0.60). A breakdown of outcomes between patients with favorable vs. unfavorable intermediate-risk disease was not provided.
Multimodal artificial intelligence (MMAI) (ArteraAI Prostate) ^{b,7-11}	AI-Pathology	A post-hoc meta-analysis of eight phase III randomized trials (NRG/RTOG 9202, 9408, 9413, 9902, 9910, 0126, 0415, and 0521) with MMAI scores have been reported. ^{7,8} There were a total of 630 patients with NCCN intermediate-risk disease who primarily received RT alone or RT+ST-ADT. Among patients with NCCN intermediate-risk disease, those with MMAI low-risk disease had 5-year and 10-year DM rates of 1% and 4%, respectively. In contrast, those with MMAI intermediate- and high-risk tumors had 5- and 10-year DM rates of 7% and 40%. MMAI was independently prognostic for DM, PCSM, and death after DM. These results suggest that the benefit of short-term ADT in NCCN intermediate-risk prostate cancer is likely to be smaller in patients with MMAI low-risk disease than in patients with MMAI intermediate- or high-risk disease. Neither a breakdown of favorable vs. unfavorable intermediate risk, nor a breakdown of outcomes after RT alone vs. RT+ST-ADT were provided.

DM = distant metastases; MFS = metastasis-free survival; PCSM = prostate cancer-specific mortality; ST-ADT = short-term ADT

^a In the absence of prospective trials, caution is warranted if using these prognostic tools to influence treatment decisions. The Panel awaits future trials that confirm the initial results described here.

^b There is also an MMAI predictive biomarker that was validated post-hoc in RTOG 9408 to predict benefit of ST-ADT added to RT in patients with intermediate-risk prostate cancer. While promising, due to differences in tissue sampling, grading, staging, and treatment, the Panel recommends further validation prior to using this predictive biomarker to guide treatment decisions in isolation.

Note: All recommendations are category 2A unless otherwise indicated.

References

PROS-H
4 OF 6



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF RISK STRATIFICATION AND BIOMARKERS

Table 1. Advanced Prognostic Tools^a

High-Risk and Very-High-Risk Prostate Cancer		
Tool	Category	Discussion
22-gene GC ⁴⁻⁶	Gene Expression	A meta-analysis of three phase III randomized trials (NRG/TOG 9202, 9413, and 9902) that included patients with high-/very-high-risk prostate cancer were profiled post-hoc with a prespecified analysis plan. ⁵ The study demonstrated the independent prognostic effect of GC on biochemical failure, DM, MFS, PCSM, and OS. Patients with low GC scores had 10-year DM rates of 6%, compared with 26% for GC high risk. The absolute benefit of LT-ADT over ST-ADT was 11% for patients with high GC scores (NNT of 9), and 3% for patients with low GC scores (NNT of 33). These results suggest that the benefit of long-term ADT in NCCN high- and very-high-risk prostate cancer is likely to be smaller in patients with low GC scores (≤ 0.45) than in patients with high GC scores (≥ 0.60). A breakdown of outcomes between patients with high- vs. very high-risk disease was not provided.
MMAI ^{b,7-12}	AI-Pathology	Multiple post-hoc meta-analyses of patients with NCCN high- or very-high-risk disease from six randomized phase III randomized trials (NRG/TOG 9202, 9408, 9413, 9902, 9910, and 0521) with MMAI scores have been reported.^{8,10} A total of 1090 patients had NCCN high-/very-high-risk disease and were treated predominately with RT+LT-ADT (with or without chemotherapy; 68%) or RT+ST-ADT (30%), and a small number with RT alone (2%). Among patients with NCCN high-/very-high-risk disease, those with MMAI low-risk disease had 5-year rates of DM of 0% and 10-year rates of 3%. In contrast, those with MMAI intermediate- and high-risk tumors had 8% and 26% risk of metastasis at 10 years post-treatment, respectively. MMAI was independently prognostic for DM, PCSM, and death after DM. These results suggest that the benefit of ADT in NCCN high- and very-high-risk prostate cancer is likely to be smaller in patients with MMAI low-risk disease than in patients with MMAI intermediate- or high-risk disease. Neither a breakdown of high- vs. very high-risk, nor a breakdown of outcomes between cohorts receiving RT+ST-ADT vs. RT+LT-ADT, were provided. Published results from four phase III randomized trials from the STAMPEDE platform, which included 1575 patients with locally advanced, M0 disease (ie, NCCN very-high-risk disease), with post-hoc derivation of MMAI scores have been reported.¹² Overall, patients with locally advanced N0M0 disease had a 5% PCSM event rate, but when divided by MMAI quartile, there was a significant difference between the lower three quartiles (3%) vs. the highest quartile (11%). When restricting to patients receiving abiraterone, these rates were 2% vs. 5%. These results suggest that the benefit of abiraterone in NCCN very-high-risk prostate cancer is likely to be smaller in patients with MMAI low-risk disease than in patients with MMAI high-risk disease.
BCR Post-RP		
Tool	Category	Discussion
22-gene GC ^{13,14}	Gene Expression	Two phase III randomized trials post-RP were profiled post-hoc with prespecified analysis plans. NRG/TOG 9601 demonstrated the independent prognostic effect of GC on DM, PCSM, and OS, and found that for patients with lower entry PSAs (< 0.7 ng/mL), the 12-year DM rate benefit from hormone therapy for patients with GC lower risk vs. GC higher risk was 0.4% vs. 11.2%. ¹³ The SAKK 09/10 phase III trial tested post-RP lower vs. higher dose RT alone. The study demonstrated the independent prognostic effect of GC on biochemical progression, clinical progression, secondary hormone therapy, DM, and MFS. ^{c,14} These results suggest that the benefit of ADT added to RT for patients with RP recurrence planned for early secondary RT is likely to be smaller in those with low or intermediate GC scores (< 0.6) than in those with high GC scores (≥ 0.60).

DM = distant metastases; LT-ADT = long-term ADT; MFS = metastasis-free survival; NNT = number needed to treat; PCSM = prostate cancer-specific mortality; ST-ADT = short-term ADT

^a In the absence of prospective trials, caution is warranted if using these prognostic tools to influence treatment decisions. The Panel awaits future trials that confirm the initial results described here.

^b There is also an MMAI predictive biomarker that was validated post-hoc in RTOG 9408 to predict benefit of ST-ADT added to RT in patients with intermediate-risk prostate cancer. While promising, due to differences in tissue sampling, grading, staging, and treatment, the Panel recommends further validation prior to using this predictive biomarker to guide treatment decisions in isolation.

^c SAKK 09/10 combined GC low and intermediate risk due to relatively similar prognosis. NRG/TOG 9601 dichotomized patients by GC low versus intermediate and high risk. However, due to the age of the tissue from NRG/TOG 9601 (> 20 years old), there is a known shifting of GC scores, and a more contemporary distribution of score distribution would approximate closer to combining GC low and intermediate risk together.

Note: All recommendations are category 2A unless otherwise indicated.

References

PROS-H
5 OF 6



PRINCIPLES OF RISK STRATIFICATION AND BIOMARKERS REFERENCES

- ¹ Cooperberg MR, Pasta DJ, Elkin EP, et al. The University of California, San Francisco Cancer of the Prostate Risk Assessment score: a straightforward and reliable preoperative predictor of disease recurrence after radical prostatectomy. *J Urol* 2005;173:1938-1942. Erratum in: *J Urol* 2006;175:2369.
- ² Dess RT, Suresh K, Zelefsky MJ, et al. Development and validation of a clinical prognostic stage group system for nonmetastatic prostate cancer using disease-specific mortality results from the International Staging Collaboration for Cancer of the Prostate. *JAMA Oncol* 2020;6:1912-1920.
- ³ Memorial Sloan-Kettering Cancer Center. Pre-Radical Prostatectomy. Available at: https://www.mskcc.org/nomograms/prostate/pre_op.
- ⁴ Spratt DE, Liu VYT, Michalski J, et al. Genomic classifier performance in intermediate-risk prostate cancer: Results from NRG Oncology/RTOG 0126 randomized phase 3 trial. *Int J Radiat Oncol Biol Phys* 2023;117:370-377.
- ⁵ Nguyen PL, Huang HR, Spratt DE, et al. Analysis of a biopsy-based genomic classifier in high-risk prostate cancer: Meta-analysis of the NRG Oncology/Radiation Therapy Oncology Group 9202, 9413, and 9902 phase 3 randomized trials. *Int J Radiat Oncol Biol Phys* 2023;116:521-529.
- ⁶ Spratt DE, Zhang J, Santiago-Jiménez M, et al. Development and validation of a novel integrated clinical-genomic risk group classification for localized prostate cancer. *J Clin Oncol* 2018;36:581-590.
- ⁷ Esteva A, Feng J, van der Wal D, et al. Prostate cancer therapy personalization via multi-modal deep learning on randomized phase III clinical trials. *NPJ Digit Med* 2022;5:71. Erratum in: *NPJ Digit Med* 2023;6:27.
- ⁸ Tward JD, Huang HC, Esteva A, et al. Prostate cancer risk stratification in NRG Oncology phase III randomized trials using multimodal deep learning with digital histopathology. *JCO Precis Oncol* 2024;e2400145.
- ⁹ Spratt DE, Tang S, Sun Y, et al. Artificial intelligence predictive model for hormone therapy use in prostate cancer. *NEJM Evid* 2023;2:EVIDoa2300023.
- ¹⁰ Spratt DE, Liu VYT, Jia AY, et al. Meta-analysis of individual patient-level data for a multimodal artificial intelligence biomarker in high-risk prostate cancer: Results from six NRG/RTOG phase 3 randomized trials. *Eur Urol* 2024;86:369-371.
- ¹¹ Ross AE, Zhang J, Huang HC, et al. External validation of a digital pathology-based multimodal artificial intelligence architecture in the NRG/RTOG 9902 phase 3 trial. *Eur Urol Oncol* 2024;7:1024-1033.
- ¹² Parker CTA, Mendes L, Liu VYT, et al. External validation of a digital pathology-based multimodal artificial intelligence-derived prognostic model in patients with advanced prostate cancer starting long-term androgen deprivation therapy: a post-hoc ancillary biomarker study of four phase 3 randomised controlled trials of the STAMPEDE platform protocol. *Lancet Digit Health* 2025;7:100885.
- ¹³ Feng FY, Huang HC, Spratt DE, et al. Validation of a 22-gene genomic classifier in patients with recurrent prostate cancer: An ancillary study of the NRG/RTOG 9601 randomized clinical trial. *JAMA Oncol* 2021;7:544-552. Erratum in: *JAMA Oncol* 2021;7:639.
- ¹⁴ Dal Pra A, Ghadjar P, Hayoz S, et al. Validation of the Decipher genomic classifier in patients receiving salvage radiotherapy without hormone therapy after radical prostatectomy - an ancillary study of the SAKK 09/10 randomized clinical trial. *Ann Oncol* 2022;33:950-958.

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF FOCAL/SUBTOTAL THERAPY OR WHOLE GLAND ABLATIVE THERAPY

General

- Ablative therapy refers to the use of a device to destroy tissue that can be directed at part of the prostate gland (termed partial gland, hemi-gland, or focal therapy [FT]) or the whole prostate gland.
- FT includes ablative therapies and non-ablative therapies, such as partial prostatectomy and focal radiation therapy.^a
- Presently, FT is an experimental and emerging technology for the initial treatment of localized prostate cancer that lacks randomized controlled trial evidence with long-term follow-up showing its superiority or noninferiority to current recommended management strategies. As such, FT meets the criteria as an alternative therapy, or a non-standard treatment.

Newly Diagnosed or Previously Untreated Prostate Cancer

- Pathologic confirmation of prostate cancer and risk stratification is required prior to consideration of ablative or FT.
- Active surveillance is the preferred treatment for low-risk prostate cancer, and ablative and focal therapies are discouraged in this population.
- The Panel also believes that ablative FT should be discouraged in patients with high- or very-high-risk, regional, or metastatic prostate cancer outside of a clinical trial.
- There is currently insufficient comparative effectiveness evidence for FT to be recommended for patients with intermediate-risk prostate cancer, and the Panel believes there is uncertainty about the long-term efficacy and toxicity of these treatments. Therefore, ablative and focal therapies in this population should be utilized only in the context of a clinical trial.
- The following device categories are currently FDA approved or cleared for the initial treatment of prostate cancer, but randomized evidence to the superiority in long-term cancer control and/or quality of life are lacking when delivered as focal rather than whole gland therapy. These treatments may be considered as FT for patients with previously untreated, localized prostate cancer only in the context of a clinical trial:
 - ▶ External beam radiotherapy^a
 - ▶ Brachytherapy
 - ▶ Cryotherapy ablation
- The following device categories are currently not FDA approved or cleared for the treatment of prostate cancer as focal or whole gland therapy, and should only be used for patients with previously untreated, localized prostate cancer in the context of a clinical trial:
 - ▶ High-intensity focused ultrasound (HIFU)
 - ▶ Transurethral ultrasound ablation (TULSA)
 - ▶ Focal laser ablation (FLA)
 - ▶ Radiofrequency ablation (RFA)
 - ▶ Irreversible electroporation (IRE)
 - ▶ Photodynamic therapy (PDT)
 - ▶ Histotripsy
 - ▶ Partial prostatectomy

^a Although ultra-hypofractionated EBRT, also known as stereotactic body radiotherapy (SBRT) or stereotactic ablative radiotherapy (SABR), is considered by some as an ablative technology, it does not fall within the general class of ablative therapies. EBRT is recommended in multiple contexts for the treatment of prostate cancer, and when delivered to the prostate it is recommended to be delivered to the whole prostate gland (discussed within [Principles of Radiation Therapy \(PROS-J\)](#)). However, when delivered as FT it should be viewed within the context of other ablative therapies at this time given similar quality of evidence and risks of residual cancer and retreatment.

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF FOCAL/SUBTOTAL THERAPY OR WHOLE GLAND ABLATIVE THERAPY

- Patients should be counseled prior to receiving FT or whole gland ablation therapy as follows:
 - ▶ HIFU, TULSA, FLA, RFA, IRE, PDT, and histotripsy are not FDA approved or cleared for the treatment of prostate cancer.
 - ▶ Currently, no treatment delivered as FT, including radiotherapy or partial prostatectomy, have demonstrated superiority or non-inferiority to guideline-recommended therapies in randomized trials with long-term follow-up. The importance of long-term follow-up is of increased importance in this setting given the high rates of secondary therapy after FTs and potential for increased toxicity.
 - ▶ For patients with low- and intermediate-risk disease¹⁻⁷:
 - ◊ Biopsy-detected (any) cancer after FT is common within 2 years post-treatment (20%–70%).
 - ◊ Erectile dysfunction (10%–40%) and urinary incontinence (0%–20%) have been reported to occur of variable severity within 2 years post-treatment.
 - ▶ FT or whole gland ablation therapy may impact the safety and/or efficacy of subsequent local therapy.
 - ▶ Moderate toxicity (grade 2) is common, and severe toxicity (grade 3 or higher) is possible with reported rates in trials from approximately 1%–14% in the first 2 years post-FT.¹⁻⁷
 - ▶ Currently, there is no evidence-based, universally agreed upon follow-up protocol after FT. Patients often undergo serial imaging and prostate biopsies.

¹ Ehdaie B, Tempany CM, Holland F, et al. MRI-guided focused ultrasound focal therapy for patients with intermediate-risk prostate cancer: a phase 2b, multicentre study. *Lancet Oncol* 2022;23:910-918.

² Duwe G, Boehm K, Haack M, et al. Single-center, prospective phase 2 trial of high-intensity focused ultrasound (HIFU) in patients with unilateral localized prostate cancer: good functional results but ontologically not as safe as expected. *World J Urol* 2023;41:1293-1299.

³ Ganzer R, Hadaschik B, Pahernik S, et al. Prospective multicenter phase II study on focal therapy (hemiblation) of the prostate with high intensity focused ultrasound. *J Urol* 2018;199:983-989.

⁴ Ghai S, Finelli A, Corr K, et al. MRI-guided focused ultrasound focal therapy for intermediate-risk prostate cancer: final results from a 2-year phase II clinical trial. *Radiol* 2024;310:e231473.

⁵ Klotz L, Pavlovich CP, Chin J, et al. Magnetic resonance imaging-guided transurethral ultrasound ablation of prostate cancer. *J Urol* 2021;205:769-779.

⁶ Mehralivand S, George AK, Hoang AN, et al. MRI-guided focal laser ablation of prostate cancer: a prospective single-arm, single-center trial with 3 years follow-up. *Diagn Interv Radiol* 2021;27:394-400.

⁷ Eggener SE, Yousuf A, Watson S, et al. Phase II evaluation of magnetic resonance imaging guided focal laser ablation of prostate cancer. *J Urol* 2016;196:1670-1675.

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF FOCAL/SUBTOTAL THERAPY OR WHOLE GLAND ABLATIVE THERAPY

Management of Recurrence After Focal Ablative or Whole Gland Ablative Therapy

- Presently, the optimal management of residual or recurrent disease after receipt of these treatments is unknown and limited evidence exists. A multi-disciplinary discussion with experienced prostate cancer specialists is recommended.
- If there is suspicion of residual or recurrent prostate cancer, a biopsy should be performed to confirm the presence of residual or recurrent prostate cancer.
 - ▶ Histologic grading of the cancer is recommended if located in previously untreated areas of the prostate. Reliability of grading within a previously treated area of the prostate is unknown at this time, and patients should be informed of this uncertainty.
- Staging imaging should be performed based on the patient's NCCN risk group at time of recurrence (see [PROS-2](#)). However, PSA groupings as used for newly diagnosed prostate cancer may underestimate the burden of disease after prior FT or ablative therapy.
 - ▶ Interpretation of imaging (ie, assessment of cT3 disease by MRI near ablation zone) after FT or whole gland ablative therapy may be less reliable.⁸
 - ▶ Advanced risk stratification tools (eg, gene expression or digital pathology-based biomarkers) have not been adequately studied for use in the context of focal or whole gland ablative therapy and are not recommended to guide treatment decisions in this setting.
- Patients with a clinically significant recurrence after initial focal therapy should be offered radical prostatectomy or prostate RT if appropriate based on life expectancy.
- Patients should be counseled as follows:
 - ▶ There is limited evidence on how to optimally manage a FT or whole gland ablative therapy recurrence.
 - ▶ Patients are at an increased risk of toxicity from subsequent local therapy compared to the primary treatment setting.
 - ▶ It is not known if radical therapy has equivalent long-term success after FT or whole gland ablative therapy given lack of trials with long-term follow-up in this setting.
 - ▶ Secondary treatment should be reserved for patients with appropriate life expectancy and Grade Group 2 and higher recurrences detected on biopsy.

Local Recurrence After Radiation Therapy

- Patients with biopsy-proven recurrence in the prostate after prior RT and without distant metastatic disease can be considered for local secondary therapy.
- However, clinicians and patients should be aware that no local therapy currently has level 1 evidence for the management of a local recurrence after definitive RT.
- Treatments with higher quality of evidence are recommended. These are discussed in the Principles of Local Secondary Therapy Post-Radiation ([PROS-L](#)) and include select forms of FT or whole gland ablative therapy.

⁸ Lai AL, Velaga J, Tay KJ, et al. Multiparametric MRI before and after focal therapy for prostate cancer: pearls and pitfalls for the reporting radiologist. Radiol;7:e240269.



PRINCIPLES OF RADIATION THERAPY

All patients diagnosed with intermediate-, high-, very-high-, regional-risk, low-volume mCSPC, biochemically recurrent, low-volume metastatic castration-sensitive prostate cancer (mCSPC) and oligometastatic mCRPC are recommended to be evaluated by a radiation oncologist as part of a multidisciplinary discussion of treatment options.

General

• EBRT

- ▶ **Treatment Planning:** Intensity modulated RT (IMRT) is recommended over 3-dimensional conformal RT to improve dose conformity.
- ▶ **Image Guidance:** Methods to improve accuracy of treatment delivery, thereby enabling reduction in planning target volume (PTV) margins, which generally result in reduced toxicity are encouraged and may include one or more of the following:
 - ◊ Image guidance with daily 3D imaging is recommended with either a cone beam CT (CBCT) or MRI.
 - ◊ Devices that assist with optimal image guidance (fiducial markers) or reduce motion (endorectal balloons).
 - ◊ Real-time intrafraction volumetric tracking.
 - ◊ Online adaptive radiotherapy if target is near mobile organs (ie, performing a simultaneous integrated boost to a positive pelvic lymph node adjacent to bowel).
- ▶ **Beam Type:** Photon and proton RT are acceptable forms of EBRT and appear to have similar outcomes in regard to toxicity, QOL, and tumor control. Potential financial toxicity should be discussed with patients.
- ▶ **Fractionation:** An extensive list of fractionation schedules have been studied. These are grouped into three categories, conventional fractionation (1.8–2 Gy/fraction), moderate hypofractionation (>2.5 to 4 Gy/fraction), and ultra-hypofractionation (>6 Gy/fraction).¹ Common dose/fraction schedules are shown in Table 1. In general, iso-effective moderate hypofractionation dosing has demonstrated noninferior tumor control, toxicity, and QOL over conventional fractionation and is more convenient for patients and thus preferred. Similarly, ultra-hypofractionation has been demonstrated to be noninferior to both moderate hypofractionation and conventional fractionation.² Thus, use of conventionally fractionated radiotherapy is no longer preferred for localized prostate cancer.
 - ◊ Ultra-hypofractionation encompasses stereotactic body radiation therapy (SBRT), which requires precision treatment setup and required additional image guidance techniques.
 - ◊ SBRT is recommended and preferred specifically when:
 - Performing metastasis-directed radiotherapy (MDRT) (see [Principles of MDT \[PROS-M\]](#))
 - There is limited progression (eg, oligoprogression) or limited residual disease and the patient is on otherwise effective systemic therapy (eg, consolidation).
 - The lesion occurs in or immediately adjacent to a previously irradiated treatment field.
 - At physician discretion for more durable control of pain than achieved with typical palliative regimens.

Note: All recommendations are category 2A unless otherwise indicated.

[References on PROS-J 10 of 11](#)



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF RADIATION THERAPY

- **Brachytherapy**

- ▶ **Patient Selection:** Providers should consider aspects of gland size, baseline urinary symptoms, and prior procedures (ie, transurethral resection of prostate) that may increase risk of adverse effects. Neoadjuvant ADT to shrink a gland to allow treatment should be considered in context of the balance between its additional toxicity with this benefit.
- ▶ **Dose Rate:** Both low-dose rate (LDR) and high-dose rate (HDR) are acceptable to be used as either monotherapy or as a boost combined with EBRT in higher risk disease. Data indicates that HDR may have lower toxicity than LDR, and while efficacy appears similar between HDR and LDR when used as a boost combined with EBRT, it remains unknown if HDR as monotherapy provides equivalent tumor control to LDR given lack of robust randomized comparisons. Post-implant dosimetry must be performed for LDR implants to verify dosimetry.
- ▶ **Monotherapy:** Brachytherapy monotherapy has been shown to have fewer side effects than when used with EBRT without compromising tumor control for intermediate-risk disease.³
- ▶ **Combination Therapy:** EBRT with an LDR brachytherapy boost has been shown to improve biochemical control over conventionally fractionated EBRT with ADT (12 months) without a focal/micro EBRT boost.⁴ This was at the expense of an increase in high-grade toxicity.⁵ Thus, careful patient selection and contemporary planning, use of recognized organ at risk (OAR) dose constraints, and use of high-quality ultrasound and other imaging are essential.

- **Elective Nodal Irradiation (ENI)**

- ▶ ENI can be considered in select patients.
- ▶ Use of IMRT with image guidance is recommended when performing ENI. Advanced techniques that include online adaptive radiotherapy may be considered to reduce ENI planning target volume margins.
- ▶ If performing ENI, it is recommended to include the common, internal, and external iliac, pre-sacral, and obturator lymph node stations. Peri-rectal/mesorectal lymph nodes may be included if there is evidence of lymph node involvement in this nodal station. Para-aortic lymph nodes should not be included in patients who do not have evidence of para-aortic lymph node involvement.

Definitive RT

- **Biocompatible and biodegradable perirectal spacers:**

- ▶ These devices may be implanted between the prostate and rectum in patients undergoing radiation therapy to the prostate in order to displace the rectum from high radiation dose regions for the purpose of toxicity reduction. Patients with grossly apparent true posterior extraprostatic extension should not undergo perirectal spacer implantation. NCCN high-risk disease without posterior extraprostatic extension is not a contraindication to spacer use.
- ▶ These devices should be placed by practitioners who can maintain a moderate to high volume of use to ensure quality.
- ▶ Additional caution should be used in the re-irradiation setting due to potential changes in tissue planes.

Note: All recommendations are category 2A unless otherwise indicated.

[References on PROS-J 10 of 11](#)



PRINCIPLES OF RADIATION THERAPY

Definitive RT (continued)

• Low Risk

- ▶ Patients with NCCN low-risk prostate cancer are encouraged to pursue active surveillance.⁶
- ▶ Those electing treatment with RT may receive EBRT or brachytherapy as monotherapy.
- ▶ The target should include the prostate and consideration of treatment of seminal vesicles (proximal).
- ▶ Patients should not be treated with any of the following:
 - ◊ ENI
 - ◊ ADT
 - ◊ Combination brachytherapy boost with EBRT

• Favorable Intermediate Risk

- ▶ The target should include the prostate and consideration of treatment of seminal vesicles (proximal or full).
- ▶ Those electing treatment with RT may receive EBRT or brachytherapy as monotherapy.
- ▶ Patients should not be treated with any of the following:
 - ◊ ENI
 - ◊ Combination brachytherapy boost with EBRT³
- ▶ ADT is generally not recommended, but can be considered if additional risk assessments suggest aggressive tumor behavior or increased potential benefit from short-term ADT.^{7,8}

• Unfavorable Intermediate Risk

- ▶ The target should include the prostate and the seminal vesicles (proximal or full).
- ▶ RT options include either EBRT, brachytherapy monotherapy, or EBRT with a brachytherapy boost.
- ▶ Multiple randomized trials that included patients with unfavorable intermediate risk who did not demonstrate superiority of combination therapy over brachytherapy monotherapy.³
- ▶ Focal boosting with isotoxic delivery can be considered if there is sufficient provider and practice expertise to delineate the dominant intraprostatic lesion (DIL) and OARs on MRI.⁹
- ▶ ENI should not be used routinely.
- ▶ ADT (level 1 data for ST-ADT) should be used unless additional risk assessments suggest less aggressive tumor behavior, or less predicted benefit, or if medically contraindicated.^{8,10} Radiotherapy dose or modality has not clearly demonstrated the ability to obviate the benefit of ADT. ST-ADT is recommended to be used as a concurrent and adjuvant approach with RT rather than a neoadjuvant and concurrent approach.¹¹

Note: All recommendations are category 2A unless otherwise indicated.

[References on PROS-J 10 of 11](#)



PRINCIPLES OF RADIATION THERAPY

Definitive RT (continued)

• High Risk

- ▶ The target should include the prostate and the seminal vesicles (full).
- ▶ RT options include EBRT or brachytherapy boost combined with EBRT. Carefully selected patients may receive brachytherapy monotherapy.
- ▶ Focal boosting with isotoxic delivery can be considered if there is sufficient provider and practice expertise to delineate the DIL and OARs on MRI.⁹
- ▶ Clinical trials have demonstrated varied results regarding the benefit of ENI. Multiple randomized trials have not demonstrated improvement in outcomes from ENI. However, the benefit of ENI was shown in one trial of patients with high-risk prostate cancer staged primarily with PSMA-PET/CT and who had a risk of nodal involvement >20%. Currently, the use of ENI is at the discretion of the treating physician.
- ▶ ADT (level 1 data for LT-ADT 12–36 months) is recommended for patients with life expectancy >5 years or who are symptomatic unless medically contraindicated. Radiotherapy dose or modality has not clearly demonstrated the ability to obviate the benefit of LT-ADT.¹⁰

• Very-High-Risk

- ▶ The target should include the prostate and the seminal vesicles (full).
- ▶ RT options include EBRT. Carefully selected patients may receive EBRT with a brachytherapy boost.
- ▶ Focal boosting with isotoxic delivery can be considered if there is sufficient provider and practice expertise to delineate the DIL and OARs on MRI.
- ▶ Combination brachytherapy boost with EBRT should not be used routinely. The primary randomized evidence to recommend combination brachytherapy boost with EBRT comes from the ASCENDE-RT trial, which excluded patients with PSA >40 ng/mL and T3b.¹²
- ▶ ADT (level 1 data for LT-ADT 18–36 months) is recommended for patients with life expectancy >5 years or who are symptomatic unless medically contraindicated.
- ▶ Currently, the use of ENI is at the discretion of the treating physician.
- ▶ Addition of abiraterone should be used very selectively as the benefit in contemporary practice with modern staging is uncertain (see [PROS-6](#) and [PROS-G 1 of 5](#)).¹³ It is not recommended to be used routinely for patients with solely MRI defined T3a disease with low volume Gleason 8 disease. Studies validating the post-hoc subset analysis of STAMPEDE for the benefit of abiraterone in very high-risk prostate cancer using other ARPIs have not yet reported, and thus the benefit of using ARPIs with RT+LT-ADT remains uncertain in contemporary patients with very high risk N0M0 disease. Radiotherapy dose or modality has not clearly demonstrated the ability to obviate the benefit of LT-ADT.¹⁰

• Regional Disease

- ▶ EBRT is recommended to include the prostate, seminal vesicles, and pelvic lymph nodes.
- ▶ Simultaneous integrated boost to involved lymph nodes is recommended while respective adjacent OAR dose constraints.
- ▶ Use of a brachytherapy boost is not recommended in these patients.
- ▶ Addition of abiraterone is recommended (see [PROS-7](#) and [PROS-G \[1 of 5\]](#)).¹³

Note: All recommendations are category 2A unless otherwise indicated.

[References on PROS-J 10 of 11](#)



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF RADIATION THERAPY

Postoperative RT:

- **Adjuvant RT:** Adjuvant RT (aRT) involves the use of EBRT post-radical prostatectomy for patients with adverse pathologic features (ie, pT3 and/or positive surgical margins) in the setting of an undetectable postoperative PSA. This may result in some patients unnecessarily receiving aRT and experience potential additional side effects.
 - ▶ The early results and aggregate meta-analysis of three trials comparing aRT to early secondary RT (sRT) did not demonstrate a difference in biochemical recurrence for patients with generally a favorable prognosis. Thus, waiting for a detectable PSA and using an early sRT approach is recommended for most patients who undergo a RP rather than an immediate aRT approach to avoid over treatment and subsequent potential side effects.¹⁴
 - ▶ Use of aRT may be reasonable for patients with multiple adverse features and/or lymph node involvement.
 - ◊ Patients with multiple adverse features and/or lymph node involvement were underrepresented in these trials, and in the RADICALS-RT trial, the largest of the trials comparing aRT to early sRT, there was a significant reduction in prostate cancer-specific mortality with aRT and numerical trends in superior freedom-from-distant metastasis with aRT.¹⁵ This raises uncertainty to the true noninferiority of early sRT in these patients with high risk of recurrence post-RP.
 - ▶ Multiple dose/fraction schedules have been used with aRT that include both conventionally fractionated (60–64 Gy in 30–36 fractions) and moderately hypofractionated regimens (52.5 Gy in 20 fractions).¹⁶
- **Secondary RT** is the use of EBRT post-RP when a patient has a detectable PSA; a lower limit is not defined and may be below 0.2 ng/mL.
 - ▶ Early sRT at a PSA of 0.1–0.2 ng/mL is recommended for most patients who experience BCR post-RP. The exception to this is in patients who have delayed recurrence many years post-RP with very slow PSA doubling times (ie, >24 months), especially when a patient has limited life expectancy.
 - ▶ Unlike in localized prostate cancer, the use of ADT in the post-RP setting has demonstrated survival improvements in only a small subset of patients, primarily in those with high pre-sRT PSAs receiving late sRT (PSA >0.7 ng/mL).¹⁷ For patients receiving early sRT there is no clear OS benefit from the addition of ADT in any of the four published randomized trials, irrespective of ADT duration, or on meta-analysis.¹⁸ ADT does significantly delay the incidence of biochemical recurrence.
 - ▶ Use of ADT with sRT should be personalized based on pre-RT PSA, clinicopathologic risk factors, patient age, life expectancy, comorbidities, and preferences, as well as advanced risk stratification tools (see [PROS-H](#)).^{18,19}
 - ◊ Generally, ST-ADT is recommended to balance toxicity, QOL, and tumor control. However, LT-ADT may be preferred for patients receiving late sRT and/or those with multiple adverse pathologic features and/or lymph node involvement.
 - ◊ Treatment intensification with an ARPI has not demonstrated an improvement in metastasis-free survival or overall survival in this population in unselected patients.^{20,21}
 - ▶ Multiple dose/fraction schedules have been used with sRT that include both conventionally fractionated (64–70.2 Gy in 32–39 fractions) and moderately hypofractionated regimens (52.5 Gy in 20 fractions or 62.5 Gy in 25 fractions).^{16,22} The moderately hypofractionated regimens have demonstrated noninferior tumor control, toxicity, and QOL. Multiple single arm trials and early follow-up of phase II randomized trials have been reported using ultra-hypofractionated RT in the sRT setting, and generally are using 32.5–34 Gy in 5 fractions.^{23,24} These studies often leverage either intrafraction motion monitoring, online adaptive radiotherapy, and strict image guidance with substantially modified contouring of target volumes to limit dose to OARs.
 - ▶ The addition of ENI to sRT plus ST-ADT has demonstrated improvements in biochemical control.²⁵ It appears the benefit is greater in patients with higher pre-sRT PSAs, as well as in patients who did not have their pelvic lymph nodes assessed, or minimally assessed, via a pelvic lymph node dissection. Use of PSMA PET/CT has not been shown to be sensitive enough to rule out the presence of pelvic lymph node micrometastases.

Note: All recommendations are category 2A unless otherwise indicated.

[References on PROS-J 10 of 11](#)

PROS-J
5 OF 11



PRINCIPLES OF RADIATION THERAPY

RT in Advanced Disease:

• Synchronous mCSPC

▶ Treatment of the primary tumor:

- ◊ Minimizing toxicity is paramount when delivering RT to the primary tumor in patients with metastatic disease. As such, it is unclear if dose escalation (ie, brachytherapy or focal boost) improves outcomes.
- ◊ Low Volume: Treatment of the primary is recommended with EBRT.^{26,27}
 - Volume of disease should be defined according to CT and bone scan, if performed.
 - PET imaging should not be used to exclude a patient from treatment of the primary tumor.
 - The strongest data for a benefit of adding RT to the primary tumor are in patients receiving either ADT alone, ADT+ docetaxel, or ADT + abiraterone for those with ≤5 bony metastases.
- ◊ High Volume: Treatment of the primary tumor with EBRT can be considered for select patients with high-volume mCSPC (category 2B).²⁷
 - RT to the primary tumor in high volume mCSPC has demonstrated improvements in time to mCRPC and a reduction in the development of severe genitourinary adverse events when given in the context of ADT with variable use of docetaxel and/or abiraterone.

▶ Treatment of Metastases

- ◊ MDT, which typically uses SBRT, may be used for patients who have a limited burden of lymph node or osseous disease. See [Principles of MDT \(PROS-M\)](#).

• Metachronous mCSPC

- ▶ A greater level of evidence supports the role of MDT in the metachronous setting as compared to synchronous mCSPC. This commonly is delivered using SBRT, but moderately hypofractionated RT may be considered.
- ▶ Use of MDT in this setting has been shown to delay the need for ADT compared to observation,²⁸ the addition of RT to ADT has been shown to prolong a treatment free eugonadal interval as compared to ADT alone,²⁹ and the addition of ADT to RT has improved progression-free survival (PFS).³⁰
- ▶ See [Principles of MDT \(PROS-M\)](#).
- ▶ mCRPC
 - ◊ The use of MDT has been shown to prolong PFS and rPFS when used for patients with oligometastatic CRPC treated with ADT+ARPI as compared to systemic therapy alone. MDT may be considered when the goal is to prolong PFS and extend duration of systemic therapy prior to needing to switch therapies.³¹ See [Principles of MDT \(PROS-M\)](#).

Note: All recommendations are category 2A unless otherwise indicated.

[References on PROS-J 10 of 11](#)



PRINCIPLES OF RADIATION THERAPY

Radiopharmaceutical Therapy

- Radiopharmaceutical therapies for prostate cancer are suitable options for improving survival and/or rPFS in select patients with advanced castration-resistant disease. Due to prior therapy exposure, specific targets, and hematologic effects of these therapies, careful selection and sequencing strategy with other therapies is important. This section discusses the two currently FDA-approved agents in use (Ra-223, Lu-177–PSMA-617).
- Radium-223
 - ▶ Radium-223 has been shown to extend survival in patients who have CRPC with symptomatic bone metastases, but no visceral metastases.³² Radium-223 alone has not been shown to extend survival in patients with visceral metastases or bulky nodal disease (>3 to 4 cm).
 - ▶ Radium-223 is administered IV once a month for 6 months by an appropriately licensed facility, usually in nuclear medicine or radiation oncology facilities.
 - ▶ Concurrent use with enzalutamide and bone health agents may be considered given the overall survival benefit over enzalutamide alone.³³ Concurrent use with other systemic therapies should be pursued with caution.
 - ▶ Bone fracture risk: Radium-223 may increase fracture risk when given concomitantly with an ARPI without use of a bone health agent. Concomitant use of denosumab or zoledronic acid is recommended.³⁴
- Lu-177–PSMA-617¹⁶
 - ▶ In patients with PSMA-positive disease, Lu-177–PSMA-617 has been shown to improve OS in patients with progressive mCRPC previously treated with androgen receptor inhibitors and taxane chemotherapy.³⁵ It has also been shown to improve rPFS in patients who have not received taxanes with PSMA-positive mCRPC who were previously treated with an androgen receptor inhibitor compared with changing to a different androgen receptor inhibitor.
 - ▶ This agent is approved for patients with mCRPC who have been treated with ARPI therapy and have received prior taxane-based chemotherapy or are considered appropriate to delay receipt of chemotherapy.³⁶
 - ▶ Lu-177–PSMA-617 is not recommended in patients with dominant PSMA-negative lesions. PSMA-negative lesions are defined as metastatic disease that lack PSMA uptake including bone with soft tissue components ≥ 1.0 cm, lymph nodes ≥ 2.5 cm in short axis, and solid organ metastases ≥ 1.0 cm in size.
 - ▶ Use of PSMA PET/CT SUVmean and use of an FDG PET/CT scan may aid in identifying patients with lower PSMA expression and non-PSMA avid disease who will derive less or potentially no benefit from 177-Lu-PSMA-617.
 - ▶ Lu-177–PSMA-617 is typically administered IV 200 mCi (7.4 GBq) every 6 weeks for up to 6 treatments by an appropriately licensed facility.
 - ▶ Use of post-treatment SPECT or PET/CT imaging can be considered to monitor PSMA expression for adaptive dosing.
 - ▶ Because Lu-177 also emits gamma radiation, appropriate precautions should be taken to minimize exposure to personnel administering the radiopharmaceutical. Treatment rooms should be monitored for potential contamination following treatments, and patients should be provided written instructions regarding radiation safety precautions following treatment.
 - ▶ For information regarding the use of PSMA-PET to define PSMA-positive disease, see [Principles of Imaging \(PROS-E\)](#).

Note: All recommendations are category 2A unless otherwise indicated.

[References on PROS-J 10 of 11](#)

PROS-J
7 OF 11



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF RADIATION THERAPY

Table 1: Below are examples of regimens that have shown acceptable efficacy and toxicity. The optimal regimen for an individual patient warrants evaluation of comorbid conditions, voiding symptoms, and toxicity of therapy. Additional fractionation schemes may be used as long as sound oncologic principles and appropriate estimate of BED are considered.

See treatment pages and [Principles of ADT \(PROS-G\)](#) for other recommendations, including recommendations for neoadjuvant/concomitant/adjuvant ADT.

EBRT Regimen	Preferred Dose/Fraction	Definitive RT						Post-Treatment RT			Advanced Disease	
		Low	FIR	UIR	High	Very-High	Regional	Post-RP		Post-RT	Primary Tumor	Metastases
								aRT	sRT		mCSPC M0 CRPC mCRPC	MDT
Conventional	1.8–2 Gy x 37–45 fx			☼	☼	✓	✓				☼	
	1.8–2 Gy x 30–39 fx							✓	✓		☼	
Moderate Hypofractionation	3 Gy x 20 fx (preferred) ^a 2.7 Gy x 26 fx 2.5 Gy x 28 fx	☼	✓	✓	✓	✓	✓			☼	☼	☼
	2.63–2.75 Gy x 20 fx 2.5 Gy x 25 fx							✓	✓	☼	✓	☼
Ultra Hypofractionation (SBRT)	9.5 Gy x 4 fx 7.25–8 Gy x 5 fx 6 Gy x 6 fx 6.1 Gy x 7 fx	☼	✓	✓	✓	☼	☼		☼	✓	✓	✓
	9–10 Gy x 3 fx 12 Gy x 2 fx 16–24 Gy x 1 fx											✓
	6.2–6.4 Gy x 5 fx								☼			
EBRT Boost Techniques												
EBRT with simultaneous integrated boost	See footnote b.		☼	✓	✓	☼	☼		☼	☼	☼	
EBRT with sequential SBRT boost	<i>Prostate:</i> 1.8 Gy x 23–28 fx <i>Boost:</i> 6 Gy x 3 fx 9.5 Gy x 2 fx			☼	☼	☼						

(✓ Preferred; ☼ Acceptable based on clinical and medical need; Regimens shaded gray are not recommended)

Note: All recommendations are category 2A unless otherwise indicated.

[Footnotes on PROS-J 8A of 11](#)



PRINCIPLES OF RADIATION THERAPY FOOTNOTES

- ^a Based on the HYDRA individual patient meta-analysis of seven phase III randomized trials, isodose moderately hypofractionated radiotherapy (MHFRT) and dose-escalated MHFRT both have similar efficacy compared with conventionally fractionated radiotherapy (CFRT), but dose-escalated MHFRT was associated with higher toxicity. Isodose regimens (eg, 60 Gy in 20 fractions) are preferred for MHFRT for localized prostate cancer. Alternative regimens (eg, 70 Gy in 28 fractions) may be used, but patients should be counseled on potentially increased risk of toxicity.¹
- ^b EBRT to whole prostate 2.2 Gy x 35 fx plus micro-boost to MRI-dominant lesion to ≤95 Gy (fractions ≤2.7 Gy).⁹ The micro-boost technique with level 1 data was established for a modestly hypofractionated regimen but has been extrapolated reasonably to other regimens in ongoing clinical trials. Care must be taken in doing so outside of clinical trials in order to respect normal tissue toxicity risk and above all in prioritizing normal organ tolerances over micro-boost coverage.

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF RADIATION THERAPY

Table 1: Below are examples of regimens that have shown acceptable efficacy and toxicity. The optimal regimen for an individual patient warrants evaluation of comorbid conditions, voiding symptoms, and toxicity of therapy. Additional fractionation schemes may be used as long as sound oncologic principles and appropriate estimate of BED are considered.

See treatment pages and [Principles of ADT \(PROS-G\)](#) for other recommendations, including recommendations for neoadjuvant/concomitant/adjunct ADT.

Brachytherapy	Preferred Dose/Fraction	Definitive RT						Post-Treatment RT			Advanced Disease	
		Low	FIR	UIR	High	Very-High	Regional	Post-RP		Post-RT	Primary	Metastases
								aRT	sRT	sRT	mCSPC M0 CRPC mCRPC	MDT
Monotherapy												
Iodine 125	145 Gy	☀	✓	✓	☀					✓		
Palladium 103	125 Gy											
Cesium 131	115 Gy											
Iridium 192	13.5 Gy x 2 implants 9.5 Gy BID x 2 implants											
EBRT + Brachytherapy Boost (combined with 45-50.4 Gy in 25-28 fx or 37.5 Gy in 15 fx)												
Iodine 125	110–115 Gy			☀	✓	☀						
Palladium 103	90–100 Gy											
Cesium 131	85 Gy											
Iridium 192	15 Gy x 1 fx 10.75 Gy x 2 fx											

(✓ Preferred; ☼ Acceptable based on clinical and medical need; Regimens shaded gray are not recommended)

Note: All recommendations are category 2A unless otherwise indicated.

[References on PROS-J 10 of 11](#)



PRINCIPLES OF RADIATION THERAPY REFERENCES

- ¹ Kishan AU, Sun Y, Tree AC, et al. Hypofractionated radiotherapy for prostate cancer (HYDRA): an individual patient data meta-analysis of randomised trials in the MARCAP consortium. *Lancet Oncol* 2025;26:459-469.
- ² van As N, Griffin C, Tree A, et al. A Phase III randomized controlled trial of Stereotactic Body Radiotherapy in localized prostate cancer. *N Engl J Med* 2024;391:1413-1425.
- ³ Michalski JM, Winter KA, Prestidge BR, et al. Effect of brachytherapy with external beam radiation therapy versus brachytherapy alone for intermediate-risk prostate cancer: NRG oncology RTOG 0232 randomized clinical trial. *J Clin Oncol* 2023;41:4035-4044.
- ⁴ Oh J, Tyldesley S, Pai H, et al. An updated analysis of the survival endpoints of ASCENDE-RT. *Int J Rad Oncol* 2023;115:1061-1070.
- ⁵ Rodda S, Tyldesley S, Morris WJ, et al. ASCENDE-RT: an analysis of treatment-related morbidity for a randomized trial comparing a low-dose-rate brachytherapy boost with a dose-escalated external beam boost for high- and intermediate-risk prostate cancer. *Int J Rad Oncol Biol Phys* 2017;98:286-295.
- ⁶ Hamdy FC, Donovan JL, Lane A, et al. Fifteen-year outcomes after monitoring, surgery, or radiotherapy for prostate cancer. *N Engl J Med* 2023;388:1547-1558.
- ⁷ Zumsteg ZS, Spratt DE, Daskivich TJ, et al. Effect of androgen deprivation on long-term outcomes of intermediate-risk prostate cancer stratified as favorable or unfavorable: A secondary analysis of the RTOG 9408 randomized clinical trial. *JAMA Netw Open* 2020;3:e2015083.
- ⁸ Krauss DJ, Karrison T, Martinez AA, et al. Dose-escalated radiotherapy alone or in combination with short-term androgen deprivation for intermediate-risk prostate cancer: results of a phase III multi-institutional trial. *J Clin Oncol* 2023;41:3203-3216.
- ⁹ Kerkmeijer LGW, Groen VH, Pos FJ, et al. Focal boost to the intraprostatic tumor in external beam radiotherapy for patients with localized prostate cancer: results from the FLAME randomized phase III trial. *J Clin Oncol* 2021;39:787-796.
- ¹⁰ Kishan AU, Sun Y, Hartman H, et al. Androgen deprivation therapy use and duration with definitive radiotherapy for localised prostate cancer: an individual patient data meta-analysis. *Lancet Oncol* 2022;23:304-316.
- ¹¹ Spratt DE, Malone S, Roy S, et al. Prostate radiotherapy with adjuvant androgen deprivation therapy (ADT) improves metastasis-free survival compared to neoadjuvant ADT: an individual patient meta-analysis. *J Clin Oncol* 2021;39:136-144.
- ¹² Morris WJ, Tyldesley S, Rodda S, et al. Androgen suppression combined with elective nodal and dose escalated radiation therapy (the ASCENDE-RT trial): an analysis of survival endpoints for a randomized trial comparing a low-dose-rate brachytherapy boost to a dose-escalated external beam boost for high- and intermediate-risk prostate cancer. *Int J Rad Oncol Biol Phys* 2017;98:275-285.
- ¹³ Attard G, Murphy L, Clarke NW, et al. Abiraterone acetate and prednisolone with or without enzalutamide for high-risk non-metastatic prostate cancer: a meta-analysis of primary results from two randomised controlled phase 3 trials of the STAMPEDE platform protocol. *Lancet* 2022;399:447-460.
- ¹⁴ Vale CL, Fisher D, Kneebone A, et al. Adjuvant or early salvage radiotherapy for the treatment of localised and locally advanced prostate cancer: a prospectively planned systematic review and meta-analysis of aggregate data. *Lancet* 2020;396:1422-1431.
- ¹⁵ Parker CC, Petersen PM, Cook AD, et al. Timing of radiotherapy (RT) after radical prostatectomy (RP): long-term outcomes in the RADICALS-RT trial (NCT00541047). *ESMO* 2024;24:656-666.
- ¹⁶ Petersen PM, Cook AD, Sydes MR, et al. Salvage radiation therapy after radical prostatectomy: analysis of toxicity by dose-fractionation in the RADICALS-RT trial. *Int J Rad Oncol Biol Phys* 2023;117:624-629.
- ¹⁷ Dess RT, Sun Y, Jackson WC, et al. Association of presalvage radiotherapy PSA levels after prostatectomy with outcomes of long-term antiandrogen therapy in men with prostate cancer. *JAMA Oncol* 2020;6:735-743.
- ¹⁸ Burdett S, Fisher DJ, Tierney JF, et al. Duration of androgen suppression with postoperative radiotherapy (DADSPORT) for nonmetastatic prostate cancer: a collaborative systematic review and meta-analysis of aggregate data. [online ahead of print] *Eur Urol* 2025;doi:10.1016/j.eururo.2025.05.013.
- ¹⁹ Feng FY, Huang HC, Spratt DE, et al. Validation of a 22-gene genomic classifier in patients with recurrent prostate cancer: an ancillary study of the NRG/RTOG 9601 randomized clinical trial. *JAMA Oncol* 2021;7:544-552.
- ²⁰ Posadas EM, Gay HA, Rodgers JP, et al. Intensification of ADT with enzalutamide in high-risk patients with biochemical relapse following radical prostatectomy undergoing salvage radiation: Initial results from RTOG 3506 (STEEL). *J Clin Oncol* 2024;42:5101.

Note: All recommendations are category 2A unless otherwise indicated.

Continued

**PROS-J
10 OF 11**



PRINCIPLES OF RADIATION THERAPY REFERENCES

- ²¹ Nguyen PL, Kollmeier M, Rathkopf DE, et al. FORMULA-509: A multicenter randomized trial of post-operative salvage radiotherapy (SRT) and 6 months of GnRH agonist with or without abiraterone acetate/prednisone (AAP) and apalutamide (Apa) post-radical prostatectomy (RP). *J Clin Oncol* 2023;41:303.
- ²² Buyyounouski MK, Pugh SL, Chen RC, et al. Noninferiority of hypofractionated vs conventional postprostatectomy radiotherapy for genitourinary and gastrointestinal symptoms: the NRG-GU003 Phase 3 randomized clinical trial. *JAMA Oncol* 2024;10:584-591.
- ²³ Nagar H, Diven MA, Rippon B, et al. A randomized controlled phase 2 trial comparing salvage radiotherapy for prostate cancer delivered in 4 versus 2 weeks (SHORTER): acute genitourinary and gastrointestinal patient-reported outcomes at a single institution. [online ahead of print] *Eur Urol Oncol* 2025;doi:10.1016/j.euo.2025.05.014.
- ²⁴ Ma TM, Ballas LK, Wilhalme H, et al. Quality-of-life outcomes and toxicity profile among patients with localized prostate cancer after radical prostatectomy treated with stereotactic body radiation: the SIMITAR multicenter phase 2 trial. *Int J Rad Oncol Biol Phys* 2023;115:142-152.
- ²⁵ Pollack A, Karrison TG, Balogh AG, et al. The addition of androgen deprivation therapy and pelvic lymph node treatment to prostate bed salvage radiotherapy (NRG Oncology/RTOG 0534 SPPORT): an international, multicentre, randomised phase 3 trial. *Lancet* 2022;399:1886-1901.
- ²⁶ Parker CC, James ND, Brawley CD, et al. Radiotherapy to the prostate for men with metastatic prostate cancer in the UK and Switzerland: Long-term results from the STAMPEDE randomised controlled trial. 2022;19:e1003998.
- ²⁷ Bossi A, Foulon S, Maldonado X, et al. Efficacy and safety of prostate radiotherapy in de novo metastatic castration-sensitive prostate cancer (PEACE-1): a multicentre, open-label, randomised, phase 3 study with a 2 × 2 factorial design. *Lancet* 2024;404:2065-2076.
- ²⁸ Deek MP, Van der Eecken K, Suter P, et al. Long-term outcomes and genetic predictors of response to metastasis-directed therapy versus observation in oligometastatic prostate cancer: analysis of STOMP and ORIOLE trials. *J Clin Oncol* 2022;40:3377-3382.
- ²⁹ Tang C, Sherry AD, Haymaker C, et al. Addition of metastasis-directed therapy to intermittent hormone therapy for oligometastatic prostate cancer: the EXTEND phase 2 randomized clinical trial. *JAMA Oncol* 2023;9:825-834.
- ³⁰ Marvaso G, Corrao G, Zaffaroni M, et al. ADT with SBRT versus SBRT alone for hormone-sensitive oligorecurrent prostate cancer (RADIOA): a randomised, open-label, phase 2 clinical trial. *Lancet Oncol* 2025;26:300-311.
- ³¹ Francolini G, Allegra AG, Detti B, et al. Stereotactic body radiation therapy and abiraterone acetate for patients affected by oligometastatic castrate-resistant prostate cancer: a randomized phase II trial (ARTO). *J Clin Oncol* 2023;41:5561-5568.
- ³² Parker C, Nilsson S, Heinrich D, et al. Alpha emitter radium-223 and survival in metastatic prostate cancer. *NEJM* 2013;369:213-223.
- ³³ Tombal B, Choudhury A, Saad F, et al. Enzalutamide plus radium-223 in metastatic castration-resistant prostate cancer: results of the EORTC 1333/PEACE-3 trial. [online ahead of print] *ESMO* 2025;doi: <https://doi.org/10.1016/j.annonc.2025.05.011>
- ³⁴ Smith M, Parker C, Saad F, et al. Addition of radium-223 to abiraterone acetate and prednisone or prednisolone in patients with castration-resistant prostate cancer and bone metastases (ERA 223): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet Oncol* 2019;20:408-419.
- ³⁵ Sartor O, de Bono J, Chi KN, et al. Lutetium-177-PSMA-617 for metastatic castration-resistant prostate cancer. *N Engl J Med* 2021;385:1091-1103.
- ³⁶ Morris MJ, Castellano D, Herrmann K, et al. 177Lu-PSMA-617 versus a change of androgen receptor pathway inhibitor therapy for taxane-naïve patients with progressive metastatic castration-resistant prostate cancer (PSMAfore): a phase 3, randomised, controlled trial. *Lancet* 2024;404:1227-1239.

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF SURGERY

Pelvic Lymph Node Dissection

- For patients undergoing RP:
 - ▶ Pelvic lymph node dissection (PLND) can be considered in patients with favorable intermediate-risk prostate cancer.
 - ▶ PLND is recommended in patients with unfavorable intermediate, high, very-high-risk, and regional prostate cancer.
 - ▶ A PLND can be excluded in patients with low predicated probability of nodal metastases by nomograms, although some patients with lymph node metastases will be missed. There is no single evidence-based threshold for performing PLND. Based on the risk of complications with PLND and extra time to perform the procedure, the published thresholds range from 2% to 7%.¹⁻⁴
 - ▶ A patient who is above the threshold for performing a PLND, but has a negative PSMA PET scan should still undergo PLND. In two studies, the sensitivity of PSMA PET for pelvic lymph node involvement among patients undergoing RP and PLND was low (about 40%), and the negative predictive value was about 81%.^{5,6} Thus, basing the decision to perform PLND on a negative PSMA PET scan could result in missing 19% of patients with positive lymph nodes.
 - ▶ PLND can be performed using an open, laparoscopic, or robotic technique and can provide staging and prognostic information.
- Extended PLND provides more complete staging and may cure some patients with microscopic metastases; therefore, an extended PLND is recommended when PLND is performed.
- An extended PLND includes removal of all node-bearing tissue from an area bound by the external iliac vein anteriorly, the pelvic sidewall laterally, the bladder wall medially, the floor of the pelvis posteriorly, Cooper's ligament distally, and the internal iliac artery proximally.

Radical Prostatectomy

- RP is an appropriate therapy for any patient not on an active surveillance program with clinically localized prostate cancer that can be completely excised surgically, who has a life expectancy of ≥ 10 years, and who has no serious comorbid conditions that would contraindicate an elective operation.
- High-volume surgeons in high-volume centers generally provide better outcomes.
- Blood loss can be substantial with RP, but can be reduced by using laparoscopic or robotic assistance or by careful control of the dorsal vein complex and periprostatic vessels when performed as open surgery.
- Urinary incontinence can be reduced by preservation of urethral length beyond the apex of the prostate and avoiding damage to the distal sphincter mechanism. Bladder neck preservation may decrease the risk of incontinence. Anastomotic strictures increase the risk of long-term incontinence.
- Recovery of erectile function is directly related to age at RP, preoperative erectile function, and the degree of preservation of the cavernous nerves. Replacement of resected nerves with nerve grafts has not been shown to be beneficial. Early restoration of erections may improve late recovery.

Secondary Radical Prostatectomy

- Secondary RP is an option for highly selected patients with local recurrence after external beam RT (EBRT), brachytherapy, or cryotherapy in the absence of metastases, but the morbidity (ie, incontinence, loss of erection, anastomotic stricture) is high and the operation should be performed by surgeons who are experienced with secondary RP.

Note: All recommendations are category 2A unless otherwise indicated.

[References on PROS-K 2 of 2](#)

PROS-K
1 OF 2



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF SURGERY REFERENCES

- ¹ Cagiannos I, Karakiewicz P, Eastham JA, et al. A preoperative nomogram identifying decreased risk of positive pelvic lymph nodes in patients with prostate cancer. *J Urol* 2003;170:1798-1803.
- ² Briganti A, Larcher A, Abdollah F, et al. Updated nomogram predicting lymph node invasion in patients with prostate cancer undergoing extended pelvic lymph node dissection: the essential importance of percentage of positive cores. *Eur Urol* 2012;61:480-487.
- ³ Gandaglia G, Ploussard G, Valerio M, et al. A novel nomogram to identify candidates for extended pelvic lymph node dissection among patients with clinically localized prostate cancer diagnosed with magnetic resonance imaging-targeted and systematic biopsies. *Eur Urol* 2019;75:506-514.
- ⁴ Gandaglia G, Martini A, Ploussard G, et al; EAU-YAU Prostate Cancer Working Group. External validation of the 2019 Briganti nomogram for the identification of prostate cancer patients who should be considered for an extended pelvic lymph node dissection. *Eur Urol* 2020;78:138-142.
- ⁵ Hope TA, Eiber M, Armstrong WR, et al. Diagnostic accuracy of 68Ga-PSMA-11 PET for pelvic nodal metastasis detection prior to radical prostatectomy and pelvic lymph node dissection: A multicenter prospective phase 3 imaging trial. *JAMA Oncol* 2021;7:1635-1642.
- ⁶ Pienta KJ, Gorin MA, Rowe SP, et al. A phase 2/3 prospective multicenter study of the diagnostic accuracy of prostate specific membrane antigen PET/CT with 18F-DCFPyL in prostate cancer patients (OSPReY). *J Urol* 2021;206:52-61.

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF LOCAL SECONDARY THERAPY POST-RADIATION

Local Secondary Therapy for Recurrent Prostate Cancer After Definitive Radiotherapy:

- Patients with biopsy-proven recurrence in the prostate after prior RT and without distant metastatic disease can be considered for local therapy. Monitoring is also an option (see [PROS-10](#)). ADT may also be included, but it should not be reflexively ordered.
- The Panel recommends that patients receive multidisciplinary counseling about the risks and benefits of each of these options in the context of the available comparative literature on this topic.^{1,2}
- Local therapy options for patients with recurrence in the prostate include^{a,b,c}:
 - Cryotherapy
 - High-intensity focused ultrasound (HIFU)
 - Irreversible electroporation (IRE) (category 2B)
 - Reirradiation
 - RP + PLND
- Reirradiation options include LDR brachytherapy, HDR brachytherapy, and SBRT.¹⁻⁷
- There is no consensus as to the most appropriate reirradiation volume, and there are published experiences for both focal/partial and whole gland reirradiation. The Panel recommends that patients receiving local therapy for RT recurrence are treated within the context of clinical trials when available and/or at experienced centers.

^a Reirradiation with LDR brachytherapy, HDR brachytherapy, and SBRT is supported by phase II trials with >36 months of median follow-up and should be strongly considered as an option in patients with >2 years interval from prior radiation who do not have ongoing moderate or higher grade radiation-associated toxicities. Phase II data with >36 months of median follow-up are available, albeit to a more limited scope, for secondary RP, cryotherapy, and HIFU.

^b Currently, all phase II and/or prospectively accrued registries reporting oncologic outcomes with >36 months of follow-up have pursued whole gland treatments. Focal therapy for prostate-only recurrences is the subject of active investigation. See [Principles of Focal/Subtotal Therapy or Whole Gland Ablative Therapy \(PROS-I\)](#).

^c There are no prospective data to guide the use and duration of ADT with reirradiation for prostate-only recurrences. The Panel recommends extrapolation from the definitive treatment setting to suggest that adding ADT or prolonging it in this setting likely offers an improvement in biochemical control. See [Principles of Radiation Therapy \(PROS-J\)](#).

Note: All recommendations are category 2A unless otherwise indicated.

[References on PROS-L 2 of 2](#)



PRINCIPLES OF LOCAL SECONDARY THERAPY POST-RADIATION REFERENCES

- ¹ Ingrosso G, Becherini C, Lancia A, et al. Nonsurgical salvage local therapies for radiorecurrent prostate cancer: A systematic review and meta-analysis. *Eur Urol Oncol* 2020;3:183-197.
- ² Valle LF, Lehrer EJ, Markovic D, et al. A systematic review and meta-analysis of local salvage therapies after radiotherapy for prostate cancer (MASTER). *Eur Urol* 2021;80:280-292.
- ³ Crook J, Rodgers JP, Pisansky TM, et al. Salvage low-dose-rate prostate brachytherapy: Clinical outcomes of a phase 2 trial for local recurrence after external beam radiation therapy (NRG Oncology/RTOG 0526). *Int J Radiat Oncol Biol Phys* 2022;112:1115-1122.
- ⁴ Yamada Y, Kollmeier MA, Pei X, et al. A phase II study of salvage high-dose-rate brachytherapy for the treatment of locally recurrent prostate cancer after definitive external beam radiotherapy. *Brachytherapy* 2014;13:111-116.
- ⁵ Fuller DB, Chen RC, Crabtree T, et al. 10 year prostate stereotactic body radiotherapy outcomes: Relapse-free survival, PSA kinetics and toxicity from a pooled analysis of two multi-institutional trials. *Int J Radiat Oncol* 2020;108:e857-e858.
- ⁶ Bergamin S, Eade T, Kneebone A, et al. Interim results of a prospective prostate-specific membrane antigen-directed focal stereotactic reirradiation trial for locally recurrent prostate cancer. *Int J Radiat Oncol Biol Phys* 2020;108:1172-1178.
- ⁷ Pasquier D, Martinage G, Janoray G, et al. Salvage stereotactic body radiation therapy for local prostate cancer recurrence after radiation therapy: A retrospective multicenter study of the GETUG. *Int J Radiat Oncol Biol Phys* 2019;105:727-734.

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF METASTASIS-DIRECTED THERAPY (MDT)

General

- MDT is the treatment of metastatic sites with a local therapy to improve oncologic outcomes, not simply to provide symptom palliation. MDT has been studied primarily with radiotherapy and with highly selected use of surgical resection in the form of lymph node dissections. MDT is not recommended to be performed with other local therapies.
- MDT is mostly commonly delivered in the form of MDRT, which specifically refers to the use of higher than palliative dose radiotherapy to provide durable local control of the areas targeted. This may be for the intended purpose of delaying the initiation of systemic therapy, improving PFS, radiographic PFS, or overall survival. Similar to systemic therapies in the polymetastatic setting, MDRT is uncommonly curative and should be viewed as a form of cytotoxic therapy, and it may need to be repeated.

Disease State

- The settings for which MDT may have utility with variable evidence quality are shown in Table 1:

Table 1.

Treatment Regimen	mCSPC		mCRPC	
	Synchronous (de novo) Oligometastatic	Metachronous Oligorecurrent ^a	Oligometastatic	Oligoprogressive
MDT	☼	✓	✓	☼

(✓ Preferred; ☼ Acceptable based on clinical and medical need)

^a Limited data suggest that lymphadenectomy may be beneficial for select patients with pelvic nodal recurrence after radical prostatectomy if they do not have evidence of distant metastases.

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF METASTASIS-DIRECTED THERAPY (MDT)

Number of Metastatic Sites

- The number of metastatic sites to define oligometastatic disease remains an evolving space and is impacted by sensitivity of imaging used. Early studies limited metastatic sites to 1–3 or 1–5 by CT, MRI, or bone scan.^{1–4} More recent studies have included up to 10 metastatic sites by PSMA-PET imaging (NCT04787744, NCT06150417, NCT03721341). The upper limit is not clearly established and is both a function of oncologic limitations and technical limitations of treating numerous metastatic sites.
 - ▶ If CT, MRI, or bone scan is used, >5 metastasis is generally an exclusion for MDT.
 - ▶ If PSMA-PET is used, >10 metastatic sites is generally an exclusion for MDT.
- Lymph nodes in the same drainage chain (eg, two adjacent para-aortic lymph nodes) are to be considered 1 metastatic site and would be treated together with one radiation therapy isocenter or plan.
- Discrete, non-contiguous bony metastases should be counted separately even if treated with one isocenter (eg, two discrete femur metastases).
- The goal of MDT is generally to treat all metastatic sites, which may include regional lymph nodes and potentially treatment of the primary tumor if untreated or evidence of local recurrence. The primary tumor in this setting should be counted as a site.

Location of Metastatic Sites

- The location of metastatic sites have generally consisted of lymph node and osseous metastatic sites, but some studies include lung and liver metastases. Brain metastases are generally not included in studies of MDRT in prostate cancer. MDT should not be used to delay systemic therapy when there is visceral organ involvement.
- Pelvic lymph nodes are also considered metastatic sites in the context of MDRT for metachronous mCSPC or mCRPC.

¹ Marvaso G, Corrao G, Zaffaroni M, et al. ADT with SBRT versus SBRT alone for hormone-sensitive oligorecurrent prostate cancer (RADIOA): a randomised, open-label, phase 2 clinical trial. *Lancet Oncol* 2025;26:300-311.

² Tang C, Sherry AD, Haymaker C, et al. Addition of metastasis-directed therapy to intermittent hormone therapy for oligometastatic prostate cancer: the EXTEND phase 2 randomized clinical trial. *JAMA Oncol* 2023;9:825-834.

³ Deek, MP, Van der Eecken K, Sutera P, et al. Long-term outcomes and genetic predictors of response to metastasis-directed therapy versus observation in oligometastatic prostate cancer: analysis of STOMP and ORIOLE trials. *J Clin Oncol* 2022;40:3377-3382.

⁴ Francolini G, Allegra AG, Detti B, et al. Stereotactic body radiation therapy and abiraterone acetate for patients affected by oligometastatic castrate-resistant prostate cancer: a randomized phase II trial (ARTO). *J Clin Oncol* 2023;41:5561-5568.

Note: All recommendations are category 2A unless otherwise indicated.



PRINCIPLES OF METASTASIS-DIRECTED THERAPY (MDT)

Radiotherapy Details (Also see [Principles of Radiation Therapy](#))

- Use of SBRT (also known in this setting as stereotactic ablative radiotherapy [SABR]) is most commonly recommended for MDRT. Established dose/fraction regimens should be used with strict OAR constraints. Daily 3D image guidance is required and motion management is recommended for thoracic and abdominal sites if relevant. Online adaptive radiotherapy, if available, may be preferred if treatment of a metastatic site is near an OAR with frequent day-to-day setup uncertainty. Appropriate clinical target volume (CTV) and PTV margins should be applied to account for microscopic spread and setup uncertainty.
- When treating a lymph node metastasis, one should generally include an elective nodal volume to the involved chain with a simultaneous integrated boost to the radiographic or biopsy confirmed lymph node. If lymph node(s) are in the pelvis, elective nodal irradiation fields should be considered, either ipsilateral or bilateral.
- Dose composites with special medical physics consults are required to account for prior radiotherapy or overlap of multiple treatment plans.
- If treatment involves more than two treatment plans simultaneously, consideration of splitting the treatment course into multiple sessions of SBRT may be required due to patient tolerance and safety (eg, 4 treatment plans for 4 metastatic sites may need to be delivered as 2 concurrent treatment plans as course 1, and then the other 2 treatment plans are delivered concurrently at a later date as a separate course of SBRT). Repeat simulation may be required based on potential for anatomy changes or use of alternative immobilization for other treatments sites.

Concurrent Systemic Therapy with MDRT

- Synchronous (de novo) oligometastatic mCSPC:
 - ▶ Systemic therapy, most commonly ADT+ARPI, and treatment of the primary tumor with RT is recommended for most patients. The concurrent addition of MDRT may be considered in select patients. For patient tolerance, it is reasonable to asynchronously treat the primary and metastatic sites in separate treatment courses.
- Metachronous oligorecurrent mCSPC:
 - ▶ Oligorecurrent disease detected primarily by PET imaging may be treated by MDRT alone or concurrently with short-term ADT (6 months). The addition of ADT improved clinical and biochemical PFS, but may simply delay the detection of recurrence rather than improve durable control of the disease.¹
 - ▶ Trials have not been reported showing a benefit of the addition of triplet chemotherapy or an ARPI in the setting of MDRT+ADT.
- Oligometastatic mCRPC:
 - ▶ First-line treatment with ADT and an ARPI ± MDRT have demonstrated improvements in PFS and radiographic PFS from the addition of MDRT.⁴ MDRT should not be used without recommended mCRPC systemic therapies.
- Oligoprogressive or oligopersistent mCRPC
 - ▶ For oligoprogressive or oligopersistent mCRPC, MDRT may be delivered concurrently with mCRPC systemic therapy.
 - ▶ MDRT should not be used concurrently with PARP inhibition (alone or in combination with abiraterone or enzalutamide) or chemotherapy outside of a clinical trial.
 - ▶ Use of MDRT may be considered for oligoprogression sequentially after approved systemic mCRPC treatments.

¹ Marvaso G, Corrao G, Zaffaroni M, et al. ADT with SBRT versus SBRT alone for hormone-sensitive oligorecurrent prostate cancer (RADIOA): a randomised, open-label, phase 2 clinical trial. *Lancet Oncol* 2025;26:300-311.

⁴ Francolini G, Allegra AG, Detti B, et al. Stereotactic body radiation therapy and abiraterone acetate for patients affected by oligometastatic castrate-resistant prostate cancer: a randomized phase II trial (ARTO). *J Clin Oncol* 2023;41:5561-5568.

Note: All recommendations are category 2A unless otherwise indicated.



PRINCIPLES OF NON-HORMONAL SYSTEMIC THERAPY

- An FDA-approved biosimilar is an appropriate substitute for any recommended systemic biologic therapy in the NCCN Guidelines.

Non-Hormonal Systemic Therapy for M1 Castration-Sensitive Prostate Cancer

- Triplet therapy with ADT, certain ARPIs, and docetaxel is an option for certain patients with mCSPC who are fit for chemotherapy based on phase 3 studies:
 - ▶ ADT with docetaxel and abiraterone was compared to ADT alone or with docetaxel in an open-label, randomized, phase 3 study. Radiographic PFS was longer in patients who received abiraterone than in those who did not. The populations receiving the triplet and doublet therapies experienced similar rates of neutropenia, febrile neutropenia, fatigue, and neuropathy, although grade ≥ 3 adverse events occurred in 63% of patients who received the triplet combination compared with 52% of those receiving ADT and docetaxel.
 - ▶ ADT with docetaxel and darolutamide was compared with ADT with docetaxel and placebo in a randomized phase 3 trial. OS, time to CRPC, skeletal event-free survival, and time to initiation of subsequent systemic antineoplastic therapy were improved in the patients who received darolutamide. Adverse events of any grade, grade 3–5 adverse events, and serious adverse events occurred at similar incidence levels between the two arms. Many of these were known effects of docetaxel. Exceptions were rash (16.6% vs. 13.5%) and hypertension (13.7% vs. 9.2%), which are known effects of androgen receptor pathway inhibitors and were more frequent in the darolutamide group.
 - ▶ An open-label, randomized, phase 3 trial compared ADT with enzalutamide to ADT with a first-generation antiandrogen in this setting. Concurrent docetaxel was allowed and used for stratification. OS was improved with the use of enzalutamide over first-generation anti-androgen regardless of the addition of docetaxel. The most common grade ≥ 3 adverse events were febrile neutropenia associated with docetaxel use (6% in both groups), fatigue (1% in the control group vs. 6% in the enzalutamide group), and hypertension (6% vs. 10%). The incidence of grade 1–3 memory impairment was 4% versus 13%.
- The use of myeloid growth factors should follow the [NCCN Guidelines for Hematopoietic Growth Factors](#), based on risk of neutropenic fever.
- An additional option for patients with BRCA2-mutated mCSPC is the use of ADT with niraparib and abiraterone.
 - ▶ ADT with niraparib and abiraterone was compared to ADT plus abiraterone doublet therapy in a randomized phase 3 trial. Most patients had high-volume and/or synchronous disease. rPFS was improved in the BRCAm population; OS data are immature. Grade 3/4 adverse events were more common in the niraparib arm (75% vs. 59%) as were treatment-emergent adverse events leading to death (14 participants vs. 7 participants). The most common grade 3/4 adverse events were anemia (29.1% in the niraparib group versus 4.6% in the control group) and hypertension (26.5% versus 18.4%).
 - ▶ Niraparib/abiraterone can be used following the completion of docetaxel in patients receiving triplet therapy or following EBRT to the primary tumor.
 - ▶ The use of this combination in patients with low-volume disease is controversial because 78% of the patients in the trial had high-volume disease and there were very few patients with low-volume metachronous disease in particular. This regimen adds considerable toxicity, and patients with low-volume disease should be informed that the benefit in this setting is not clear.
 - ▶ The fine-particle (category 2B; other recommended option) or standard formulation of abiraterone can be given with single-agent niraparib as a substitute for the combination niraparib/abiraterone tablet.

Note: All recommendations are category 2A unless otherwise indicated.



PRINCIPLES OF NON-HORMONAL SYSTEMIC THERAPY

Non-Hormonal Systemic Therapy for M1 CRPC

• **Chemotherapy**

▶ **Docetaxel with concurrent steroid**

- ◊ Concurrent steroid includes daily prednisone, which may be omitted on the day of chemotherapy administration when dexamethasone is given.
- ◊ Every-3-week docetaxel with concurrent steroid is the preferred first-line chemotherapy treatment based on phase 3 clinical trial data for patients with symptomatic mCRPC. Adverse events associated with docetaxel include neutropenia, leukopenia, febrile neutropenia, neutropenic infections, fluid retention, hypersensitivity reaction, hepatic function impairment, neuropathy, and other low-grade adverse events (eg, fatigue, nausea, vomiting, alopecia, diarrhea).
- ◊ Only regimens utilizing docetaxel on an every-3-week schedule demonstrated beneficial impact on survival. The duration of therapy should be based on the assessment of benefit and toxicities. In the pivotal trials establishing survival advantage of docetaxel-based chemotherapy, patients received up to 10 cycles of treatment if no progression and no prohibitive toxicities were noted.
- ◊ Docetaxel retreatment can be attempted after progression on an ARPI and docetaxel in patients with mCRPC whose cancer has not demonstrated definitive evidence of progression on prior docetaxel therapy in either the castration-sensitive setting or the mCRPC setting.
- ◊ Rare patients may have exposure to docetaxel without prior ARPI (ie, in the setting of EBRT to the primary tumor for low-volume synchronous mCSPC). Docetaxel rechallenge is not recommended for such patients in the pre-ARPI mCRPC setting.

▶ **Cabazitaxel with concurrent steroid**

- ◊ Concurrent steroid includes daily prednisone, which may be omitted on the day of chemotherapy administration when dexamethasone is given.
- ◊ Patients who are not candidates for docetaxel or who are intolerant of docetaxel should be considered for cabazitaxel with concurrent steroid, based on results that suggest clinical activity of cabazitaxel in mCRPC. Cabazitaxel was associated with lower rates of peripheral neuropathy than docetaxel, particularly at 20 mg/m² (12% vs. 25%) and may be appropriate in patients with pre-existing mild peripheral neuropathy. Current data do not support greater efficacy of cabazitaxel over docetaxel.
- ◊ Cabazitaxel at 25 mg/m² with concurrent steroid has been shown in a randomized phase 3 study (TROPIC) to prolong OS, PFS, PSA response, and radiologic response when compared with mitoxantrone and prednisone and is FDA approved in the post-docetaxel second-line setting. Toxicity at this dose was significant and included febrile neutropenia, severe diarrhea, fatigue, nausea/vomiting, anemia, thrombocytopenia, sepsis, and renal failure. A recent trial, PROSELICA, compared cabazitaxel 25 mg/m² every 3 weeks to 20 mg/m² every 3 weeks. Cabazitaxel 20 mg/m² had less toxicity; febrile neutropenia, diarrhea, and fatigue were less frequent. Cabazitaxel at 20 mg/m² had a significantly lower PSA response rate but non-significantly lower radiographic response rate and non-significantly shorter PFS and OS (13.4 vs. 14.5 months) compared to 25 mg/m².
- ◊ Cabazitaxel at 25 mg/m² with concurrent steroid improved radiographic PFS and reduced the risk of death compared with abiraterone or enzalutamide in patients with prior docetaxel treatment for mCRPC in the CARD study.
- ◊ Biweekly cabazitaxel at 16 mg/m² with prophylactic G-CSF significantly reduced the risk of neutropenia/neutropenic complications compared to the 25 mg/m² dose with G-CSF in patients ≥65 years in a phase 3 trial. Clinical outcomes were comparable between the 2 groups.
- ◊ No chemotherapy regimen to date has demonstrated improved survival or quality of life after cabazitaxel, and trial participation should be encouraged.

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF NON-HORMONAL SYSTEMIC THERAPY

Non-Hormonal Systemic Therapy for M1 CRPC (continued)

• **Chemotherapy**

▶ **Cabazitaxel/carboplatin with concurrent steroid**

- ◊ Concurrent steroid includes daily prednisone, which may be omitted on the day of chemotherapy administration when dexamethasone is given.
- ◊ Cabazitaxel 20 or 25 mg/m² plus carboplatin AUC 4 mg/mL per minute with growth factor support can be considered for fit patients with aggressive variant mCRPC (ie, visceral metastases, low PSA and bulky disease, high lactate dehydrogenase (LDH), high CEA, lytic bone metastases, NEPC histology) or unfavorable genomics (defects in at least 2 of *PTEN*, *TP53*, and *RB1*). The most common grade 3–5 adverse events were fatigue, anemia, neutropenia, and thrombocytopenia. Corn PG, et al. *Lancet Oncol* 2019;20:1432-1443.
 - Cabazitaxel starting dose can be either 20 mg/m² or 25 mg/m² for patients with mCRPC whose cancer has progressed despite prior docetaxel chemotherapy. Cabazitaxel 25 mg/m² with concurrent steroid may be considered for healthy patients who wish to be more aggressive. Growth factor support may be needed with either dose.

▶ **Mitoxantrone with prednisone**

- ◊ Mitoxantrone with prednisone may provide palliation but has not been shown to extend survival in two randomized trials. Adverse events associated with mitoxantrone are similar to docetaxel, but with lower rates of grade 3 or 4 neutropenic fevers, cardiovascular events, nausea and vomiting, metabolic disturbances, and neurologic events.

▶ **Increasing PSA should not be used as the sole criteria for progression. Assessment of response should incorporate clinical and radiographic criteria.**

▶ **See [NCCN Guidelines for Hematopoietic Growth Factors](#) for recommendations on growth factor support.**

• **PARP Inhibitors with or without ARPIs**

- ▶ Inactivation of HRR genes has been associated with response to PARP inhibitor (PARPi) therapy, and a number of FDA-approved agents and combinations are now available ([Table 1](#) and [Discussion](#)). Although there are multiple specific gene alterations included in the FDA labels for specific PARPi and combinations (see [Table 1](#)), the response rates have been most pronounced for patients with *BRCA2* mutation-associated prostate cancers. Treatment responses have been variable for other genes alterations, and this information should be taken into account in decisions about treatment sequencing and patient counseling. For example, responses have been less robust for patients whose cancers carry ATM mutations, and other recommended regimen options may be favored until more data are available on a gene-by-gene level.
- ▶ Loss of *BRCA2* may be especially associated with response to PARPi therapy compared to other HRR gene alterations.
- ▶ There has been heterogeneity of response to olaparib for non-*BRCA* mutations based on the specific gene mutation ([Discussion](#)).
- ▶ The fine-particle (category 2B; other recommended option) or standard formulation of abiraterone can be given with single-agent niraparib as a substitute for the combination niraparib/abiraterone tablet.
- ▶ The fine-particle abiraterone (category 2B; other recommended option) may be given with olaparib as a substitute for standard abiraterone.

Note: All recommendations are category 2A unless otherwise indicated.



PRINCIPLES OF NON-HORMONAL SYSTEMIC THERAPY

Non-Hormonal Systemic Therapy for M1 CRPC (continued)

• **Immunotherapy**

▶ **Sipuleucel-T**

- ◊ Sipuleucel-T is only for asymptomatic or minimally symptomatic patients with no liver metastases, life expectancy >6 months, and ECOG performance status 0–1.
- ◊ Sipuleucel-T is not recommended for patients with small cell prostate cancer/NEPC.
- ◊ Sipuleucel-T has been shown in a phase 3 clinical trial to extend mean survival from 21.7 months in the control arm to 25.8 months in the treatment arm, which constitutes a 22% reduction in mortality risk.
- ◊ Sipuleucel-T is well-tolerated; common complications include chills, pyrexia, and headache.
- ▶ **Pembrolizumab** is an option for certain patients with MSI-H/dMMR mCRPC ([PROS-18](#)).
- ◊ Pembrolizumab may cause severe, life-threatening immune-mediated adverse reactions, which may include but are not limited to: pneumonitis, colitis, hepatitis, myocarditis, endocrinopathies, exfoliative dermatologic conditions, renal failure and nephritis, and ocular toxicities. See [NCCN Guidelines for Management of Immunotherapy-Related Toxicities](#).
- ◊ Limited data suggest that pembrolizumab may be associated with some benefit in patients with mCRPC and TMB ≥10 mut/mB. Lenis AT, et al. Clin Canc Res 2024;30:3894-3903.
- ◊ Pembrolizumab and berahyaluronidase alfa-pmph subcutaneous injection may be substituted for IV pembrolizumab. Pembrolizumab and berahyaluronidase alfa-pmph has different dosing and administration instructions compared to IV pembrolizumab.

• **Other Targeted Agents**

- ▶ Pan-cancer, tumor-agnostic treatments can be considered for patients with actionable mutations.

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

PRINCIPLES OF NON-HORMONAL SYSTEMIC THERAPY

Table 1. PARP Inhibitors with or without ARPIs

Treatment	FDA-Approved Disease State Indication	Biomarker Pathogenic variant or mutation in gene (germline and/or somatic)	Clinical Trial
Olaparib	mCRPC that has progressed following prior treatment with enzalutamide or abiraterone.	<i>BRCA2, BRCA1, ATM, BARD1, BRIP1, CDK12, CHEK1, CHEK2, FANCL, PALB2, RAD51B, RAD51C, RAD51D, or RAD54L</i>	PROfound ¹
Rucaparib	mCRPC that has been treated with androgen receptor-directed therapy and taxane-based chemotherapy	<i>BRCA2, BRCA1</i>	TRITON2 ^{2,3}
Olaparib/ Abiraterone	mCRPC	<i>BRCA2, BRCA1</i>	PROpel ⁴
Talazoparib/ Enzalutamide	mCRPC	<i>BRCA2, BRCA1, ATM, ATR, CDK12, CHEK2, FANCA, MLH1, MRE11A, NBN, PALB2, or RAD51C</i>	TALAPRO-2 ^{5,6}
Niraparib/ Abiraterone	mCRPC	<i>BRCA2, BRCA1</i>	MAGNITUDE ^{7,8}

¹ de Bono JD, Mateo J, Fizazi K, et al. Olaparib for metastatic castration-resistant prostate cancer. *NEJM* 2020;382:2091-2102.

² Abida W, Patnaik A, Campbell D, et al. Rucaparib in men with metastatic castration resistant prostate cancer harboring a BRCA1 or BRCA2 gene alteration. *J Clin Oncol* 2020;38:3763-3772.

³ Abida W, Campbell D, Patnaik A, et al. Rucaparib for the treatment of metastatic castration-resistant prostate cancer associated with a DNA damage repair gene alteration: final results from the Phase 2 TRITON2 study. *Eur Urol* 2023;84:321-330.

⁴ Saad F, Clarke NW, Oya M, et al. Olaparib plus abiraterone versus placebo plus abiraterone in metastatic castration-resistant prostate cancer (PROpel): final prespecified overall survival results of a randomised, double-blind, phase 3 trial. *Lancet Oncol* 2023;24:1094-1108.

⁵ Agarwal N, Azad AA, Carles J, et al. Talazoparib plus enzalutamide in men with first-line metastatic castration-resistant prostate cancer (TALAPRO-2): a randomised, placebo-controlled, phase 3 trial. *Lancet* 2023;402:291-303.

⁶ Fizazi K, Azad AA, Matsubara N, et al. First-line talazoparib with enzalutamide in HRR-deficient metastatic castration-resistant prostate cancer: the phase 3 TALAPRO-2 trial. 2023;30:257-264.

⁷ Chi KN, Rathkopf D, Smith MR, et al. Niraparib and abiraterone acetate for metastatic castration-resistant prostate cancer. *J Clin Oncol* 2023;41:339-3351.

⁸ Chi KN, Sandhu S, Smith MR, et al. Niraparib plus abiraterone acetate with prednisone in patients with metastatic castration-resistant prostate cancer and homologous recombination repair gene alterations: second interim analysis of the randomized phase III MAGNITUDE trial. 2023;34:772-782.

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

American Joint Committee on Cancer (AJCC) TNM Staging System For Prostate Cancer (8th ed., 2017)

Table 1. Definitions for T, N, M

Clinical T (cT)

T Primary Tumor

- TX** Primary tumor cannot be assessed
- T0** No evidence of primary tumor
- T1** Clinically inapparent tumor that is not palpable
 - T1a** Tumor incidental histologic finding in 5% or less of tissue resected
 - T1b** Tumor incidental histologic finding in more than 5% of tissue resected
 - T1c** Tumor identified by needle biopsy found in one or both sides, but not palpable
- T2** Tumor is palpable and confined within prostate
 - T2a** Tumor involves one-half of one side or less
 - T2b** Tumor involves more than one-half of one side but not both sides
 - T2c** Tumor involves both sides
- T3** Extraprostatic tumor that is not fixed or does not invade adjacent structures
 - T3a** Extraprostatic extension (unilateral or bilateral)
 - T3b** Tumor invades seminal vesicle(s)
- T4** Tumor is fixed or invades adjacent structures other than seminal vesicles such as external sphincter, rectum, bladder, levator muscles, and/or pelvic wall.

Pathological T (pT)

T Primary Tumor

- T2** Organ confined
- T3** Extraprostatic extension
 - T3a** Extraprostatic extension (unilateral or bilateral) or microscopic invasion of bladder neck
 - T3b** Tumor invades seminal vesicle(s)
- T4** Tumor is fixed or invades adjacent structures other than seminal vesicles such as external sphincter, rectum, bladder, levator muscles, and/or pelvic wall

Note: There is no pathological T1 classification.

Note: Positive surgical margin should be indicated by an R1 descriptor, indicating residual microscopic disease.

N Regional Lymph Nodes

- NX** Regional lymph nodes cannot be assessed
- N0** No positive regional nodes
- N1** Metastases in regional node(s)

M Distant Metastasis

- M0** No distant metastasis
- M1** Distant metastasis
 - M1a** Nonregional lymph node(s)
 - M1b** Bone(s)
 - M1c** Other site(s) with or without bone disease

Note: When more than one site of metastasis is present, the most advanced category is used. M1c is most advanced.

Used with permission of the American College of Surgeons, Chicago, Illinois. The original source for this information is the AJCC Cancer Staging Manual, Eighth Edition (2017) published by Springer International Publishing.



NCCN Guidelines Version 5.2026

Prostate Cancer

Table 2. AJCC Prognostic Groups

Group	T	N	M	PSA (ng/mL)	Grade Group
Stage I	cT1a-c	N0	M0	PSA <10	1
	cT2a	N0	M0	PSA <10	1
	pT2	N0	M0	PSA <10	1
Stage IIA	cT1a-c	N0	M0	PSA ≥10 <20	1
	cT2a	N0	M0	PSA ≥10 <20	1
	pT2	N0	M0	PSA ≥10 <20	1
	cT2b	N0	M0	PSA <20	1
	cT2c	N0	M0	PSA <20	1
Stage IIB	T1-2	N0	M0	PSA <20	2
Stage IIC	T1-2	N0	M0	PSA <20	3
	T1-2	N0	M0	PSA <20	4
Stage IIIA	T1-2	N0	M0	PSA ≥20	1-4
Stage IIIB	T3-4	N0	M0	Any PSA	1-4
Stage IIIC	Any T	N0	M0	Any PSA	5
Stage IVA	Any T	N1	M0	Any PSA	Any
Stage IVB	Any T	Any N	M1	Any PSA	Any

Histopathologic Type

This classification applies to adenocarcinomas and squamous carcinomas, but not to sarcoma or transitional cell (urothelial) carcinoma of the prostate. Adjectives used to describe histologic variants of adenocarcinomas of prostate include mucinous, signet ring cell, ductal, and neuroendocrine, including small cell carcinoma. There should be histologic confirmation of the disease.

Definition of Histologic Grade Group (G)

Recently, the Gleason system has been compressed into so-called Grade Groups.

Grade Group	Gleason Score	Gleason Pattern
1	≤6	≤3+3
2	7	3+4
3	7	4+3
4	8	4+4, 3+5, 5+3
5	9 or 10	4+5, 5+4, 5+5

Note: When either PSA or Grade Group is not available, grouping should be determined by T category and/or either PSA or Grade Group as available.

Used with permission of the American College of Surgeons, Chicago, Illinois. The original source for this information is the AJCC Cancer Staging System Manual, Eighth Edition (2017) published by Springer International Publishing.



NCCN Guidelines Version 5.2026

Prostate Cancer

ABBREVIATIONS

3D	three-dimensional	DCE	dynamic contrast-enhanced	IGRT	image-guided radiation therapy
ACE-27	Adult Comorbidity Evaluation-27 Index	DEXA	dual-energy x-ray absorptiometry	IMRT	intensity modulated RT
ADT	androgen deprivation therapy	DIL	dominant intraprostatic lesion	IRE	irreversible electroporation
AI	artificial intelligence	DM	distant metastases	IRF	intermediate risk factor
ARPI	androgen receptor pathway inhibitor	dMMR	mismatch repair deficient	ITT	intention to treat
aRT	adjuvant RT	DRE	digital rectal examination	LDH	lactate dehydrogenase
ASTRO	American Society for Radiation Oncology	DWI	diffusion-weighted imaging	LDR	low dose rate
AUC	area under the curve	EBRT	external beam radiation therapy	LHRH	luteinizing hormone-releasing hormone
BCR	biochemical recurrence	ECG	electrocardiogram	LT-ADT	long-term androgen deprivation therapy
BED	biologically effective dose	ECHO	echocardiogram	mCRPC	metastatic castration-resistant prostate cancer
BMI	body mass index	ENI	elective nodal irradiation	mCSPC	metastatic castration-sensitive prostate cancer
CBRT	cone beam CT	FDG	fluorodeoxyglucose	MDT	metastasis-directed therapy
CEA	carcinoembryonic antigen	FFS	failure-free survival	MDRT	metastasis-directed radiotherapy
CHIP	clonal hematopoiesis of indeterminate potential	FIR	favorable intermediate-risk	MFS	metastasis-free survival
CLR	correct localization rate	FLA	focal laser ablation	MMAI	multimodal artificial intelligence
CRPC	castration-resistant prostate cancer	FRAX	Fracture Risk Assessment Tool	MMR	mismatch repair
CSPC	castration-sensitive prostate cancer	FT	focal therapy	mpMRI	multiparametric MRI
ctDNA	circulating tumor DNA	GC	genomic classifier	MSI	microsatellite instability
CVD	cardiovascular disease	HDR	high dose rate	MSI-H	microsatellite instability-high
		HIFU	high-intensity focused ultrasound		
		HRD	homologous recombination deficiency		
		HRR	homologous recombination repair		

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

ABBREVIATIONS

NEPC	neuroendocrine prostate cancer	rh	radiohybrid
NNT	number needed to treat	RFA	radiofrequency ablation
		ROI	region of interest
OAR	organ at risk	RP	radical prostatectomy
OS	overall survival	rPFS	radiographic progression-free survival
		RTOG	Radiation Therapy Oncology Group
PARPi	PARP inhibitor		
PCSM	prostate cancer-specific mortality		
PCSS	prostate cancer-specific survival		
PDT	photodynamic therapy	SABR	stereotactic ablative radiotherapy
PFS	progression-free survival	SBRT	stereotactic body radiation therapy
PLND	pelvic lymph node dissection	sHR	subdistribution hazard ratio
PNRT	prophylactic nodal radiotherapy	SPECT	single-photon emission computed tomography
PPV	positive predictive value	SRE	skeletal-related event
PSA	prostate-specific antigen	sRT	secondary RT
PSADT	prostate-specific antigen doubling time	ST-ADT	short-term androgen deprivation therapy
PSMA	prostate-specific membrane antigen		
PTV	planning target value	TMB	tumor mutational burden
		TULSA	transurethral ultrasound ablation
QOL	quality of life	VUS	variant of uncertain significance

Note: All recommendations are category 2A unless otherwise indicated.



NCCN Guidelines Version 5.2026

Prostate Cancer

NCCN Categories of Evidence and Consensus	
Category 1	Based upon high-level evidence (≥1 randomized phase 3 trials or high-quality, robust meta-analyses), there is uniform NCCN consensus (≥85% support of the Panel) that the intervention is appropriate.
Category 2A	Based upon lower-level evidence, there is uniform NCCN consensus (≥85% support of the Panel) that the intervention is appropriate.
Category 2B	Based upon lower-level evidence, there is NCCN consensus (≥50%, but <85% support of the Panel) that the intervention is appropriate.
Category 3	Based upon any level of evidence, there is major NCCN disagreement that the intervention is appropriate.

All recommendations are category 2A unless otherwise indicated.

NCCN Categories of Preference	
Preferred	Interventions that are based on superior efficacy, safety, and evidence; and, when appropriate, affordability.
Other recommended	Other interventions that may be somewhat less efficacious, more toxic, or based on less mature data; or significantly less affordable for similar outcomes.
Useful in certain circumstances	Other interventions that may be used for selected patient populations (defined with recommendation).

All recommendations are considered appropriate.



NCCN Guidelines Version 5.2026

Prostate Cancer

Discussion

Table of Contents

This discussion corresponds to the NCCN Guidelines for Prostate Cancer. Last updated: January 23, 2026.

Overview	MS-2
Guidelines Update Methodology	MS-2
Literature Search Criteria	MS-2
Sensitive/Inclusive Language Usage	MS-3
Initial Prostate Cancer Diagnosis	MS-3
Estimates of Life Expectancy	MS-3
Quality-of-Life and Shared Decision-Making	MS-4
Prostate Cancer Genetics	MS-4
Inequities Based on Race and Ethnicity	MS-7
Risk Stratification for Clinically Localized Disease	MS-8
NCCN Risk Groups	MS-8
Nomograms	MS-10
Advanced Risk Stratification Tools	MS-10
Initial Clinical Assessment and Staging Evaluation	MS-12
Imaging Techniques	MS-13
Multiparametric MRI (mpMRI)	MS-13
PET Imaging	MS-14
Risks of Imaging	MS-16
Observation	MS-17
Active Surveillance	MS-18
Radical Prostatectomy	MS-23
Radiation Therapy	MS-25
Comparison of Treatment Options for Localized Disease	MS-31
Other Local Therapies	MS-32
Disease Monitoring	MS-32
Post-Radical Prostatectomy Approaches	MS-33
Early Secondary or Adjuvant Treatment for Adverse Features After Radical Prostatectomy	MS-33
Adjuvant or Early Secondary Therapy for pN1	MS-34
Secondary Treatment for PSA Persistence or Biochemical Recurrence After Radical Prostatectomy	MS-34
Post-Radiation Recurrence	MS-35
Second Biochemical Recurrence	MS-36
Androgen Deprivation Therapy	MS-37
ADT for Initial Treatment of Clinically Localized (N0,M0) Disease	MS-38
ADT for Initial Treatment of Regional Disease	MS-39
Palliative ADT	MS-39

Primary ADT for mCSPC	MS-39
Intermittent Versus Continuous ADT	MS-40
Adverse Effects of Traditional ADT	MS-41
Management of mCSPC	MS-43
Doublet Therapies for mCSPC	MS-43
Triplet Therapies for mCSPC	MS-46
PARP Inhibitor Combination for mCSPC	MS-48
EBRT to the Primary Tumor in Synchronous Low-Volume M1 Disease	MS-48
MDT for Oligometastatic CSPC	MS-49
Progression to and Management of CRPC	MS-50
Secondary Hormone Therapy for CRPC	MS-51
Abiraterone Acetate in mCRPC	MS-52
Enzalutamide in M0 and M1 CRPC	MS-53
Apalutamide in M0 CRPC	MS-54
Darolutamide in M0 CRPC	MS-55
Other Secondary Hormone Therapies	MS-55
Chemotherapy, Immunotherapy, and Targeted Therapy for mCRPC	MS-55
Docetaxel	MS-56
Cabazitaxel	MS-56
Cabazitaxel/Carboplatin	MS-57
Sipuleucel-T	MS-58
Pembrolizumab	MS-58
Mitoxantrone	MS-60
Treatment Options for Patients with DNA Repair Gene Mutations	MS-60
Radiopharmaceuticals for mCRPC	MS-65
Lutetium Lu 177 vipivotide tetraxetan	MS-65
Radium-223	MS-66
MDT for mCRPC	MS-67
Small Cell/Neuroendocrine Prostate Cancer	MS-67
Additional Treatment Options for Bone Metastases	MS-68
Considerations for Visceral Metastases	MS-69
Sequencing of Therapy in CRPC	MS-69
Summary	MS-70
Table 1: Selected Active Surveillance Experiences with Large Patient Cohorts	MS-71
References	MS-72



NCCN Guidelines Version 5.2026

Prostate Cancer

Overview

An estimated 313,780 new cases of prostate cancer will be diagnosed in the United States in 2025, accounting for 30% of new cancer cases in men.¹ It is the most common cancer in men in the United States, who currently have a 1 in 8 lifetime risk of developing prostate cancer.¹ The incidence of prostate cancer declined by approximately 40% from 2007 to 2014, but since that time has increased at a rate of 3% annually. These trends largely reflect the changes in prostate-specific antigen (PSA) screening recommendations. The decrease in PSA screening that followed the 2012 USPSTF recommendations against routine testing was associated with a rise in the diagnosis of regional and metastatic disease.²⁻¹⁰

Researchers further estimate that prostate cancer will account for 11% of male cancer deaths in the United States in 2025, with an estimated 35,770 deaths.¹ The age-adjusted death rate from prostate cancer declined by 52% from 1993 to 2017, but the death rate has become more stable in recent years, with a 0.5% annual decrease from 2012 through 2022.¹ For all stages combined, the 5-year relative survival rate for prostate cancer is 97%.¹ The comparatively low death rate suggests that increased public awareness with earlier detection and treatment has affected mortality from this prevalent cancer, but is also complicated by screening-related lead-time bias and detection of indolent cancers. Maintenance of this low death rate is threatened by the rising prostate cancer incidence and diagnosis of advanced disease.

Unfortunately, large inequities exist in incidence of and mortality from prostate cancer across racial and ethnic groups. The incidence rate in Black individuals is 67% higher than in white individuals, with prostate cancer accounting for 44% of cancer diagnoses in Black men and a 1 in 6 lifetime risk of a prostate cancer diagnosis.¹¹ Black individuals are also more likely to be diagnosed with more aggressive disease and are less

likely to have had a PSA test within the past year.¹¹ The mortality rate from prostate cancer in this population is two to four times higher than all other racial and ethnic groups; prostate cancer accounts for 17% of male cancer deaths in the United States.^{1,11} However, the overall prognosis by race appears similar when patients are treated with the same guideline-concordant care.¹²

Use of PSA for early detection of potentially fatal prostate cancer coupled with the use of imaging and the consideration of risk calculators and/or biomarkers to improve the specificity of screening should decrease the risk of overdetected (see the NCCN Guidelines for Prostate Cancer Early Detection, available at www.NCCN.org). This reduced overdetected along with the use of active surveillance in appropriate patients should reduce overtreatment AND preserve the relatively low rates of prostate cancer mortality.

Guidelines Update Methodology

The complete details of the Development and Update of the NCCN Guidelines are available at www.NCCN.org.

Literature Search Criteria

Prior to the update of the NCCN Guidelines for Prostate Cancer, an electronic search of the PubMed database was performed to obtain key literature in prostate cancer published since the previous Guidelines update, using the search term “prostate cancer.” The PubMed database was chosen because it remains the most widely used resource for medical literature and indexes peer-reviewed biomedical literature.¹³

The search results were narrowed by selecting studies in humans published in English. Results were confined to the following article types: Clinical Trial, Phase III; Clinical Trial, Phase IV; Guideline; Practice Guideline; Randomized Controlled Trial; Meta-Analysis; Systematic



NCCN Guidelines Version 5.2026

Prostate Cancer

Reviews; and Validation Studies. The data from key PubMed articles as well as articles from additional sources deemed as relevant to these Guidelines as discussed by the Panel during the Guidelines update have been included in this version of the Discussion section. Recommendations for which high-level evidence is lacking are based on the Panel's review of lower-level evidence and expert opinion.

Sensitive/Inclusive Language Usage

NCCN Guidelines strive to use language that advances the goals of equity, inclusion, and representation.¹⁴ NCCN Guidelines endeavor to use language that is person-first; not stigmatizing; anti-racist, anti-classist, anti-misogynist, anti-ageist, anti-ableist, and anti-weight-biased; and inclusive of individuals of all sexual orientations and gender identities. NCCN Guidelines incorporate non-gendered language, instead focusing on organ-specific recommendations. This language is both more accurate and more inclusive and can help fully address the needs of individuals of all sexual orientations and gender identities. NCCN Guidelines will continue to use the terms men, women, female, and male when citing statistics, recommendations, or data from organizations or sources that do not use inclusive terms. Most studies do not report how sex and gender data are collected and use these terms interchangeably or inconsistently. If sources do not differentiate gender from sex assigned at birth or organs present, the information is presumed to predominantly represent cisgender individuals. NCCN encourages researchers to collect more specific data in future studies and organizations to use more inclusive and accurate language in their future analyses.

Initial Prostate Cancer Diagnosis

Initial suspicion of prostate cancer is based on an abnormal digital rectal exam (DRE) or an elevated PSA level. A separate NCCN Guidelines Panel has written guidelines for prostate cancer early detection (see the NCCN Guidelines for Prostate Early Detection, available at

www.NCCN.org). Definitive diagnosis requires biopsies of the prostate, usually performed by a urologist using a needle under transrectal ultrasound (TRUS) guidance. A pathologist assigns a Gleason primary and secondary grade to the biopsy specimen. Clinical staging is based on the TNM (tumor, node, metastasis) classification from the AJCC Staging Manual, Eighth Edition.¹⁵ NCCN treatment recommendations are based on risk stratification that includes TNM staging rather than on AJCC prognostic grouping.

Pathology synoptic reports (protocols) are useful for reporting results from examinations of surgical specimens; these reports assist pathologists in providing clinically useful and relevant information. The NCCN Guidelines Panel favors pathology synoptic reports from the College of American Pathologists (CAP) that comply with the Commission on Cancer (CoC) requirements.¹⁶

Estimates of Life Expectancy

Estimates of life expectancy have emerged as a key determinant of primary treatment, particularly when considering observation. Life expectancy can be estimated for groups of individuals, but it is difficult to extrapolate these estimates to an individual patient. Life expectancy can be estimated using the Social Security Administration Life Insurance Tables,¹⁷ the Memorial Sloan Kettering Male Life Expectancy tool,¹⁸ or the University of California San Francisco (UCSF) Lee Schonberg Index¹⁹ and adjusted for individual patients by adding or subtracting 50% based on whether one believes the patient is in the healthiest quartile or the unhealthiest quartile, respectively.²⁰ As an example, the Social Security Administration Life Expectancy for a 65-year-old American male is 17.5 years. If judged to be in the upper quartile of health, a life expectancy of 26.3 years is assigned. If judged to be in the lower quartile of health, a life expectancy of 8.8 years is assigned. Thus, treatment recommendations could change dramatically using the NCCN Guidelines if a 65-year-old



NCCN Guidelines Version 5.2026

Prostate Cancer

patient was assessed to be in excellent health or if their health is in decline.

Quality-of-Life and Shared Decision-Making

Patient quality of life (QOL, also referred to as health-related quality of life [HRQOL]) should be taken into consideration when discussing and deciding upon management strategies for all types of cancer. In regards to prostate cancer, addressing QOL is especially important due to early detection and increased survival rates.²¹ The management of localized prostate cancer can have risks and side effects related to radiation, surgery, and hormone therapy—including functional effects on urinary, bowel, and sex organs.²²⁻²⁷ Some side effects have been linked to lasting psychological effects; for example an association has been shown between erectile dysfunction and depressive symptoms.^{28,29} Additionally, the use of androgen deprivation therapy (ADT) is associated with 23% higher rates of depression compared to patients who did not receive ADT.²⁹ Use of a standardized patient-reported outcomes instrument, such as the abbreviated Expanded Prostate Cancer Index Composite (EPIC-26), can aid in the assessment of baseline urinary, bowel, and sexual function and serve as a basis for shared-decision making.²¹

Shared decision-making should be collaborative and informative. This approach is especially applicable when multiple treatment options are considered clinically acceptable and when some treatment options have greater potential for adverse effects on QOL than others. When patients have actively participated in making decisions about their medical care, they generally report that they feel empowered, have better rates of adherence to the medical plan, and are less likely to display regrets about the decisions that were made.³⁰ When shared-decision making is utilized, patients can make informed decisions with more realistic expectations and greater satisfaction over the long term.

Prostate Cancer Genetics

Family history of prostate cancer raises the risk of prostate cancer.³¹⁻³⁴ In addition, prostate cancer has been associated with hereditary breast and ovarian cancer (HBOC) syndrome (due to germline mutations in homologous DNA repair genes) and Lynch syndrome (resulting from germline mutations in DNA mismatch repair [MMR] genes).³⁴⁻³⁹ In fact, approximately 11% of patients with prostate cancer and at least 1 additional primary cancer carry germline mutations associated with increased cancer risk.⁴⁰ Therefore, the Panel recommends a thorough review of personal and family history for all patients with prostate cancer at the time of initial diagnosis.^{41,42}

The presence of germline mutations has implications for family genetic counseling, cancer risk syndromes, and assessment of personal risk for subsequent cancers. Some patients with prostate cancer and their families may be at increased risk for breast and ovarian cancer, melanoma, and pancreatic cancer (HBOC); colorectal cancers (Lynch syndrome); and other cancer types. While DNA repair defects carry a worse prognosis for those with metastatic prostate cancer, it does not appear to be prognostic in localized disease.⁴³ Finally, there are possible treatment implications for patients with DNA repair defects with advanced disease (see *Treatment Options for Patients with DNA Repair Gene Mutations*, below).

Prostate cancer is often associated with somatic mutations that occur in the tumor but not in the germline. Both germline and tumor mutations are discussed herein.

Homologous DNA Repair Genes

Somatic mutations in DNA repair pathway genes occur in up to 19% of localized prostate tumors and 23% of metastatic CRPC (mCRPC) tumors, with most mutations found in *BRCA2* and *ATM*.^{44,45} These tumor mutations are often associated with germline mutations. For example, 42% of



NCCN Guidelines Version 5.2026

Prostate Cancer

patients with mCRPC and somatic mutations in *BRCA2* were found to carry the mutation in their germlines.⁴⁴ In localized prostate cancer, that number was 60%.⁴⁵

Overall, germline DNA repair mutations have been reported with the lowest frequencies seen in patients with lower-risk localized prostate cancer (1.6%–3.8%), higher frequencies in those with higher-risk localized disease (6%–8.9%), and the highest frequencies in those with metastatic disease (7.3%–16.2%).^{44,46–52} One study found that 11.8% of patients with metastatic prostate cancer have germline mutations in 1 of 16 DNA repair genes: *BRCA2* (5.3%), *ATM* (1.6%), *CHEK2* (1.9%), *BRCA1* (0.9%), *RAD51D* (0.4%), *PALB2* (0.4%), *ATR* (0.3%), and *NBN*, *PMS2*, *GEN1*, *MSH2*, *MSH6*, *RAD51C*, *MRE11A*, *BRIP1*, or *FAM175A*.⁵¹

An additional study showed that 9 of 125 patients with high-risk, very-high-risk, or metastatic prostate cancer (7.2%) had pathogenic germline mutations in *MUTYH* (4), *ATM* (2), *BRCA1* (1), *BRCA2* (1), and *BRIP1* (1).⁴⁸ In this study, the rate of metastatic disease among those with a mutation identified was high (28.6%, 2 of 7 patients). Although having a relative with breast cancer was associated with germline mutation identification ($P = .035$), only 45.5% of the mutation carriers in the study had mutations that were concordant with their personal and family history. Another study also found that a family history of breast cancer increased the chances of identifying a germline DNA repair gene mutation in patients with prostate cancer (OR, 1.89; 95% CI, 1.33–2.68; $P = .003$).⁵³ In a study of an unselected cohort of 3607 patients with a personal history of prostate cancer who had germline genetic testing based on clinician referral, 11.5% had germline mutations in *BRCA2*, *CHEK2*, *ATM*, *BRCA1*, or *PALB2*.⁵⁴

More than 2% of Ashkenazi Jews carry germline mutations in *BRCA1* or *BRCA2*, and these carriers have a 16% chance (95% CI, 4%–30%) of developing prostate cancer by the age of 70.⁵⁵ In a study of 251 unselected Ashkenazi Jewish patients with prostate cancer, 5.2% had

germline mutations in *BRCA1* and *BRCA2*, compared with 1.9% of control Ashkenazi Jewish males.⁵⁶

Germline *BRCA1* or *BRCA2* mutations have been associated with an increased risk for prostate cancer in numerous reports.^{38,39,56–66} In particular, *BRCA2* mutations have been associated with a 2- to 6-fold increase in the risk for prostate cancer, whereas the association of *BRCA1* mutations and increased risks for prostate cancer are less consistent.^{38,39,56,58,60,65,67,68} In addition, limited data suggest that germline mutations in *ATM*, *PALB2*, and *CHEK2* increase the risk of prostate cancer.^{69–72} Furthermore, prostate cancer in individuals with germline *BRCA* mutations (*BRCAm*) appears to occur earlier, and in patients with metastatic prostate cancer, has a more aggressive phenotype, and is associated with significantly reduced survival times than in patients without these mutations.^{43,67,73–79}

DNA Mismatch Repair Genes

Tumor mutations in *MLH1*, *MSH2*, *MSH6*, and *PMS2* may result in tumor microsatellite instability (MSI) and deficient MMR (dMMR; detected by immunohistochemistry) and are sometimes associated with germline mutations and Lynch syndrome. Patients with Lynch syndrome may have an increased risk for prostate cancer. In particular, studies show an increased risk for prostate cancer in patients who are older and have germline *MSH2* mutations.^{80,81}

In a study of >15,000 patients with cancer treated at Memorial Sloan Kettering Cancer Center who had their tumor and matched normal DNA sequenced and tumor MSI status assessed, approximately 5% of 1048 patients with prostate cancer had MSI-high (MSI-H) or MSI-indeterminate tumors, 5.6% of whom were found to have Lynch syndrome (0.29% of patients with prostate cancer).³⁵ In another prospective case series, the tumors of 3.1% of 1033 patients with prostate cancer demonstrated MSI-



NCCN Guidelines Version 5.2026

Prostate Cancer

H/dMMR status, and 21.9% of these patients had Lynch syndrome (0.68% of the total population).⁸² In a study of an unselected cohort of 3607 patients with a personal history of prostate cancer who had germline genetic testing based on clinician referral, 1.7% had germline mutations in *PMS2*, *MLH1*, *MSH2*, or *MSH6*.⁵⁴

Effect of Intraductal/Cribriform or Ductal Histology

Ductal prostate carcinomas are rare, accounting for approximately 1.3% of prostate carcinomas.⁸³ Intraductal prostate cancer may be more common, especially in higher risk groups, and may be associated with a poor prognosis.⁸⁴ It is important to note that there is significant overlap in diagnostic criteria and that intraductal, ductal, and invasive cribriform features may coexist in the same biopsy. By definition, intraductal carcinoma includes cribriform proliferation of malignant cells as long as they remain confined to a preexisting gland that is surrounded by basal cells. These features are seen frequently with an adjacent invasive cribriform component and would be missed without the use of basal cell markers.

Limited data suggest that acinar prostate adenocarcinoma with invasive cribriform pattern, intraductal carcinoma of prostate (IDC-P), or ductal adenocarcinoma component have increased genomic instability.⁸⁵⁻⁸⁸ In particular, tumors with these histologies may be more likely to harbor somatic MMR gene alterations than those with adenocarcinoma histology.⁸⁸⁻⁹⁰ In addition, limited data suggest that germline homologous DNA repair gene mutations may be more common in prostate tumors of ductal or intraductal origin^{91,92} and that intraductal histology is common in germline *BRCA2* mutation carriers with prostate cancer.⁹³ Overall, the Panel believes that the data connecting histology and the presence of genomic alterations are stronger for intraductal than ductal histology at this time.

Genetic Testing Recommendations

Germline Testing Based on Family History, Histology, and Risk Groups

The Panel recommends inquiring about family and personal history of cancer and known germline variants at time of initial diagnosis. Germline testing should be considered in appropriate individuals where it is likely to impact the prostate cancer treatment and clinical trial options, management of risk of other cancers, and/or potential risk of cancer in family members. Criteria for germline testing from the following NCCN Guidelines (available at www.NCCN.org) should be reviewed at time of initial diagnosis and, if relevant, at recurrence:

- The NCCN Guidelines for Genetic/Familial High-Risk Assessment: Breast, Ovarian, Pancreatic, and Prostate (page CRIT-6)
- The NCCN Guidelines for Genetic/Familial High-Risk Assessment: Colorectal, Endometrial, and Gastric (page HRS-3).

If criteria are met, multigene testing is recommended (see GENE-1 in the NCCN Guidelines for Genetic/Familial High-Risk Assessment: Breast, Ovarian, Pancreatic, and Prostate).

Genetic counseling resources and support are critical, and post-test genetic counseling is recommended if a germline mutation (pathogenic/likely pathogenic variant) is identified. Cascade testing for relatives is critical to inform the risk for familial cancers in all relatives. Post-test genetic counseling is recommended if there is a positive family history but no pathogenic variant OR if only germline variants of uncertain significance (VUS) are identified. This is to ensure accurate understanding of family implications and review indications for additional testing and/or follow up (including clinical trials of reclassification). Resources are available to check the known pathologic effects of genomic variants (eg, <https://brcaexchange.org/about/app>; <https://www.ncbi.nlm.nih.gov/clinvar/>). Information regarding germline mutations in patients with metastatic



NCCN Guidelines Version 5.2026

Prostate Cancer

disease can be used to inform future treatments or to determine eligibility for clinical trials.

Somatic Tumor Testing Based on Risk Groups

Tumor testing recommendations are as follows:

1. Tumor testing for somatic homologous recombination gene mutations (eg, *BRCA1*, *BRCA2*, *ATM*, *PALB2*, *FANCA*, *RAD51D*, *CHEK2*, *CDK12*) can be considered in patients with regional (N1) prostate cancer and is recommended for those with metastatic disease.
2. Tumor testing for MSI-H or dMMR can be considered in patients with regional or metastatic castration-sensitive prostate cancer (CSPC) and is recommended in the mCRPC setting.
3. Tumor mutational burden (TMB) testing is recommended in patients with mCRPC.

The Panel strongly recommends a metastatic biopsy for histologic and molecular evaluation. When unsafe or unfeasible, plasma ctDNA assay is an option, preferably collected during biochemical (PSA) and/or radiographic progression in order to maximize diagnostic yield. Caution is needed when interpreting ctDNA-only evaluation due to potential interference from clonal hematopoiesis of indeterminate potential (CHIP), which can result in a false-positive biomarker signal.⁹⁴

If MSI testing is performed, testing using an NGS assay validated for prostate cancer is preferred.⁹⁵⁻⁹⁷ If MSI-H or dMMR is found, the patient should be referred for genetic counseling to assess for the possibility of Lynch syndrome. MSI-H or dMMR indicate eligibility for pembrolizumab for patients with mCRPC (see *Pembrolizumab*, below).

Post-test genetic counseling is also recommended if pathogenic/likely pathogenic somatic mutations in any gene that has clinical implications are also identified in germline (eg, *BRCA1*, *BRCA2*, *ATM*, *PALB2*, *CHEK2*,

MLH1, *MSH2*, *MSH6*, *PMS2*). Virtually none of the NGS tests is designed or validated for germline assessment. Therefore, over-interpretation of germline findings should be avoided. If a germline mutation is suspected, the patient should be recommended for genetic counseling and follow-up dedicated germline testing.

Inequities Based on Race and Ethnicity

A CDC analysis of population-based cancer registries found that the incidence of prostate cancer was 67% higher in Black individuals than in white individuals in the United States.¹¹ Data also show that Black Americans are more likely to be diagnosed with more advanced disease and less likely to receive PSA screening.^{11,98-101} Furthermore, the mortality rate from prostate cancer in the Black American population is two to four times higher than all other racial and ethnic groups.^{1,11} Moreover, Black American patients with low-risk or favorable intermediate-risk prostate cancer have an increase in all-cause mortality after treatment, potentially due to cardiovascular complications after ADT.¹⁰²

Significant inequities also exist for other racial and ethnic groups. For example, Hispanic patients are more likely than non-Hispanic white patients to present with higher risk localized or metastatic prostate cancer.^{103,104} Furthermore, Hispanic individuals are less likely than non-Hispanic white individuals to receive treatment and more likely to experience treatment delays.^{103,104} Similarly, Asian American, Native Hawaiian, and Pacific Islander individuals are more likely to be diagnosed at a higher risk group than white individuals.¹⁰⁵

The main drivers for these racial and ethnic inequities are lower rates of prostate cancer screening, diagnostic procedures, and overall access to health care; treatment disparities; inadequate health insurance coverage; and other factors related to social determinants of health (eg, environmental exposures, patient and physician behaviors, delays in



NCCN Guidelines Version 5.2026

Prostate Cancer

diagnosis).^{98,100,106-112} For example, an analysis of SEER data from 2004 to 2011 determined that Black and Hispanic individuals had a decreased likelihood of receiving treatment than white individuals even when diagnosed with high-risk disease.¹¹³ Importantly, data show that Black patients who receive equitable prostate cancer treatment have similar outcomes to white patients.^{12,114-116}

Risk Stratification for Clinically Localized Disease

Unlike in some cancers where treatments for early-stage disease are guided by predictive biomarkers, such as ER/PR or HER2 status in breast cancer, in localized prostate cancer treatment is based largely on risk stratification of prognosis.

NCCN and other risk classification schemas, such as CARPA¹¹⁷ or STAR-CAP,¹¹⁸ are prognostic and have not been shown to be predictive of benefit to a specific treatment. Thus, common decision points, such as conservative management versus radical therapy, radiotherapy alone versus the addition of short-term ADT, the use of radiotherapy with short-term versus long-term ADT, or the addition of abiraterone acetate/prednisone to radiotherapy plus long-term ADT are based on clinical trial inclusion criteria, expert opinion, and estimates of absolute benefit and harm from a given therapy in the context of NCCN risk groups.

There are newer risk classification schemas that have been shown to outperform NCCN risk groups,^{118,119} as well as tools (ie, imaging, gene expression biomarkers) that together improve risk stratification. These tools should not be ordered reflexively. They are recommended only when they will have the ability to change management. Improved risk stratification can better identify patients who may derive greater or lesser absolute benefit from a given treatment.

NCCN Risk Groups

The NCCN Guidelines have, for many years, incorporated a risk stratification scheme that uses a minimum of stage, Gleason grade, and PSA to assign patients to risk groups. Risk group stratification has been published widely and validated, and provides a better basis for treatment recommendations than clinical stage alone.¹²⁰⁻¹²²

Prostate cancer grading continues to evolve. The grading system was updated during the 2014 International Society of Urological Pathology (ISUP) Consensus Conference.¹²³ Several changes were made to the assignment of Gleason pattern based on pathology. The new system assigns Grade Groups from 1 to 5, derived from the Gleason score.

- Grade Group 1: Gleason score ≤ 6 ; only individual discrete well-formed glands
- Grade Group 2: Gleason score $3+4=7$; predominantly well-formed glands with lesser component of poorly formed/fused/cirribriform glands
- Grade Group 3: Gleason score $4+3=7$; predominantly poorly formed/fused/cirribriform glands with lesser component of well-formed glands
 - For cases with $>95\%$ poorly formed/fused/cirribriform glands or lack of glands on a core or at radical prostatectomy, the component of $<5\%$ well-formed glands is not factored into the grade.
- Grade Group 4: Gleason score $4+4=8$; $3+5=8$; $5+3=8$
 - Only poorly formed/fused/cirribriform glands; or
 - Predominantly well-formed glands and lesser component lacking glands (poorly formed/fused/cirribriform glands can be a more minor component); or
 - Predominantly lacking glands and lesser component of well-formed glands (poorly formed/fused/cirribriform glands can be a more minor component)



NCCN Guidelines Version 5.2026

Prostate Cancer

- Grade Group 5: Gleason score 9–10; lack gland formation (or with necrosis) with or without poorly formed/fused/cribriform glands
 - For cases with >95% poorly formed/fused/cribriform glands or lack of glands on a core or at radical prostatectomy, the component of <5% well-formed glands is not factored into the grade.

The new Grade Group system was initially validated in two separate cohorts of patients treated for prostate cancer with either radical prostatectomy or radiation: one with >26,000 patients and one with 5880 patients.^{124,125} Both studies found that Grade Groups predicted the risk of recurrence after primary treatment with modest discrimination. For instance, in the larger study, the 5-year biochemical recurrence-free progression probabilities after radical prostatectomy for Grade Groups 1 through 5 were 96% (95% CI, 95–96), 88% (95% CI, 85–89), 63% (95% CI, 61–65), 48% (95% CI, 44–52), and 26% (95% CI, 23–30), respectively. The separation between Grade Groups was less pronounced in the radiation therapy (RT) cohort, likely because of increased use of ADT in the higher risk groups. In another study of the new ISUP Grade Group system, all-cause mortality and prostate cancer-specific mortality were higher in patients in Grade Group 5 than in those in Grade Group 4.¹²⁶ Additional studies have supported the validity of this new system.¹²⁷⁻¹³²

The NCCN Panel has accepted the Grade Group system to inform better treatment discussions compared to those using Gleason score. Patients remain divided into low-, intermediate-, high-, and very-high-risk groups. If a patient meets the criteria for intermediate risk, they are then further risk stratified into favorable and unfavorable subsets based on the number of intermediate risk factors, grade group, and the percentage of positive biopsy cores, as delineated in the algorithm above.

The NCCN Guidelines Panel recognized that heterogeneity exists within each risk group.¹¹⁸ For example, an analysis of 12,821 patients showed that those assigned to the intermediate-risk group by clinical stage (T2b–T2c) had a lower risk of recurrence than those categorized according to Gleason score (7) or PSA level (10–20 ng/mL).¹³³ A similar trend of superior recurrence-free survival was observed in patients placed in the high-risk group by clinical stage (T3a) compared to those assigned by Gleason score (8–10) or PSA level (>20 ng/mL), although it did not reach statistical significance. Other studies have reported differences in outcomes in the high-risk group depending on risk factors or primary Gleason pattern.^{134,135} Evidence also shows heterogeneity in the low-risk group, with PSA levels and percent positive cores affecting pathologic findings after radical prostatectomy.^{136,137}

In a retrospective study, 1024 patients with intermediate-risk prostate cancer were treated with radiation with or without neoadjuvant and concurrent ADT.¹³⁸ Multivariate analysis revealed that primary Gleason pattern 4, number of positive biopsy cores ≥50%, and presence of >1 intermediate-risk factors (IRFs; ie, T2b–c, PSA 10–20 ng/mL, Gleason score 7) were significant predictors of increased incidence of distant metastasis. The authors used these factors to separate the patients into unfavorable and favorable intermediate-risk groups and determined that the unfavorable intermediate-risk group had worse PSA recurrence-free survival and higher rates of distant metastasis and prostate cancer-specific mortality than the favorable intermediate-risk group. The use of active surveillance in patients with favorable intermediate-risk prostate cancer is discussed below (see *Active Surveillance in Favorable Intermediate Risk*). The NCCN Panel has included the separation of the intermediate risk group into favorable and unfavorable subsets in their risk stratification scheme.



NCCN Guidelines Version 5.2026

Prostate Cancer

Nomograms

A nomogram is a predictive instrument that takes a set of input data (variables) and makes predictions about an outcome. Nomograms may potentially predict more accurately for the individual patient than risk groups, because they utilize granular prognostic variables. The Partin tables are an early example of a nomogram for patients with clinically localized prostate cancer.¹³⁹⁻¹⁴² The tables give the probability (95% CI) that a patient with a certain clinical stage, Gleason score, and PSA will have a cancer of each pathologic stage. Nomograms can be used to inform treatment decision-making for patients contemplating active surveillance,¹⁴³⁻¹⁴⁵ radical prostatectomy,¹⁴⁶⁻¹⁴⁹ neurovascular bundle preservation¹⁵⁰⁻¹⁵² or omission of pelvic lymph node dissection (PLND) during radical prostatectomy,¹⁵³⁻¹⁵⁶ brachytherapy,^{146,157-159} or external beam RT (EBRT).^{146,160} Biochemical progression-free survival (PFS) can be reassessed postoperatively using age, diagnostic serum PSA, and pathologic grade and stage.^{146,161-163} Potential success of adjuvant or post-recurrence RT after unsuccessful radical prostatectomy can be assessed using a nomogram.^{146,164}

None of the current models predicts with perfect accuracy, and only some of these models predict metastasis^{145,146,161,165,166} and cancer-specific death.^{147,149,167-169} Given the competing causes of mortality, many patients who sustain PSA recurrence will not live long enough to develop clinical evidence of distant metastases or to die from prostate cancer. Not all PSA recurrences are clinically relevant; thus, PSADT may be a more useful measure of risk of death.¹⁷⁰ The NCCN Guidelines Panel recommends that NCCN risk groups be used to begin the discussion of options for the treatment of clinically localized prostate cancer and that nomograms be used to provide additional and more individualized information.

Advanced Risk Stratification Tools

The Panel strongly advocates the maximum utilization of all routine clinical, pathologic, and patient information to risk stratify patients. This includes the information that forms the basis of the NCCN risk groups, as well as other routine information that can be derived from standard tests, such as PSA density, PSA velocity, percentage or absolute pattern 4, and intraductal and cribriform architecture. Other tools have been developed that, variably, have been shown to be superior to standard clinical tools at estimating prognosis, and these are referred to as advanced risk stratification tools. These currently include multigene expression tests and digital pathology-based artificial intelligence tests, as discussed in more detail below.

These advanced risk stratification tools should only be considered when they have the potential ability to change disease management; they should not be ordered reflexively. Importantly, the Panel notes that there are risks to using tools that lack robust validation to change management decisions, as they may drive patients or providers to choose inappropriate options. For example, in one trial, 200 patients with very low to favorable-intermediate risk prostate cancer (excluding those with grade group 2 and >3 positive cores and limiting inclusion of those with PSA 10–20 ng/mL to only those with PSA density <0.15) were randomized to standard counseling with or without the 17-gene genomic prostate score (GPS) assay.¹⁷¹ Patients with lower health literacy who received the GPS test were 7-fold less likely to choose active surveillance compared to controls. Therefore, if advanced tools are used, the Panel recommends only tools that have high-quality, long-term clinical trial data, ideally across multiple clinical trials.

Currently, the Panel recommends using tests with high-quality, prospective validation. For predictive tools, ideally this validation is from dedicated prospective biomarker-focused trials (eg, NRG GU006).



NCCN Guidelines Version 5.2026

Prostate Cancer

Alternatively, this can be from post-hoc assessment of the biomarker in completed prospective trials, ideally multiple such trials. For prognostic tools, the Panel generally recommends the use of prognostic tests that are validated in well-designed prospective studies with clinically meaningful endpoints based on disease settings that guide a specific treatment indication based on a specific score or result. These studies can be either prospective integral or integrated clinical trial(s) (eg, NRG GU009, NRG GU010) or post-hoc correlative analyses of prospective trials.

Tumor Multigene Expression Testing

Gene testing of a tumor offers the potential of added insight into the biologic behavior of a cancer that could thereby aid in clinical decision-making.

Several tissue-based molecular assays have been developed to improve risk stratification. The 22-gene genomic classifier (GC; Decipher) is discussed in more detail in the *Principles of Risk Stratification* in the algorithm above; others are discussed below. The training and development of each assay is distinct, the genes analyzed are unique, and there is heterogeneity in the robustness of validation. It is important to understand that companies may make invalid or inaccurate claims about these tests, which can create confusion for patients and providers. Furthermore, the resulting test score, such as low or high risk, may be discordant between different tests due to multiple factors.

It is clear that use of tissue-based molecular assays will often change decisions about disease management.¹⁷²⁻¹⁷⁵ However, what is unclear from such studies is if the change in management improved patient outcomes or was appropriate. Furthermore, it is increasingly clear that each assay should be evaluated independently because they do not capture the same biology. For example, a study of over 50,000 patients was performed to compare three commonly used gene expression

signatures via creation of derived signatures for each test, and the correlation between the signatures was poor (R^2 , 0.32–0.36).¹⁷⁶

While multiple gene expression tests have made claims to improve the safety and/or efficacy of active surveillance, such an improvement has not been born out in prospective studies. For example, evaluation of diagnostic biopsy tissue from patients enrolled in the Canary PASS multicenter active surveillance cohort suggested that results of a molecular assay were not associated with adverse pathology in combination with clinical variables nor was there an association with upgrading in surveillance biopsies.¹⁷⁷ While this study utilized GPS, no other gene expression or digital pathology-based tool has positive prospective validation in this setting.

Currently, the above-mentioned validation criteria for prognostic biomarker tests has been reached by the 22-gene GC prognostic assay, which has reported outcomes from multiple post-hoc analyses of randomized trials.¹⁷⁸⁻¹⁸¹ The relevant disease settings and more details for this tool are available in the *Principles of Risk Stratification* in the algorithm above. There are multiple biomarker-directed randomized trials utilizing the 22-gene GC assay that have completed enrollment or are ongoing (eg, NRG GU006, 009, 010).

Multiple retrospective studies suggest that the 17-gene GPS tool (previously called Oncotype Dx) may be prognostic for patients with localized prostate cancer.^{182,183} GPS was also studied in the prospective Canary PASS active surveillance cohort with post-hoc biomarker analysis.¹⁷⁷ GPS results were obtained from 432 patients, 101 of whom underwent radical prostatectomy after an initial period of active surveillance. The authors concluded that adding GPS to a model containing PSA density and diagnostic grade group did not significantly improve adverse pathology stratification over the clinical variables alone (HR, 1.17; 95% CI, 1.00–1.43; $P = .066$). Additionally, there was no



NCCN Guidelines Version 5.2026

Prostate Cancer

association observed between GPS and subsequent biopsy upgrade ($P = 0.48$). Event rates and sample size may have impacted the results. Additionally, as previously referenced, the GPS test was evaluated in a randomized trial and demonstrated that it decreased the relative odds of choosing active surveillance by approximately 50% with variable statistical significance depending on analysis method ($P = .029$ when excluding participants with inadequate biopsy specimens who did not receive a planned GPS result; $P = .067$ for all patients in an intention-to-treat analysis).¹⁷¹

Similarly, multiple retrospective institutional and tumor registry studies suggest that the 31-gene cell cycle progression test (CCP; Prolaris) may have prognostic value for patients with localized prostate cancer.¹⁸⁴⁻¹⁸⁶ No prospective trials have been published to date.

Numerous other tests have been developed with variable evidence quality that are currently outside of the Discussion presented here.

Digital Pathology-Based Tools

More recently a new class of advanced risk stratification tool has been developed that utilizes artificial intelligence (AI) to analyze digital pathology slides from a patient's prostate biopsy or prostatectomy specimen. Analogous to the multigene prognostic assays, these tools are primarily focused to improve risk stratification. The multimodal AI (MMAI) test (ArteraAI Prostate) is one such test, which gained FDA approval in July 2025. The MMAI test was validated in multiple post-hoc analyses of completed randomized trials.¹⁸⁷⁻¹⁸⁹

The MMAI test also has developed a potential predictive model to guide the use of short-term ADT. The predictive model was validated in RTOG 9408, a randomized trial of low-dose radiotherapy with or without 4 months of ADT, and was shown to identify patients more likely to benefit from ADT.¹⁹⁰ A similar, but unique model also was developed that may predict

the benefit of short-term versus long-term ADT and validated in RTOG 9202.¹⁹¹ Given the considerable changes in staging, grading, and treatment since the start of RTOG 9202 and 9408 (>30 years ago), it is unclear if the predictive models still work in contemporarily treated patients, and further validation is recommended by the Panel.

The relevant disease settings and more details for this tool are available in the *Principles of Risk Stratification* in the algorithm above.

Initial Clinical Assessment and Staging Evaluation

For patients with a life expectancy of ≤ 5 years without clinical symptoms who have low- and intermediate-risk prostate cancer, further imaging and treatment should be delayed until symptoms develop, at which time imaging can be performed and ADT should be given. Those with a life expectancy ≤ 5 years who fall into the high- or very-high-risk groups should undergo bone imaging and, if indicated by nomogram prediction of lymph node involvement, pelvic +/- abdominal imaging.

For symptomatic patients and those with unfavorable intermediate-risk, high-risk, and very-high-risk prostate cancer and a life expectancy of >5 years, bone and soft tissue imaging is appropriate:

- PSMA-PET/CT or PSMA-PET/MRI can be considered for bone and soft tissue (full body) imaging.
 - Because of the increased sensitivity and specificity of PSMA-PET tracers for detecting micrometastatic disease compared to CT, MRI, and bone scan at both initial staging and biochemical recurrence, PSMA-PET/CT or PSMA-PET/MRI can serve as a more effective front-line imaging tool for these patients.
- Bone imaging can be achieved by conventional technetium-99m-MDP bone scan.



NCCN Guidelines Version 5.2026

Prostate Cancer

- Plain films, CT, MRI, PSMA-PET/CT or PSMA PET/MRI, or PET/CT or PET/MRI with F-18 sodium fluoride, C-11 choline, F-18 fluciclovine can be considered for equivocal results on initial bone imaging.
- Soft tissue imaging of the pelvis, abdomen, and chest can include chest CT and abdominal/pelvic CT or abdominal/pelvic MRI. Multiparametric MRI (mpMRI) is preferred over CT for pelvic staging.

mpMRI may detect larger and clinically significant prostate cancer (Grade Group ≥ 2), extracapsular extension, and seminal vesicle invasion (T staging) and is preferred over CT for abdominal/pelvic staging. mpMRI has been shown to be equivalent to CT scan for pelvic lymph node evaluation.

See *Imaging Techniques* below for a more detailed discussion.

Imaging Techniques

Imaging techniques are useful for staging and for detecting metastases and tumor recurrence. Current clinical imaging techniques for prostate cancer include conventional radiography (ie, x-rays), ultrasound, CT, MRI, single photon emission computed tomography (SPECT, scintigraphy), and PET. Some of these modalities have the ability to assess both anatomy and tumor function/biology. For example, functional MR sequences can be added to conventional anatomic MR sequences in a clinical examination such as diffusion-weighted imaging (DWI) to assess tumor cellularity or MR spectroscopy (MRS) to assess tumor metabolism.

Different modalities can also be merged to maximize prostate cancer assessment. For example, the functional information obtained with PET can be combined with the spatial and anatomic information from either CT (ie, PET/CT) or MRI (ie, PET/MRI) to inform about the locations of tumor foci for diagnosis or therapy response. Another example of the advantage of combining modalities is MR-ultrasound fusion guided biopsy (eg, MR-

TRUS), where MRI datasets containing information on suspicious lesions identified by the radiologist are used by the urologist to navigate ultrasound-guided biopsies of the prostate for more accurate diagnosis.¹⁹² More details on each technique are outlined in the algorithm under *Principles of Imaging*, above.

Multiparametric MRI (mpMRI)

The use of mpMRI in the staging and characterization of prostate cancer has increased in the last few years. mpMRI examinations typically include three sequences: T2-weighted imaging, DWI, and dynamic contrast enhancement (DCE) imaging. There has been increased interest in biparametric imaging that excludes the use of gadolinium contrast in prostate MRI examinations; however, more data are needed to identify the risk groups who would benefit most from this approach.¹⁹³ In general, it is recommended that mpMRI be performed on a 3 Tesla (3T) magnetic strength MRI scanner. This is the highest strength scanner in routine clinical use and provides the best possible evaluation of prostate cancer.

Additional instrumentation can be used, such as an endorectal coil (ERC) to improve image quality. If a lower strength (eg, 1.5T) MRI is required for a patient because of indwelling medical device incompatibility with 3T MRI, an ERC is recommended. Use of ERC in routine prostate imaging is controversial. Current data suggest that a 3T exam with ERC may not be significantly better than a 3T exam without ERC. Moreover, there may not be a significant difference in image interpretation between a 1.5T with ERC and 3T without ERC.¹⁹⁴ The use of ERC in prostate MRI also introduces new problems into the clinical workflow including patient discomfort, prostate distortion, increased scanner time and expense, and requirement of someone experienced to place the ERC.

Evidence supports the implementation of mpMRI in several aspects of prostate cancer management.¹⁹² *First*, mpMRI helps detect larger and/or



NCCN Guidelines Version 5.2026

Prostate Cancer

more clinically significant cancers (ie, Grade Group ≥ 2).¹⁹⁵ mpMRI has been incorporated into MRI-TRUS fusion-targeted biopsy protocols, which has led to an increase in the diagnosis of high-grade cancers with fewer biopsy cores, while reducing detection of low-grade and insignificant cancers.¹⁹⁶⁻¹⁹⁸ In fact, a recently published clinical study identified that MRI-targeted biopsy synergized with conventional systematic biopsy to identify more clinically significant cancers.¹⁹⁹ *Second*, mpMRI aids in better assessment of extracapsular extension (T staging), with high negative predictive values (NPVs) in patients with low-risk disease.²⁰⁰ mpMRI results may inform decision-making regarding nerve-sparing operation.²⁰¹ *Third*, mpMRI is equivalent to CT scan for staging of pelvic lymph nodes.^{202,203} *Finally*, mpMRI outperforms bone scan and targeted x-rays for detection of bone metastases, with a sensitivity of 98% to 100% and specificity of 98% to 100% (vs. sensitivity of 86% and specificity of 98%–100% for bone scan plus targeted x-rays).²⁰⁴

PET Imaging

The use of PET/CT or PET/MRI imaging using tracers other than F-18 fluorodeoxyglucose (FDG) for staging of small-volume recurrent or metastatic prostate cancer has rapidly expanded in recent years.¹⁹² Currently, there are several PET tracers that are FDA approved for use in patients with prostate cancer: Ga-68 PSMA-11 (PSMA-HBED-CC), F-18 piflufolastat (DCFPyL), F-18 flutufolastat PSMA (rh-PSMA-7.3), C-11 choline, F-18 fluciclovine, and F-18 sodium fluoride. Although these tracers are approved for the evaluation of patients with biochemical recurrence, the PSMA tracers are also approved for patients at initial staging with suspected metastatic disease. Tracer distribution in patients with prostate cancer can be imaged with either PET/CT or PET/MRI modalities. Although CT and MRI are equivalent in the assessment of lymphadenopathy, PET/MRI has the added advantage over PET/CT with enhanced tissue contrast that is especially important in evaluation of pelvic anatomy and prostate cancer assessment. Table 1 in the *Principles of*

Imaging above summarizes the FDA-cleared PET imaging tracers studied in prostate cancer.

PSMA-PET refers to a growing body of radiopharmaceuticals that target prostate specific membrane antigen (PSMA) on the surface of prostate cells. Because of the high density of PSMA receptors on the surface of cancer cells relative to adjacent prostate, PSMA-PET has the advantage of high signal-to-noise relative to adjacent tissues. The mechanistic role of androgen receptor signaling in PSMA regulation is still being investigated, as multiple reports in animals and humans suggest that androgen modulation can affect PSMA expression and may even be dichotomous in patients with castration-sensitive versus castrate-resistant disease.²⁰⁵⁻²⁰⁷ There are multiple PSMA radiopharmaceuticals at various stages of investigation. At this time, the NCCN Guidelines only recommend the currently FDA-approved PSMA tracers: F-18 piflufolastat, F-18 flutufolastat, and Ga-68 PSMA-11. PSMA-PET/CT or PSMA-PET/MRI can be considered as an alternative to standard imaging of bone and soft tissue for initial staging, the detection of biochemically recurrent disease, and as workup for progression with bone scan plus CT or MRI for the evaluation of bone, pelvis, and abdomen.

Studies suggest that PSMA-PET imaging has a higher sensitivity than C-11 choline or F-18 fluciclovine PET imaging, especially at very low PSA levels.²⁰⁸⁻²¹⁵ The reported sensitivity and specificity for PSMA-11 PET/CT in the detection of nodal involvement in primary staging of patients with intermediate-, high-, and very-high-risk disease is 40% and 95%, respectively.²¹⁶ The patient-level positive predictive value (PPV) in detection of lesions in patients with biochemical recurrence (BCR) is 92%.²¹⁷ Similarly, the reported sensitivity and specificity for piflufolastat PET/CT in the detection of nodal involvement in primary staging of patients with unfavorable intermediate-, high-, and very-high-risk disease are 31% to 42% and 96% to 99%, respectively.^{218,219} The patient-level



NCCN Guidelines Version 5.2026

Prostate Cancer

correct localization rate (CLR; patient-level PPV validated by anatomic lesion co-localization) for piflufolastat PET/CT is 85% to 87%.²²⁰ Thus, PSMA-11 and piflufolastat are considered equivalent. F-18 flutemetastat PSMA is part of a novel class of tracers referred to as radiohybrid (rh) ligands. These rh ligands have the unique advantage of offering two binding sites for radionuclides (ie, F-18 or Ga-68), which increases its flexibility in imaging. In addition, the presence of a chelator in these rh ligands also allows for chelation of Lu-177 for its use as a theranostic as well as imaging agent.

Because of the increased sensitivity and specificity of PSMA-PET tracers for detecting micrometastatic disease compared to CT, MRI, and bone scan at both initial staging and biochemical recurrence, PSMA PET/CT or PSMA-PET/MRI can serve as a more effective front-line imaging tool for these patients.

PET/CT or PET/MRI detect small-volume disease in bone and soft tissues.^{221,222} The reported sensitivity and specificity of C-11 choline PET/CT in restaging patients with biochemical recurrence ranges from 32% to 93% and from 40% to 93%, respectively.²²³⁻²³² The reported sensitivity and specificity of F-18 fluciclovine PET/CT ranges from 37% to 90% and from 40% to 100%, respectively.^{229,233,234} A prospective study compared F-18 fluciclovine and C-11 choline PET/CT scans in 89 patients, and agreement was 85%.²²⁹ Thus, choline and fluciclovine are considered equivalent in the evaluation of patients with biochemical recurrence. The Panel believes that F-18 fluciclovine PET/CT or PET/MRI or C-11 choline PET/CT or PET/MRI may be used in patients with biochemical recurrence after primary treatment for further soft tissue and/or bone evaluation after bone scan, chest CT, and abdominal/pelvic CT or abdominal/pelvic MRI.

The use of these PET tracers can lead to changes in clinical management. The FALCON trial showed that results of F-18 fluciclovine PET/CT in 104 patients with biochemical recurrence after definitive therapy resulted in a

change in disease management for 64% of patients.²³⁵ In addition, the LOCATE trial demonstrated that fluciclovine frequently changed disease management plans for patients with biochemical recurrence.²³⁶ In a similar fashion, data also show that PSMA-PET has the ability to change radiation treatment planning in 53% (N = 45) of patients with high- and very-high-risk prostate cancer using PSMA-11 as well as change disease management in over half of a prospective cohort of 635 patients with BCR.^{237,238} However, whether changes to treatment planning because of PET tracers have an impact on long-term survival remains to be studied.

The increasing use of PSMA-PET has identified the potential for considerable biological diversity among disease foci within a given individual with prostate cancer, especially in the mCRPC setting. This heterogeneity can be detected with a combination of PSMA-PET and fluorodeoxyglucose (FDG)-PET. Initial data suggest that metastases with PSMA-negative/FDG-positive mismatches may exist in patients with mCRPC undergoing Lu-177-PSMA-617 radioligand therapy and that patients with these mismatches may have worse outcomes. To overcome the limitations of PSMA-PET, the Panel currently recommends the use of MRI or a dedicated high resolution diagnostic contrast-enhanced CT in those with a negative PSMA-PET scan, because the non-contrast CT component of PSMA-PET/CT is insufficient to detect visceral metastatic disease, especially in the liver, due to its high physiological uptake.

F-18 sodium fluoride targets osteoblast activity where the fluoride is deposited into new bone formation, thus limiting use of this agent to the detection of osseous metastases. Fluoride PET/CT has greater sensitivity than standard bone scintigraphy in the detection of bone metastases, with 77% to 94% sensitivity, 92% to 99% specificity, and 82% to 97% PPV.²³⁹ However, emerging evidence indicates that other tracers such as PSMA are at least equivalent to fluoride in the detection of osseous metastases with the added advantage of soft tissue metastasis detection.²⁴⁰



NCCN Guidelines Version 5.2026

Prostate Cancer

Histologic or radiographic confirmation of involvement detected by PET imaging is recommended whenever feasible due to the presence of false positives. Although false positives exist, literature suggests that these are outweighed by the increase in true positives detected by PET relative to bone scintigraphy. To reduce the false-positive rate, physicians should consider the intensity of PSMA-PET uptake and correlative CT findings in the interpretation of scans. Several reporting systems have been proposed but have not been validated or widely used.^{241,242} Moreover, although PET imaging may change treatment,²³⁶ it may not change oncologic outcome. Earlier detection of bone metastatic disease, for instance, may result in earlier use of newer and more expensive therapies, which may not improve oncologic outcomes or OS.

Risks of Imaging

Medical imaging is a critical tool in the evaluation and comprehensive care of patients with malignancy. However, as with any medical procedure, imaging is not without risk. Some of these risks are concrete and tangible, while others are less clear. Risks associated with imaging include exposure to ionizing radiation, adverse reaction to contrast media, false-positive scans, and overdetection. Inappropriate use of imaging also has been identified as a significant contributor to health care costs in the United States and worldwide. Therefore, imaging should be performed only when medically appropriate, and in a manner that reduces risk (eg, minimizing radiation dose). An algorithmic approach to the use of imaging, such as by NCCN and the Appropriateness Criteria developed by the American College of Radiology,²⁴³ can assist in medical decision-making.

Exposure to Ionizing Radiation

Deterministic and stochastic are two types of effects from exposure to ionizing radiation by x-ray, CT, or PET/CT. Deterministic effects are those that occur at a certain dose level and include events such as cataracts and radiation burns. No effect is seen below the dose threshold. Medical

imaging is always performed almost below the threshold for deterministic effects. Stochastic effects tend to occur late, increase in likelihood as dose increases, and have no known lower “safe” limit. The major stochastic effect of concern in medical imaging is radiation-induced malignancy. Unfortunately, no direct measurements are available to determine risk of cancer arising from one or more medical imaging events, so risks are calculated using other models (such as from survivors of radiation exposure). The literature is conflicting about the precise risk of secondary malignancies in patients undergoing medical imaging procedures. There is a small but finite risk of developing secondary malignancies because of medical imaging procedures, and the risk is greatest in young patients. However, the absolute risk of fatal malignancy arising from a medical imaging procedure is very low and is difficult to detect given the prevalence of cancer in the population and the multiple factors that contribute to oncogenesis.²⁴⁴ Efforts should be made to minimize dose from these procedures, which begin with judicious use of imaging only when justified by the clinical situation. Harm may arise from not imaging a patient, from disease non-detection or erroneous staging.

Adverse Reaction to Contrast Media

Many imaging studies use contrast material delivered by oral, intravenous, or rectal routes. The use of contrast material may improve study performance, but reactions to contrast material may occur and they should therefore be used only when warranted. Some patients develop adverse reactions to iodinated intravenous contrast material. Most reactions are mild cutaneous reactions (eg, urticaria, pruritus) but occasionally severe reactions can be life-threatening (bronchospasm or anaphylaxis). The risk of severe reaction is low with non-ionic contrast materials.²⁴⁵ Both iodinated CT contrast material and gadolinium-based MR contrast materials can be problematic in patients with reduced renal function. Gadolinium MR contrast media, in particular, is contraindicated in patients with acute renal failure or stage V chronic kidney disease (glomerular



NCCN Guidelines Version 5.2026

Prostate Cancer

filtration rate [GFR] <15).²⁴⁵ Patients in this category are significantly more likely to develop nephrogenic systemic fibrosis (NSF). Centers performing imaging studies with contrast materials should have policies in place to address the use of contrast in these patients.

False-Positive Scans and Overdetection

Every imaging test has limitations for sensitivity, specificity, and accuracy that involve both the nature of the imaging modality as well as the interpretation by the physician. Harm can arise when a tumor or tumor recurrence is not detected (ie, false negative), but harm to the patient and added expense to the medical system also can result from false-positive scans. Extensive workup of imaging findings that may be benign or indolent (ie, overdetection) can lead to significant patient anxiety, additional and unnecessary imaging, and invasive procedures that carry their own risks for adverse outcomes.

Accurate and medically relevant interpretation of imaging studies requires familiarity and expertise in the imaging modality, attention to detail in image review, knowledge of tumor biology, and familiarity with treatment options and algorithms. Challenging cases are best addressed through direct communication, either physician-to-physician or in a multidisciplinary tumor board setting.

The increased false positive detection of benign rib lesions with PSMA PET/CT is well established. If these are called metastatic foci then patients may be inappropriately recategorized from localized prostate cancer where they may receive curative-intent local therapy, to now receive doublet systemic therapy. Thus, expert interpretation of PSMA PET/CT scans is essential to avoid misclassification.

Observation

Observation involves monitoring the course of prostate cancer with a history and physical exam no more often than every 12 months (without

surveillance biopsies) until symptoms develop or are thought to be imminent. If patients under observation become symptomatic, an assessment of disease burden can be performed, and treatment or palliation can be considered. Observation thus differs from active surveillance. The goal of observation is to maintain quality of life (QOL) by avoiding noncurative treatment when prostate cancer is unlikely to cause mortality or significant morbidity. The main advantage of observation is avoidance of possible side effects of routine biopsies and unnecessary definitive therapy or ADT. However, patients may develop urinary retention or pathologic fracture without prior symptoms or increasing PSA level.

Observation is applicable to patients who are older or in declining health with comorbidity that are more likely than prostate cancer to be the cause of death. Johansson and colleagues²⁴⁶ observed that only 13% of patients developed metastases 15 years after diagnosis of T0–T2 disease and only 11% had died from prostate cancer. Because prostate cancer will not be treated for cure for patients with shorter life expectancies, observation for as long as possible is a reasonable option based on physician discretion. When symptoms develop or are imminent, patients can begin palliative treatment.

The SPCG-4 and PIVOT trials compared the use of observation versus radical prostatectomy (RP) in patients with localized prostate cancer.^{247–249} Results from SPCG-4 did not demonstrate a significant reduction in prostate cancer-specific mortality with RP compared to observation in patients with low-risk disease.²⁴⁷ In PIVOT, there was no difference in all-cause mortality between these two management strategies in those with low-risk disease, strongly implying that treatment of low-risk prostate cancer should generally be avoided. RP was superior to observation for patients with intermediate-risk disease. However, RP did not significantly improve death from any cause in those with high-risk disease in either trial.^{247,249}



NCCN Guidelines Version 5.2026

Prostate Cancer

Active Surveillance

The NCCN Guidelines Panel remains concerned about the problems of overtreatment related to the increased frequency of diagnosis of prostate cancer from widespread use of PSA for early detection or screening (see the NCCN Guidelines for Prostate Cancer Early Detection, available at www.NCCN.org). The debate about whether it is necessary to diagnose and treat every individual who has prostate cancer is fueled by the high prevalence of prostate cancer upon autopsy of the prostate²⁵⁰; the high frequency of positive prostate biopsies in individuals with normal DREs and serum PSA values²⁵¹; the contrast between the incidence and mortality rates of prostate cancer; and the need to treat an estimated 37 patients with screen-detected prostate cancer^{252,253} or 100 patients with low-risk prostate cancer²⁵⁴ to prevent one death from the disease.

Several large, population-based prostate cancer screening trials have shown that properly applied early detection protocols can reduce prostate cancer mortality (see the NCCN Guidelines for Prostate Cancer Early Detection, available at www.NCCN.org). However, wide-spread screening has led to overdiagnosis of clinically insignificant prostate cancer (grade group 1). A growing body of high-level evidence supports the use of mpMRI to reduce unnecessary biopsies in individuals with an elevated PSA who are not likely to have clinically significant prostate cancer. Its use should decrease the risk of overdiagnosis while maintaining the detection of potentially fatal prostate cancer. These improvements coupled with the use of active surveillance in appropriate patients should reduce overtreatment while preserving the relatively low rates of prostate cancer mortality.

Models of prostate cancer detection and progression from the early 2000's estimated that 23% to 42% of all U.S. screen-detected cancers were overtreated²⁵⁵ and that PSA detection was responsible for up to 12.3 years of lead-time bias.²⁵⁶ Encouragingly, the proportion of patients with low-risk

prostate cancer choosing active surveillance has increased over time, and was at approximately 60% in 2021.²⁵⁷⁻²⁵⁹ This trend has also been observed in the Veterans Affairs Integrated Health Care System, in which active surveillance increased from 2005 to 2015: from 4% to 39% of those <65 years and from 3% to 41% of those ≥65 years.²⁶⁰

Active surveillance involves actively monitoring the course of the disease with the expectation to deliver curative therapy if the cancer progresses. Unlike observation, active surveillance is mainly applicable to patients with sufficient life expectancy who may benefit from early detection of cancer progression and radical treatment. Because these patients have a longer life expectancy, they should be followed closely, and treatment should start promptly should the cancer progress so as not to miss the chance for cure.

Early versions of active surveillance were termed active monitoring. This approach differs from active surveillance as it primarily only relies on PSA monitoring, whereas active surveillance typically involved serial prostate biopsies and potential for serial imaging.

The ProtecT study, which randomized 1643 patients with localized prostate cancer to active monitoring, radical prostatectomy, or RT, found no significant difference in the primary outcome of prostate cancer mortality at a median of 10 years follow-up.²⁶¹ Of 17 prostate cancer deaths (1% of study participants), 8 were in the active surveillance group, 5 were in the surgery group, and 4 were in the radiation group ($P = .48$ for the overall comparison). However, a 12.2% absolute increase in the rate of disease progression and a 3.4% absolute increase in the rate of metastases or prostate cancer death were seen in the active surveillance group.^{261,262} Approximately 23% of participants had Gleason scores 7–10, and 5 of 8 deaths in the active surveillance group were in this subset. Longer follow-up has been reported with a median follow-up of 15-years, and overall survival remains similar across the 3 groups.²⁶³ However,



NCCN Guidelines Version 5.2026

Prostate Cancer

metastases were more common in the active monitoring group (9.4% vs. ~5% in the surgery and RT groups), primarily driven by the patients with intermediate-risk disease who were included in the study. By 15-years, approximately 70% of patients in the active monitoring arm had received treatment.

An updated analysis from the ProtecT trial of 712 patients that had pathology available for re-review was reported.²⁶⁴ The authors demonstrated that 13.1% of patients had cribriform-positive disease, which significantly increased the risk of developing metastases (HR, 3.61; 95% CI, 1.60–8.11). RT with neoadjuvant ADT significantly reduced metastasis risk (HR, 0.35; 95% CI, 0.16–0.78; $P = .04$), whereas surgery did not significantly improve long-term outcomes compared with active monitoring (15-year cumulative incidence in patients with cribriform-positive disease: 26% for surgery, 25% for active monitoring, and 8% for RT plus ADT).

Patient-reported outcomes were compared among the 3 groups of ProtecT.^{25,265} The surgery group experienced the greatest negative effect on sexual function and urinary continence, whereas the RT group had a slight decrease in bowel function. It has been noted that the RT utilized in the trial was 3D radiotherapy without image guidance or use of hydrogel spacers.²⁶⁶ In addition, studies have shown that active surveillance does not adversely impact psychological well-being or QOL.^{25,267-271}

Several large active surveillance cohort studies have shown that >50% of those eligible for active surveillance may safely avoid treatment, and thus the possible associated side effects of treatment, for at least 10 years (see *Table 1* below).] For example, in one study, 55% of the population remained untreated at 15 years.²⁷² Although a proportion of patients on active surveillance will eventually undergo treatment, the delay does not appear to impact cure rates, and numerous studies have shown that active surveillance can be a safe option for many patients.²⁷²⁻²⁸² In fact, a 2015 meta-analysis of 26 active surveillance cohort studies that included 7627

patients identified only 8 prostate cancer deaths and 5 cases of metastasis.²⁸³

For low-risk prostate cancer, active surveillance is preferred.

The favorable intermediate risk grouping was not developed to identify optimal active surveillance, but rather to identify patients with intermediate-risk disease who can safely receive RT alone without ADT. The favorable-intermediate-risk group includes patients that are well suited for active surveillance, such as those with Grade group 1 and PSA <10 ng/mL, with a low PSA density. However, it also includes patients with multiple cores of Grade group 2 disease with cribriform features and a high PSA density. Patients should be counseled that unselected Grade group 2 intermediate-risk prostate cancer has been shown to have unacceptably high risks of metastasis when managed by observation, active monitoring, or active surveillance.²⁸⁴ Patients with low-volume disease (ie, 1–2 positive cores), low percent or absolute Gleason pattern 4, lack of cribriform features, and low PSA density are likely better candidates for active surveillance among patients with Grade group 2 disease.²⁶⁴

Therefore, for select patients with favorable intermediate-risk disease, a recommendation for active surveillance must be based on careful individualized weighing of several factors: life expectancy, general health condition, disease characteristics, potential side effects of treatment, and patient preference. Shared decision-making, after appropriate counseling on the risks and benefits of the various options, is critical. Data in these settings are discussed under *Patient Selection*, below.

The Panel believes there is an urgent need for further clinical research regarding the criteria for recommending active surveillance, the criteria for reclassification on active surveillance, and the schedule for active surveillance especially as it pertains to prostate biopsies, which pose an increasing burden.



NCCN Guidelines Version 5.2026

Prostate Cancer

Patient Selection

NCCN risk groups are the main determinant used to select patients who can safely defer treatment.

Active Surveillance in Low-Risk Disease

Panel consensus is that active surveillance is preferred for most patients with low-risk prostate cancer and a life expectancy ≥ 10 years based on the data discussed above. However, the Panel recognizes that there is heterogeneity across the low-risk group, and that some factors may be associated with an increased probability of near-term grade reclassification including high PSA density, a high number of positive cores (eg, ≥ 3), high gene expression risk (from tissue-based molecular tumor analysis).²⁸⁵⁻²⁸⁷ Of note, core involvement in the major active surveillance cohort studies was generally low (see *Table 1* below) and very few patients with grade group 2 disease were included. Therefore, in very select patients with low-risk prostate cancer, upfront treatment with radical prostatectomy or RT may be appropriate based on shared decision-making with the patient. Patient anxiety should be addressed so that it is not a deterrent to the appropriate use of active surveillance.

Active Surveillance in Favorable Intermediate-Risk Disease

The literature on outcomes of active surveillance in patients with intermediate-risk prostate cancer is limited because few prospective studies of active surveillance included patients with intermediate-risk prostate cancer (see *Table 1*, below).

In prospective studies that included patients with intermediate-risk disease, favorable prostate cancer-specific survival rates of 94% to 100% were seen for the full cohorts.^{272,274,276} However, with extended follow-up, the Toronto group has demonstrated inferior metastasis-free survival for patients with intermediate-risk prostate cancer (15-year metastasis-free survival for cases of Gleason 6 or less with PSA < 10 ng/mL, 94%;

Gleason 6 or less with PSA 10–20 ng/mL, 94%; Gleason 3+4 with PSA 20 ng/mL or less, 84%; and Gleason 4+3 with PSA 20 ng/mL or less, 63%).²⁸⁸ Furthermore, there was an unacceptably high 15-year distant metastasis rate (16.4%) that largely occurred between 10 to 15 years after diagnosis.²⁸⁸ Likewise, Canary PASS included a small subset of patients with grade group 2 disease, but only 2 patients had 15 years of follow-up. Two patients with grade group 2 prostate cancer developed metastatic disease between 9 and 12 years,²⁸⁹ with an estimated 15-year risk of distant metastasis of approximately 10%.²⁸⁴ Thus, 5 to 10 years of follow-up may be insufficient to assess active surveillance safety.

Overall, the Panel interpreted these data to show that a subset of patients with favorable intermediate-risk prostate cancer and life expectancy > 10 years may be considered for active surveillance, especially with grade group 1 disease. However, the precise inclusion criteria and follow-up protocols need continued refinement. Patients must understand that a significant proportion of those clinically staged as having favorable intermediate-risk prostate cancer may have higher risk disease.²⁹⁰⁻²⁹³ Particular consideration to active surveillance may be appropriate for those patients with a low percentage of Gleason pattern 4 cancer, low tumor volume, low PSA density, and/or low genomic risk (from tissue-based molecular tumor analysis), but should be approached with caution, include informed decision-making, and use close monitoring for progression. How to optimize active surveillance for patients with grade group 2 prostate cancer remains an active area of investigation.

Confirmatory Testing

Confirmatory testing is critical to the process of active surveillance because a subset of patients will be found to no longer be candidates for active surveillance after confirmatory testing. Active surveillance starts after confirmatory testing. Since an initial prostate biopsy may underestimate tumor grade or volume, confirmatory testing is strongly



NCCN Guidelines Version 5.2026

Prostate Cancer

recommended within the first 6 to 12 months of diagnosis for patients who are considering active surveillance.

Before starting on an active surveillance program, mpMRI with calculation of PSA density should be considered to confirm candidacy for active surveillance if not performed during initial workup.²⁹⁴ Patients with PI-RADS 4 or 5 on mpMRI have an increased risk of biopsy progression during active surveillance.²⁹⁵

In patients with low- and favorable-intermediate-risk disease, molecular tumor analysis has not been shown in prospective studies to clearly improve long-term outcomes for patients receiving active surveillance. One study examined the role of molecular tumor analysis for predicting upgrading on surveillance biopsy or the presence of adverse pathology on eventual radical prostatectomy in patients in an active surveillance cohort.¹⁷⁷ In this study, results of the molecular testing did not significantly improve risk stratification over the use of clinical variables alone.

If results of mpMRI and/or molecular testing are concerning, a repeat biopsy may be appropriate.

Early confirmatory testing may not be necessary in patients who have had a complete workup including mpMRI prior to diagnostic biopsy. However, all patients should undergo a confirmatory prostate biopsy within 1 to 2 years of their diagnostic biopsy.

Active Surveillance Program

The current NCCN recommendations for the active surveillance program include PSA no more often than every 6 months unless clinically indicated; DRE no more often than every 12 months unless clinically indicated; repeat prostate biopsy no more often than every 12 months unless clinically indicated; and consideration of repeat mpMRI no more often than every 12 months unless clinically indicated. mpMRI may be particularly

useful to exclude the presence of anterior cancer when PSA levels increase and systematic prostate biopsies remain negative.²⁹⁶

Results of a study of 211 patients with Grade Group 1 prostate cancer who had initial and repeat mpMRIs and PSA monitoring suggest that a negative initial mpMRI predicts a low risk of Gleason upgrading by systematic biopsy.²⁹⁷ In addition, PSA velocity was significantly associated with subsequent progression in those with an initial negative mpMRI. In contrast, those with high-risk visible lesions on mpMRI before initiation of active surveillance had an increased risk of progression. A meta-analysis of 43 studies found the sensitivity and NPV for mpMRI to be 0.81 and 0.78, respectively.²⁹⁸ An analysis of patients in Canary PASS found that mpMRI had an NPV and PPV for detecting Grade Group ≥ 2 cancer of 83% and 31%, respectively.²⁹⁹ Another study found the NPV of mpMRI to be 80%.³⁰⁰

In patients with a suspicious lesion on mpMRI, MRI-US fusion biopsy improves the detection of higher grade (Grade Group ≥ 2) cancers. Early experience supports the utilization of mpMRI in biopsy protocols to better risk stratify patients under active surveillance.³⁰¹⁻³⁰³ However, more recent studies have shown that a significant proportion of high-grade cancers are detected with systematic biopsy and not targeted biopsy in patients on active surveillance.³⁰⁴⁻³⁰⁶

Whereas the intensity of surveillance may be tailored on an individual basis (eg, based on life expectancy and risk of reclassification), most patients should have prostate biopsies incorporated as part of their monitoring, but no more often than every 12 months, because PSA kinetics may not be reliable for predicting progression. Repeat biopsy is useful to determine whether higher Gleason grade exists, which may influence prognosis and hence the decision to continue active surveillance or proceed to definitive local therapy.³⁰⁷ A repeat prostate biopsy should also be considered if the prostate exam changes, if mpMRI (if done)



NCCN Guidelines Version 5.2026

Prostate Cancer

suggests more aggressive disease, or if PSA increases. However, literature suggests that as many as 5% of patients undergoing prostate biopsy will suffer an adverse event, including serious ones such as infections requiring hospitalization.³⁰⁸⁻³¹⁰ Therefore, many clinicians choose to wait 2 years for a biopsy if there are no signs of progression.

Patients should be transitioned to observation (see *Observation*, above) when life expectancy is <10 years and they have low risk disease.

Considerations for Treatment of Patients on Active Surveillance

Reliable parameters of prostate cancer progression await the results of ongoing clinical trials. PSADT is not considered reliable enough to be used alone to detect disease progression.³¹¹ If repeat biopsy shows Grade Group ≥ 3 disease, or if tumor is found in a greater number of biopsy cores or in a higher percentage of a given biopsy core, cancer progression may have occurred. Grade reclassification on repeat biopsy is the most common factor influencing a change in management approach from active surveillance to treatment. Other factors affecting decisions to actively treat patients on active surveillance include increase in tumor volume and a rise in PSA density. Considerations for a change in management strategy should be made in the context of the patient's life expectancy. Patient anxiety over continued active surveillance should be addressed if treatment does not appear warranted.

Each of the major active surveillance series has used different criteria for reclassification.^{272,275,278-281,312-315} Reclassification criteria were met by 23% of patients with a median follow-up of 7 years in the Toronto experience,³¹³ 36% of patients with a median follow-up of 5 years in the Johns Hopkins experience,²⁷⁵ and 16% of patients with a median follow-up of 3.5 years in the University of California, San Francisco (UCSF) experience²⁸¹ (also see *Table 1*, below). Uncertainty regarding reclassification criteria and the desire to avoid missing an opportunity for cure drove several reports that

dealt with the validity of commonly used reclassification criteria. The Toronto group demonstrated that a PSA trigger point of PSADT <3 years could not be improved upon by using a PSA threshold of 10 or 20, PSADT calculated in various ways, or PSA velocity >2 ng/mL/y.³¹⁶ The Johns Hopkins group used biopsy-demonstrated reclassification to Gleason pattern 4 or 5 or increased tumor volume on biopsy as their criteria for reclassification. Of 290 patients on an annual prostate biopsy program, 35% demonstrated reclassification at a median follow-up of 2.9 years.³¹⁷ Neither PSADT (area under the curve [AUC], 0.59) nor PSA velocity (AUC, 0.61) was associated with prostate biopsy reclassification. Both groups have concluded that PSA kinetics cannot replace regular prostate biopsy, although treatment of most patients who demonstrate reclassification on prostate biopsy prevents evaluation of biopsy reclassification as a criterion for treatment or reduction of survival. Treatment of all patients who developed Gleason pattern 4 on annual prostate biopsies has thus far resulted in only two prostate cancer deaths among 1298 patients (0.15%) in the Johns Hopkins study.²⁷⁵ However, it remains uncertain whether treatment of all who progressed to Gleason pattern 4 was necessary. Studies remain in progress to identify the best trigger points when interventions with curative intent may still be successful.

The patients in the Toronto cohort were followed for a median of 6.8 years.³¹³ Of the 30% (n = 145) of patients who progressed, 8% had an increase in Gleason grade, 14% had a PSADT <3 years, 1% developed a prostate nodule, and 3% were treated because of anxiety. One hundred thirty-five of these 145 patients were treated: 35 by radical prostatectomy, 90 by external beam RT (EBRT) with or without ADT, and 10 with ADT alone. The 10- and 15-year actuarial cause-specific survival rates for the entire cohort were 98.1% and 94.3%, respectively.²⁷² Only 15 of 993 (1.5%) patients had died of prostate cancer, an additional 13 patients (1.3%) had developed metastatic disease, and only 36.5% of the cohort



NCCN Guidelines Version 5.2026

Prostate Cancer

had received treatment by 10 years. In an analysis of 592 patients enrolled in this cohort who had ≥ 1 repeat prostate biopsies, 31.3% of cases were upgraded. Fifteen percent of upgraded cases were upgraded to Gleason ≥ 8 , and 62% of total upgraded cases proceeded to active treatment.³¹⁸ Another analysis of this cohort revealed that metastatic disease developed in 13 of 133 patients with Gleason 7 disease (9.8%) and 17 of 847 patients with Gleason ≤ 6 disease (2.0%).³¹⁹ PSADT and the number of positive scores were also predictors of increased risk for the development of metastatic disease.

In comparison, among 192 patients on active surveillance who underwent delayed treatment at a median of 2 years after diagnosis in the Johns Hopkins experience, 5-year biochemical PFS was 96% for those who underwent radical prostatectomy and 75% for those who underwent radiation.³¹⁵ The two groups were similar by pathologic Gleason grade, pathologic stage, and margin positivity. All patients treated by radical prostatectomy after progression on active surveillance had freedom from biochemical progression at a median follow-up of 37.5 months, compared to 97% of those in the primary radical prostatectomy group at a median follow-up of 35.5 months. A later publication from this group showed that 23 of 287 patients who were treated after active surveillance (8%) experienced biochemical recurrence, and the rate was independent of the type of treatment.²⁷⁵ Several studies have shown that delayed radical prostatectomy does not increase the rates of adverse pathology.^{279,320-322}

Radical Prostatectomy

Radical prostatectomy is an appropriate option for patients whose cancer appears clinically localized to the prostate and can safely undergo general anesthesia and an operation. Because of potential perioperative morbidity, radical prostatectomy should generally be reserved for patients whose life expectancy is ≥ 10 years.

Radical prostatectomy was compared to watchful waiting in a randomized trial of 695 patients with early-stage prostate cancer (mostly T2) in the SPCG-4 trial.^{247,323} With a median follow-up of 12.8 years, those assigned to the radical prostatectomy group had significant improvements in disease-specific survival, OS, and risk of metastasis and local progression.³²³ The reduction in mortality was confirmed at 18-year follow-up, with an absolute difference of 11%.²⁴⁷ Overall, 8 patients needed to be treated to avert one death; that number fell to 4 for patients < 65 years of age. Longer follow-up results were also reported, in which the cumulative incidence of death from prostate cancer was 19.6% and 31.3% in the radical prostatectomy and watchful waiting groups, respectively, at 23 years, with a mean increase of 2.9 years of life in the radical prostatectomy group.²⁴⁸ However, in the small subset of patients with high-risk disease, there was no improvement in survival between treatment groups. A report of 30-year follow-up of this trial showed that the absolute difference in death due to prostate cancer between groups increased over time.³²⁴ Radical prostatectomy resulted in a 48% lower risk of death from prostate cancer and a mean of 2.2 life-years gained. The authors concluded that treating six patients would result in one prostate cancer death averted. The results of this trial offer high-quality evidence to support radical prostatectomy alone as a treatment option for patients with intermediate-risk clinically localized prostate cancer.

While the SPCG-4 and PIVOT trials comparing observation to surgery did not clearly demonstrate an improvement in outcomes among those with high-risk disease, these trials enrolled few patients and are not adequately powered to rigorously assess the role of surgery in high-risk disease.^{248,249} Trials that have utilized surgery in high-risk disease generally demonstrate a 70% to 80% risk of recurrence and thus surgery should be viewed as one component of a multi-modality approach.³²⁵ For example, in the CALGB 90203 trial of patients who received surgery in the control arm,



NCCN Guidelines Version 5.2026

Prostate Cancer

69% received adjuvant or secondary treatment, most commonly radiotherapy with or without ADT.

There remains no high-level evidence for the use of radical prostatectomy for patients with NCCN very-high-risk, regional/node-positive, or metastatic disease. Ideally surgery is used in the context of a clinical trial or when radiotherapy plus ADT is contraindicated. Surgery should be discussed as one component of a planned multimodality approach in these patients. There is an ongoing trial, SPCG-15, which aims to compare RP versus RT plus ADT in patients with high-risk prostate cancer.³²⁶

Radical prostatectomy is also a treatment option for select patients experiencing a local recurrence after primary EBRT, but morbidity (incontinence, erectile dysfunction, bladder neck contracture, and rectal injury) remain significantly higher than when radical prostatectomy is used as initial therapy.^{327,328} Patient selection is important, and post-RT recurrence radical prostatectomy should only be performed by highly experienced surgeons.

Operative Techniques and Adverse Effects

The retropubic and the perineal approaches are reasonable to perform a radical prostatectomy. Surgeons who perform a high-volume of operations in high-volume centers generally achieve superior outcomes.^{329,330} Laparoscopic and robot-assisted radical prostatectomy are commonly used and are considered comparable to conventional approaches in experienced hands.³³¹⁻³³³ In a cohort study using SEER Medicare-linked data on 8837 patients, minimally invasive compared to open radical prostatectomy was associated with shorter length of hospital stay, less need for blood transfusions, and fewer surgical complications, but rates of incontinence and erectile dysfunction were higher.³³⁴ A second large study reported no difference in overall complications, readmission, and

additional cancer therapies between open and robot-assisted radical prostatectomy, although the robotic approach was associated with higher rates of genitourinary complications and lower rates of blood transfusion.³³⁵ Oncologic outcomes of a robotic versus open approach were similar when assessed by use of additional therapies³³⁴ or rate of positive surgical margins,³³⁶ although longer follow-up is necessary. A meta-analysis of 19 observational studies (n = 3893) reported less blood loss and lower transfusion rates with minimally invasive techniques than with open operation.³³⁶ Risk of positive surgical margins was the same. Two more recent meta-analyses showed a statistically significant advantage in favor of a robotic approach compared to an open approach in 12-month urinary continence³³⁷ and potency recovery.³³⁸ Early results from a randomized controlled phase 3 study comparing robot-assisted laparoscopic radical prostatectomy and open radical retropubic prostatectomy in 326 patients were published in 2016.^{339,340} Urinary function and sexual function scores and rates of postoperative complications did not differ significantly between the groups at 6, 12, and 24 months after surgery. Rates of positive surgical margins were similar, based on a superiority test (10% in the open group vs. 15% in the robotic group). Assessment of oncologic outcomes from this trial will be limited because postoperative management and additional cancer therapies were not standardized between the groups.³³⁹

Return of urinary continence after radical prostatectomy may be improved by preserving the urethra beyond the prostatic apex and by avoiding damage to the distal sphincter mechanism. Bladder neck preservation may allow more rapid recovery of urinary control.³⁴¹ Anastomotic strictures that increase the risk of long-term incontinence are less frequent with modern surgical techniques. Recovery of erectile function is related directly to the degree of preservation of the cavernous nerves, age at surgery, and preoperative erectile function. Improvement in urinary and sexual function has been reported with nerve-sparing techniques.^{342,343}



NCCN Guidelines Version 5.2026

Prostate Cancer

Replacement of resected nerves with nerve grafts does not appear to be effective for patients undergoing wide resection of the neurovascular bundles.³⁴⁴ The ability of mpMRI to detect extracapsular extension can aid in decision-making in nerve-sparing surgery.²⁰¹

Common adverse events after RP include urinary incontinence, erectile dysfunction, penile shortening, and urethral or bladder neck contracture or stricture. The rates of occurrence vary by patient population (eg, by age and risk group), surgeon skill, and surgical technique (eg, unilateral vs. bilateral nerve sparing). In the ProtecT trial and PACE-A, which both randomized patients to RP versus RT, RP was associated with 40% to 60% risk of erectile dysfunction and 20% to 50% risk of urinary incontinence requiring at least one pad per day.^{25,345}

Pelvic Lymph Node Dissection

PLND can be considered with a radical prostatectomy for patients with favorable intermediate, unfavorable intermediate, high, or very-high-risk prostate cancer. If performed, it is recommended to be an extended PLND.^{346,347} An extended PLND includes removal of all node-bearing tissue from an area bounded by the external iliac vein anteriorly, the pelvic side wall laterally, the bladder wall medially, the floor of the pelvis posteriorly, Cooper's ligament distally, and the internal iliac artery proximally. Removal of more lymph nodes using the extended technique has been associated with increased likelihood of finding lymph node metastases, thereby providing more complete staging.³⁴⁸⁻³⁵⁰ A survival advantage with more extensive lymphadenectomy has been suggested by several studies, possibly due to elimination of microscopic metastases,^{349,351-353} although definitive proof of oncologic benefit is lacking.³⁵⁴ PLND can be performed safely laparoscopically, robotically, or as an open procedure, and complication rates should be similar among the three approaches.

Radiation Therapy

RT techniques used in prostate cancer can be described generally by the following: delivery method (ie, internal vs. EBRT), source of radiation (ie, photon vs. proton), planning technique (ie, 3D vs. intensity modulated RT [IMRT]), image guidance (none vs. cone beam CT [CBCT] or MRI), adaptation (ie, online adaptive treatment), and fractionation (ie, conventional vs. moderate hypofractionation vs. stereotactic body radiotherapy).

External Beam Radiation Therapy

Over the past several decades, EBRT techniques have evolved to allow higher doses of radiation to be administered safely. EBRT can be delivered either with photons, also called x-rays, or proton beam. Photons are used in >95% of cases in the United States.

Proton Therapy

Proton therapy and x-ray–based therapies like IMRT can deliver highly conformal doses to the prostate. Proton-based therapies will deliver less low-dose radiation to some of the surrounding normal tissues not immediately adjacent to the prostate, such as muscle, bone, vessels, and fat. These tissues do not routinely contribute to the morbidity of prostate radiation and are relatively resilient to radiation injury; therefore, the benefit of decreased dose to these types of normal, non-critical tissues has not been clearly proven.

Only one randomized trial has been performed, the PartiQOL trial, and results are reported in abstract only.³⁵⁵ This trial randomized 450 patients with low- and intermediate-risk prostate cancer to either proton or photons. After a median follow-up of 5-years, there were no significant differences in patient-reported quality of life, physician reported toxicity, or biochemical recurrence. Thus, both treatments appear to offer excellent tumor control and good safety profile.



NCCN Guidelines Version 5.2026

Prostate Cancer

The costs associated with proton beam facility construction and proton beam treatment are high compared to the expense of building and using the more common photon linear accelerator-based practice.³⁵⁶ These higher costs may translate into higher charges for protons, a point that should be considered when tumor control and side effects appear similar between photon and proton beam therapy for prostate cancer, especially if these charges are incurred by the patient.

The NCCN Panel believes no clear evidence supports a benefit or decrement to proton therapy over IMRT for either treatment efficacy or long-term toxicity. Conventionally fractionated or moderately hypofractionated prostate proton therapy can be considered a reasonable alternative to x-ray–based regimens at clinics with appropriate technology, physics, and clinical expertise. Insufficient data is present to recommend their delivery with ultra-hypofractionation.

Photon (X-ray) Beam

Treatment planning in the 1980s used two-dimensional planning techniques. By the 1990s this was largely replaced by three-dimensional planning. In the late 1990s IMRT planning techniques were increasingly utilized, and as of 2010 nearly all centers utilize IMRT planning. This technique has been shown to improve dose conformality.

In parallel, improvements in image guidance, also referred to as image guided RT (IGRT), have occurred. The purpose of IGRT is to ensure the planned dose is delivered as intended. Given the potential changes in anatomy day to day due to bladder and rectal filling, IGRT is essential to ensure accurate alignment of the prostate and target volumes and avoidance of organs at risk. IGRT techniques often include use of implanted fiducials, electromagnetic targeting and tracking, or endorectal balloon, which have variably been shown to decrease complications and allow margin reduction.³⁵⁷ More recently, the use of real-time tracking and online adaptive radiotherapy have also demonstrated the ability to improve

dosimetry and enable margin reduction, which may result in reduced toxicity.³⁵⁸

The advancements in treatment planning and IGRT have enabled greater comfort and confidence to test increasingly hypofractionated regimens. Historically, conventionally fractionated RT was utilized with 1.8 to 2.0 Gy per fraction over 8 to 9 weeks of daily RT. This schema has been compared in numerous randomized trials to moderately hypofractionated RT in 2.5 to 4 Gy/fraction, which has been shown to be noninferior when using iso-effective doses.³⁵⁹⁻³⁶⁸ Recently, the HYDRA individual patient meta-analyses of these trials demonstrated that 60 Gy in 20 fractions, or regimens with a similar iso-effective dose, should be preferred.³⁶⁹ Reduction from >40 treatments to 20 treatments reduces both financial toxicity and increases patient convenience.

The relatively slow proliferation rate of prostate cancer is reflected in a low α/β ratio, most commonly reported between 1 and 4.³⁷⁰ Because the α/β ratio for prostate cancer is similar to or lower than the surrounding tissues responsible for most of the toxicity reported with radiation, appropriately designed radiation treatment fields and schedules using ultra hypofractionated regimens were hypothesized to result in similar cancer control rates without increased risk of late toxicity.

SBRT is a technique that delivers highly conformal, high-dose radiation in five or fewer treatment fractions, which are safe to administer only with precise, image-guided delivery.³⁷¹ Numerous prospective studies demonstrated safety and efficacy of SBRT in the setting of localized prostate cancer.³⁷² Two large randomized phase 3 trials have reported results. HYPO-RT-PC trial enrolled 1200 patients with intermediate- and high-risk prostate cancer and demonstrated noninferiority of 42.7 Gy in seven fractions to 78.0 Gy in 39 fractions with respect to FFS.³⁷³ Likewise, PACE-B compared 5-fraction SBRT with a dose of 40 Gy to the prostate and 36.25 Gy to the planning target volume versus moderate



NCCN Guidelines Version 5.2026

Prostate Cancer

hypofractionated RT with 62 Gy in 20 fractions.³⁷⁴ Results showed noninferiority in tumor control and generally similar safety profiles.

SBRT regimens should utilize IMRT planning and daily IGRT. SBRT is an appropriate standard RT method at clinics with appropriate technology, physics, and clinical expertise.

The Panel lists fractionation schemes that have shown acceptable efficacy and toxicity within the *Principles of Radiation Therapy*.

Relative contraindications or considerations to EBRT include prior pelvic irradiation, active inflammatory disease of the rectum, permanent indwelling Foley catheter, very low bladder capacity, chronic moderate or severe diarrhea, and bladder outlet obstruction requiring a suprapubic catheter.

Hydrogel Per-Rectal Spacers

Biomaterials have been developed, tested, and FDA approved to serve as spacer materials when inserted between the rectum and prostate.^{375,376} In a randomized phase 3 multicenter clinical trial of patients undergoing image-guided IMRT (IG-IMRT), the risk of late (3-year) grade 2 or higher, physician-recorded rectal complications declined from 5.7% to 0% in the control versus hydrogel spacer group.³⁷⁷ The control group had a significant reduction in bowel QOL decline. No significant differences in urinary adverse events were noted in those receiving hydrogel placement versus controls. Results of a secondary analysis of this trial suggest that use of a perirectal spacer may decrease the sexual side effects of radiation.³⁷⁸ Spacer implantation may be associated with rare complications such as rectum perforation and urethral damage.^{379,380} Retrospective data also support its use in similar patients undergoing brachytherapy. Overall, the Panel believes that biocompatible and biodegradable perirectal spacer materials may be implanted between the prostate and rectum in patients undergoing RT in order to displace the

rectum from high radiation dose regions. Patients with obvious rectal invasion or visible posterior extension should not undergo perirectal spacer implantation.

EBRT for Low-Risk Disease

Active surveillance is preferred for low-risk disease. If EBRT is to be used, it should be given as monotherapy without a brachytherapy boost, without nodal irradiation, and without ADT. Ideally it is given with SBRT or moderately hypofractionated RT.

EBRT for Intermediate-Risk Disease

The favorable/unfavorable distinction within the intermediate-risk group should be used to help guide treatment for these patients. Generally, patients with favorable intermediate-risk disease should be treated with EBRT alone without a brachytherapy boost, without nodal irradiation, and without ADT.³⁸¹ Use of SBRT or moderately hypofractionated RT are preferred. Patients with unfavorable intermediate-risk disease generally should be considered for the addition of short-term ADT.^{382,383} Studies have shown that the addition of a focal EBRT boost may further improve biochemical control in these patients.³⁸⁴ Similarly, the addition of a brachytherapy boost may improve biochemical control, at the expense of increased toxicity.³⁸⁵ Trials to date that have enrolled patients with intermediate-risk disease have not shown improvements in disease control from the use of elective nodal irradiation (ENI), and have generally seen an increase in toxicity.³⁸⁶

EBRT for Patients with High-Risk or Very-High-Risk Disease

EBRT represents the only local therapy for patients with high- or very-high-risk disease that has been shown to improve overall survival when compared to ADT alone.³⁸⁷⁻³⁹⁰ Furthermore, the addition of ADT, specifically long-term ADT of >18 months, has been shown to improve overall survival as compared to RT alone or RT with short-term ADT.³⁸²



NCCN Guidelines Version 5.2026

Prostate Cancer

There are mixed results regarding the benefit of ENI in this patient population. Three multicenter cooperative group trials have not demonstrated improvement in outcomes from ENI; GETUG-01, RTOG 9413, and recently in abstract form RTOG 0924.^{386,391,392} However, one small single center trial from India, POP-RT, showed that pelvic radiation can improve biochemical FFS and DFS compared with prostate-only radiation in patients with high- and very-high-risk prostate cancer.³⁹³ With the recent results of NRG/RTOG 0924 (ClinicalTrials.gov ID NCT01368588) demonstrating no improvement in distant metastasis, prostate cancer-specific mortality, or overall survival from ENI in an approximately 2500 patient trial of unfavorable-intermediate and high-risk prostate cancer, the role of ENI remains unclear.

More recently, the STAMPEDE trial has reported a post-hoc analysis of the benefit of adding abiraterone acetate/prednisone to EBRT plus long-term ADT in patients with NCCN very-high-risk prostate cancer.³⁹⁴ Results demonstrated improvement in metastasis-free survival from the addition of abiraterone at the expense of increased grade ≥ 3 side effects. Notably, the patients enrolled in this trial had markedly higher rates of metastasis and death from prostate cancer than those observed in other trials of high-risk prostate cancer conducted by NRG, EORTC, or TROG. Given some patients had PSAs >1000 ng/mL on the STAMPEDE trial but were called non-metastatic, it remains unclear the true value of the addition of abiraterone in contemporary very-high-risk disease. ENZARAD, a randomized trial of patients with high- and very-high-risk and clinical node positive disease, recently reported results in abstract form and demonstrated that the addition of the APRI enzalutamide to RT+ADT did not improve metastasis-free survival or overall survival.³⁹⁵ Ongoing trials utilizing other ARPIs have not yet reported their results, such as ATLAS (NCT02531516).

EBRT for Node-Positive Disease

EBRT with neoadjuvant, concurrent, and/or adjuvant ADT and abiraterone is the preferred option for patients with clinical N1 disease based upon the previously discussed STAMPEDE trial.³⁹⁴ Other options in this setting include EBRT with ADT, ADT alone or with abiraterone, and, for select patients, RP plus PLND (see *Radical Prostatectomy*, above).

Post-Recurrence EBRT

The use of EBRT in patients with biochemical recurrence is discussed under *Post-Radical Prostatectomy Approaches*, below.

Brachytherapy

Brachytherapy involves placing radioactive sources into the prostate tissue. Brachytherapy has been used traditionally for low-risk disease because earlier studies found it less effective than EBRT for high-risk disease.^{121,396} However, increasing evidence suggests that technical advancements in brachytherapy may provide a role for contemporary brachytherapy in higher-risk localized and locally advanced prostate cancer.³⁹⁷

There are currently two methods for prostate brachytherapy: low dose-rate (LDR) and high dose-rate (HDR). LDR brachytherapy consists of placement of permanent seed implants in the prostate. The short range of the radiation emitted from these low-energy sources allows delivery of adequate dose levels to the cancer within the prostate, with excessive irradiation of the bladder and rectum avoided. Post-implant dosimetry should be performed to document the quality of an LDR implant.³⁹⁸ HDR brachytherapy, which involves temporary insertion of a radiation source, is a newer approach.

Two groups have observed a lower risk of urinary frequency, urgency, and rectal pain with HDR brachytherapy compared with LDR brachytherapy (permanent seed implant).^{399,400} Vargas and colleagues⁴⁰¹ reported that



NCCN Guidelines Version 5.2026

Prostate Cancer

HDR brachytherapy results in a lower risk of erectile dysfunction than LDR brachytherapy. Commonly prescribed doses for LDR and HDR brachytherapy are listed in the *Principles of Radiation Therapy* within the algorithm above.

For patients with very large or very small prostates, symptoms of bladder outlet obstruction (high International Prostate Symptom Score), or a previous transurethral resection of the prostate (TURP), seed implantation may be more difficult. These patients also have an increased risk of side effects. Neoadjuvant ADT may be used to shrink the prostate to an acceptable size; however, increased toxicity is expected from ADT, and prostate size may not decline in some patients. The potential toxicity of ADT must be weighed against the possible benefit of target reduction.

Ideally, the accuracy of brachytherapy treatment should be verified by daily prostate localization with techniques of IGRT: CT, ultrasound, implanted fiducials, or electromagnetic targeting/tracking. Endorectal balloons may be used to improve prostate immobilization. Perirectal spacer materials (discussed under *External Beam Radiation Therapy*, above) may be used when the previously mentioned techniques are insufficient to reduce side effects due to anatomic geometry or other patient-related factors (eg, medication usage, comorbid conditions). Patients with obvious rectal invasion or visible cT3 and posterior extension should not undergo perirectal spacer implantation.

Brachytherapy Alone for Localized Disease

Active surveillance is preferred for low-risk disease, but brachytherapy alone is an option for select patients with low-risk disease. Brachytherapy alone is an option for patients with favorable intermediate-risk and for carefully selected patients with unfavorable intermediate-risk prostate cancer, depending on life expectancy. Patients with high-risk cancers are not generally considered candidates for brachytherapy alone. Either LDR or HDR brachytherapy can be used in this setting; however, there are

concerns regarding the efficacy of both single fraction HDR and, more recently, two fraction HDR as monotherapy.⁴⁰²

There are no reported trials comparing SBRT or other contemporary EBRT methods to brachytherapy. ASCENDE-SBRT is comparing a brachytherapy boost versus SBRT.⁴⁰³

Brachytherapy Boost

LDR or HDR brachytherapy can be added as a boost to EBRT plus ADT in patients with unfavorable intermediate- or high-risk prostate cancer being treated with curative intent and for carefully selected patients with very-high-risk disease. Combining EBRT and brachytherapy allows dose escalation. This combination has demonstrated improved biochemical control over EBRT plus ADT alone in randomized trials, but with higher toxicity.⁴⁰⁴⁻⁴⁰⁶

The randomized ASCENDE-RT trial compared two methods of dose escalation in 398 patients with intermediate- or high-risk prostate cancer: dose-escalated EBRT boost to 78 Gy or LDR brachytherapy boost.³⁸⁵ All patients were initially treated with 12 months of ADT and pelvic EBRT to 46 Gy. An ITT analysis found that the primary endpoint of biochemical PFS was 86% versus 75% at 7 years for the LDR boost versus EBRT boost arms (log-rank $P < .001$). Toxicity was higher in the brachytherapy arm, with the cumulative incidence of grade 3 genitourinary events at 5 years of 18.4% for brachytherapy boost and 5.2% for EBRT boost ($P < .001$).⁴⁰⁷ A trend for increased gastrointestinal toxicity with brachytherapy boost was also seen (cumulative incidence of grade 3 events at 5 years, 8.1% vs. 3.2%; $P = .12$). However, at 6-year follow-up, health-related QOL was similar between the groups in most domains, except that physical and urinary function scales were significantly worse in the LDR arm.⁴⁰⁸ Whereas the toxicity is increased with the use of brachytherapy boost, this and other randomized controlled trials have not shown an improvement in OS or cancer-specific survival.⁴⁰⁹



NCCN Guidelines Version 5.2026

Prostate Cancer

The optimal duration of ADT with EBRT plus brachytherapy is unclear. The TROG RADAR trial randomized patients to receive 6 versus 18 months of ADT, and a subset of participants received EBRT with an HDR boost.⁴¹⁰ In the subset who received EBRT plus HDR, the use of 18 months of ADT improved overall survival compared to 6 months. More recently, the TRIP/TRIGU0907 randomized trial of EBRT plus LDR compared 6 versus 30 months of ADT.⁴¹¹ The study found no significant difference in biochemical progression between arms, both were <10% at 7-year follow-up.

To address concerns for increased toxicity incidence associated with brachytherapy boost from historical trial data, careful patient selection and contemporary planning associated with lesser toxicity, such as use of recognized organ at risk dose constraints, use of high-quality ultrasound and other imaging, and prescription of dose as close as possible to the target without excessive margins, should be implemented.

Post-Recurrence Brachytherapy

Brachytherapy can be considered in patients with biochemical recurrence after EBRT. Results of a phase 2 study of post-recurrence HDR brachytherapy after EBRT included relapse-free survival, distant metastases-free survival, and cause-specific survival rates of 68.5%, 81.5%, and 90.3%, respectively, at 5 years.⁴¹² Toxicities were mostly grade 1 and 2 and included gastrointestinal toxicity and urethral strictures, and one case of Grade 3 urinary incontinence. In another prospective phase 2 trial, the primary endpoint of grade ≥3 late treatment-related gastrointestinal and genitourinary adverse events at 9 to 24 months after post-recurrence brachytherapy was below the unacceptable threshold, at 14%.⁴¹³

Data on the use of brachytherapy after permanent brachytherapy are limited, but the Panel agrees that it can be considered for carefully selected patients. Decisions regarding the use of brachytherapy in the

recurrent-disease setting should consider comorbidities, extent of disease, and potential complications. Brachytherapy in this setting is best performed at high-volume centers.

Radiation for Distant Metastases

EBRT can be used for multiple purposes in the setting of metastatic disease. Historically, it was only used as a means of palliation, often for bone pain, obstruction, or bleeding. A short course of 8 Gy x 1 is as effective for palliation as, and less costly than, 30 Gy in 10 fractions.⁴¹⁴ In a randomized trial of 898 patients with bone metastases, grade 2–4 acute toxicity was observed less often in the 8-Gy arm (10%) than in the 30-Gy arm (17%) ($P = .002$); however, the retreatment rate was higher in the 8-Gy group (18%) than in the 30-Gy group (9%) ($P < .001$).⁴¹⁵ In another study of 425 patients with painful bone metastases, a single dose of 8 Gy was noninferior to 20 Gy in multiple fractions in terms of overall pain response to treatment.⁴¹⁶ The SCORAD randomized trial did not show noninferiority for ambulatory status of single-fraction 8-Gy EBRT to 20 Gy in 5 fractions.⁴¹⁷

Eight Gy as a single dose is as effective for short-term (ie, 12 week) pain palliation at any bony site as longer courses of radiation, but re-treatment rates are higher. Other regimens (eg, 30 Gy in 10 fractions) may be used as alternative palliative dosing depending on clinical scenario. Neither regimen has demonstrated clear long-term pain or cancer control and are intended for transient pain control palliation.

EBRT may also be used as an oncologic treatment and not solely for the palliation of symptoms in the metastatic setting. The topic of metastasis-directed radiation therapy is discussed in detail below (see *MDT for Oligometastatic CSPC* and *MDT for mCRPC*).



NCCN Guidelines Version 5.2026

Prostate Cancer

Comparison of Treatment Options for Localized Disease

Multiple randomized trials have compared RT to RP. The ProtecT trial demonstrated that, with a median follow-up of 15 years, patients with low and intermediate risk prostate cancer had similar rates of developing metastasis or death.^{263,265} A post-hoc subset analysis of the ProtecT trial demonstrated that patients with cribriform features on active monitoring had significantly reduced rates of distant metastasis when treated with RT plus ADT, but not with RP.²⁶⁴

The ProtecT trial demonstrated that urinary incontinence rates and impotence rates were significantly higher with RP than RT. There were small but significant increases in bowel side effects with RT as compared to surgery. The RT utilized on this trial was 3D conformal radiotherapy without contemporary image guidance or rectal spacer use.

More recently, results from PACE-A, which included patients with intermediate-risk prostate cancer receiving either SBRT versus RP, were reported.³⁴⁵ No spacer gels were utilized. The study demonstrated that, with more modern RT, there were smaller differences in patient-reported bowel quality of life as compared to surgery. Furthermore, those receiving surgery were >3-fold more likely to develop impotence, and >8-fold more likely to develop urinary incontinence requiring ≥1 pads per day.

Recently, a cross-trial comparison using two phase 3 randomized trials was performed.⁴¹⁸ Trials were matched by era, country, and randomization. NRG/RTOG 0521 randomized patients with high-risk prostate cancer to RT plus 24 months of ADT with or without 6 cycles of docetaxel, and CALGB randomized patients with high-risk prostate cancer to surgery with or without 6 cycles of docetaxel with concurrent ADT in the neoadjuvant setting. The patients in the RT trial were generally older and had more aggressive prognostic features. This study demonstrated that patients receiving RT with ADT had significantly reduced risk of

developing distant metastasis than patients treated with a surgical approach, but there were no significant differences in death from prostate cancer. As this was not a true randomized trial of the two treatment approaches, confounding may impact the results.

Numerous non-randomized studies have been reported that should be interpreted with caution given the high risk for selection bias and residual confounding. Barocas et al compared radical prostatectomy, EBRT, and active surveillance in 2550 patients and found that, after 3 years, radical prostatectomy was associated with a greater worsening in urinary and sexual function than either EBRT or active surveillance.⁴¹⁹ However, radical prostatectomy was also associated with an improvement in urinary irritative symptoms that was not observed with radiation or active surveillance. Health-related QOL measures including bowel and hormonal function were similar among the groups, as was disease-specific survival.

Chen et al compared radical prostatectomy, EBRT, and brachytherapy against active surveillance in 1141 patients.²³ As in the Barocas study, radical prostatectomy was associated with greater declines in sexual and urinary function than other treatments at 3 months. In this study, EBRT was associated with worse short-term bowel function, and both EBRT and brachytherapy were associated with worsened urinary obstructive and irritative symptoms. By 2 years, however, differences among the groups compared with active surveillance were insignificant. Results of a systematic review showed similar findings to these studies.⁴²⁰

Another study examined patient-reported outcomes in >2000 patients with localized prostate cancer managed by radical prostatectomy, brachytherapy, EBRT with or without ADT, or active surveillance.²⁴ By 5 years, most functional differences were minimal between management approaches. However, radical prostatectomy was associated with worse incontinence in the full cohort and with worse sexual function in those with



NCCN Guidelines Version 5.2026

Prostate Cancer

unfavorable intermediate-, high-, or very-high-risk disease than in those treated with EBRT and ADT.

Other Local Therapies

Many therapies have been investigated for the treatment of localized prostate. Cryotherapy was investigated as a potential whole gland treatment option. A study by Donnelly and colleagues included 244 patients with T2 or T3 disease who were randomly assigned to either cryotherapy or EBRT.⁴²¹ In this small trial, all patients received neoadjuvant ADT. Patients who received cryotherapy reported poorer sexual function.⁴²² For patients with locally advanced cancer, cryoablation was associated with a lower 8-year biochemical progression-free rate compared to EBRT in a small trial of 62 patients.⁴²³

Other ablative therapies (eg, high-intensity focused ultrasound [HIFU], transurethral ultrasound ablation [TULSA], irreversible electroporation [IRE], photodynamic therapy [PDT]) are not recommended as primary therapy for localized prostate cancer due to lack of long-term randomized data comparing these treatments to radiation or radical prostatectomy.⁴²⁴⁻⁴³⁰ Single arm phase I or II trials have shown high rates of residual or recurrent cancer (20%–70%) and clinically meaningful toxicity (ie, incontinence, obstruction, erectile dysfunction). Furthermore, monitoring strategies appear to be compromised (ie, MRI accuracy) and secondary treatment may carry higher risk of complications. These ablative treatments can be utilized in the context of a clinical trial, ideally a randomized trial compared them to a current standard treatment option. Companies and institutions may make non-evidenced-based claims to the evidence, efficacy, and safety of focal therapy.

At this time, the only local therapy options that can be considered are cryotherapy, HIFU, and IRE (category 2B) in the setting of non-metastatic

RT recurrence due to the overall lack of high-quality evidence in this setting for any modality. Also see *Post-Radiation Recurrence*, below.

Disease Monitoring

Please refer to the NCCN Guidelines for Survivorship (available at www.NCCN.org) for recommendations regarding common consequences of cancer and cancer treatment (eg, anxiety, depression, trauma, and distress; hormone-related symptoms; sexual dysfunction) and on the promotion of physical activity, weight management, and proper immunizations in cancer survivors. Prostate-cancer specific survivorship recommendations pertaining to bone health and cardiovascular disease are included in the algorithm above and are discussed in more detail in the section on *Adverse Effects of Traditional ADT*, below.

Patients After Initial Definitive Therapy

For patients initially treated with curative intent, serum PSA levels should be measured every 6 to 12 months for the first 5 years and then annually. PSA testing every 3 months may be better for patients at high risk of recurrence. When prostate cancer recurred after radical prostatectomy, Pound and colleagues found that 45% of patients experienced recurrence within the first 2 years, 77% within the first 5 years, and 96% by 10 years.⁴³¹

Patients with N1 CSPC on ADT

The intensity of clinical monitoring for patients on ADT for N1 CSPC is determined by the response to initial ADT, EBRT, or both. Follow-up evaluation of these patients should include physical examination and PSA measurement every 3 to 6 months. Imaging for symptoms or increasing PSA. The relative risk for bone metastasis or death increases as PSADT falls. Bone imaging should be performed more frequently in patients with shorter PSADTs.⁴³²



NCCN Guidelines Version 5.2026

Prostate Cancer

Patients with Localized Disease Under Observation

Patients with localized disease on observation follow the same monitoring recommendations as patients with N1 CSPC who are on ADT.

Workup for Progression

Castrate levels of testosterone should be documented if clinically indicated in patients with signs of progression, with adjustment of ADT as necessary. If serum testosterone levels are <50 ng/dL, the patient should undergo disease workup with bone and soft tissue imaging (see *Imaging Techniques* above for more details).

Post-Radical Prostatectomy Approaches

Patients who have undergone radical prostatectomy should be monitored as described above. However, some patients will have adverse pathologic features, positive lymph nodes, or biochemical persistence or recurrence. Options for these patients include adjuvant radiation (the use of EBRT immediately following radical prostatectomy in the setting of an undetectable postoperative PSA) with or without ADT, secondary radiation (the use of EBRT after radical prostatectomy when PSA is detectable) with or without ADT, or monitoring. Decisions about when to initiate post-radical prostatectomy radiation and whether to include ADT for such patients are complex. The Panel recommends use of nomograms and consideration of age and comorbidities, clinical and pathologic information, PSA levels, and PSADT to individualize treatment discussion. The various scenarios are discussed in more detail below.

Bone and soft tissue imaging and prostate bed biopsy should be considered in patients with PSA persistence/recurrence after RP if their life expectancy is >5 years. The utility of imaging for patients with an early biochemical recurrence after radical prostatectomy depends on disease risk before operation and pathologic stage, Gleason grade, PSA, and PSADT after recurrence. Patients with low- and intermediate-risk disease

and low postoperative serum PSA levels have a very low risk of positive bone scans or CT scans.^{433,434} In a series of 414 bone scans performed in 230 patients with biochemical recurrence after radical prostatectomy, the rate of a positive bone scan for patients with PSA >10 ng/mL was only 4%.⁴³⁵ In another study, the probability of a positive bone scan for a patient not on ADT after radical prostatectomy was <5% unless the PSA increased to 40 to 45 ng/mL.⁴³⁶ Thus bone imaging is appropriate when patients develop symptoms or when PSA levels are increasing rapidly. A prostate bed biopsy can also be considered. However, PSMA PET/CT imaging has increased sensitivity at low PSA levels post-RP, and approximately 30% to 40% of patients will have detectable disease on PSMA PET/CT at PSA levels of 0.2 to 0.5 ng/mL.

Early Secondary or Adjuvant Treatment for Adverse Features After Radical Prostatectomy

Patients with undetectable PSA levels and without lymph node metastases following radical prostatectomy can be considered for immediate adjuvant treatment if adverse pathologic features are present (ie, positive margins, seminal vesicle invasion, extracapsular extension). However, monitoring these patients is the preferred option (category 1). Waiting for a detectable and rising PSA or PSA ≥0.1 ng/mL before initiating secondary radiation is preferred in this setting, since this approach avoids over-treatment and subsequent potential side effects and does not appear to compromise oncologic outcomes.^{437,438}

In the RADICALS-RT trial, 1396 patients with adverse features after radical prostatectomy were randomized to receive immediate adjuvant treatment or to be monitored with a policy to initiate secondary treatment if PSA levels reached 0.1 ng/mL or rose three consecutive times.⁴³⁹ Participants were followed for a median 4.9 years and no differences were seen in 5-year biochemical PFS and freedom from non-protocol hormone therapy. There was no statistically significant difference in the primary



NCCN Guidelines Version 5.2026

Prostate Cancer

endpoint of freedom from distant metastasis between the arms at 10 years (93% in the adjuvant arm vs. 90% in the early secondary therapy arm; 95% CI, 0.43–1.07; $P = .095$) and there was no difference in OS.⁴³⁸ Importantly, urinary incontinence and grade 3–4 urethral strictures were more frequent in the adjuvant therapy group.

The GETUG-AFU 17 trial and the TROG 08.03/ANZUP RAVES trial were both terminated early for unexpectedly low event rates, but similarly found no evidence of oncologic benefit with increased risk of genitourinary toxicity and erectile dysfunction when an adjuvant therapy approach was used.^{440,441} One small randomized trial in Finland ($n = 126$) saw a significant improvement in 10-year survival for biochemical recurrence with the use of adjuvant therapy in patients with positive margins or extracapsular extension (HR, 0.26; 95% CI, 0.14–0.48; $P < .001$).⁴⁴² However, there was no difference in OS, and grade 3 adverse events were significantly higher in the adjuvant arm.

Several meta-analyses have also shown that adjuvant radiotherapy improved biochemical control, but not overall survival, in patients with adverse features and that the approach leads to a higher incidence of adverse events, though some noted that careful patient selection is warranted.^{437,443–445} Real world evidence supports the safety of this approach for most patients.^{446,447}

The Panel notes immediate adjuvant radiation may be reasonable for patients with multiple adverse features, a population that was under-represented in the above-mentioned clinical trials. True noninferiority of the delayed, early secondary therapy approach has not been demonstrated in patients with a high risk of recurrence after surgery.

Adjuvant or Early Secondary Therapy for pN1

Patients with positive lymph nodes found during or after radical prostatectomy have several disease management options to consider.

Immediate adjuvant treatment with EBRT with or without ADT is one approach for these patients based on historic data.⁴⁴⁸ However, retrospective data show that initial observation may be safe in some patients with pN1 disease at radical prostatectomy, because 28% of a cohort of 369 patients remained free from biochemical recurrence at 10 years.⁴⁴⁹ Therefore, another option is monitoring with consideration of early secondary treatment for a detectable and rising PSA or PSA >0.1 ng/mL. However, RADICALS-RT, GETUG-AFU 17, and TROG 08.03/ANZUP RAVES have very few pN1 patients that were enrolled.^{439–441}

An analysis of the National Cancer database identified 8074 patients with lymph node metastases after radical prostatectomy.⁴⁵⁰ The results indicate that immediate adjuvant EBRT and ADT improved OS compared to monitoring for those who also had adverse pathologic features but not for those without these features. Therefore, use of early secondary radiotherapy can be considered for patients with pN1 disease, with careful patient selection and monitoring.

The addition of abiraterone to EBRT with ADT can also be considered for patients with positive lymph after surgery (category 2B). However, these patients comprised $<5\%$ of patients on the STAMPEDE trial, and caution is warranted with extrapolation of data from cN1 disease.³⁹⁴

Secondary Treatment for PSA Persistence or Biochemical Recurrence After Radical Prostatectomy

The timing of secondary treatment for patients whose only evidence of cancer after radical prostatectomy is a persistent or increasing PSA is influenced by PSA levels, PSADT, and the underlying comorbidities of the patient. PSADT should be calculated to inform nomogram use and counseling for patients with PSA persistence or recurrence after radical prostatectomy. Early secondary treatment (at a PSA of 0.1–0.2 ng/mL) is recommended for most patients who experience a biochemical recurrence after RP. However, patients who have a delayed recurrence (eg, many



NCCN Guidelines Version 5.2026

Prostate Cancer

years post-RP) with very slow PSA doubling times (ie, >24 months) can consider delaying secondary treatment, especially if they have limited life expectancy.

For those with PSA persistence or recurrence after RP without evidence of M1 disease, secondary treatment with EBRT is the preferred approach. Large retrospective cohort studies support the use of EBRT in the setting of biochemical recurrence, because it is associated with decreased all-cause mortality and increased prostate cancer-specific survival.^{451,452}

Numerous phase 3 randomized trials have evaluated the role of hormone therapy with secondary EBRT in this setting, as well as a harmonized aggregate meta-analysis.⁴⁵³⁻⁴⁵⁷ In general, these studies have demonstrated that 6 to 24 months of hormone therapy improved outcomes primarily for patients receiving late secondary RT (PSA levels >0.5 ng/mL), but a smaller or no benefit is clear at lower PSA levels. For example, RTOG 9601 demonstrated an OS benefit with 2 years of bicalutamide with secondary EBRT, but enrolled patients primarily receiving late secondary EBRT.⁴⁵⁸ In contrast, the RADICALS-HD trial randomized patients to post-operative EBRT with or without 6 months of ADT and enrolled mostly adjuvant or early secondary EBRT patients. They showed that ADT did not improve MFS or OS. The DADSPORT meta-analysis did not show an OS benefit for the addition of ADT in patients receiving early secondary RT, although an MFS and PCSS benefit were seen.⁴⁵⁹

Based on these data, short-term ADT is generally recommended to balance toxicity, QOL, and tumor control. However, long-term ADT may be preferred for patients receiving late secondary RT and/or those with multiple adverse pathologic features or lymph node involvement.

There remains no randomized data to support the addition of an ARPI to EBRT and ADT. However, abiraterone is an option for patients with lymph

node metastases (category 2B), based on extrapolation from STAMPEDE.³⁹⁴

Monitoring patients without evidence of M1 disease in the setting of PSA persistence or a rising PSA post-RP is another option, especially if the risk of rapid progression appears to be low based on factors such as PSA levels, PSADT, and results of nomograms and with consideration of patient age and comorbidities. It is important to note that some patients have detectable PSA after radical prostatectomy due to benign prostate tissue in the prostate fossa. They have low stable PSAs and a very low risk of prostate cancer progression.^{460,461} Serial PSA measurements can be helpful for stratifying patients at highest risk of progression and metastases.

Post-Radiation Recurrence

The 2006 Phoenix definition was revised by ASTRO and the RTOG in Phoenix: 1) PSA rise by 2 ng/mL or more above the nadir PSA is the standard definition for biochemical recurrence after EBRT with or without hormonal therapy; and 2) A recurrence evaluation should be considered when PSA has been confirmed to be increasing after radiation even if the rise above nadir is not yet 2 ng/mL, especially in candidates for additional local therapy who are young and healthy.⁴⁶² Retaining a strict version of the ASTRO definition allows comparison with a large existing body of literature. Rapid increase of PSA may warrant evaluation (prostate biopsy) prior to meeting the Phoenix definition, especially in younger or healthier patients.

Workup for PSA recurrence or a positive DRE after RT for a patients with a life expectancy >5 years typically includes PSADT calculation, bone and soft tissue imaging, and consideration of a prostate/seminal vesicle biopsy. Biopsy is particularly useful if the staging workup does not reveal



NCCN Guidelines Version 5.2026

Prostate Cancer

metastatic disease and local recurrence is suspected. The junction of the seminal vesicle and prostate is a common site of recurrence.

Local radiation recurrences are most responsive to additional therapy when PSA levels at the time of treatment are low (<5 ng/mL). Options for therapy for those without evidence of distant metastases include monitoring with physical exam, PSA every 3 to 6 months, and imaging for symptoms or increasing PSA. For those deemed to be at higher risk of rapid progression, treatment options are ADT and local secondary therapy with or without ADT. For those with positive lymph nodes, abiraterone can be added. Treatment should be individualized based on the patient's risk of progression and the potential benefits and risks involved with each option.

Local secondary therapy options in the setting of RT recurrence include cryotherapy,⁴⁶³ HIFU,⁴⁶⁴⁻⁴⁶⁹ IRE^{470,471} (category 2B), reirradiation, and RP with PLND. The Panel recommends that patients receive multidisciplinary counseling about the risks and benefits of each of these options in the context of the available comparative literature on this topic.^{472,473} Reirradiation options include LDR brachytherapy, HDR brachytherapy, and SBRT based on phase II and other prospective and retrospective studies.^{412,474-476} Radical prostatectomy after RT recurrence can result in long-term disease control, but is often associated with impotence and urinary incontinence.⁴⁷⁷ Overall, there is limited evidence for these approaches, and the Panel recommends that patients receiving local therapy for RT recurrence are treated within the context of clinical trials when available and/or at experienced centers.

Second Biochemical Recurrence

Patients who have been treated for recurrence after definitive therapy for localized prostate cancer may continue to experience a rise in PSA levels without developing metastases. The Panel refers to this disease setting as

second biochemical recurrence. A staging workup with imaging is recommended. For patients without evidence of lymph node or distant metastases, risk stratification is essential to determine treatment options and is based mainly on PSA levels and PSADT. The definition of high-risk second biochemical recurrence differs between clinical trials but is generally considered to include patients with a PSADT ≤9 months and other adverse prognostic features.

For individuals deemed to be a low risk of rapid disease progression, monitoring until diagnosis of metastatic disease is preferred. ADT monotherapy is also an option, and PSA levels and PSADT should be considered when deciding whether to begin ADT.

For patients deemed to be at high risk of rapid disease progression, treatment intensification can be considered based on results of two trials as discussed below. Monitoring and ADT are equally reasonable options for these patients, and shared decision-making is key.

In the phase 3 EMBARK trial, 1068 patients with prostate cancer and high-risk biochemical recurrence were randomly assigned into one of three groups using a 1:1:1 ratio.⁴⁷⁸ The “combination group” received enzalutamide (160 mg/day) plus leuprolide, the “leuprolide-alone group” received placebo plus leuprolide, and the “monotherapy group” received enzalutamide alone. Eligibility for the trial included the following high-risk criteria: M0 by CT, MRI, or bone scan; PSADT ≤9 months; PSA ≥2 ng/mL above nadir after RT or ≥1 ng/mL after radical prostatectomy with or without postoperative RT; and patients were not considered a candidate for pelvic-directed therapy. Patients were monitored for a median follow-up time of 60.7 months. The trial demonstrated that the primary endpoint, metastasis-free survival, was significantly improved for enzalutamide plus leuprolide (HR for metastasis or death, 0.42; 95% CI, 0.30–0.61; $P < .001$) and for enzalutamide alone (HR, 0.63; 95% CI, 0.46–0.87; $P = .005$) when



NCCN Guidelines Version 5.2026

Prostate Cancer

compared to leuprolide alone. The most common adverse events associated with combination therapy and enzalutamide monotherapy were hot flashes and fatigue. Enzalutamide monotherapy was also significantly associated with gynecomastia (45% compared with 8% to 9% in the combination and leuprolide alone groups), nipple pain (15% compared with 1%–3%), and breast tenderness (14% compared with 1%). Based on these results, the Panel includes enzalutamide with or without leuprolide as options for patients with high-risk biochemical recurrence after maximal pelvic therapy in the absence of metastatic disease.

In the phase 3 PRESTO trial, 503 patients with biochemically recurrent prostate cancer were randomly assigned into one of three groups using a 1:1:1 ratio.⁴⁷⁹ The three arms of the study included: ADT alone (control arm A), ADT plus apalutamide (experimental arm B), or ADT plus apalutamide plus abiraterone acetate/prednisone (AAP; experimental arm C). The dose of apalutamide was 240mg/day. To be eligible for the trial, patients had prostate cancer with biochemical recurrence after radical prostatectomy and the following high-risk criteria: PSADT $\leq$ 9 months; PSA $\geq$ 0.5 ng/mL; and prior adjuvant or secondary RT or not considered a candidate for RT. The planned duration of treatment was 52 weeks. The primary endpoint of PSA progression-free survival (PSA-PFS) was met in patients in both experimental arms. PSA progression was defined as serum PSA $>$ 0.2 ng/mL after completion of treatment. At the first interim analysis (median follow-up of 21.3 and 21.5 months in arms B and C, respectively), both experimental arms had significant prolongation of PSA-PFS compared to the control arm (ADT plus apalutamide, 24.9 months; ADT plus apalutamide plus AAP, 26.0 months; ADT alone, 20.3 months). For the comparison of ADT with and without apalutamide, the HR was 0.52 (95% CI, 0.35–0.77; $P = .00047$), and for the comparison of ADT plus apalutamide/AAP the HR was 0.48 (95% CI, 0.32–0.71; $P = .00008$). There was no difference in median time to testosterone recovery across treatment arms. The most prevalent treatment-associated adverse events

across treatment arms were hot flashes (78%), fatigue (55%), injection site reaction (33%), and hypertension (31%). Hypertension was the most common grade $\geq$ 3 adverse event, experienced in 7%, 8%, and 19% of patients in arms A, B, and C, respectively. Even though the study was not designed to directly compare the two experimental arms, there was not shown to be further clinical benefit with the addition of AAP to apalutamide with ADT. Based on the results of this study, the Panel includes apalutamide plus ADT as a category 2B option for patients with non-metastatic disease and high-risk of biochemical recurrence after radical prostatectomy who have received maximal pelvic therapy.

Androgen Deprivation Therapy

ADT is administered as primary systemic therapy for regional or advanced disease and as neoadjuvant/concomitant/adjuvant therapy in combination with radiation in patients with unfavorable-intermediate or higher risk localized prostate cancers. It is also used as palliative therapy for symptomatic patients with a life expectancy $\leq$ 5 years.

Placing patients with early prostate cancer on ADT should not be routine practice.^{480,481} Bicalutamide monotherapy is not recommended to be used and is recommended when used to be in combination with ADT as per the FDA label.

Medical castration (ie, LHRH agonist or antagonist) and surgical castration (ie, bilateral orchiectomy) are equally effective options for ADT. Castrate levels of serum testosterone ($<$ 50 ng/dL; $<$ 1.7 nmol/L) should be achieved, because low nadir serum testosterone levels were shown to be associated with improved cause-specific survival in the PR-7 study.⁴⁸² Patients who do not achieve adequate suppression of serum testosterone ($<$ 50 ng/dL) with medical or surgical castration can be considered for additional hormonal manipulations (with antiandrogens, LHRH antagonists, or steroids), although the clinical benefit remains uncertain. Monitoring



NCCN Guidelines Version 5.2026

Prostate Cancer

testosterone levels 12 weeks after first dose of LHRH therapy and upon increase in PSA should be considered.

The LHRH antagonist, relugolix, has been studied as ADT in patients with localized and advanced prostate cancer.^{483,484} In the HERO study, 622 patients received relugolix (120 mg orally once daily) and 308 received leuprolide (injections every 3 months) for 48 weeks. Sustained castrate levels of testosterone (<50 ng per deciliter) through 48 weeks was superior with relugolix over leuprolide (96.7%; 95% CI, 94.9–97.9 vs. 88.8% [95% CI, 84.6–91.8]; $P < .001$ for superiority). Time to castration and testosterone recovery were both faster in the relugolix arm. The incidence of major adverse cardiovascular events was 2.9% in the relugolix arm and 6.2% in the leuprolide arm (HR, 0.46; 95% CI, 0.24–0.88).

ADT for Initial Treatment of Clinically Localized (N0,M0) Disease

ADT should not be used as monotherapy in clinically localized prostate cancer unless there is a contraindication to definitive local therapy, such as life expectancy less than 5 years and comorbidities (see *Palliative ADT*, below).

In the clinically localized setting, ADT using an LHRH agonist—alone or with a first-generation antiandrogen—or an LHRH antagonist can be used as a neoadjuvant, concurrent, and/or adjuvant to EBRT in patients with unfavorable intermediate-, high-, or very-high-risk prostate cancer, as described in more detail below.

ADT used as neoadjuvant treatment before radical prostatectomy is strongly discouraged outside of a clinical trial.

Neoadjuvant, Concurrent, and/or Adjuvant ADT with EBRT for Unfavorable Intermediate-Risk Disease

The MARCAP consortium performed an individual patient meta-analysis of nearly every randomized trial assessing the use and prolongation of ADT with EBRT in localized prostate cancer.³⁸² They demonstrated that the addition of 4 to 6 months of ADT improved OS for patients with intermediate- and high-risk disease. Additionally, the prolongation of adjuvant ADT for a total duration of 18 to 36 months further improved OS for patients primarily with high-risk prostate cancer. However, the prolongation of neoadjuvant ADT for multiple months did not significantly improve biochemical control, MFS, or OS. Furthermore, post-hoc analyses of RTOG 9408 and subgroup analyses of RTOG 0815 demonstrate that patients with multiple intermediate-risk features have a greater absolute benefit from ADT than those with a single intermediate-risk factor.^{381,383}

Thus, the Panel recommends 4 to 6 months of ADT when EBRT is given to patients as initial treatment of unfavorable intermediate-risk prostate cancer.

Two randomized phase 3 trials assessed the optimal sequencing of ADT with EBRT: RTOG 9413 and the Ottawa 0101 trials.⁴⁸⁵ These trials were pooled in an individual patient meta-analysis and demonstrated that concurrent/adjuvant ADT significantly improved biochemical control, distant metastasis, and MFS over a neoadjuvant/concurrent approach. Thus, concurrent/adjuvant ADT is preferred over neoadjuvant ADT in this setting.

Neoadjuvant, Concurrent, and/or Adjuvant ADT with EBRT for High-Risk or Very-High-Risk Disease

ADT combined with EBRT is an effective treatment for patients with high-risk or very-high-risk disease, as discussed in the *Radiation Therapy* section above. Combination therapy was consistently associated with



NCCN Guidelines Version 5.2026

Prostate Cancer

improved disease-specific survival and OS compared to single-modality treatment in randomized phase 3 studies.^{382,388,389,486-488}

Evidence favors long-term over short-term ADT for patients with high- and very-high-risk disease. MARCAP performed an individual patient level meta-analysis of every randomized trial that compared short-term versus long-term ADT with radiotherapy.³⁸² This study demonstrated that long-term ADT significantly improved MFS and OS over short-term ADT, with a greater absolute benefit in patients with high-/very high-risk disease as compared with intermediate-risk disease.

While the TROG RADAR trial demonstrated that in patients who received EBRT+HDR brachytherapy boost had an OS benefit from 18 months of ADT versus 6 months, the ASCENDE-RT trial of EBRT+LDR brachytherapy boost utilized only 12 months as their fixed duration of ADT.^{385,410} Thus, it may be reasonable for select patients with high-risk disease (not very high-risk) to use 12 months of ADT if receiving EBRT and LDR brachytherapy per the ASCENDE-RT trial regimen.

ADT for Initial Treatment of Regional Disease

One option for patients with regional disease is EBRT with 2 to 3 years of neoadjuvant/concurrent/adjuvant ADT with or without abiraterone. Neoadjuvant/concurrent/adjuvant ADT options are an LHRH agonist, an LHRH agonist with a first-generation antiandrogen, or an LHRH antagonist. Abiraterone should not be co-administered with an antiandrogen.

Data regarding the treatment of cN1 disease is limited. Exploratory analysis of STAMPEDE suggested that there may be benefit to the use of RT in this setting.⁴⁸⁹ Failure-free survival was better in those for whom RT was planned than for those receiving ADT alone (HR, 0.48; 95% CI, 0.29–0.79).

Another option for the treatment of patients initially diagnosed with node-positive disease is primary ADT. Primary ADT options are orchiectomy, an LHRH agonist, an LHRH agonist with a first-generation antiandrogen, or an LHRH antagonist; or orchiectomy, LHRH agonist, or LHRH antagonist with abiraterone.

Palliative ADT

Palliative ADT can be given to patients with a life expectancy of ≤ 5 years who have regional or metastatic prostate cancer. Palliative ADT also can be given to patients with disease progression during observation, usually when symptoms develop or when changes in PSA levels suggest that symptoms are imminent. The options in this setting are orchiectomy, LHRH agonist, or LHRH antagonist.

Primary ADT for mCSPC

ADT is strongly recommended in combination therapy for mCSPC (see Management of mCSPC, below). The use of ADT monotherapy in this setting is discouraged unless there are clear contraindications to combination therapy. If ADT monotherapy is given, intermittent ADT can be considered to reduce toxicity. Close monitoring of PSA and testosterone levels and possibly imaging is required when using intermittent ADT, especially during off-treatment periods, and patients may need to switch to continuous ADT upon signs of disease progression. ADT options for mCSPC include orchiectomy, LHRH agonist, or LHRH antagonist with various ARPIs and/or docetaxel. Metastasis-directed therapy or EBRT to the primary tumor are also options for certain patients. Specific recommendations depend on prior therapy and volume of disease as described in more detail below (see *Management of mCSPC*).

In patients with overt metastases in weight-bearing bone who are at risk of developing symptoms associated with the flare in testosterone with initial LHRH agonist alone, antiandrogen therapy should precede or be



NCCN Guidelines Version 5.2026

Prostate Cancer

coadministered with LHRH agonist for at least 7 days to diminish ligand binding to the androgen receptor.^{490,491} LHRH antagonists rapidly and directly inhibit the release of androgens, unlike LHRH agonists that initially stimulate LHRH receptors prior to hypogonadism. Therefore, no initial flare is associated with these agents and coadministration of antiandrogen is unnecessary.

Intermittent Versus Continuous ADT

ADT is associated with substantial side effects, which generally increase with the duration of treatment. Intermittent ADT is an approach based on the premise that cycles of androgen deprivation followed by re-exposure may delay “androgen independence,” reduce treatment morbidity, and improve QOL.^{492,493} Some patients who have no ADT-related morbidity may find the uncertainty of intermittent ADT not worthwhile. Intermittent ADT requires close monitoring of PSA and testosterone levels, especially during off-treatment periods, and patients may need to switch to continuous therapy upon signs of disease progression.

Intermittent ADT in Non-Metastatic Disease

The Canadian-led PR.7 trial was a phase 3 trial of intermittent versus continuous ADT in patients with non-metastatic prostate cancer who experienced biochemical recurrence after primary or post-recurrence EBRT.⁴⁹⁴ One thousand three hundred eighty-six patients with PSA >3 ng/mL were randomly assigned to intermittent ADT or continuous ADT. At a median follow-up of 6.9 years, the intermittent approach was noninferior to continuous ADT with respect to OS (8.8 vs. 9.1 years, respectively; HR, 1.02; 95% CI, 0.86–1.21). More patients died from prostate cancer in the intermittent ADT arm (120 of 690 patients) than in the continuous ADT arm (94 of 696 patients), but this was balanced by more non-prostate cancer deaths in the continuous ADT arm. Physical function, fatigue, urinary problems, hot flashes, libido, and erectile dysfunction showed modest improvement in the intermittent ADT group. The population of participants

was heterogeneous, so it remains unclear which of these asymptomatic patients benefited from treatment. It is possible that many of these patients could have delayed ADT without harm. The population had a low disease burden, and 59% of deaths in the trial were not related to prostate cancer. Follow-up longer than 6.9 years may be required for disease-specific deaths to out-balance deaths by other causes.

The multinational European ICELAND trial randomized 702 participants with locally advanced or biochemically recurrent prostate cancer to continuous or intermittent ADT.⁴⁹⁵ Clinical outcomes, which included time to PSA progression, PSA PFS, OS, mean PSA levels over time, QOL, and adverse events, were similar between the arms.

A 2015 meta-analysis identified 6 randomized controlled trials comparing continuous with intermittent ADT in patients with locally advanced prostate cancer and found no difference in mortality and progression and an advantage of the intermittent approach in terms of QOL and adverse effects.⁴⁹⁶

Therefore, the Panel believes that patients receiving ADT for M0 recurrent disease can consider intermittent ADT based on shared decision making.

Intermittent ADT in Metastatic Disease

Hussain and colleagues⁴⁹⁷ conducted the SWOG (Southwest Oncology Group) 9346 trial to compare intermittent and continuous ADT in patients with metastatic disease. After 7 months of induction ADT, 1535 patients whose PSA dropped to 4 ng/mL or below (thereby demonstrating androgen sensitivity) were randomized to intermittent or continuous ADT. At a median follow-up of 9.8 years, median survival was 5.1 years for the intermittent ADT arm and 5.8 years for the continuous ADT arm. The HR for death with intermittent ADT was 1.10 with a 90% CI between 0.99 and 1.23, which exceeded the prespecified upper boundary of 1.20 for noninferiority. The authors stated that the survival results were



NCCN Guidelines Version 5.2026

Prostate Cancer

inconclusive, and that a 20% greater mortality risk with the intermittent approach cannot be ruled out. The study demonstrated better erectile function and mental health in patients receiving intermittent ADT at 3 months, but the difference became insignificant thereafter, most likely due to contamination on the intermittent arm of patients who may have returned to continuous ADT at the prespecified time points. A secondary analysis of SWOG 9346 showed that intermittent ADT did not reduce endocrine, bone, or cognitive events, whereas it led to an unexpected increase in the incidence of ischemic and thrombotic events.⁴⁹⁸

In a post-hoc stratification analysis of the trial, patients with minimal disease had a median survival of 5.4 years when receiving intermittent ADT versus 6.9 years when receiving continuous ADT (HR, 1.19; 95% CI, 0.98–1.43).⁴⁹⁷ The median survival was 4.9 years in the intermittent ADT arm compared to 4.4 years in the continuous ADT arm for patients with extensive disease (HR, 1.02; 95% CI, 0.85–1.22). These subgroup analyses are hypothesis-generating.

A population-based analysis that included 9772 patients ≥66 years with advanced prostate cancer showed that intermittent ADT reduced the risks of total serious cardiovascular events by 36%, heart failure by 38%, and pathologic fracture by 48%, compared with continuous ADT.⁴⁹⁹ Furthermore, several meta-analyses of randomized controlled trials reported no difference in survival between intermittent ADT and continuous ADT,⁵⁰⁰⁻⁵⁰² although another analysis concluded that the non-inferiority of intermittent to continuous ADT in terms of survival has not been clearly demonstrated.⁵⁰³ Still, the intermittent approach leads to marked improvement in QOL compared to the continuous approach in most studies, and the Panel believes that intermittent ADT should be strongly considered in appropriate patients with metastatic disease.

Adverse Effects of Traditional ADT

ADT has a variety of adverse effects including hot flashes, vasomotor instability, loss of libido, erectile dysfunction, shrinkage of penis and testicles, loss of muscle mass and strength, fatigue, anemia, breast enlargement and tenderness/soreness, depression and mood swings, hair loss, osteoporosis, greater incidence of clinical fractures, obesity, insulin resistance, alterations in lipids, and greater risk for diabetes, acute kidney injury, and cardiovascular disease.⁵⁰⁴⁻⁵⁰⁶ The intensity and spectrum of these side effects vary greatly. In general, the side effects of continuous ADT increase with the duration of treatment. In addition, some forms of ADT may result in lower risk than others. For example, post-hoc analysis of a randomized trial of LHRH antagonist versus LHRH agonist found lower risk of cardiac events in patients with existing cardiac disease treated with LHRH antagonist.⁵⁰⁷ In addition, relugolix was associated with a lower risk of major adverse cardiovascular events than leuprolide in the phase 3 HERO study, although the FDA considered these results in HERO to be exploratory and therefore did not allow for these data to be included in the prescribing information for relugolix.⁴⁸³ Results from the randomized controlled PRONOUNCE trial suggest that there is no difference in cardiovascular safety between LHRH agonists and antagonists.⁵⁰⁸ However, all patients had cardiovascular risk factors optimized or addressed in both arms, which does not reflect real world practice.

Some evidence suggests that orchiectomy may be safer than an LHRH agonist. The heart and T lymphocytes have receptors for LHRH. Therefore, LHRH agonists may affect cardiac contractility, vascular plaque stability, and inflammation.⁵⁰⁹ Four hundred twenty-nine patients with metastatic prostate cancer who underwent orchiectomy were compared to 2866 patients who received LHRH agonist between 1995 and 2009.⁵¹⁰ Orchiectomy was associated with lower risk of fracture, peripheral arterial disease, and cardiac-related complications, although risk was similar for



NCCN Guidelines Version 5.2026

Prostate Cancer

diabetes, deep vein thrombosis, pulmonary embolism, and cognitive disorders.

Evidence suggests that a link between ADT and cognitive decline, dementia, or future Alzheimer's disease may exist, although data are inconsistent, the risk is low, and the link remains to be proven.⁵¹¹⁻⁵¹⁸ Some data suggests that the risk of cognitive effects may be increased with the addition of a second-generation ARPI.⁵¹⁹

Patients and their medical providers should be advised about these risks prior to treatment. Many side effects of ADT are reversible or can be avoided or mitigated. For example, physical activity can counter many of these symptoms and should be recommended (see NCCN Guidelines for Survivorship, available at www.NCCN.org). Specific recommendations for bone and cardiovascular health are included in the *Principles of Survivorship in Prostate Cancer* in the algorithm above and in the text that follows.

Bone Health During ADT

Medical or surgical ADT is associated with greater risk for osteoporosis and clinical fractures. In large population-based studies, for example, ADT was associated with a 21% to 54% relative increase in fracture risk.⁵²⁰⁻⁵²²

Longer treatment duration conferred greater fracture risk. Age and comorbidity also were associated with higher fracture incidence. In a population-based cohort of 3295 patients, surgical castration was associated with a significantly lower risk of fractures than medical castration using an LHRH agonist (HR, 0.77; 95% CI, 0.62–0.94; $P = .01$).⁵⁰⁹ ADT increases bone turnover and decreases bone mineral density,⁵²³⁻⁵²⁶ a surrogate for fracture risk in patients with non-metastatic disease. Bone mineral density of the hip and spine decreases by approximately 2% to 3% per year during initial therapy. Most studies have reported that bone mineral density continues to decline steadily during long-term therapy. ADT significantly decreases muscle mass,⁵²⁷ and

treatment-related sarcopenia appears to contribute to frailty and increased risk of falls in patients who are older.

The NCCN Guidelines Panel recommends screening and treatment for osteoporosis according to guidelines for the general population from the Bone Health and Osteoporosis Foundation, as outlined in the algorithm above.⁵²⁸

Earlier randomized controlled trials demonstrated that bisphosphonates increase bone mineral density, a surrogate for fracture risk, during ADT.⁵²⁹⁻⁵³¹ In 2011, the FDA approved denosumab as a treatment to prevent bone loss and fractures during ADT. Denosumab binds to and inhibits the receptor activator of NF- κ B ligand (RANKL) to blunt osteoclast function and delay generalized bone resorption and local bone destruction. Approval was based on a phase 3 study that randomized 1468 patients with non-metastatic prostate cancer undergoing ADT to either biannual denosumab or placebo. At 24 months, denosumab increased bone mineral density by 6.7% and reduced fractures (1.5% vs. 3.9%) compared to placebo.⁵³² Denosumab also was approved for prevention of SREs in patients with bone metastasis (see *Chemotherapy, Immunotherapy, and Targeted Therapy*).

Currently, treatment with denosumab, zoledronic acid, or alendronate is recommended when the absolute fracture risk warrants drug therapy, as outlined in the algorithm above. A baseline DEXA scan before start of therapy and a follow-up DEXA scan after 1 year of therapy is recommended by the International Society for Clinical Densitometry to monitor response. Use of biochemical markers of bone turnover is not recommended. There are no existing guidelines on the optimal frequency of vitamin D testing.



NCCN Guidelines Version 5.2026

Prostate Cancer

Diabetes and Cardiovascular Disease

Cardiovascular disease and diabetes are leading causes of morbidity and mortality in the general population. Cardiovascular disease is also a leading cause of death in patients with prostate cancer.⁵³³ In a landmark population-based study, ADT was associated with higher incidence of diabetes and cardiovascular disease.⁵³⁴ After controlling for other variables, which included age and comorbidity, ADT with an LHRH agonist was associated with increased risk for new diabetes (HR, 1.44; $P < .001$), coronary artery disease (HR, 1.16; $P < .001$), and myocardial infarction (HR, 1.11; $P = .03$). More recent data show that the main cause for the increased risk in cardiovascular disease in this population is the use of ADT and ARPIs.^{535,536}

Importantly, increased physical activity may decrease the symptoms and cardiovascular side effects of patients treated with ADT.⁵³⁷ Management of modifiable cardiovascular disease risk factors can also decrease the risk of subsequent cardiovascular events.

ADT may also prolong the QT/QTc interval. Providers should consider whether the benefits of ADT outweigh the potential risks in patients with congenital long QT syndrome, congestive heart failure, and frequent electrolyte abnormalities, and in patients taking drugs known to prolong the QT interval. Electrolyte abnormalities should be corrected, and periodic monitoring of electrocardiograms and electrolytes should be considered.

Based on the observed adverse metabolic effects of ADT and the association between ADT and higher incidence of diabetes and cardiovascular disease, screening for and intervention to prevent/treat diabetes and cardiovascular disease are recommended for patients receiving ADT, as described in the *Principles of Survivorship in Prostate Cancer*, in the algorithm above.

Management of mCSPC

ADT with treatment intensification is strongly recommended for patients with metastatic castration-sensitive prostate cancer (mCSPC). The use of ADT monotherapy in this setting is discouraged unless there are clear contraindications to combination therapy. Treatment intensification options include doublet therapy of ADT with an androgen receptor pathway inhibitor (ARPI; abiraterone, apalutamide, darolutamide, or enzalutamide); triplet therapy of ADT with docetaxel and one of the same ARPIs; metastasis-directed therapy (MDT) for oligometastases (see *MDT for Oligometastatic CSPC*, below); ADT with niraparib and abiraterone for *BRCA2*-mutated tumors; or ADT with EBRT to the primary tumor with or without docetaxel, abiraterone, apalutamide, or enzalutamide for low-metastatic burden (see *EBRT to the Primary Tumor in Low-Metastatic-Burden M1 Disease*, above). The specific recommended therapy options vary depending on whether the metastases were diagnosed in the synchronous or metachronous setting and on whether disease is oligometastatic, low-volume metastatic, or high-volume metastatic.

The data supporting doublet or triplet therapy in this setting are discussed below. For some of the combinations recommended by the Panel, supporting data are limited. They are included based on extrapolation from the studies of other agents, since the Panel considers the four ARPIs with approval in prostate cancer to be generally interchangeable.

Doublet Therapies for mCSPC

Abiraterone Acetate in mCSPC

In February 2018, the FDA approved abiraterone in combination with prednisone for mCSPC. This approval was based on two randomized phase 3 clinical trials of abiraterone and low-dose prednisone plus ADT in patients with newly diagnosed metastatic prostate cancer or high-risk or node-positive disease (STAMPEDE and LATITUDE) that demonstrated improved OS over ADT alone.⁵³⁸



NCCN Guidelines Version 5.2026

Prostate Cancer

In LATITUDE, 1199 patients with high-risk mCSPC were randomized to abiraterone with prednisone 5 mg once daily or matching placebos. High-risk disease was defined as at least two of the following: Gleason score 8–10, ≥ 3 bone metastases, and visceral metastases.⁵³⁸ Efficacy was demonstrated at the first interim analysis, and the trial was unblinded. The primary endpoint of OS was met and favored abiraterone (HR, 0.62; 95% CI, 0.51–0.76; $P < .0001$). Estimated 3-year OS rates improved from 49% to 66% at 30-month follow-up. Secondary endpoints were improved and included delayed castration-resistant radiographic progression (from median 14.8–33.2 months), PSA progression (7.4–33.2 months), time to pain progression, and initiation of chemotherapy. After the first interim analysis, 72 patients crossed over from placebo to abiraterone. Final OS analysis of LATITUDE after a median follow-up of 51.8 months showed median OS was significantly longer in the abiraterone group than in the placebo group (53.3 vs. 36.5 months; HR, 0.66; 95% CI, 0.56–0.78; $P < .0001$).⁵³⁹

Adverse events were higher with abiraterone and prednisone but were generally mild in nature and largely related to mineralocorticoid excess (ie, hypertension, hypokalemia, edema), hormonal effects (ie, fatigue, hot flashes), and liver toxicity.⁵³⁸ Cardiac events, such as atrial fibrillation, were rare but slightly increased with abiraterone. The overall discontinuation rate due to side effects was 12%. Patient-reported outcomes were improved with the addition of abiraterone, with improvements in pain intensity progression, fatigue, functional decline, prostate cancer-related symptoms, and overall health-related QOL.⁵⁴⁰ A limitation of this trial is that only 27% of placebo-treated patients received abiraterone or enzalutamide at progression, and only 52% of these patients received any life-prolonging therapy.⁵³⁸

The second randomized trial (STAMPEDE) of 1917 patients with CSPC demonstrated similar OS benefits.⁵⁴¹ However, unlike LATITUDE,

STAMPEDE eligibility permitted patients with high-risk N0,M0 disease (2 of 3 high-risk factors: stage T3/4, PSA >40 , or Gleason score 8–10; $n = 509$), or N1,M0 disease (pelvic nodal metastases; $n = 369$) in addition to patients with metastatic disease, who made up the majority of patients ($n = 941$). Most patients were newly diagnosed, while a minority had recurrent, high-risk, or metastatic disease after local therapy ($n = 98$). Thus, STAMPEDE was a heterogeneous mix of patients with high-risk, non-metastatic, node-positive, or metastatic disease. In patients with M1 disease, treatment with abiraterone plus prednisone was continued until progression. In patients with N1 or M0 disease, 2 years of abiraterone plus prednisolone was used if curative-intent EBRT was utilized. OS was improved in the overall population (HR, 0.63; 95% CI, 0.5–0.76; $P < .0001$) and in the M1 and N1 subsets, without any heterogeneity of treatment effect by metastatic status. The survival benefit of abiraterone was larger in patients <70 years of age than those ≥ 70 years (HR, 0.94 vs. HR, 0.51). Patients ≥ 70 years also suffered increased toxicities, which suggests heterogeneity in clinical benefits by age and comorbidity. The secondary endpoint of FFS, which included PSA recurrence, was improved overall (HR, 0.29; $P < .0001$) and in all subgroups regardless of M1 (HR, 0.31), N1 (HR, 0.29), or M0 (HR, 0.21) status. No heterogeneity for FFS was observed based on subgroups or by age. In this trial, subsequent life-prolonging therapy was received by 58% of those in the control group, which included 22% who received abiraterone and 26% who received enzalutamide. Thus, these data reflect a survival advantage of initial abiraterone in newly diagnosed patients compared with deferring therapy to the CRPC setting.

Adverse events in STAMPEDE were similar to that reported in LATITUDE, but were increased in patients ≥ 70 years, with higher incidences of grade 3–5 adverse events with abiraterone (47% vs. 33%) and 9 versus 3 treatment-related deaths. Severe hypertension or cardiac disorders were noted in 10% of patients and grade 3–5 liver toxicity in 7%, which



NCCN Guidelines Version 5.2026

Prostate Cancer

illustrates the need for blood pressure and renal and hepatic function monitoring.

Taken together, these data led the NCCN Panel to recommend abiraterone with 5-mg once-daily prednisone as a treatment option with ADT for patients with newly diagnosed mCSPC (category 1). Alternatively, the fine-particle formulation of abiraterone can be used with 4 mg methylprednisolone PO twice daily (category 2B; see *Abiraterone Acetate in mCRPC*, below).

The standard formulation of abiraterone can be given at 250 mg/day and administered following a low-fat breakfast as an alternative to the dose of 1000 mg/day after an overnight fast (see *Abiraterone Acetate in mCRPC*, below).⁵⁴² The cost savings may reduce financial toxicity and improve adherence in those who will not take or cannot afford the standard dose.

Apalutamide in mCSPC

The double-blind phase 3 TITAN clinical trial randomized 1052 patients with mCSPC to ADT with apalutamide (240 mg/day) or placebo.⁵⁴³

Participants were stratified by Gleason score at diagnosis, geographic region, and previous docetaxel treatment. The median follow-up was 22.7 months. Both primary endpoints were met: radiographic PFS (68.2% vs. 47.5% at 24 months; HR for radiographic progression or death, 0.48; 95% CI, 0.39–0.60; $P < .001$) and OS (82.4% vs. 73.5% at 24 months; HR for death, 0.67; 95% CI, 0.51–0.89; $P = .005$). Adverse events that were more common with apalutamide than with placebo included rash, hypothyroidism, and ischemic heart disease. Health-related QOL was maintained during treatment.⁵⁴⁴ At final analysis of TITAN, median OS was improved with apalutamide plus ADT compared with ADT alone after a median follow-up of 44 months (not reached vs. 52.2 months; HR, 0.65; 95% CI, 0.53–0.79; $P < .001$).⁵⁴⁵

Apalutamide is a category 1 option for patients with mCSPC. The FDA approved this indication in September 2019.

Enzalutamide in mCSPC

The open-label randomized phase 3 ENZAMET clinical trial compared enzalutamide (160 mg/day) plus ADT (LHRH analog or surgical castration) with a first-generation antiandrogen (bicalutamide, nilutamide, or flutamide) plus ADT in 1125 patients with mCSPC.⁵⁴⁶ Stratification was by volume of disease, planned use of early docetaxel, planned use of bone antiresorptive therapy, comorbidity score, and trial site. The primary endpoint of OS was met at the first interim analysis with median follow-up of 34 months (HR for death, 0.67; 95% CI, 0.52–0.86; $P = .002$). Enzalutamide also improved secondary endpoints, such as PFS using PSA levels and clinical PFS. An additional analysis was triggered at 470 deaths.⁵⁴⁷ After a median follow-up of 68 months, the 5-year OS rate was again lower in the first-generation antiandrogen group than in the enzalutamide group (HR, 0.70; 95% CI, 0.58–0.84; $P < .0001$). The median OS was not reached.

In the double-blind randomized phase 3 ARCHES clinical, 1150 patients with mCSPC were randomized to receive ADT with either enzalutamide (160 mg/day) or placebo. Participants were stratified by disease volume and prior docetaxel use. The primary endpoint was radiographic PFS, which was improved in the enzalutamide group after a median follow-up of 14.4 months (19.0 months vs. not reached; HR, 0.39; 95% CI, 0.30–0.50; $P < .001$).⁵⁴⁸ At the final, prespecified OS analysis, median OS was not met in either group, but a 34% reduction in the risk of death was observed in those receiving enzalutamide versus placebo (HR, 0.66; 95% CI, 0.53–0.81; $P < .001$).⁵⁴⁹ This result could be an underestimate of the effect of enzalutamide, since approximately 32% of the patients assigned placebo crossed over to enzalutamide after unblinding.



NCCN Guidelines Version 5.2026

Prostate Cancer

The safety of enzalutamide in these trials was similar to that seen in previous trials in the castration-resistant setting. Adverse events associated with enzalutamide in these trials included fatigue, seizures, and hypertension.^{546,548}

Enzalutamide is a category 1 option for patients with mCSPC. The FDA approved this indication in December 2019.

Darolutamide in mCSPC

The phase 3 ARANOTE trial assessed darolutamide with ADT compared with placebo and ADT in 669 patients with mCSPC.⁵⁵⁰ The primary endpoint of radiological PFS was improved in the darolutamide arm compared with the placebo arm (HR, 0.54; 95% CI, 0.41–0.71; $P < .0001$). The benefit was consistent across the low- and high-volume subgroups. Some of the secondary endpoints were also met, including delayed time to mCRPC and time to pain progression. However, a significant improvement in OS was not evident in the current follow-up (HR, 0.81; 95% CI, 0.59–1.12).

The FDA approved this indication for darolutamide in June 2025. The Panel include darolutamide with ADT as an option for patients with low- and high-volume mCSPC. Because an OS benefit has not been demonstrated for darolutamide doublet therapy, it is not a category 1 recommendation at this time.

Docetaxel in mCSPC

Docetaxel with ADT has been studied as an upfront option for patients with mCSPC in two phase 3 trials (ECOG 3805/CHAARTED and STAMPEDE).^{551,552} CHAARTED randomized 790 patients with mCSPC to docetaxel (75 mg/m² IV every 3 weeks x 6 doses) plus ADT or ADT alone.⁵⁵² After a median follow-up of 53.7 months, the patients in the combination arm experienced a longer OS than those in the ADT arm (57.6 vs. 47.2 months; HR, 0.72; 95% CI, 0.59–0.89; $P = .002$).⁵⁵³

Subgroup analysis showed that the survival benefit was more pronounced in the 65% of participants with high-volume disease (HR, 0.63; 95% CI, 0.50–0.79; $P < .001$). Patients with low metastatic burden in CHAARTED did not derive a survival benefit from the inclusion of docetaxel (HR, 1.04; 95% CI, 0.70–1.55; $P = .86$).

The STAMPEDE trial, a multi-arm, multi-stage phase 3 trial, included patients with both M0 and M1 CSPC.⁵⁵¹ The results in the M1 population confirmed the survival advantage of adding docetaxel (75 mg/m² IV every 3 weeks x 6 doses) to ADT seen in the CHAARTED trial. In STAMPEDE, extent of disease was not evaluated in the 1087 patients with metastatic disease, but the median OS for all patients with M1 disease was 5.4 years in the ADT-plus-docetaxel arm versus 3.6 years in the ADT-only arm (a difference of 1.8 years between groups compared with a 1.1-year difference in CHAARTED).

Patients with low metastatic burden did not have definitively improved survival outcomes in the ECOG CHAARTED study or a similar European trial (GETUG-AFU 15).^{552,554,555} Furthermore, the triplet options of ADT with docetaxel and either abiraterone or darolutamide showed improved OS over ADT with docetaxel (see below). The Panel therefore does not include docetaxel with ADT as an option for patients with mCSPC. Rather, patients with high-volume mCSPC who are fit for chemotherapy should be considered for triplet therapy.

Triplet Therapies for mCSPC

Data from the PEACE-1 and ARASENS trials indicate that triplet therapies of ADT with docetaxel and an ARPI—either abiraterone or darolutamide—improve OS over ADT with docetaxel in patients with high-volume metastatic CSPC.^{556,557} These trials are discussed below. Both of these combinations are included as category 1, preferred options for patients with high-volume mCSPC, and their use is encouraged for patients with



NCCN Guidelines Version 5.2026

Prostate Cancer

high-volume disease who are fit for chemotherapy. However, the Panel notes that no studies have compared doublet therapies (ADT plus an ARPI; discussed above) to triplet therapies. Therefore, doublet therapies are also suitable options for patients with high-volume metastatic disease.

These triplet combinations are also included as options in the low-volume, synchronous CSPC setting based on results of a meta-analysis, although their use in this setting is controversial and should be reserved for patients who desire aggressive treatment.⁵⁵⁸ No data support the use of triplet therapy in low-volume metachronous mCSPC, and they are not recommended in this setting.

Docetaxel Plus Abiraterone in CSPC

PEACE-1 was an international, open-label, randomized, phase 3 study conducted in seven European countries.⁵⁵⁶ Using a 2 × 2 factorial design, 1173 patients with de novo metastatic prostate cancer were randomized at a 1:1:1:1 ratio to standard of care (ADT alone or with docetaxel), standard of care with RT, standard of care with abiraterone, or standard of care with radiation and abiraterone. The two primary endpoints of the trial were radiographic PFS and OS. Adjusted Cox regression modelling showed no interaction between abiraterone and RT, so data were pooled for the analysis of abiraterone efficacy. Consistent with results of older studies, at a median follow-up of 3.5 years, radiographic PFS was longer in patients who received abiraterone than in those that did not (HR, 0.54; 99.9% CI, 0.41–0.71; $P < .0001$) as was OS (HR, 0.82; 95.1% CI, 0.69–0.98; $P = .030$). An OS benefit with abiraterone was also seen in the subset of patients with high metastatic burden as defined by CHAARTED criteria (HR, 0.77 ;95% CI, 0.62–0.96), but was not seen in those with low metastatic burden (HR, 0.93 ;95% CI, 0.69–1.28).

As part of the analysis, the efficacy of abiraterone was assessed in the population that received docetaxel. As in the overall population, radiographic PFS (HR, 0.50; 99.9% CI, 0.34–0.71; $P < .0001$) and OS

(HR, 0.75; 95.1% CI, 0.59–0.95; $P = .017$) were longer in those receiving all three therapies compared with those only receiving ADT and docetaxel. The populations receiving the triplet and doublet therapies experienced similar rates neutropenia, febrile neutropenia, fatigue, and neuropathy, although grade ≥3 adverse events occurred in 63% of patients who received the triplet combination compared with 52% of those receiving ADT and docetaxel.

Based on these data, the Panel includes this regimen as a category 1 option for patients with high-volume mCSPC.

Docetaxel Plus Darolutamide in CSPC

The international, phase 3 trial ARASENS trial, the second phase 3 trial evaluating a triplet therapy, randomized 1306 patients with mCSPC to receive ADT and docetaxel with either darolutamide or matching placebo.⁵⁵⁷ The primary endpoint, OS, was improved in the darolutamide group at 4 years (62.7%; 95% CI, 58.7–66.7) compared with the placebo group (50.4%; 95% CI, 46.3–54.6). The risk of death was lower in the darolutamide group by about 32% (HR, 0.68; 95% CI, 0.57–0.80; $P < .001$). The addition of darolutamide also showed significant benefits over placebo for secondary efficacy endpoints, including time to CRPC (HR, 0.36; 95% CI, 0.30–0.42; $P < .001$), skeletal event-free survival (HR, 0.61; 95% CI, 0.52–0.72; $P < .001$), and time to initiation of subsequent systemic antineoplastic therapy (HR, 0.39; 95% CI, 0.33–0.46; $P < .001$). Subgroup analysis showed a similar improvement in OS in those with high-volume disease as defined by CHAARTED criteria as in the overall population.⁵⁵⁹ An OS benefit was not observed in those with low metastatic burden (HR, 0.68; 95% CI, 0.41–1.13), but median survival was not reached in either arm.

Adverse events of any grade, grade 3 to 5 adverse events, and serious adverse events occurred at similar incidence levels between the two arms. Many of these were known effects of docetaxel. The most frequent



NCCN Guidelines Version 5.2026

Prostate Cancer

adverse events were alopecia (40.5% of patients in the darolutamide arm vs. 40.6% with placebo), neutropenia (39.3% vs. 38.8%), fatigue (33.1% vs. 32.9%), and anemia (27.8% vs. 25.1%). Exceptions were rash (16.6% vs. 13.5%) and hypertension (13.7% vs. 9.2%), which are known effects of androgen receptor pathway inhibitors and were more frequent in the darolutamide group.

The FDA approved this indication in August 2022, and the Panel includes this regimen as a category 1 option for patients with high-volume mCSPC.

PARP Inhibitor Combination for mCSPC

Niraparib Plus Abiraterone in BRCA2-Mutated mCSPC

ADT with PARP inhibitors alone or with ARPIs have been options for patients with HRR-mutated mCRPC for several years (see *Treatment Options for Patients with DNA Repair Gene Mutations*, below). The randomized, double-blind, phase 3 AMPLITUDE trial assessed ADT with niraparib and abiraterone compared with ADT plus abiraterone doublet therapy in 696 patients with homologous recombination repair (HRR)-deficient mCSPC.⁵⁶⁰ Most patients had high-volume disease (78%) and/or synchronous disease (87%) and most had *BRCA2* (42%) or *BRCA1* (7.2%) gene alterations. Sixteen percent of patients had received docetaxel. After a median follow-up of 30.8 months, the primary endpoint of median radiographic PFS (rPFS) was met in the *BRCA* subset (not reached vs. 26.0 months; HR, 0.52; 95% CI, 0.37–0.72; $P < .0001$) as well as in the full intention-to-treat population (not reached vs. 29.5 months; HR, 0.63; 95% CI, 0.49–0.80; $P = 0.0001$). This effect was largely driven by the *BRCA2*-mutated subset. OS data are immature, and the trial did not include formal cross-over to PARP inhibition for patients in the abiraterone group. Roughly one-third of participants in the control group received a PARP inhibitor following progression to mCRPC at the time of analysis.

Grade 3/4 adverse events were more common in the niraparib arm (75% vs. 59%) as were treatment-emergent adverse events leading to death (14 participants vs. 7 participants). The most common grade 3/4 adverse events were anemia (29.1% in the niraparib group versus 4.6% in the control group) and hypertension (26.5% versus 18.4%).

The FDA approved the combination of niraparib plus abiraterone with ADT for patients with *BRCA2*-mutated mCSPC in December 2025. It is included in these Guidelines as an option for patients with high-volume, synchronous or metachronous, mCSPC. It is also included for patients with synchronous or metachronous low-volume mCSPC (category 2B), but the Panel cautions that this regimen adds considerable toxicity and the benefit in this population is not yet clear due to the low numbers of such patients included in the AMPLITUDE trial. The panel notes that the niraparib plus abiraterone combination can be used following the completion of docetaxel in patients receiving triplet therapy or following EBRT to the primary tumor.

EBRT to the Primary Tumor in Synchronous Low-Volume M1 Disease

Patients with newly diagnosed, low-volume metastatic prostate cancer can be considered for ADT with EBRT to the primary tumor based on results from the randomized controlled phase 3 STAMPEDE trial.⁵⁶¹ In this multicenter, international study, 2061 patients were randomized to lifelong ADT with or without EBRT to the primary tumor (either 55 Gy in 20 daily fractions over 4 weeks or 36 Gy in 6 weekly fractions over 6 weeks). The primary outcome of OS by intention-to-treat (ITT) analysis was not met (HR, 0.92; 95% CI, 0.80–1.06; $P = .266$), but EBRT improved the secondary outcome of failure-free survival (FFS; HR, 0.76; 95% CI, 0.68–0.84; $P < .0001$). In a pre-planned subset analysis, outcomes of patients with high-metastatic burden (defined as visceral metastases; ≥ 4 bone metastases with ≥ 1 outside the vertebral bodies or pelvis; or both) and



NCCN Guidelines Version 5.2026

Prostate Cancer

those with low-metastatic burden (all others) were determined. EBRT improved OS (adjusted HR, 0.68; 95% CI, 0.52–0.90), prostate cancer-specific survival (adjusted HR, 0.65; 95% CI, 0.47–0.90), FFS (adjusted HR, 0.59; 95% CI, 0.49–0.72), and PFS (adjusted HR, 0.78; 95% CI, 0.63–0.98) in patients with low-metastatic burden, but not in patients with high metastatic burden. Long-term results have been reported, confirming the benefit of RT to the primary tumor in the setting of ADT with or without docetaxel.⁵⁶²

Abiraterone may be added to ADT with EBRT to the primary tumor in patients with low-volume synchronous metastases based on results of the PEACE-1, open-label, randomized trial.⁵⁶³ In this trial, participants were randomized 1:1:1:1 to ADT alone or with docetaxel (standard of care, SOC), SOC with abiraterone, SOC with radiation to the prostate, or SOC with abiraterone and radiation to the prostate. Results demonstrated that the addition of RT to the primary tumor in patients with low-volume disease treated with abiraterone led to improvements in median radiographic PFS (4.4 vs. 7.5 years). RT to the primary tumor also reduced rates of serious genitourinary toxicity regardless of disease volume, and time to castration resistance was delayed in the full population. Thus, some patients with high-volume disease may also benefit from RT to the primary tumor.

In PEACE-1, the benefits of RT to the primary tumor were only seen in patients receiving abiraterone, not in those receiving ADT alone or with docetaxel.⁵⁶³ However, in a secondary analysis of the STAMPEDE trial, the benefits of RT on OS and FFS in patients with low-volume disease were seen regardless of planned docetaxel use.⁵⁶⁴ The Panel therefore includes the addition of docetaxel to ADT and EBRT to the primary tumor as a category 2B recommendation for patients with low-volume synchronous mCSPC.

MDT for Oligometastatic CSPC

Treatment of metastatic sites with local therapy with the intent to improve oncologic outcomes (eg, delaying the initiation of systemic therapy or ADT; improving PFS, radiographic PFS, or OS) is known as metastasis-directed therapy, or MDT. MDT has been primarily studied as metastasis-directed RT (MDRT) and with the highly selected use of surgical lymph node dissection. MDRT is delivered at a higher-than-palliative dose to provide durable local control of the areas targeted and is the most used form of MDT.

MDT is used in patients with oligometastatic disease. The number of metastatic sites to define oligometastatic disease remains an evolving space and is impacted by the sensitivity of the imaging modality used. Early studies included patients with 1 to 3 or 1 to 5 metastatic sites by CT, MRI, or bone scan.⁵⁶⁵⁻⁵⁶⁸ More recent studies allow for up to 10 metastatic sites by PSMA-PET imaging (NCT04787744, NCT06150417, NCT03721341). The upper limit is not clearly established and is both a function of oncologic limitations and technical limitations of treating numerous metastatic sites. It should be noted that the goal of MDT is generally to treat all metastatic sites, which may include regional lymph nodes and, potentially, the primary tumor if untreated or if there is evidence of local recurrence. The primary tumor in this setting should be counted as a site. The Panel notes that general exclusion limits of >5 and >10 metastases by CT, MRI, or bone scan or by PSMA-PET imaging, respectively, are appropriate.

The best evidence supporting the use of MDT in the CSPC setting comes from randomized phase 2 studies in patients with metachronous oligorecurrent disease.^{565,568-570} These trials mostly used MDRT and showed that the approach improved ADT-free survival or PFS over monitoring or ADT. For example, the ORIOLE trial included 54 previously treated patients with 1 to 3 metastases by conventional imaging who



NCCN Guidelines Version 5.2026

Prostate Cancer

were randomized to receive MDRT or observation.⁵⁷⁰ Median PFS was better in the MDRT group than in the observation group (not reached vs. 5.8 months; HR, 0.30; 95% CI, 0.11–0.81; $P = .002$), and the treatment was well tolerated. STOMP randomized 62 patients with biochemically recurrent CSPC and ≤ 3 metastases to surveillance or to MDT with surgery or RT.⁵⁶⁹ The median ADT-free survival was improved in the MDT arm after a median 3-year follow-up (13 vs. 21 months; HR, 0.60; 80% CI, 0.40–0.90; log-rank $P = .11$). In a combined, longer-term analysis of ORIOLE and STOMP, median PFS was longer with MDT compared with observation (pooled HR, 0.44; 95% CI, 0.29–0.66; $P < .001$) after a median follow-up of 52.5 months.⁵⁶⁵

The EXTEND trial included a more heterogeneous group of patients, as 24 of the 87 enrolled patients had no prior definitive therapy to the prostate and 7 patients had mCRPC.⁵⁶⁸ Participants had ≤ 5 metastases amenable to MDRT and were randomized to MDRT with intermittent ADT or to intermittent ADT alone. After a median follow-up of 22 months, median PFS was improved in the MDRT/ADT group compared with the ADT-only group (not reached vs. 15.8 months; HR, 0.25; 95% CI, 0.12–0.55; $P < .001$). Analysis of a separate basket of participants in EXTEND who received continuous ADT have also been reported.⁵⁷¹ Results showed that the inclusion of MDT improved the primary endpoint of PFS improved in the participants who received continuous ADT (47 vs. 22 months; HR, 0.50; one-sided $P = .036$) and in the combined group of intermittent or continuous ADT (36 months vs. 17 months; HR, 0.45; $P < .001$).

The SABR-COMET phase 2 study, which enrolled patients with breast, lung, colorectal, and prostate cancers who had a controlled primary tumor and 1 to 5 metastases amenable to MDRT, showed an improvement on OS with an MDT approach.⁵⁷² SABR-COMET included 16 patients with prostate cancer; 14 were randomized to the MDRT arm, and 2 were randomized to receive palliative RT. Patients in both arms

received palliative systemic therapy as appropriate. After a median follow-up of 51 months, improvements were seen in 5-year OS rate (17.7% vs. 42.3%; 95% CI, 0.28–0.56; stratified log-rank $P = .006$) in the total population of 99 patients. A post-hoc sensitivity analysis was used to address the imbalance in the distribution of patients with prostate cancer between the two arms of the study. When patients with prostate cancer were excluded from the analysis, the 5-year OS rate continued to trend in favor of the MDRT group (16.2% vs. 33.1%; stratified log-rank test $P = .085$).

The benefit of adding ADT to MDRT in patients with oligorecurrent CSPC was assessed in the randomized, phase 2 RADIOSA trial.⁵⁶⁷ The 105 enrolled patients were randomized to 6 months of ADT with MDRT or MDRT alone. After a median follow-up of 31 months, the median clinical PFS was improved in the group receiving ADT (32.2 vs. 15.1 months; HR, 0.43; 95% CI, 0.26–0.72; $P = .001$).

Based on these data, the Panel recommends MDT with or without ADT as an option for patients with metachronous oligometastatic CSPC. These patients may alternatively be treated with ADT plus systemic therapy for low-volume mCSPC with or without concurrent MDT.

There is currently no randomized evidence in the synchronous oligometastatic setting, just single-arm prospective trials and retrospective cohorts. However, the Panel believes that concurrent MDT with recommended systemic therapy can be considered in select patients with synchronous oligometastatic disease.

Progression to and Management of CRPC

Most advanced disease eventually stops responding to traditional ADT and is categorized as castration-resistant (also known as castration-recurrent). CRPC is defined as prostate cancer that progresses clinically, radiographically, or biochemically despite castrate levels of serum



NCCN Guidelines Version 5.2026

Prostate Cancer

testosterone (<50 ng/dL).⁵⁷³ Patients whose disease progresses to CRPC during primary ADT should receive a laboratory assessment to assure a castrate level of testosterone (<50 ng/dL; <1.7 nmol/L). Imaging tests may be indicated to monitor for signs of distant metastases. Factors affecting the frequency of imaging include individual risk, age, overall patient health, PSA velocity, and Gleason grade.

For patients who develop CRPC, ADT with orchiectomy or an LHRH agonist or antagonist should be continued to maintain castrate serum levels of testosterone (<50 ng/dL).

Patients with CRPC and no signs of distant metastasis on conventional imaging studies (M0) can consider monitoring with continued ADT if PSADT is >10 months (preferred), because these patients will have a relatively indolent disease history.⁵⁷⁴ Secondary hormone therapy with continued ADT is an option mainly for patients with shorter PSADT (≤10 months) as described below.

For patients who develop mCRPC, metastatic lesion biopsy is recommended, as is MSI/MMR testing, if not previously performed. If MSI-H or dMMR is found, referral to genetic counseling should be made to assess for the possibility of Lynch syndrome. These patients should also have germline and tumor testing to check for mutations in homologous recombination repair (HRR) genes (eg, *BRCA1*, *BRCA2*, *ATM*, *PALB2*, *FANCA*, *RAD51D*, *CHEK2*, *CDK12*) if not done previously.⁵⁷⁵ This information may be used for genetic counseling, cascade germline testing for family members, early use of platinum chemotherapy, and understanding eligibility for biomarker-directed treatments or clinical trials. TMB testing is also recommended for patients with mCRPC.

ADT is continued in patients with mCRPC while additional therapies, including secondary hormone therapies, chemotherapies, immunotherapies, radiopharmaceuticals, and/or targeted therapies, are

applied sequentially or concurrently, as discussed in the sections that follow; all patients should receive best supportive care. The Panel defined treatment options for patients with mCRPC based on previous exposure to ARPIs (abiraterone, enzalutamide, darolutamide, or apalutamide) and docetaxel. Abiraterone given as part of neoadjuvant/concomitant/adjuvant ADT with EBRT is not considered prior ARPI therapy.

The decision to initiate therapy in the CRPC setting after disease progression on one or more treatments should be based on the available high-level evidence of safety, efficacy, and tolerability of these agents and the application of this evidence to an individual patient. Prior exposures to therapeutic agents should be considered. Data to inform the optimal sequence for delivery of these agents in patients with mCRPC is evolving (see *Sequencing of Therapy in CRPC*, below). Choice of therapy is based largely on clinical considerations, which include patient preferences, prior treatment, presence or absence of visceral disease, symptoms, and potential side effects.

NCCN recommends that patients being treated for CRPC be closely monitored with radiologic imaging (ie, CT, bone imaging), PSA tests, and clinical exams for evidence of progression. Therapy should be continued until clinical progression or intolerability, with consideration of the fact that even in cases where PSA remains undetectable, bone imaging may reveal progression.^{576,577} The sequential use of these agents is recommended in patients who remain candidates for further systemic therapy. The Panel also notes that Pan-cancer, tumor-agnostic treatments can be considered for patients with mCRPC who have actionable mutations. Clinical trial and best supportive care are additional options.

Secondary Hormone Therapy for CRPC

Research has shown enhancement of autocrine and/or paracrine androgen synthesis in the tumor microenvironment of patients receiving



NCCN Guidelines Version 5.2026

Prostate Cancer

ADT.^{578,579} Androgen signaling consequent to non-gonadal sources of androgen in CRPC refutes earlier beliefs that CRPC was resistant to further hormone therapies. The development of novel ARPIs demonstrating efficacy in the non-metastatic CRPC and mCRPC settings dramatically changed the paradigm of CRPC treatment over the past two decades.

Abiraterone Acetate in mCRPC

In April 2011, the FDA approved the androgen synthesis inhibitor, abiraterone, in combination with low-dose prednisone, for the treatment of patients with mCRPC who have received prior chemotherapy containing docetaxel. This FDA approval in the post-docetaxel, mCRPC setting was based on the results of a phase 3, randomized, placebo-controlled trial (COU-AA-301) in patients with mCRPC previously treated with docetaxel-containing regimens.^{580,581} Patients were randomized to receive either abiraterone 1000 mg orally once daily (n = 797) or placebo once daily (n = 398), and both arms received daily prednisone. In the final analysis, median survival was 15.8 versus 11.2 months in the abiraterone and placebo arm, respectively (HR, 0.74; 95% CI, 0.64–0.86; $P < .0001$).⁵⁸¹ Time to radiographic progression, PSA decline, and pain palliation also were improved by abiraterone.^{581,582}

FDA approval in the pre-docetaxel setting occurred in December 2012, and was based on the randomized phase 3 COU-AA-302 trial of abiraterone and prednisone (n = 546) versus prednisone alone (n = 542) in patients with asymptomatic or minimally symptomatic, mCRPC.⁵⁸³ Most participants in this trial were not taking narcotics for cancer pain and none had visceral metastatic disease or prior ketoconazole exposure. The coprimary endpoint of radiographic PFS was improved from 8.3 to 16.5 months with abiraterone (HR, 0.53; $P < .001$). OS was improved at final analysis with a median follow-up of 49.2 months (34.7 vs. 30.3 months; HR, 0.81; 95% CI, 0.70–0.93; $P = .003$).⁵⁸⁴ Key secondary endpoints of

time to symptomatic deterioration, time to chemotherapy initiation, time to pain progression, and PSA PFS improved significantly with abiraterone treatment; PSA declines (62% vs. 24% with >50% decline) and radiographic responses (36% vs. 16% RECIST responses) were more common.

The most common adverse reactions with abiraterone/prednisone (>5%) were fatigue (39%); back or joint discomfort (28%–32%); peripheral edema (28%); diarrhea, nausea, or constipation (22%); hypokalemia (17%); hypophosphatemia (24%); atrial fibrillation (4%); muscle discomfort (14%); hot flushes (22%); urinary tract infection; cough; hypertension (22%, severe hypertension in 4%); urinary frequency and nocturia; dyspepsia; or upper respiratory tract infection. The most common adverse drug reactions that resulted in drug discontinuation were increased aspartate aminotransferase and/or alanine aminotransferase (11%–12%), or cardiac disorders (19%, serious in 6%).

Based on the studies described here, abiraterone is a category 1, preferred option for mCRPC without prior ARPI therapy. It can also be considered in patients with mCRPC following progression on another ARPI, although other therapies are preferred in this setting.

In May 2018, the FDA approved a novel, fine-particle formulation of abiraterone, in combination with methylprednisolone, for the treatment of patients with mCRPC. In studies of healthy males, this formulation at 500 mg was shown to be bioequivalent to 1000 mg of the originator formulation.^{585,586} In a phase 2 therapeutic equivalence study, 53 patients with mCRPC who were not treated previously with abiraterone, enzalutamide, radium-223, or chemotherapy (docetaxel for mCRPC completed ≥ 1 year prior to enrollment was allowed) were randomized to 500 mg daily of the new, fine-particle formulation plus 4 mg methylprednisolone orally twice daily or to 1000 mg of the originator formulation daily plus 5 mg prednisone orally twice daily.⁵⁸⁷



NCCN Guidelines Version 5.2026

Prostate Cancer

Bioequivalence of these doses was confirmed based on serum testosterone levels, PSA response, and abiraterone pharmacokinetics. The rates of total and grade 3/4 adverse events were similar between the arms, with musculoskeletal and connective tissue disorders occurring more frequently in the originator-treated patients (37.9% vs. 12.5%). The Panel believes that the fine-particle formulation of abiraterone can be used instead of the original formulation of abiraterone in the treatment of patients with mCRPC (category 2A).

Abiraterone should be given with concurrent steroid (either oral prednisone 5 mg twice daily or oral methylprednisolone 4 mg twice daily, depending on which formulation is given) to abrogate signs of mineralocorticoid excess that can result from treatment. These signs include hypertension, hypokalemia, and peripheral edema. Thus, monitoring of liver function, potassium and phosphate levels, and blood pressure readings on a monthly basis is warranted during abiraterone therapy. Symptom-directed assessment for cardiac disease also is warranted, particularly in patients with pre-existing cardiovascular disease.

A randomized phase 2 noninferiority study of 75 patients with mCRPC compared 1000 mg/day abiraterone after an overnight fast with 250 mg/day after a low-fat breakfast.⁵⁴² The primary endpoint was log change in PSA, with secondary endpoints of PSA response ($\geq 50\%$) and PFS. The primary endpoint favored the low-dose arm (log change in PSA, -1.59 vs. -1.19), as did the PSA response rate (58% vs. 50%), with an equal PFS of 9 months in both arms. Noninferiority of the low dose was established according to the predefined criteria. Therefore, abiraterone can be given at 250 mg/day administered following a low-fat breakfast, as an alternative to the dose of 1000 mg/day after an overnight fast in patients who will not take or cannot afford the standard dose. The cost savings may reduce financial toxicity and improve adherence. Food impacts absorption unpredictably; therefore, side effects should be monitored and standard

dosing (1000 mg on empty stomach) utilized if excess toxicity is observed on modified dosing (250 mg with food).

Abiraterone with Dexamethasone in mCRPC

Switching from prednisone to dexamethasone 0.5 mg/day can be considered for patients with mCRPC with disease progression on either formulation of abiraterone. Trials show improved PSA responses and PFS and acceptable safety using this strategy.^{588,589}

The SWITCH study was a single-arm, open-label, phase 2 study of this approach with 26 enrolled patients.⁵⁸⁸ The primary endpoint, the proportion of patients with a PSA decline $\geq 30\%$ in 6 weeks, was 46.2%. No significant toxicities were observed, and two radiologic responses were seen. In another study, 48 consecutive patients with mCRPC, with disease progression on abiraterone with prednisone, were switched to abiraterone with 0.5 mg/day dexamethasone.⁵⁸⁹ The primary endpoint of median PFS was 10.35 months, and PSA levels decreased or stabilized in 56% of patients after switching to dexamethasone.

Enzalutamide in M0 and M1 CRPC

In August 2012, the FDA approved enzalutamide, a next-generation antiandrogen, for treatment of patients with mCRPC who had received prior docetaxel chemotherapy. Approval was based on the results of the randomized, phase 3, placebo-controlled AFFIRM trial.^{590,591} AFFIRM randomized 1199 patients to enzalutamide (160 mg daily) or placebo in a 2:1 ratio and the primary endpoint was OS. Median survival was improved with enzalutamide from 13.6 to 18.4 months (HR, 0.63; $P < .001$). Survival was improved in all subgroups analyzed. Secondary endpoints were also improved significantly, which included the proportion of patients with $>50\%$ PSA decline (54% vs. 2%), radiographic response (29% vs. 4%), radiographic PFS (8.3 vs. 2.9 months), and time to first skeletal-related event (SRE) (16.7 vs. 13.3 months). QOL measured using validated



NCCN Guidelines Version 5.2026

Prostate Cancer

surveys was improved with enzalutamide compared to placebo. Adverse events were mild, and included fatigue (34% vs. 29%), diarrhea (21% vs. 18%), hot flushes (20% vs. 10%), headache (12% vs. 6%), and seizures (0.6% vs. 0%). The incidence of cardiac disorders did not differ between the arms. Patients in the AFFIRM study were maintained on LHRH agonist/antagonist therapy and could receive bone supportive care medications. The seizure risk in the enzalutamide FDA label was 0.9% versus 0.6% in the manuscript.^{590,592}

Another phase 3 trial studied enzalutamide in the pre-chemotherapy setting. The PREVAIL study randomly assigned 1717 patients with chemotherapy-naïve metastatic prostate cancer to daily enzalutamide or placebo.^{593,594} The study was stopped early due to benefits shown in the treatment arm. Compared to the placebo group, the enzalutamide group showed improved median PFS (20.0 vs. 5.4 months) and median OS (35.3 vs. 31.3 months). Improvements in all secondary endpoints were also observed (eg, the time until chemotherapy initiation or first SRE).

Thus, enzalutamide represents a category 1, preferred treatment option for patients with mCRPC without prior ARPI therapy. It can also be considered in patients with mCRPC with prior exposure to another ARPI, although other therapies are preferred in this setting.

The randomized, double-blind, placebo-controlled phase 3 PROSPER trial assessed the use of enzalutamide in 1401 patients with non-metastatic CRPC.⁵⁹⁵ Patients with PSADT ≤ 10 months were stratified according to PSADT (< 6 vs. ≥ 6 months) and use of bone-sparing agents and randomized 2:1 to enzalutamide (160 mg/day) plus ADT or placebo plus ADT. Enzalutamide improved the primary endpoint of metastasis-free survival over placebo (36.6 vs. 14.7 months; HR for metastasis or death, 0.29; 95% CI, 0.24–0.35; $P < .0001$). Median OS was longer in the enzalutamide group than in the placebo group (67.0 vs. 56.3 months; HR for death, 0.73; 95% CI, 0.61–0.89; $P = 0.001$).⁵⁹⁶ Adverse events included

fatigue (33% vs. 14%), hypertension (12% vs. 5%), major adverse cardiovascular events (5% vs. 3%), and mental impairment disorders (5% vs. 2%). Patient-reported outcomes from PROSPER indicate that enzalutamide delayed pain progression, symptom worsening, and decrease in functional status, compared with placebo.⁵⁹⁷

The FDA expanded its approval for enzalutamide to include patients with non-metastatic CRPC in July 2018, and the Panel believes that patients with M0 CRPC can be offered enzalutamide, if PSADT is ≤ 10 months (category 1, preferred option).

Patients receiving enzalutamide have no restrictions for food intake and concurrent prednisone is permitted but not required.⁵⁹⁰

Apalutamide in M0 CRPC

The FDA approved apalutamide for treatment of patients with non-metastatic CRPC in February 2018. This approval was based on the phase 3 SPARTAN trial of 1207 patients with M0 CRPC and PSADT ≤ 10 months.⁵⁹⁸ Participants were stratified according to PSADT (> 6 vs. ≤ 6 months), use of bone-sparing agents, and the presence of metastatic pelvic lymph nodes (N0 vs. N1). After a median follow-up of 20.3 months, apalutamide at 240 mg/day with ADT improved the primary endpoint of metastasis-free survival over placebo with ADT (40.5 vs. 16.2 months; HR for metastasis or death, 0.28; 95% CI, 0.23–0.35; $P < .001$). Adverse events included rash (24% vs. 5.5%), fracture (11% vs. 6.5%), and hypothyroidism (8% vs. 2%). In a prespecified exploratory analysis of SPARTAN, health-related QOL was maintained in both the apalutamide and placebo groups.⁵⁹⁹

After a median follow-up of 52 months, final OS analysis showed that participants in SPARTAN experienced an improved median OS with apalutamide versus placebo (73.9 vs. 59.9 months; HR, 0.78; 95% CI, 0.64–0.96; $P = .016$).⁶⁰⁰ This longer OS reached prespecified statistical



NCCN Guidelines Version 5.2026

Prostate Cancer

significance, even though 19% of participants crossed over from placebo to apalutamide.

Apalutamide is a category 1, preferred option for patients with M0 CRPC if PSADT is ≤ 10 months.

Darolutamide in M0 CRPC

The FDA approved darolutamide for treatment of patients with non-metastatic CRPC in July 2019. The phase 3 ARAMIS study randomized 1509 patients with M0 CRPC and PSADT ≤ 10 months 2:1 to darolutamide (600 mg twice daily) or placebo.⁶⁰¹ Participants were stratified according to PSADT (>6 vs. ≤ 6 months) and the use of osteoclast-targeted agents. The median follow-up time was 17.9 months. Darolutamide improved the primary endpoint of metastasis-free survival compared to placebo (40.4 vs. 18.4 months; HR for metastasis or death, 0.41; 95% CI, 0.34–0.50; $P < .001$).

Patients in the placebo group of ARAMIS crossed over to darolutamide ($n = 170$) or received other life-prolonging therapy ($n = 137$). Final analysis occurred after a median follow-up time of 29.0 months. The risk of death was 31% lower in the darolutamide group than in the placebo group (HR for death, 0.69; 95% CI, 0.53–0.88; $P = .003$).⁶⁰² OS at 3 years was 83% (95% CI, 80–86) in the darolutamide group compared with 77% (95% CI, 72–81) in the placebo group. Adverse events that occurred more frequently in the treatment arm included fatigue (12.1% vs. 8.7%), pain in an extremity (5.8% vs. 3.2%), and rash (2.9% vs. 0.9%). The incidence of fractures was similar between darolutamide and placebo (4.2% vs. 3.6%).⁶⁰¹

Darolutamide is a category 1, preferred option for patients with M0 CRPC if PSADT is ≤ 10 months.

Other Secondary Hormone Therapies

Other options for secondary hormone therapy include a first-generation antiandrogen, antiandrogen withdrawal, corticosteroid, or ketoconazole (adrenal enzyme inhibitor) with hydrocortisone.^{603–605} However, none of these strategies has been shown to prolong survival in randomized clinical trials. In the mCRPC setting, these options should only be used for select patients who are not candidates for other recommended mCRPC therapies.

A randomized phase 2 trial, TRANSFORMER, compared the effect of bipolar androgen therapy (BAT) with that of enzalutamide on PFS in 195 patients with asymptomatic, mCRPC with prior progression on abiraterone.⁶⁰⁶ BAT involves rapid cycling between high and low serum testosterone to disrupt the adaptive upregulation of the androgen receptor that occurs with low testosterone levels. Patients in the BAT arm received testosterone cypionate 400 mg intramuscularly once every 28 days. The PFS was 5.7 months in both arms (HR, 1.14; 95% CI, 0.83–1.55; $P = .42$). Crossover was allowed after disease progression, and OS was similar between the groups. BAT resulted in more favorable patient-reported QOL. The Panel awaits more data on this approach.

Chemotherapy, Immunotherapy, and Targeted Therapy for mCRPC

Research has expanded the therapeutic options for patients with mCRPC. In addition to the hormonal and radiopharmaceutical therapies described in other sections, options include chemotherapy, immunotherapy, and targeted therapy. As noted above, selection of therapy depends on patient preferences, prior treatment exposures, the presence or absence of symptoms, the location of metastases, the presence of certain biomarkers, and consideration of potential side effects.



NCCN Guidelines Version 5.2026

Prostate Cancer

Docetaxel

Docetaxel was FDA-approved for mCRPC in May 2004. Two randomized phase 3 studies evaluated docetaxel-based regimens in symptomatic or rapidly progressive CRPC (TAX 327 and SWOG 9916).⁶⁰⁷⁻⁶⁰⁹ TAX 327 compared docetaxel (every 3 weeks or weekly) plus prednisone to mitoxantrone plus prednisone in 1006 patients.⁶⁰⁸ Every-3-week docetaxel resulted in higher median OS than mitoxantrone (18.9 vs. 16.5 months; $P = .009$). This survival benefit was maintained at extended follow-up.⁶⁰⁹ The SWOG 9916 study showed improved survival with docetaxel when combined with estramustine compared to mitoxantrone plus prednisone.⁶⁰⁷

Thus, docetaxel is a category 1 option for treatment of docetaxel-naïve mCRPC. It is the preferred option post-ARPI in patients without prior docetaxel exposure.

The standard regimen is 75 mg/m² every 3 weeks. An alternative to every-3-week docetaxel is a biweekly regimen of 50 mg/m². This regimen is based on a large randomized phase 2 trial of 346 patients with mCRPC randomized to either every-2-week docetaxel or every-3-week docetaxel, each with maintenance of ADT and prednisone.⁶¹⁰ Patients treated with the every-2-week regimen survived an average of 19.5 months compared to 17.0 months with the every-3-week regimen ($P = .015$). Time to progression and PSA decline rate favored every-2-week therapy. Tolerability was improved with every-2-week docetaxel; febrile neutropenia rate was 4% versus 14% and other toxicities and overall QOL were similar.

The duration of docetaxel therapy should be based on the assessment of benefit and toxicities. Treatment with ≥ 8 cycles of docetaxel may be associated with better OS than fewer cycles in the mCRPC setting.⁶¹¹

Retrospective analysis from the GETUG-AFU 15 trial suggests that docetaxel may benefit some patients with CRPC who received docetaxel

in the CSPC setting.⁶¹² Thus, the Panel believes that docetaxel can be given as a rechallenge after progression on an ARPI in the mCRPC setting if the patient's cancer did not demonstrate definitive evidence of progression on prior docetaxel therapy in either the castration-sensitive setting or the mCRPC setting.

Adverse events associated with docetaxel include neutropenia, leukopenia, febrile neutropenia, neutropenic infections, fluid retention, hypersensitivity reaction, hepatic function impairment, neuropathy, and other low-grade adverse events (eg, fatigue, nausea, vomiting, alopecia, diarrhea).

Cabazitaxel

In June 2010, the FDA approved cabazitaxel, a semi-synthetic taxane derivative, for patients with mCRPC previously treated with a docetaxel-containing regimen. An international randomized phase 3 trial (TROPIC) randomized 755 patients with progressive mCRPC to receive cabazitaxel 25 mg/m² or mitoxantrone 12 mg/m², each with daily prednisone.⁶¹³ A 2.4-month improvement in OS was demonstrated with cabazitaxel compared to mitoxantrone (HR, 0.72; $P < .0001$). The improvement in survival was balanced against a higher toxic death rate with cabazitaxel (4.9% vs. 1.9%), which was due, in large part, to differences in rates of sepsis and renal failure. Febrile neutropenia was observed in 7.5% of patients treated with cabazitaxel versus 1.3% of patients treated with mitoxantrone. The incidences of severe diarrhea (6%), fatigue (5%), nausea/vomiting (2%), anemia (11%), and thrombocytopenia (4%) also were higher in patients treated with cabazitaxel, which indicated the need for vigilance and treatment or prophylaxis in this setting to prevent febrile neutropenia. The survival benefit was sustained at an updated analysis with a median follow-up of 25.5 months.⁶¹⁴ Furthermore, results of a post-hoc analysis of this trial suggested that the occurrence of grade ≥ 3 neutropenia after



NCCN Guidelines Version 5.2026

Prostate Cancer

cabazitaxel treatment was associated with improvements in both PFS and OS.⁶¹⁵

The multicenter CARD study was a randomized, open-label clinical trial that compared cabazitaxel with either abiraterone or enzalutamide in 255 patients with mCRPC who had previously received docetaxel and either abiraterone or enzalutamide.⁶¹⁶ Cabazitaxel at 25 mg/m² with concurrent steroid improved the primary endpoint of radiographic PFS (8.0 vs. 3.7 months; HR, 0.54; $P < .0001$) and reduced the risk of death (13.6 vs. 11.0 months; HR, 0.64; $P = .008$) compared with abiraterone or enzalutamide in these patients. Cabazitaxel was also associated with an increased rate of pain response and delayed time to pain progression and SREs.⁶¹⁷

The phase 3 open-label, multinational, noninferiority PROSELICA study compared 20 mg/m² cabazitaxel with 25 mg/m² cabazitaxel in 1200 patients with mCRPC who progressed on docetaxel.⁶¹⁸ The lower dose was found to be noninferior to the higher dose for median OS (13.4 months [95% CI, 12.19–14.88] vs. 14.5 months [95% CI, 13.47–15.28]), and grade 3/4 adverse events were decreased (39.7% vs. 54.5%). In particular, grade ≥3 neutropenia rates were 41.8% and 73.3% for the lower and higher dose groups, respectively.

Results from the phase 3 FIRSTANA study suggest that cabazitaxel has clinical activity in patients with chemotherapy-naïve mCRPC.⁶¹⁹ Median OS, the primary endpoint, was similar between 20 mg/m² cabazitaxel, 25 mg/m² cabazitaxel, and 75 mg/m² docetaxel (24.5 months, 25.2 months, and 24.3 months, respectively). Cabazitaxel was associated with lower rates of peripheral sensory neuropathy than docetaxel, particularly at 20 mg/m² (12% vs. 25%). However, the Panel does not currently recommend cabazitaxel in docetaxel-naïve patients.

Based on these data, cabazitaxel is included in these Guidelines as a category 1, preferred option after exposure to docetaxel and an ARPI in

patients with mCRPC. Cabazitaxel at 20 mg/m² every 3 weeks, with or without growth factor support, is the recommended dose for fit patients. Cabazitaxel at 25 mg/m² may be considered for healthy patients who opt for more aggressive treatment. Biweekly cabazitaxel at 16 mg/m² with prophylactic G-CSF is an option for patients ≥65 years based on results from the phase 3 CABASTY trial.⁶²⁰ Using the lower dose significantly reduced the risk of neutropenia/neutropenic complications compared to the 25 mg/m² dose with G-CSF. Clinical outcomes were comparable between the 2 groups.

Cabazitaxel should be given with concurrent steroids (daily prednisone or dexamethasone on the day of chemotherapy). Physicians should follow current guidelines for prophylactic white blood cell growth factor use, particularly in this heavily pretreated population. In addition, supportive care should include antiemetics (prophylactic antihistamines, H2 antagonists, and corticosteroids prophylaxis) and symptom-directed antidiarrheal agents. Cabazitaxel was tested in patients with hepatic dysfunction in a small, phase I, dose-escalation study.⁶²¹ Cabazitaxel was tolerated in patients with mild to moderate hepatic impairment. However, cabazitaxel should not be used in patients with severe hepatic dysfunction. Cabazitaxel should be stopped upon clinical disease progression or intolerance.

Cabazitaxel/Carboplatin

Cabazitaxel 20 mg/m² plus carboplatin AUC 4 mg/mL per minute with growth factor support can be considered post-ARPI for fit patients with aggressive variant mCRPC (visceral metastases, low PSA and bulky disease, high lactate dehydrogenase [LDH], high carcinoembryonic antigen [CEA], lytic bone metastases, and neuroendocrine prostate cancer [NEPC] histology) or unfavorable genomics (defects in at least 2 of *PTEN*, *TP53*, and *RB1*). This recommendation is based on a phase 1–2, open label, randomized study.⁶²² In the phase 2 portion, 160 patients were



NCCN Guidelines Version 5.2026

Prostate Cancer

randomized to receive cabazitaxel alone or with carboplatin, and the primary endpoint was investigator-assessed PFS. In the ITT population, median PFS was 4.5 months in the cabazitaxel arm versus 7.3 months in the cabazitaxel/carboplatin arm (HR, 0.69; 95% CI, 0.50–0.95; $P = .018$). The most common grade 3–5 adverse events (fatigue, anemia, neutropenia, and thrombocytopenia) were all more common in the combination arm. Post-hoc analyses showed that patients with aggressive variant disease had a longer median PFS in the combination arm than the cabazitaxel arm (7.5 vs. 1.7 months; $P = .017$). Patients without aggressive variant tumors, on the other hand, had similar median PFS regardless of treatment (6.5 vs. 6.3 months; $P = .38$).

Sipuleucel-T

In April 2010, sipuleucel-T became the first in a new class of cancer immunotherapeutic agents to be approved by the FDA. This autologous cancer “vaccine” involves collection of the white blood cell fraction-containing, antigen-presenting cells from each patient; exposure of the cells to the prostatic acid phosphatase-granulocyte macrophage colony-stimulating factor (PAP-GM-CSF recombinant fusion protein); and subsequent reinfusion of the cells. The pivotal study was a phase 3, multicenter, randomized, double-blind trial (D9902B).⁶²³ Five hundred twelve patients with minimally symptomatic or asymptomatic mCRPC were randomized 2:1 to receive sipuleucel-T or placebo. Eighteen-point two percent of patients had received prior chemotherapy, which included docetaxel; eligibility requirements included no chemotherapy for 3 months and no steroids for 1 month prior to enrollment. Median survival in the vaccine arm was 25.8 months compared to 21.7 months in the control arm. In a subset analysis, both those who did and those who did not receive prior chemotherapy benefited from sipuleucel-T treatment. Sipuleucel-T treatment resulted in a 22% reduction in mortality risk (HR, 0.78; 95% CI, 0.61–0.98; $P = .03$). Common complications included mild

to moderate chills (54.1%), pyrexia (29.3%), and headache (16.0%), which usually were transient.

A prospective registry of patients with mCRPC, PROCEED, enrolled 1976 patients from 2011 to 2017, who were followed for a median of 46.6 months.⁶²⁴ The safety and tolerability of sipuleucel-T were consistent with previous findings, and the median OS was 30.7 months (95% CI, 28.6–32.2 months).

Sipuleucel-T is included in these Guidelines as an option for asymptomatic patients with mCRPC regardless of prior ARPI or docetaxel, although the Panel notes that the data supporting its use post-docetaxel is limited. Benefit of sipuleucel-T has not been reported in patients with visceral metastases and is not recommended if visceral metastases are present. Sipuleucel-T is also not recommended for patients with small cell prostate cancer/NEPC. The Panel prefers that sipuleucel-T be used as a therapy for asymptomatic or minimally symptomatic patients with mCRPC, so that disease burden is lower and immune function is potentially more intact. Patients should have good performance level (ECOG 0–1), estimated life expectancy >6 months, and no liver metastases. Clinicians and patients should be aware that the usual markers of benefit (decline in PSA and improvement in bone or CT scans) are not seen. Therefore, benefit to the individual patient cannot be ascertained using currently available testing.

Pembrolizumab

The FDA approved the tumor-agnostic use of pembrolizumab, an anti-PD-1 antibody, for treatment of patients with unresectable or metastatic MSI-H or dMMR solid tumors who have progressed on prior treatment and who have no satisfactory alternative treatment options in May 2017. This approval was based on the treatment of 149 patients across five clinical studies involving MSI-H or dMMR colorectal ($n = 90$) or non-colorectal ($n = 59$) cancer for an objective response rate of 40% (59/149).⁵⁹² All patients



NCCN Guidelines Version 5.2026

Prostate Cancer

received ≥ 1 prior regimen. Among the non-colorectal cohorts, two patients had mCRPC: one achieved a partial objective response, and the other achieved stable disease for >9 months.

Outcomes of additional patients with mCRPC treated with pembrolizumab have been reported.^{82,625-629} In an early study, 10 patients with CRPC and non-visceral metastases (bone = 7; lymph nodes = 2; bone and liver = 1) who had disease progression on enzalutamide were treated with pembrolizumab and enzalutamide.⁶²⁵ Some of the patients also had experienced disease progression on additional therapies (docetaxel for CSPC, abiraterone, and/or sipuleucel-T). Three of the 10 patients showed a near complete PSA response. Two of these three patients had radiographically measurable disease and achieved a partial radiographic response (including a response in liver metastases). Of the remaining patients, three showed stable disease, and four displayed no evidence of clinical benefit. Genetic analysis of biopsy tissue revealed that one patient whose disease showed PSA response had an MSI-H tumor, whereas the other patient with responsive disease and two with non-responsive disease did not. The nonrandomized phase Ib KEYNOTE-028 trial included 23 patients with advanced, progressive prostate cancer, of whom 74% had received ≥ 2 previous therapies for metastatic disease.⁶²⁷ The objective response rate by investigator review was 17.4% (95% CI, 5.0%–38.8%), with four confirmed partial responses. Eight patients (34.8%) had stable disease. Treatment-related adverse events occurred in 61% of patients after a median follow-up of 7.9 months; 17% of the cohort experienced grade 3/4 events (ie, grade 4 lipase increase, grade 3 peripheral neuropathy, grade 3 asthenia, grade 3 fatigue).

KEYNOTE-199 was a multi-cohort, open-label phase II study in 258 patients with mCRPC and prior treatment with docetaxel and at least one ARPI that assessed pembrolizumab in patients regardless of MSI status.⁶³⁰ Cohorts 1 and 2 included patients with programmed cell death

ligand 1 (PD-L1)–positive (n = 133) and PD-L1–negative (n = 66) prostate cancer, respectively. Cohort 3 included those with bone-predominant disease with positive or negative PD-L1 expression (n = 59). The primary endpoint of overall response rate (ORR) was 5% (95% CI, 2%–11%) in cohort 1 and 3% (95% CI, $<1\%$ to 11%) in cohort 2. Responses were durable (range, 1.9 to ≥ 21.8 months).

The most common adverse events from pembrolizumab are fatigue, pruritus, diarrhea, anorexia, constipation, nausea, rash, fever, cough, dyspnea, and musculoskeletal pain. Pembrolizumab also may be associated with immune-mediated side effects, which include colitis, hepatitis, endocrinopathies, pneumonitis, or nephritis.

Based on the available data, the Panel includes pembrolizumab as an option for patients with MSI-H or dMMR mCRPC (category 2B). The prevalence of MMR deficiency in metastatic CRPC is estimated at 2% to 5%,^{44,626,631} and testing for MSI-H or dMMR can be performed using DNA testing or immunohistochemistry. If tumor MSI-H or dMMR is identified, the Panel recommends referral to genetic counseling for consideration of germline testing for Lynch syndrome.

In June 2020, the FDA granted accelerated approval for pembrolizumab's tumor-agnostic use in patients with unresectable or metastatic TMB-high (TMB-H) [≥ 10 mutations/megabase (mut/Mb)] solid tumors that have progressed following prior treatment and who have no satisfactory alternative treatment options. Results from prospective biomarker analysis of the multicohort, non-randomized, open-label, phase 2 KEYNOTE-158 trial support this approval, but this trial did not include any patients with prostate cancer.⁶³² One retrospective study found that 1.5% of patients with prostate cancer had TMB-H tumors.⁶³¹ Of those patients, 8 had received an immune checkpoint inhibitor; 4 patients (50%) experienced a reduction in PSA of $\geq 50\%$ that lasted at least 1 week. The Panel therefore



NCCN Guidelines Version 5.2026

Prostate Cancer

notes that pembrolizumab may be associated with some benefit in patients with mCRPC and TMB ≥ 10 mut/Mb.

Mitoxantrone

Two randomized trials assessed the role of mitoxantrone in patients with mCRPC.^{633,634} Although there was no improvement in OS, palliative responses and improvements in QOL were seen with mitoxantrone.

Mitoxantrone can be used for palliation in symptomatic patients with mCRPC who cannot tolerate other therapies after disease progression on prior docetaxel and an ARPI.

Treatment Options for Patients with DNA Repair Gene Mutations

Early studies suggest germline and somatic mutations in HRR genes (eg, *BRCA1*, *BRCA2*, *ATM*, *PALB2*, *FANCA*, *RAD51D*, *CHEK2*) may be predictive of the clinical benefit of poly-ADP ribose polymerase (PARP) inhibitors.⁶³⁵⁻⁶³⁷ PARP inhibitors are oral agents that exert their activity through the concept of synthetic lethality.⁶³⁸ PARP inhibitor therapy options are discussed below.

DNA repair defects have also been reported to be predictive for sensitivity to platinum agents in CRPC and other cancers.⁶³⁹⁻⁶⁴³ Platinum agents have shown some activity in patients with CRPC without molecular selection.⁶⁴⁴ Studies of platinum agents in patients with CRPC that have DNA repair gene mutations are needed.

Results of one study suggested that patients with mCRPC and germline mutations in DNA repair genes may have better outcomes if treated with abiraterone or enzalutamide than with taxanes.⁵² However, it should be noted that the response in patients with mCRPC and HRR gene mutations to standard therapies is similar to the response in patients without mutations.^{645,646}

Patients with *CDK12* mutations tend to have aggressive disease, with high rates of metastases and short OS. Their disease also does not respond well to hormonal therapy, PARP inhibitors, or taxanes. Two large, multi-institutional, retrospective studies have shown that 11% to 33% of patients with mCRPC and *CDK12* mutations experienced disease response to PD-1 inhibitors (ie, nivolumab, pembrolizumab), some with durable responses.^{647,648} There are also limited data from phase 2 trials indicating that ipilimumab plus nivolumab may have some activity against *CDK12*-mutated mCRPC.^{649,650} The Panel awaits more data on the use of PD-1 inhibition in patients with *CDK12* mutations.

Olaparib

Preliminary clinical data using olaparib suggested favorable activity of this agent in patients with HRR gene mutations, but not in those without HRR mutations.^{636,637,651} The phase 3 PROfound study was a randomized trial evaluating olaparib 300 mg twice daily versus physician's choice of abiraterone or enzalutamide in patients with mCRPC and progression on at least one novel hormonal agent (abiraterone or enzalutamide) and up to one prior taxane agent (permitted but not required).⁶⁵² Patients were required to have a somatic or germline HRR gene mutation, and were allocated to one of two cohorts: cohort A comprised patients with *BRCA1/2* or *ATM* mutations, and cohort B comprised patients with a mutation in at least one of 12 other HRR genes (*BARD1*, *BRIP1*, *CDK12*, *CHEK1*, *CHEK2*, *FANCL*, *PALB2*, *PPP2R2A*, *RAD51B*, *RAD51C*, *RAD51D*, or *RAD54L*). The primary endpoint of improving radiographic PFS with olaparib versus abiraterone/enzalutamide was met in cohort A (HR, 0.34; 95% CI, 0.25–0.47; $P < .001$), and radiographic PFS was also superior in the entire study population encompassing cohorts A+B (HR, 0.49; 95% CI, 0.38–0.63; $P < .001$).

In addition, final analysis of PROfound showed that OS was improved with olaparib versus abiraterone/enzalutamide in cohort A (HR, 0.69; 95% CI,



NCCN Guidelines Version 5.2026

Prostate Cancer

0.50–0.97; $P = .02$), despite the fact that 86 of 131 patients (66%) crossed over to olaparib after disease progression in the control arm.⁶⁵³

The Panel notes that there may be heterogeneity of response to olaparib based on which gene has a mutation. Efficacy in PROfound appears to be driven by the cohort of patients with at least one alteration in *BRCA2*, *BRCA1*, or *ATM*, and in particular by patients with *BRCA2* or *BRCA1* mutations based on exploratory gene-by-gene analysis.⁶⁵³ Patients with *BRCA2* mutations in PROfound experienced an OS benefit with olaparib (HR, 0.59; 95% CI, 0.37–0.95), whereas the HR for OS in patients with *ATM* mutations was 0.93 (95% CI, 0.53–1.75).⁶⁵³ Furthermore, there were few patients in PROfound with mutations in some of the genes. For example, only 4 patients had *BRIP1* mutations (2 in olaparib arm and 2 in control arm), 2 patients had *RAD51D* mutations (both in olaparib arm), and no patients had *RAD51C* mutations.⁶⁵² Patients with *PPP2R2A* mutations in PROfound experienced an unfavorable risk-benefit profile.

As a result of the favorable efficacy data from the PROfound trial, the FDA approved olaparib (300 mg twice daily) in May 2020 for use in patients with mCRPC and deleterious or suspected deleterious germline or somatic HRR gene mutations in at least one of 14 genes (*BRCA1*, *BRCA2*, *ATM*, *BARD1*, *BRIP1*, *CDK12*, *CHEK1*, *CHEK2*, *FANCL*, *PALB2*, *RAD51B*, *RAD51C*, *RAD51D*, or *RAD54L*) and who had previously received treatment with enzalutamide or abiraterone.

Adverse events that may occur with olaparib treatment include anemia (including that requiring transfusion), fatigue, nausea or vomiting, anorexia, weight loss, diarrhea, thrombocytopenia, creatinine elevation, cough, and dyspnea. Rare but serious side effects may include thromboembolic events (including pulmonary emboli), drug-induced pneumonitis, and a theoretical risk of myelodysplasia or acute myeloid leukemia.⁶⁵²

Since some patients in PROfound had prior taxane therapy, olaparib use might be reasonable in patients with mCRPC before or after docetaxel treatment. The Panel therefore recommends olaparib as an option for patients with mCRPC, previous ARPI, and an HRRm regardless of prior docetaxel therapy. The HRR genes to be considered for use of olaparib are *BRCA1*, *BRCA2*, *ATM*, *BARD1*, *BRIP1*, *CDK12*, *CHEK1*, *CHEK2*, *FANCL*, *PALB2*, *RAD51B*, *RAD51C*, *RAD51D* and *RAD54L*. Olaparib is included in the Guidelines as a category 1 recommendation for BRCAm mCRPC and is a preferred option for these patients if they have not yet received docetaxel.

Any commercially available analytically and clinically validated somatic tumor and germline assays can be used to identify patients for treatment. The Panel strongly recommends a metastatic biopsy for histologic and molecular evaluation; a plasma circulating tumor DNA (ctDNA) assay can be used if a metastatic biopsy is unsafe or not feasible.

Careful monitoring of complete blood counts and hepatic and renal function, along with type and screens and potential transfusion support and/or dose reductions as needed for severe anemia or intolerance are recommended during olaparib therapy.

Rucaparib

Rucaparib is another PARP inhibitor approved for use in patients with mCRPC. This agent received accelerated FDA approval in May 2020 based on the preliminary favorable data from the TRITON2 clinical trial. In that open-label, single-arm, phase 2 trial, patients with mCRPC harboring a deleterious or suspected deleterious germline or somatic *BRCA1* or *BRCA2* mutation, who had previously received therapy with an ARPI plus one taxane chemotherapy, were treated with rucaparib 600 mg twice daily.⁶⁵⁴ The primary endpoint of TRITON2 was the objective response rate in patients with measurable disease, and was 43.5% (95% CI, 31.0%–56.7%) in this *BRCA1/2*-mutated population. Median radiographic



NCCN Guidelines Version 5.2026

Prostate Cancer

PFS, a key secondary endpoint, was 9.0 months (95% CI, 8.3–13.5 months). The most common adverse events were asthenia/fatigue, nausea, and anemia/decreased hemoglobin, with grade ≥ 3 anemia/decreased hemoglobin in 25.2% of participants. Final analysis of TRITON2 confirmed results of the earlier analysis.⁶⁵⁵

In the randomized phase 3 TRITON3 study, patients with mCRPC and a germline or somatic *BRCA1/2* or *ATM* mutation who have previously received an ARPI but no chemotherapy for mCRPC were randomized 2:1 to rucaparib versus physician's choice of therapy (abiraterone, enzalutamide, or docetaxel).⁶⁵⁶ The primary endpoint of TRITON3, the median duration of imaging-based PFS, was significantly longer at 62 months in the group of 270 participants assigned to receive rucaparib than in the 135 participants who received a control medication (10.2 vs. 6.4 months; HR, 0.61; 95% CI, 0.47–0.80; $P < .001$). This effect was also seen in the 201 patients and 101 patients in each group with a *BRCa*m (11.2 vs. 6.4 months; HR, 0.50; 95% CI, 0.36–0.69). For those with *ATM* mutations, an exploratory analysis suggested a possible improvement as well (8.1 vs. 6.8 months; HR, 0.95; 95% CI, 0.59–1.52). As in TRITON2, the most frequent adverse events with rucaparib were fatigue and nausea.

The Panel recommends rucaparib as an option for patients with mCRPC, a *BRCA1* or *BRCA2* mutation, and prior treatment with an ARPI. It is a category 1, preferred option pre-docetaxel. Rucaparib should not be used in patients with HRR gene mutations other than *BRCA1/2*.⁶⁵⁷ Adverse events that may occur with rucaparib include anemia (including that requiring transfusion), fatigue, asthenia, nausea or vomiting, anorexia, weight loss, diarrhea or constipation, thrombocytopenia, increased creatinine, increased liver transaminases, and rash. Rare but serious side effects of rucaparib include a theoretical risk of myelodysplasia or acute myeloid leukemia, as well as fetal teratogenicity.^{654,657}

The preferred method of selecting patients for rucaparib treatment is somatic and germline analysis of *BRCA1* and *BRCA2* from a metastatic biopsy. A ctDNA sample can be used if biopsy is unsafe or not feasible.

As with olaparib, careful monitoring of complete blood counts and hepatic and renal function, along with type and screens and potential transfusion support and/or dose reductions as needed for severe anemia or intolerance are recommended during treatment with rucaparib.

Olaparib Plus Abiraterone

Pre-clinical data suggest that PARP-1 promotes androgen receptor activity.⁶⁵⁸ Additional pre-clinical data show that androgen receptor inhibitors can down-regulate DNA repair genes, creating a situation similar to that of HRR mutation.^{659,660} These results suggest that the combination of PARP inhibition with androgen receptor inhibition may have an enhanced antitumor effect and that this effect may not be limited to patients with HRR mutations. In fact, a randomized phase 2 trial showed that the combination of abiraterone with olaparib increased radiographic PFS over abiraterone and placebo in patients with mCRPC regardless of HRR status (ITT population: HR, 0.65; 95% CI, 0.44–0.97; $P = .034$).⁶³⁷

The PROpel trial was an international, double-blind, phase 3 trial comparing abiraterone and olaparib with abiraterone and placebo in 796 patients with mCRPC regardless of HRR mutation status.⁶⁶¹ Prior docetaxel in the localized or mCSPC setting was allowed, but patients were untreated for CRPC. The primary endpoint, imaging-based PFS by investigator assessment in the ITT population, was significantly longer in the abiraterone/olaparib group than in the abiraterone/placebo group (24.8 vs. 16.6 months; HR, 0.66; 95% CI, 0.54–0.81; $P < .001$). HRR mutations were identified in tumors of 226 patients; 552 patients did not have HRR tumor mutations. The HR for the primary endpoint in those with HRR mutations was 0.50 (95% CI, 0.34–0.73). The safety profile of the olaparib/abiraterone combination was as expected based on the known



NCCN Guidelines Version 5.2026

Prostate Cancer

safety profiles of the individual drugs, with the most common adverse events being anemia, fatigue/asthenia, and nausea.

Final OS data from PROpel showed that OS was not significantly improved with the abiraterone/olaparib combination therapy in the full cohort after a median follow up of approximately 36.5 months (42.1 vs 34.7 months; HR, 0.81; 95% CI, 0.67–1.00; $P = .054$).⁶⁶² In a post-hoc exploratory analysis, the BRCAm population saw an OS benefit, with a median OS of 23.0 months in the abiraterone arm and not reached in the combination arm (HR, 0.29; 95% CI, 0.14–0.56). A smaller OS benefit was seen in the HRRm group overall, and no OS benefit was evident in the non-HRRm and the non-BRCAm/other HRRm subgroups.

In May 2023, the FDA approved the combination of olaparib with abiraterone for the treatment of adult patients with BRCAm mCRPC. Based on the results of PROpel, olaparib/abiraterone is included in the NCCN Guidelines as an option for patients with mCRPC and a pathogenic *BRCA1* or *BRCA2* mutation (germline and/or somatic) who have not yet received an ARPI (category 1).

Talazoparib Plus Enzalutamide

Talazoparib is another PARP inhibitor; it has had an FDA indication in breast cancer. The open-label, international phase 2 TALAPRO-1 trial included 127 patients with an HRR mutation and progressive, mCRPC, all of whom received at least one dose of talazoparib.⁶⁶³ The objective response rate after a median follow-up of 16.4 months was 29.8% (95% CI, 21.2–39.6). The most common grade 3–4 treatment-emergent adverse events were anemia (31%), thrombocytopenia (9%), and neutropenia (8%).

As noted above (see *Olaparib Plus Abiraterone*), pre-clinical data suggest that the PARP inhibition combined with androgen receptor inhibition may have an enhanced antitumor effect that may not be limited to those with

HRR mutations. The randomized, double-blind, phase 3 TALAPRO-2 study compared enzalutamide plus talazoparib with enzalutamide plus placebo in 805 patients with untreated mCRPC.⁶⁶⁴ HRR gene alteration status and treatment with docetaxel and/or abiraterone in the castration-sensitive setting were used to stratify the randomization. The primary endpoint was radiographic PFS in the ITT population. At the planned primary analysis, median radiographic PFS was not reached (95% CI, 27.5 months–not reached) for the talazoparib group and was 21.9 months (95% CI, 16.6–25.1) for the control group (HR, 0.63; 95% CI, 0.51–0.78; $P < .0001$).

HRR mutations were present in 21% of TALAPRO-2 participants, with *BRCA* alterations being the most common.⁶⁶⁴ The HR for radiographic PFS in the HRR-deficient subgroup was more strongly in favor of the talazoparib combination than in the HRR-proficient/unknown population (0.46 [95% CI, 0.30–0.70; $P = .0003$] vs. 0.70 [95% CI, 0.54–0.89; $P = .0039$]). Among HRR mutations, talazoparib conferred a 77% lower risk of radiographic progression or death in those with tumor mutations in *BRCA1* or *BRCA2* (HR, 0.23; 95% CI, 0.10–0.53; $P = .0002$), whereas the corresponding reduction was 34% (HR, 0.66; 95% CI, 0.39–1.12; $P = .12$) in those with non-*BRCA* HRR alterations.

Prior therapy also affected the radiographic PFS outcomes in this trial.⁶⁶⁴ In the 179 participants in TALAPRO-2 who had received docetaxel in earlier disease settings, the HR for radiographic PFS was 0.51 (95% CI, 0.32–0.81; $P = .0034$). In the small population of 50 participants in the ITT population who had received prior novel hormonal therapy, the corresponding HR was non-significant at 0.57 (95% CI, 0.28–1.16; $P = .12$).

Final results from an HRRm-only cohort of TALAPRO-2 at a median follow-up of 44.2 months showed that median OS was improved with the combination compared with enzalutamide alone (45.1 vs. 31.1 months;



NCCN Guidelines Version 5.2026

Prostate Cancer

HR, 0.62; 95% CI, 0.48–0.81; two-sided $P = .0005$).⁶⁶⁵ Median radiographic PFS was also improved with the talazoparib/enzalutamide group compared with the enzalutamide group (30.7 vs. 12.3 months; HR, 0.47; 95% CI, 0.36–0.61; $P < .0001$).

The safety profile of enzalutamide plus talazoparib was consistent with the known safety profiles of the individual drugs, with the most common adverse events in those who received talazoparib being anemia, neutropenia, and fatigue. However, hematologic adverse events were of higher grades and occurred more frequently than would be expected with talazoparib alone. Overall, the combination had significant toxicity, with dose interruption due to adverse events in 75% of participants in the talazoparib group compared with 23% in the placebo group. Dose reductions due to adverse events occurred in 56% and 7% of the talazoparib and placebo groups, respectively.

Based on the results from the first TALAPRO-2 cohort, the FDA approved talazoparib plus enzalutamide for HRRm mCRPC in June 2023. The Panel includes talazoparib plus enzalutamide as a category 1 treatment option for patients with mCRPC and a pathogenic mutation (germline and/or somatic) in one of certain HRR and other DNA repair genes (*BRCA1*, *BRCA2*, *ATM*, *ATR*, *CDK12*, *CHEK2*, *FANCA*, *MLH1*, *MRE11A*, *NBN*, *PALB2*, or *RAD51C*) who have not yet had treatment with an ARPI. Use of talazoparib/enzalutamide for those who have received prior ARPI therapy without prior docetaxel is controversial (category 2B) because a benefit of this combination over use of a PARP inhibitor alone has not been shown in this setting, but responses are likely.

Niraparib Plus Abiraterone

Another PARP inhibitor, niraparib, has also been studied in combination with androgen inhibition in the setting of mCRPC. The randomized, double-blinded phase 3 MAGNITUDE trial compared niraparib plus abiraterone to placebo plus abiraterone in 423 patients with mCRPC and

HRR mutations and an additional 247 patients without HRR mutations.⁶⁶⁶ Prior chemotherapy and novel hormonal therapy were allowed in the mCSPC or M0 CRPC settings, and were received by 3.1% and 20.1% of the total HRRm cohort, respectively.

The primary endpoint of MAGNITUDE was radiographic PFS. After a median follow-up of 18.6 months, radiographic PFS was improved for those receiving niraparib in the HRRm group overall (16.5 vs. 13.7 months; HR, 0.73; 95% CI, 0.56–0.96; $P = .022$) as well as in the BRCAm subgroup (16.6 vs. 10.9 months; HR, 0.53; 95% CI, 0.36–0.79; $P = .001$). However, radiographic PFS was not improved in the subgroup of patients with non-*BRCA* HRR mutations (HR, 0.99; 95% CI, 0.68–1.44). For the cohort without HRR mutations, fertility was declared based on prespecified criteria. The secondary endpoints of time to symptomatic progression and time to initiation of cytotoxic chemotherapy were improved with the combination therapy in the HRRm and BRCAm cohorts.

A second interim analysis of MAGNITUDE included a prespecified, inverse probability censoring weighting analysis of OS, which was designed to account for the receipt of subsequent therapies, including PARP inhibitors.⁶⁶⁷ Results of this analysis suggest that there may be an OS benefit for the combination therapy (HR, 0.54; 95% CI, 0.33–0.90; nominal $P = .0181$).

The incidence of grade 3/4 adverse events was higher with the combination of niraparib plus abiraterone compared with placebo and abiraterone (67.0% vs. 46.4%).⁶⁶⁶ Anemia (28.3% vs. 7.6%) and hypertension (14.6% vs. 12.3%) were the most reported grade ≥3 adverse events. Overall, the combination was tolerable and QOL was maintained.

Based on these results, the FDA approved niraparib plus abiraterone for the treatment of patients with BRCAm mCRPC in August 2023. The Panel includes niraparib plus abiraterone as a treatment option for patients with



NCCN Guidelines Version 5.2026

Prostate Cancer

mCRPC and a pathogenic *BRCA1* or *BRCA2* mutation (germline and/or somatic) who have not yet had treatment in the setting of mCRPC. This is a category 1 recommendation for those without prior ARPI. Use of niraparib/abiraterone for those who have received a prior ARPI without prior docetaxel is controversial (category 2B) because a benefit of this combination over use of a PARP inhibitor alone has not been shown in this setting.

Radiopharmaceuticals for mCRPC

Lutetium Lu 177 vipivotide tetraxetan

Lu-177-PSMA-617 is a radiopharmaceutical that is administered intravenously and is indicated for PSMA-positive mCRPC that has been treated with androgen receptor pathway inhibition and taxane-based chemotherapy. The active moiety is a radionuclide that delivers radiation to PSMA-expressing and surrounding cells, which induces DNA damage and leads to cell death. The approval of Lu-177-PSMA-617 was based on the international, open-label phase III VISION trial of 831 patients with mCRPC and PSMA-positive metastatic lesions. Patients in VISION were previously treated with at least one androgen receptor-directed therapy and one or two taxane-based chemotherapy regimens.⁶⁶⁸ Patients had at least one PSMA-positive metastatic lesion and no PSMA-negative lesions determined by Ga-68 labeled PSMA-11 PET/CT imaging. Patients were randomized in a 2:1 ratio to receive standard of care (abiraterone, enzalutamide, bisphosphonates, RT, denosumab, and/or glucocorticoids) and Lu-177-PSMA-617 (7.4 GBq or 200 mCi every 6 weeks for 4–6 cycles) or standard of care alone.

The median OS was improved in the Lu-177-PSMA-617 group compared to the control group (15.3 vs. 11.3 months; HR, 0.62; 95% CI, 0.52–0.74; $P < .001$). Similarly, the median PFS was improved in the Lu-177-PSMA-617 group compared to the control group (8.7 vs. 3.4 months; HR, 0.40; 99.2% CI, 0.29–0.57; $P < .001$). The incidence of grade ≥ 3 adverse events

(particularly anemia, thrombocytopenia, lymphopenia, and fatigue) was significantly higher in the Lu-177-PSMA-617 group compared to the control group.⁶⁶⁸

The FDA approved Lu-177-PSMA-617 in the post-ARPI setting in March 2022.

Another randomized controlled phase 3 trial, PSMAfore, assessed the efficacy of Lu-177-PSMA-617 pre-docetaxel in 468 patients with mCRPC who experienced disease progression on an ARPI.⁶⁶⁹ The primary endpoint of rPFS in the ITT population was improved with Lu-177-PSMA-617 compared with a change in ARPI. At a median of 24.1 months after randomization, median radiographic PFS was 11.6 months in the Lu-177-PSMA-617 group versus 5.6 months in the control group (HR 0.49; 95% CI, 0.39–0.61). There was no difference in OS, but OS data are difficult to interpret because patients were allowed to crossover from an ARPI change to Lu-177-PSMA-617 upon radiographic progression, and 57% of patients in the control arm did so. Importantly, there were fewer grade 3–5 toxicities in Lu-177-PSMA-617 arm.

Based on the results of the PSMAfore trial, the FDA expanded the indication for Lu-177-PSMA-617 to include adult patients with PSMA-positive mCRPC who have progressed after ARPI therapy and are considered appropriate candidates for delaying taxane-based chemotherapy.

The NCCN Panel recommends Lu-177-PSMA-617 as a treatment option for patients with one or more PSMA-positive lesion and/or metastatic disease that is predominately PSMA-positive and with no dominant PSMA-negative metastatic lesions who have been treated previously with an ARPI and a taxane-based chemotherapy (category 1) or who have prior ARPI therapy and are considered appropriate to delay taxane-based chemotherapy.



NCCN Guidelines Version 5.2026

Prostate Cancer

PSMA-negative lesions are defined as metastatic disease that lacks PSMA uptake including bone with soft tissue components ≥ 1.0 cm, lymph nodes ≥ 2.5 cm in short axis, and solid organ metastases ≥ 1.0 cm in size. The Panel notes that Ga-68 PSMA-11, F-18 piflufolastat PSMA, or F-18 flutufolastat PSMA can be used to identify patients eligible for treatment with Lu-177–PSMA-617.

Radium-223

In May 2013, the FDA approved radium-223 dichloride, an alpha particle-emitting radioactive agent. This first-in-class radiopharmaceutical was approved for treatment of mCRPC in patients with symptomatic bone metastases and no known visceral metastatic disease. Approval was based on clinical data from a multicenter, phase 3, randomized trial (ALSYMPCA) that included 921 patients with symptomatic CRPC, two or more bone metastases, and no known visceral disease.⁶⁷⁰ Fifty-seven percent of the patients received prior docetaxel and all patients received best supportive care. Patients were randomized in a 2:1 ratio to 6 monthly radium-223 intravenous injections or placebo. Compared to placebo, radium-223 significantly improved OS (median 14.9 vs. 11.3 months; HR, 0.70; 95% CI, 0.058–0.83; $P < .001$) and prolonged time to first SRE (median 15.6 vs. 9.8 months). Preplanned subset analyses showed that the survival benefit of radium-223 was maintained regardless of prior docetaxel use.⁶⁷¹ ITT analyses from ALSYMPCA showed that radium-223 also may reduce the risk of symptomatic SREs.⁶⁷² Grade 3/4 hematologic toxicity was low (3% neutropenia, 6% thrombocytopenia, and 13% anemia), likely due to the short range of radioactivity.⁶⁷⁰ Fecal elimination of the agent led to generally mild non-hematologic side effects, which included nausea, diarrhea, and vomiting. Radium-223 was associated with improved or slower decline of QOL in ALSYMPCA.⁶⁷³

The multicenter, international, double-blind, placebo-controlled, phase 3 ERA 223 trial randomized patients with bone-metastatic chemotherapy-

naïve CRPC to abiraterone with or without radium-223.⁶⁷⁴ The patients were asymptomatic or mildly symptomatic. The primary endpoint of symptomatic skeletal event-free survival in the ITT population was not met. In fact, the addition of radium-223 to abiraterone was associated with an increased frequency of bone fractures compared with placebo.

The randomized PEACE-3 trial also compared radium-223 with or without an ARPI in patients who were ARPI-naïve.⁶⁷⁵ Radium-223 with enzalutamide was compared with enzalutamide therapy alone in 446 patients with mildly symptomatic mCRPC. The use of bone-protecting agents (denosumab or zoledronic acid) was made mandatory following results from ERA 223. The primary endpoint of radiological PFS was improved in the combination arm compared with enzalutamide alone (16.4 vs. 19.4 months; HR, 0.69; 95% CI, 0.54–0.87; $P = .0009$). At a preplanned interim OS analysis, median OS was also improved with the addition of radium-223 (35.0 vs. 42.3 months; HR, 0.69; 95% CI 0.52–0.90; $P = .0031$). Grade ≥ 3 adverse events occurred more commonly in the combination arm (65.6% vs. 55.8%) and included hypertension (in 34% of the combination arm), fatigue (6%), fracture (5%), anemia (5%), and neutropenia (5%). Fractures were also more common in the combination arm (24.3% vs. 13.4%).

In an earlier safety analysis of PEACE-3, the cumulative incidence of fractures at 1.5 years in patients who received a bone-protecting agent was 2.8% in participants receiving radium-223 plus enzalutamide and 3.9% in those receiving enzalutamide alone.⁶⁷⁶ In the absence of bone agents, these numbers were 45.9% and 22.3%, respectively. This result suggests that radium-223 combined with an ARPI may be safe if preventive administration of a bone agent is used.

Radium-223 is a category 1 option to treat symptomatic bone metastases without visceral metastases in patients with mCRPC regardless of prior therapy. Radium-223 plus enzalutamide is included as an option for



NCCN Guidelines Version 5.2026

Prostate Cancer

patients with bone-metastatic CRPC without prior exposure to an ARPI. Hematologic evaluation should be performed according to the FDA label before treatment initiation and before each subsequent dose.⁵⁹² Radium-223 given in combination with chemotherapy (such as docetaxel) outside of a clinical trial has the potential for additive myelosuppression.⁵⁹² Its use in combination with docetaxel or any other systemic therapy except ADT or enzalutamide should be pursued with caution. It should not be used in patients with visceral metastases. All patients receiving radium-223 should be given concomitant denosumab or zoledronic acid.

MDT for mCRPC

Although most data supporting the use of MDT for oligometastatic prostate cancer is in the oligorecurrent CSPC setting, as discussed in detail above (see *MDT for Oligometastatic CSPC*), MDT has also been studied in the oligoprogressive CRPC. As noted above, the EXTEND trial included 7 patients with oligoprogressive CRPC, although results of MDT in this subset specifically were not reported.⁵⁶⁸

ARTO is a phase 2 study that included 157 patients with CRPC and 1 to 3 metastatic lesions who were randomized to receive abiraterone alone or abiraterone with MDRT.⁵⁶⁶ The rate of biochemical response (defined as a $\geq 50\%$ decrease in PSA levels at 6 months compared to baseline), which was the primary endpoint, was 92% in the MDRT group compared with 68.3% in the control group (OR, 4.22; 95% CI, 2.12–8.38; $P < .001$). PFS, a secondary endpoint was also improved with the use of MDRT (HR, 0.35; 95% CI, 0.21–0.57; $P < .001$). A subgroup analysis of ARTO further suggested that MDRT for oligoprogressive mCRPC may result in similar PFS as second-line systemic therapy.⁶⁷⁷

Initial results from the phase 2 GROUQ-PCS-9 were presented at the 2025 ASCO Genitourinary Cancers Symposium.⁶⁷⁸ MDRT added to ADT and enzalutamide in oligometastatic CRPC (with 1–5 metastases) led to

improvements in radiological PFS, biochemical PFS, and time to next line of therapy compared to ADT with enzalutamide alone.

The Panel includes MDT with mCRPC systemic therapy as an option for patients with oligometastatic or oligoprogressive CRPC regardless of prior therapy.

Small Cell/Neuroendocrine Prostate Cancer

De novo small cell carcinoma in untreated prostate cancer occurs rarely and is very aggressive.⁶⁷⁹ Treatment-associated small cell prostate cancer/NEPC that occurs in patients with mCRPC is more common.⁶⁸⁰ In a multi-institution prospective series of 202 consecutive patients with mCRPC, all of whom underwent metastatic biopsies, small cell/neuroendocrine histology was present in 17% of patients.⁶⁸⁰ Patients with small cell/neuroendocrine tumors and prior abiraterone and/or enzalutamide had a shorter OS when compared with those with adenocarcinoma and prior abiraterone and/or enzalutamide (HR, 2.02; 95% CI, 1.07–3.82). Genomic analysis showed that DNA repair mutations and small cell/neuroendocrine histology were almost mutually exclusive.

Small cell/neuroendocrine carcinoma of the prostate should be considered in patients with disease that no longer responds to ADT and who test positive for metastases. These relatively rare tumors are associated with low PSA levels despite large metastatic burden and visceral disease.⁶⁸¹ Those with initial Grade Group 5 are especially at risk. Biopsy of accessible metastatic lesions to identify patients with small cell/neuroendocrine histomorphologic features is recommended in patients with mCRPC.

These patients may be treated with cytotoxic chemotherapy (ie, cisplatin/etoposide, carboplatin/etoposide, docetaxel/carboplatin, cabazitaxel/carboplatin).^{622,682,683} Physicians should consult the NCCN Guidelines for Small Cell Lung Cancer for additional options in the first and



NCCN Guidelines Version 5.2026

Prostate Cancer

subsequent lines of therapy (available at www.NCCN.org), because the behavior of small cell/neuroendocrine carcinoma of the prostate is similar to that of small cell carcinoma of the lung.

Additional Treatment Options for Bone Metastases

In a multicenter study, 643 patients with CRPC and asymptomatic or minimally symptomatic bone metastases were randomized to intravenous zoledronic acid every 3 weeks or placebo.⁶⁸⁴ At 15 months, fewer patients in the zoledronic acid 4-mg group than patients in the placebo group had SREs (33% vs. 44%; $P = .02$). An update at 24 months also revealed an increase in the median time to first SRE (488 vs. 321 days; $P = .01$).⁶⁸⁵ No significant differences were found in OS. Other bisphosphonates have not been shown to be effective for prevention of disease-related skeletal complications.

The randomized TRAPEZE trial used a 2 X 2 factorial design to compare clinical PFS (pain progression, SREs, or death) as the primary outcome in 757 patients with bone-metastatic CRPC treated with docetaxel alone or with zoledronic acid, 89Sr, or both.⁶⁸⁶ The bone-directed therapies had no statistically significant effect on the primary outcome or on OS in unadjusted analysis. However, adjusted analysis revealed a small effect for 89Sr on clinical PFS (HR, 0.85; 95% CI, 0.73–0.99; $P = .03$). For secondary outcomes, zoledronic acid improved the SRE-free interval (HR, 0.78; 95% CI, 0.65–0.95; $P = .01$) and decreased the total SREs (424 vs. 605) compared with docetaxel alone.

Denosumab was compared to zoledronic acid in a randomized, double-blind, placebo-controlled study in patients with CRPC.⁶⁸⁷ The absolute incidence of SREs was similar in the two groups; however, the median time to first SRE was delayed by 3.6 months by denosumab compared to zoledronic acid (20.7 vs. 17.1 months; $P = .0002$ for noninferiority; $P = .008$ for superiority). The rates of important SREs with denosumab were

similar to zoledronic acid and included spinal cord compression (3% vs. 4%), need for radiation (19% vs. 21%), and pathologic fracture (14% vs. 15%). Treatment-related toxicities reported for zoledronic acid and denosumab were similar and included hypocalcemia (more common with denosumab 13% vs. 6%), arthralgias, and osteonecrosis of the jaw (ONJ, 1%–2% incidence).

Therefore, denosumab every 4 weeks (category 1, preferred) or zoledronic acid every 3 to 4 weeks is recommended for patients with CRPC and bone metastases to prevent or delay disease-associated SREs. SREs include pathologic fractures, spinal cord compression, operation, or EBRT to bone. The optimal duration of zoledronic acid or denosumab in patients with CRPC and bone metastases remains unclear. A multi-institutional, open-label, randomized trial in 1822 patients with bone-metastatic prostate cancer, breast cancer, or multiple myeloma found that zoledronic acid every 12 weeks was noninferior to zoledronic acid every 4 weeks.⁶⁸⁸ In the every-12-week and every-4-week arms, 28.6% and 29.5% experienced at least 1 SRE within 2 years of randomization, respectively.

Use of zoledronic acid in patients with CSPC and bone metastases is not associated with lower risk for SREs.⁶⁸⁹ Therefore, the routine use of these agents in bone-metastatic CSPC is not recommended. Bone antiresorptive agents should, however, be used for SRE prevention in patients with CSPC if they have treatment-related bone loss (see *Principles of Survivorship, Bone Health in Prostate Cancer*, in the algorithm above).

Oral hygiene, baseline dental evaluation for high-risk individuals, and avoidance of invasive dental surgery during therapy are recommended to reduce the risk of ONJ.⁶⁹⁰ Most, but not all, patients who develop ONJ have preexisting dental problems.⁶⁹¹ If invasive dental surgery is necessary, therapy should be deferred until the dentist confirms that the patient has healed completely from the dental procedure. Supplemental



NCCN Guidelines Version 5.2026

Prostate Cancer

calcium and vitamin D are recommended to prevent hypocalcemia in patients receiving either denosumab or zoledronic acid.

Monitoring of creatinine clearance is required to guide dosing of zoledronic acid. Zoledronic acid should be dose reduced in patients with impaired renal function (estimated creatinine clearance 30–60 mL/min) and held for creatinine clearance <30 mL/min. Denosumab may be administered to patients with impaired renal function or even patients on hemodialysis; however, the risk for severe hypocalcemia and hypophosphatemia is greater, and the dose, schedule, and safety of denosumab have not yet been defined. A single study of 55 patients with creatinine clearance <30 mL/min or on hemodialysis evaluated the use of 60-mg-dose denosumab.⁵⁹² Hypocalcemia should be corrected before starting denosumab, and serum calcium monitoring is required for denosumab and recommended for zoledronic acid, with repletion as needed.

Radium-223 is a category 1 option to treat symptomatic bone metastases in patients with mCRPC without visceral metastases (see *Radium-223*, above). The use of palliative RT is also an option.

Clinical research on the prevention or delay of disease spread to bone continues. A phase 3 randomized trial of 1432 patients with non-metastatic CRPC at high risk of bone involvement showed that denosumab delayed bone metastasis by 4 months compared to placebo.⁶⁹² OS was not improved, and the FDA did not approve denosumab for the prevention of bone metastases.

Considerations for Visceral Metastases

The Panel defines visceral metastases as those occurring in the liver, lung, adrenal gland, peritoneum, or brain. Soft tissue/lymph node sites are not considered visceral metastases. In general, there are fewer data on treatment of patients with CRPC and visceral metastases than for those without visceral metastases.

Sequencing of Therapy in CRPC

The number of treatment options for patients with CRPC has expanded rapidly over the past several years. Although the optimal sequence of therapies remains undefined, some data that can help with treatment selection in some cases continues to emerge.

After abiraterone or enzalutamide, data suggest that giving the alternate ARPI may not be the optimal strategy considering the availability of other treatment options, including chemotherapy. The CARD trial, for instance, showed that treatment with cabazitaxel significantly improved clinical outcomes over enzalutamide or abiraterone in patients with mCRPC who had been previously treated with docetaxel and the alternate hormonal therapy (abiraterone or enzalutamide).⁶¹⁶ Furthermore, data suggest cross-resistance between abiraterone and enzalutamide.⁶⁹³⁻⁶⁹⁶ Results of a randomized, open-label, phase 2, crossover trial suggest that the sequence of abiraterone followed by enzalutamide may be more efficacious than the reverse.⁶⁹⁷

Some data inform the sequencing of therapies in patients with PSMA-positive mCRPC. The multicenter, unblinded, randomized phase 2 TheraP trial compared PSA response after Lu-177-PSMA-617 versus cabazitaxel in 200 patients with PSMA-positive mCRPC who previously received docetaxel.⁶⁹⁸ Prior androgen receptor-directed therapy was permitted. Among the ITT population, the PSA response rate was 66% in the Lu-177-PSMA-617 arm compared with 37% in the cabazitaxel arm (difference 29%; 95% CI, 16–42; $P < .0001$). These numbers were 66% and 44%, respectively, in those who received treatment (difference 23%; 95% CI, 9–37; $P = .0016$). Furthermore, grade 3–4 adverse events were less frequent in the Lu-177-PSMA-617 arm than in the cabazitaxel arm (33% vs. 53%). Results from the phase 3 PSMAfore trial as discussed above (see *Lutetium Lu 177 vipivotide tetraxetan*) showed that Lu-177-PSMA-617



NCCN Guidelines Version 5.2026

Prostate Cancer

improved rPFS compared with switching to a different ARPI in docetaxel-naïve patients.⁶⁶⁹

Data for patients with HRRm mCRPC are more limited, but comparative effectiveness research suggests that olaparib may result in superior radiographic PFS than cabazitaxel in patients with *BRCA1* or *BRCA2* mutations and prior treatment with docetaxel.⁶⁹⁹ Furthermore, data from PROfound and TRITON3 suggest that a PARP inhibitor is preferred over a different ARPI in patients with BRCAm mCRPC and prior ARPI exposure.^{652,656}

No chemotherapy regimen has demonstrated improved survival or QOL after cabazitaxel or cabazitaxel/carboplatin in patients with mCRPC of adenocarcinoma histology, although several systemic agents other than mitoxantrone have shown palliative and radiographic response benefits in clinical trials (ie, carboplatin, cyclophosphamide, doxorubicin, vinorelbine, carboplatin/etoposide, docetaxel/carboplatin, gemcitabine/oxaliplatin, paclitaxel/carboplatin).⁷⁰⁰⁻⁷⁰⁹ No survival benefit for any these combination regimens over sequential single-agent regimens has been demonstrated,

and toxicity is higher. Treatment with these regimens could be considered after an informed discussion between the physician and an individual patient about treatment goals and risks/side effects and alternatives, which must include best supportive care. In patients not able to receive life prolonging therapy, prednisone and dexamethasone at low doses may provide palliative benefits.⁷¹⁰ Participation in a clinical trial is encouraged.

Summary

The intention of these guidelines is to provide a framework on which to base treatment decisions. Prostate cancer is a complex disease, with many controversial aspects of management and with limited data to support some of the treatment recommendations. Several variables (including adjusted life expectancy, disease characteristics, predicted outcomes, and patient preferences) must be considered by the patient and physician to tailor prostate cancer therapy for the individual patient.



NCCN Guidelines Version 5.2026

Prostate Cancer

Table 1: Selected Active Surveillance Experiences with Large Patient Cohorts

Cohort	Toronto ^{272,313,319}	Johns Hopkins ^{275, 312,314,711}	UCSF ²⁸¹		Canary PASS ²⁸⁹	Cooley/Catalona Meta-Dataset ²⁷⁷	PRIAS ²⁷⁸
			Initial Cohort	Newer Cohort			
Number of patients	993	1298	321	810	2155	6775	5302
Median age (y)	68	66	63	62	63	64	66
Core involvement and risk groups	% of cohort with ≤2 positive cores, 69 25% IR, D'Amico criteria	Median # positive cores, 1	Mean % positive cores, 20.3%	Not available	Median % positive cores, 12.5% 17% NCCN IR/HR	% of cohort with ≤2 positive cores, 77.6	% of cohort with ≤2 positive cores, 99
Median follow-up (months)	77	60	43	60	86	80	120
Conversion to treatment*	36.5% (10-y) 45% (15-y)	50% (10-y)	24% (3-y)	40% (5-y)	49% (10-y)	33% (6.7-y)	52% (5-y) 73% (10-y)
Systemic progression	6.6% systemic progression in IR group	0.15% distant metastases	0% distant metastases	0.1%	0.5% distant metastases	0.4%	0.2%
Lymph node involvement and/or metastasis	3.1% (1.8% distant metastases; 1.3% pos. lymph nodes)	0.08% pos. lymph nodes	0.2% pos. lymph nodes		0.5% pos. lymph nodes		
Cancer-specific survival	98% (10-y)	99.9% (10-y)	100% (5-y)	100% (5-y)	99.9% (10-y)	99.8% (6.7-y)	>99% (10-y)
Overall survival	80% (10-y)	93% (10-y)	98% (10-y)	98% (5-y)	94.3% (10-y)	—	—
*Reason for conversion to treatment (% of entire cohort)							
Gleason grade change	9.5%	15.1%	38%	—	43% (10-y)	49%	34% (5-y) 41% (20-y) ^a
PSA increase	11.7%	—	26%	—	—	8.5%	—
Tumor volume increase	—	—	—	—	—	7.2%	—
Personal choice	1.6%	8%	8%	—	—	5% (anxiety)	5%

IR = intermediate risk; HR = high risk.

^aProtocol-based reclassification (included change in Gleason grade, number of positive cores, or cT stage)



NCCN Guidelines Version 5.2026

Prostate Cancer

References

1. Siegel RL, Kratzer TB, Giaquinto AN, et al. Cancer statistics, 2025. *CA Cancer J Clin* 2025;75:10–45. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39817679>.
2. Moyer VA. Screening for prostate cancer: U.S. Preventive Services Task Force recommendation statement. *Annals of internal medicine* 2012;157:120–134. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22801674>.
3. Barocas DA, Mallin K, Graves AJ, et al. Effect of the USPSTF grade D recommendation against screening for prostate cancer on incident prostate cancer diagnoses in the United States. *J Urol* 2015;194:1587–1593. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26087383>.
4. Drazer MW, Huo D, Eggener SE. National prostate cancer screening rates after the 2012 US Preventive Services Task Force recommendation discouraging prostate-specific antigen-based screening. *J Clin Oncol* 2015;33:2416–2423. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26056181>.
5. Etzioni R, Gulati R. Recent trends in PSA testing and prostate cancer incidence: A look at context. *JAMA Oncol* 2016;2:955–956. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27010657>.
6. Fedewa SA, Ward EM, Brawley O, Jemal A. Recent patterns of prostate-specific antigen testing for prostate cancer screening in the United States. *JAMA Intern Med* 2017;177:1040–1042. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28437537>.
7. Houston KA, King J, Li J, Jemal A. Trends in prostate cancer incidence rates and prevalence of prostate-specific antigen screening by socioeconomic status and regions in the US, 2004–2013. *J Urol* 2017;199:676–682. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28965781>.
8. Kearns JT, Holt SK, Wright JL, et al. PSA screening, prostate biopsy, and treatment of prostate cancer in the years surrounding the USPSTF recommendation against prostate cancer screening. *Cancer* 2018;124:2733–2739. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29781117>.
9. Jemal A, Fedewa SA, Ma J, et al. Prostate cancer incidence and PSA testing patterns in relation to USPSTF screening recommendations. *JAMA* 2015;314:2054–2061. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26575061>.
10. Sammon JD, Abdollah F, Choueiri TK, et al. Prostate-specific antigen screening after 2012 US Preventive Services Task Force recommendations. *JAMA* 2015;314:2077–2079. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26575066>.
11. Saka AH, Giaquinto AN, McCullough LE, et al. Cancer statistics for African American and Black people, 2025. *CA Cancer J Clin* 2025;75:111–140. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39976243>.
12. Dess RT, Hartman HE, Mahal BA, et al. Association of Black race with prostate cancer-specific and other-cause mortality. *JAMA Oncol* 2019;5:975–983. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31120534>.
13. PubMed Overview. National Institutes of Health; Available at: <https://pubmed.ncbi.nlm.nih.gov/about/>. Accessed October 24, 2025.
14. Freedman-Cass DA, Fischer T, Alpert AB, et al. The value and process of inclusion: Using sensitive, respectful, and inclusive language and images in NCCN content. *J Natl Compr Canc Netw* 2023;21:434–441. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37156485>.
15. Amin MB, Greene FL, Edge S, et al., eds. *AJCC Cancer Staging Manual* (ed 8th Edition). New York: Springer; 2017.
16. Cancer Protocol Templates. College of American Pathologists; 2025. Available at: <https://www.cap.org/protocols-and-guidelines/cancer-reporting-tools/cancer-protocol-templates#protocols>. Accessed October 24, 2025.



NCCN Guidelines Version 5.2026 Prostate Cancer

17. Social Security Administration. Period Life Table. 2025. Available at: <https://www.ssa.gov/OACT/STATS/table4c6.html>. Accessed October 24, 2025.
18. Male Life Expectancy. Memorial Sloan Kettering Cancer Center; Available at: https://www.mskcc.org/nomograms/prostate/male_life_expectancy. Accessed October 24, 2025.
19. Lee Schonberg Index. University of California San Francisco; Available at: <https://eprognosis.ucsf.edu/leeschonberg.php>. Accessed October 10, 2025.
20. Howard DH. Life expectancy and the value of early detection. J Health Econ 2005;24:891–906. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16129128>.
21. Szymanski KM, Wei JT, Dunn RL, Sanda MG. Development and validation of an abbreviated version of the expanded prostate cancer index composite instrument for measuring health-related quality of life among prostate cancer survivors. Urology 2010;76:1245–1250. Available at:
22. Sanda MG, Dunn RL, Michalski J, et al. Quality of life and satisfaction with outcome among prostate-cancer survivors. N Engl J Med 2008;358:1250–1261. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18354103>.
23. Chen RC, Basak R, Meyer AM, et al. Association between choice of radical prostatectomy, external beam radiotherapy, brachytherapy, or active surveillance and patient-reported quality of life among men with localized prostate cancer. JAMA 2017;317:1141–1150. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28324092>.
24. Hoffman KE, Penson DF, Zhao Z, et al. Patient-reported outcomes through 5 years for active surveillance, surgery, brachytherapy, or external beam radiation with or without androgen deprivation therapy for localized prostate cancer. JAMA 2020;323:149–163. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31935027>.
25. Donovan JL, Hamdy FC, Lane JA, et al. Patient-reported outcomes after monitoring, surgery, or radiotherapy for prostate cancer. N Engl J Med 2016;375:1425–1437. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27626365>.
26. Odeo S, Degu A. Factors affecting health-related quality of life among prostate cancer patients: A systematic review. J Oncol Pharm Pract 2020;26:1997–2010. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32972301>.
27. Naccarato A, Consuelo Souto S, Matheus WE, et al. Quality of life and sexual health in men with prostate cancer undergoing radical prostatectomy. Aging Male 2020;23:346–353. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30040004>.
28. Nelson CJ, Mulhall JP, Roth AJ. The association between erectile dysfunction and depressive symptoms in men treated for prostate cancer. J Sex Med 2011;8:560–566. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21155979>.
29. Klaassen Z, Arora K, Wilson SN, et al. Decreasing suicide risk among patients with prostate cancer: Implications for depression, erectile dysfunction, and suicidal ideation screening. Urol Oncol 2018;36:60–66. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28964659>.
30. Makarov DV, Chrouser K, Gore JL, et al. AUA white paper on implementation of shared decision making into urological practice. Urol Pract 2016;3:355–363. Available at:
31. Albright F, Stephenson RA, Agarwal N, et al. Prostate cancer risk prediction based on complete prostate cancer family history. Prostate 2015;75:390–398. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25408531>.
32. Bratt O, Drevin L, Akre O, et al. Family history and probability of prostate cancer, differentiated by risk category: a nationwide population-based study. J Natl Cancer Inst 2016;108. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27400876>.



NCCN Guidelines Version 5.2026

Prostate Cancer

33. Jansson F, Drevin L, Frisell T, et al. Concordance of non-low-risk disease among pairs of brothers with prostate cancer. *J Clin Oncol* 2018;36:JCO2017766907. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29652556>.
34. Beebe-Dimmer JL, Kapron AL, Fraser AM, et al. Risk of prostate cancer associated with familial and hereditary cancer syndromes. *J Clin Oncol* 2020;38:1807–1813. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32208047>.
35. Latham A, Srinivasan P, Kemel Y, et al. Microsatellite instability is associated with the presence of Lynch syndrome pan-cancer. *J Clin Oncol* 2018;37:JCO1800283. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30376427>.
36. Haraldsdottir S, Hampel H, Wei L, et al. Prostate cancer incidence in males with Lynch syndrome. *Genet Med* 2014;16:553–557. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24434690>.
37. Ryan S, Jenkins MA, Win AK. Risk of prostate cancer in Lynch syndrome: a systematic review and meta-analysis. *Cancer Epidemiol Biomarkers Prev* 2014;23:437–449. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24425144>.
38. Moran A, O'Hara C, Khan S, et al. Risk of cancer other than breast or ovarian in individuals with BRCA1 and BRCA2 mutations. *Fam Cancer* 2012;11:235–242. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22187320>.
39. Mersch J, Jackson MA, Park M, et al. Cancers associated with BRCA1 and BRCA2 mutations other than breast and ovarian. *Cancer* 2015;121:269–275. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25224030>.
40. Pilie PG, Johnson AM, Hanson KL, et al. Germline genetic variants in men with prostate cancer and one or more additional cancers. *Cancer* 2017;123:3925–3932. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28657667>.
41. Cheng HH, Sokolova AO, Schaeffer EM, et al. Germline and somatic mutations in prostate cancer for the clinician. *J Natl Compr Canc Netw* 2019;17:515–521. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31085765>.
42. Giri VN, Knudsen KE, Kelly WK, et al. Implementation of germline testing for prostate cancer: Philadelphia Prostate Cancer Consensus Conference 2019. *J Clin Oncol* 2020;38:2798–2811. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32516092>.
43. Stopsack KH, Vijai J, Conry M, et al. Germline DNA damage repair variants and prognosis of patients with high-risk or metastatic prostate cancer. *Clin Cancer Res* 2025;31:122–129. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39450704>.
44. Robinson D, Van Allen EM, Wu YM, et al. Integrative clinical genomics of advanced prostate cancer. *Cell* 2015;161:1215–1228. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26000489>.
45. Cancer Genome Atlas Research N. The molecular taxonomy of primary prostate cancer. *Cell* 2015;163:1011–1025. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26544944>.
46. Carter HB, Helfand B, Mamawala M, et al. Germline mutations in ATM and BRCA1/2 are associated with grade reclassification in men on active surveillance for prostate cancer. *Eur Urol* 2019;75:743–749. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30309687>.
47. Wu Y, Yu H, Li S, et al. Rare germline pathogenic mutations of DNA repair genes are most strongly associated with grade group 5 prostate cancer. *Eur Urol Oncol* 2020;3:224–230. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31948886>.
48. Giri VN, Obeid E, Gross L, et al. Inherited mutations in men undergoing multigene panel testing for prostate cancer: Emerging implications for personalized prostate cancer genetic evaluation. *JCO Precision Oncol* 2017;published online May 4, 2017. Available at: <http://ascopubs.org/doi/full/10.1200/PO.16.00039>.



NCCN Guidelines Version 5.2026

Prostate Cancer

49. Yadav S, Hart SN, Hu C, et al. Contribution of inherited DNA-repair gene mutations to hormone-sensitive and castrate-resistant metastatic prostate cancer and implications for clinical outcome. *JCO Precis Oncol* 2019;3. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32923857>.
50. Boyle JL, Hahn AW, Kapron AL, et al. Pathogenic germline DNA repair gene and HOXB13 mutations in men with metastatic prostate cancer. *JCO Precis Oncol* 2020;4. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32923906>.
51. Pritchard CC, Mateo J, Walsh MF, et al. Inherited DNA-repair gene mutations in men with metastatic prostate cancer. *N Engl J Med* 2016;375:443–453. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27433846>.
52. Castro E, Romero-Laorden N, Del Pozo A, et al. PROREPAIR-B: A prospective cohort study of the impact of germline DNA repair mutations on the outcomes of patients with metastatic castration-resistant prostate cancer. *J Clin Oncol* 2019;37:490–503. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30625039>.
53. Giri VN, Hegarty SE, Hyatt C, et al. Germline genetic testing for inherited prostate cancer in practice: Implications for genetic testing, precision therapy, and cascade testing. *Prostate* 2018;79:333–339. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30450585>.
54. Nicolosi P, Ledet E, Yang S, et al. Prevalence of germline variants in prostate cancer and implications for current genetic testing guidelines. *JAMA Oncol* 2019;5:523–528. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30730552>.
55. Struewing JP, Hartge P, Wacholder S, et al. The risk of cancer associated with specific mutations of BRCA1 and BRCA2 among Ashkenazi Jews. *N Engl J Med* 1997;336:1401–1408. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/9145676>.
56. Kirchhoff T, Kauff ND, Mitra N, et al. BRCA mutations and risk of prostate cancer in Ashkenazi Jews. *Clin Cancer Res* 2004;10:2918–2921. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15131025>.
57. Cancer risks in BRCA2 mutation carriers. The Breast Cancer Linkage Consortium. *J Natl Cancer Inst* 1999;91:1310–1316. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10433620>.
58. Agalliu I, Gern R, Leanza S, Burk RD. Associations of high-grade prostate cancer with BRCA1 and BRCA2 founder mutations. *Clin Cancer Res* 2009;15:1112–1120. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19188187>.
59. Ford D, Easton DF, Bishop DT, et al. Risks of cancer in BRCA1-mutation carriers. Breast Cancer Linkage Consortium. *Lancet* 1994;343:692–695. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7907678>.
60. Gallagher DJ, Gaudet MM, Pal P, et al. Germline BRCA mutations denote a clinicopathologic subset of prostate cancer. *Clin Cancer Res* 2010;16:2115–2121. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20215531>.
61. Leongamornlert D, Mahmud N, Tymrakiewicz M, et al. Germline BRCA1 mutations increase prostate cancer risk. *Br J Cancer* 2012;106:1697–1701. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22516946>.
62. Liede A, Karlan BY, Narod SA. Cancer risks for male carriers of germline mutations in BRCA1 or BRCA2: a review of the literature. *J Clin Oncol* 2004;22:735–742. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14966099>.
63. Thompson D, Easton DF. Cancer incidence in BRCA1 mutation carriers. *J Natl Cancer Inst* 2002;94:1358–1365. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/12237281>.
64. Tulinius H, Olafsdottir GH, Sigvaldason H, et al. The effect of a single BRCA2 mutation on cancer in Iceland. *J Med Genet* 2002;39:457–462. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12114473>.
65. van Asperen CJ, Brohet RM, Meijers-Heijboer EJ, et al. Cancer risks in BRCA2 families: estimates for sites other than breast and ovary. *J Med*



NCCN Guidelines Version 5.2026 Prostate Cancer

Genet 2005;42:711–719. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16141007>.

66. Lecarpentier J, Silvestri V, Kuchenbaecker KB, et al. Prediction of breast and prostate cancer risks in male BRCA1 and BRCA2 mutation carriers using polygenic risk scores. J Clin Oncol 2017;35:2240–2250. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28448241>.

67. Page EC, Bancroft EK, Brook MN, et al. Interim results from the IMPACT study: Evidence for prostate-specific antigen screening in BRCA2 mutation carriers. Eur Urol 2019;76:831–842. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31537406>.

68. Mano R, Tamir S, Kedar I, et al. Malignant abnormalities in male BRCA mutation carriers: Results from a prospectively screened cohort. JAMA Oncol 2018;4:872–874. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29710070>.

69. Helgason H, Rafnar T, Olafsdottir HS, et al. Loss-of-function variants in ATM confer risk of gastric cancer. Nat Genet 2015;47:906–910. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26098866>.

70. Erkko H, Xia B, Nikkila J, et al. A recurrent mutation in PALB2 in Finnish cancer families. Nature 2007;446:316–319. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/17287723>.

71. Naslund-Koch C, Nordestgaard BG, Bojesen SE. Increased risk for other cancers in addition to breast cancer for CHEK2*1100delC heterozygotes estimated from the Copenhagen General Population Study. J Clin Oncol 2016;34:1208–1216. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26884562>.

72. Wu Y, Yu H, Zheng SL, et al. A comprehensive evaluation of CHEK2 germline mutations in men with prostate cancer. Prostate 2018;78:607–615. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29520813>.

73. Castro E, Goh C, Olmos D, et al. Germline BRCA mutations are associated with higher risk of nodal involvement, distant metastasis, and

poor survival outcomes in prostate cancer. J Clin Oncol 2013;31:1748–1757. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23569316>.

74. Mitra A, Fisher C, Foster CS, et al. Prostate cancer in male BRCA1 and BRCA2 mutation carriers has a more aggressive phenotype. Br J Cancer 2008;98:502–507. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18182994>.

75. Na R, Zheng SL, Han M, et al. Germline mutations in ATM and BRCA1/2 distinguish risk for lethal and indolent prostate cancer and are associated with early age at death. Eur Urol 2016;71:740–747. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27989354>.

76. Narod SA, Neuhausen S, Vichodez G, et al. Rapid progression of prostate cancer in men with a BRCA2 mutation. Br J Cancer 2008;99:371–374. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18577985>.

77. Thorne H, Willems AJ, Niedermayr E, et al. Decreased prostate cancer-specific survival of men with BRCA2 mutations from multiple breast cancer families. Cancer Prev Res (Phila) 2011;4:1002–1010. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21733824>.

78. Tryggvadottir L, Vidarsdottir L, Thorgeirsson T, et al. Prostate cancer progression and survival in BRCA2 mutation carriers. J Natl Cancer Inst 2007;99:929–935. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17565157>.

79. Wei Y, Wu J, Gu W, et al. Prognostic value of germline DNA repair gene mutations in de novo metastatic and castration-sensitive prostate cancer. Oncologist 2020;25:e1042–e1050. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32190957>.

80. Dominguez-Valentin M, Sampson JR, Seppala TT, et al. Cancer risks by gene, age, and gender in 6350 carriers of pathogenic mismatch repair variants: findings from the Prospective Lynch Syndrome Database. Genet Med 2020;22:15–25. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31337882>.



NCCN Guidelines Version 5.2026 Prostate Cancer

81. Moller P, Seppala TT, Bernstein I, et al. Cancer risk and survival in path_MMR carriers by gene and gender up to 75 years of age: a report from the Prospective Lynch Syndrome Database. *Gut* 2018;67:1306–1316. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28754778>.

82. Abida W, Cheng ML, Armenia J, et al. Analysis of the prevalence of microsatellite instability in prostate cancer and response to immune checkpoint blockade. *JAMA Oncol* 2019;5:471–478. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30589920>.

83. Zhou M. High-grade prostatic intraepithelial neoplasia, PIN-like carcinoma, ductal carcinoma, and intraductal carcinoma of the prostate. *Mod Pathol* 2018;31:S71–79. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29297491>.

84. Porter LH, Lawrence MG, Ilic D, et al. Systematic review links the prevalence of intraductal carcinoma of the prostate to prostate cancer risk categories. *Eur Urol* 2017;72:492–495. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28342640>.

85. Chua MLK, Lo W, Pintilie M, et al. A prostate cancer "nimbusus": Genomic instability and SchLAP1 dysregulation underpin aggression of intraductal and cribriform subpathologies. *Eur Urol* 2017;72:665–674. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28511883>.

86. Seipel AH, Whittington T, Delahunt B, et al. Genetic profile of ductal adenocarcinoma of the prostate. *Hum Pathol* 2017;69:1–7. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28457729>.

87. Bottcher R, Kweldam CF, Livingstone J, et al. Cribriform and intraductal prostate cancer are associated with increased genomic instability and distinct genomic alterations. *BMC Cancer* 2018;18:8. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29295717>.

88. Schweizer MT, Antonarakis ES, Bismar TA, et al. Genomic characterization of prostatic ductal adenocarcinoma identifies a high prevalence of DNA repair gene mutations. *JCO Precis Oncol* 2019;3. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31123724>.

89. Antonarakis ES, Shaukat F, Isaacsson Velho P, et al. Clinical features and therapeutic outcomes in men with advanced prostate cancer and DNA mismatch repair gene mutations. *Eur Urol* 2018;75:378–382. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30337059>.

90. Antonarakis ES, Shaukat F, Isaacsson Velho P, et al. Clinical features and therapeutic outcomes in men with advanced prostate cancer and DNA mismatch repair gene mutations. *Eur Urol* 2019;75:378–382. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30337059>.

91. Isaacsson Velho P, Silberstein JL, Markowski MC, et al. Intraductal/ductal histology and lymphovascular invasion are associated with germline DNA-repair gene mutations in prostate cancer. *Prostate* 2018;78:401–407. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29368341>.

92. Taylor RA, Fraser M, Livingstone J, et al. Germline BRCA2 mutations drive prostate cancers with distinct evolutionary trajectories. *Nat Commun* 2017;8:13671. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28067867>.

93. Risbridger GP, Taylor RA, Clouston D, et al. Patient-derived xenografts reveal that intraductal carcinoma of the prostate is a prominent pathology in BRCA2 mutation carriers with prostate cancer and correlates with poor prognosis. *Eur Urol* 2015;67:496–503. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25154392>.

94. Jensen K, Konnick EQ, Schweizer MT, et al. Association of Clonal Hematopoiesis in DNA Repair Genes With Prostate Cancer Plasma Cell-free DNA Testing Interference. *JAMA Oncol* 2021;7:107–110. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33151258>.

95. Middha S, Zhang L, Nafa K, et al. Reliable pan-cancer microsatellite instability assessment by using targeted next-generation sequencing data. *JCO Precis Oncol* 2017;2017. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30211344>.



NCCN Guidelines Version 5.2026 Prostate Cancer

96. Guedes LB, Antonarakis ES, Schweizer MT, et al. MSH2 loss in primary prostate cancer. *Clin Cancer Res* 2017;23:6863–6874. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28790115>.

97. Hempelmann JA, Lockwood CM, Konnick EQ, et al. Microsatellite instability in prostate cancer by PCR or next-generation sequencing. *J Immunother Cancer* 2018;6:29. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29665853>.

98. Carthon B, Sibold HC, Pentz RD. Prostate cancer: Community education and disparities in diagnosis and treatment. *Oncologist* 2021;26:537–548. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33683758>.

99. Lillard JW, Jr., Moses KA, Mahal BA, George DJ. Racial disparities in Black men with prostate cancer: A literature review. *Cancer* 2022;128:3787–3795. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36066378>.

100. Mahal BA, Berman RA, Taplin ME, Huang FW. Prostate cancer-specific mortality across Gleason scores in Black vs nonblack men. *JAMA* 2018;320:2479–2481. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30561471>.

101. Wang M, Chi G, Bodovski Y, et al. Temporal and spatial trends and determinants of aggressive prostate cancer among Black and white men with prostate cancer. *Cancer Causes Control* 2020;31:63–71. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31732913>.

102. Kovtun KA, Chen MH, Braccioforte MH, et al. Race and mortality risk after radiation therapy in men treated with or without androgen-suppression therapy for favorable-risk prostate cancer. *Cancer* 2016;122:3608–3614. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27490845>.

103. Swami N, Baez YA, Franco I, et al. Localized prostate cancer disparities in risk group at presentation and access to treatment for Hispanic men. *Prostate Cancer Prostatic Dis* 2023;26:309–316. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35306541>.

104. Hougen HY, Swami N, Dee EC, et al. Disparities in diagnosis, treatment access, and time to treatment among Hispanic men with metastatic prostate cancer. *JCO Oncol Pract* 2023;19:645–653. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37262399>.

105. Jain B, Ng K, Santos PMG, et al. Prostate cancer disparities in risk group at presentation and access to treatment for Asian Americans, Native Hawaiians, and Pacific Islanders: A study with disaggregated ethnic groups. *JCO Oncol Pract* 2022;18:e204–e218. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34709962>.

106. Barocas DA, Grubb R, 3rd, Black A, et al. Association between race and follow-up diagnostic care after a positive prostate cancer screening test in the Prostate, Lung, Colorectal, and Ovarian Cancer Screening Trial. *Cancer* 2013;119:2223–2229. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23559420>.

107. Han Y, Rand KA, Hazelett DJ, et al. Prostate cancer susceptibility in men of African ancestry at 8q24. *J Natl Cancer Inst* 2016;108. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26823525>.

108. Vince RA, Jr., Jiang R, Bank M, et al. Evaluation of social determinants of health and prostate cancer outcomes among Black and white patients: A systematic review and meta-analysis. *JAMA Netw Open* 2023;6:e2250416. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36630135>.

109. Yamoah K, Johnson MH, Choeurng V, et al. Novel biomarker signature that may predict aggressive disease in African American men with prostate cancer. *J Clin Oncol* 2015;33:2789–2796. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26195723>.

110. Zhang H, Messing EM, Travis LB, et al. Age and racial differences among PSA-detected (AJCC stage T1cN0M0) prostate cancer in the U.S.: a population-based study of 70,345 men. *Front Oncol* 2013;3:312. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24392353>.

111. Bickell NA, Lin JJ, Abramson SR, et al. Racial disparities in clinically significant prostate cancer treatment: The potential health information



NCCN Guidelines Version 5.2026

Prostate Cancer

technology offers. J Oncol Pract 2018;14:e23–e33. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/29194001>.

112. Friedlander DF, Trinh QD, Krasnova A, et al. Racial disparity in delivering definitive therapy for intermediate/high-risk localized prostate cancer: The impact of facility features and socioeconomic characteristics. Eur Urol 2017;73:445–451. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/28778619>.

113. Moses KA, Orom H, Brasel A, et al. Racial/Ethnic Disparity in Treatment for Prostate Cancer: Does Cancer Severity Matter? Urology 2017;99:76–83. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/27667157>.

114. Riviere P, Luterstein E, Kumar A, et al. Survival of African American and non-Hispanic white men with prostate cancer in an equal-access health care system. Cancer 2020;126:1683–1690. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/31984482>.

115. Alexander M, Zhu K, Cullen J, et al. Race and overall survival in men diagnosed with prostate cancer in the Department of Defense Military Health System, 1990–2010. Cancer Causes Control 2019;30:627–635. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30997591>.

116. Halabi S, Dutta S, Tangen CM, et al. Clinical outcomes in men of diverse ethnic backgrounds with metastatic castration-resistant prostate cancer. Ann Oncol 2020;31:930–941. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/32289380>.

117. Cooperberg MR, Pasta DJ, Elkin EP, et al. The University of California, San Francisco Cancer of the Prostate Risk Assessment score: a straightforward and reliable preoperative predictor of disease recurrence after radical prostatectomy. J Urol 2005;173:1938–1942. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/15879786>.

118. Dess RT, Suresh K, Zelefsky MJ, et al. Development and validation of a clinical prognostic stage group system for nonmetastatic prostate cancer using disease-specific mortality results from the international staging collaboration for cancer of the prostate. JAMA Oncol

2020;6:1912–1920. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/33090219>.

119. Zelic R, Garmo H, Zugna D, et al. Predicting prostate cancer death with different pretreatment risk stratification tools: A head-to-head comparison in a nationwide cohort study. Eur Urol 2020;77:180–188. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31606332>.

120. D'Amico AV, Whittington R, Malkowicz SB, et al. Biochemical outcome after radical prostatectomy or external beam radiation therapy for patients with clinically localized prostate carcinoma in the prostate specific antigen era. Cancer 2002;95:281–286. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/12124827>.

121. D'Amico AV, Whittington R, Malkowicz SB, et al. Biochemical outcome after radical prostatectomy, external beam radiation therapy, or interstitial radiation therapy for clinically localized prostate cancer. JAMA 1998;280:969–974. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/9749478>.

122. D'Amico AV, Whittington R, Malkowicz SB, et al. Pretreatment nomogram for prostate-specific antigen recurrence after radical prostatectomy or external-beam radiation therapy for clinically localized prostate cancer. J Clin Oncol 1999;17:168–172. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/10458230>.

123. Epstein JI, Egevad L, Amin MB, et al. The 2014 International Society of Urological Pathology (ISUP) consensus conference on Gleason grading of prostatic carcinoma: definition of grading patterns and proposal for a new grading system. Am J Surg Pathol 2016;40:244–252. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/26492179>.

124. Epstein JI, Zelefsky MJ, Sjoberg DD, et al. A contemporary prostate cancer grading system: a validated alternative to the Gleason score. Eur Urol 2016;69:428–435. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/26166626>.

125. Loeb S, Folkvaljon Y, Robinson D, et al. Evaluation of the 2015 Gleason grade groups in a nationwide population-based cohort. Eur Urol



NCCN Guidelines Version 5.2026

Prostate Cancer

2016;69:1135–1141. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26707871>.

126. Ham WS, Chalfin HJ, Feng Z, et al. New prostate cancer grading system predicts long-term survival following surgery for Gleason score 8–10 prostate cancer. *Eur Urol* 2016;71:907–912. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27876305>.

127. Delahunt B, Egevad L, Srigley JR, et al. Validation of International Society of Urological Pathology (ISUP) grading for prostatic adenocarcinoma in thin core biopsies using TROG 03.04 'RADAR' trial clinical data. *Pathology* 2015;47:520–525. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26325671>.

128. Mathieu R, Moschini M, Beyer B, et al. Prognostic value of the new grade groups in prostate cancer: a multi-institutional European validation study. *Prostate Cancer Prostatic Dis* 2017;20:197–202. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28071673>.

129. Leapman MS, Cowan JE, Simko J, et al. Application of a prognostic Gleason grade grouping system to assess distant prostate cancer outcomes. *Eur Urol* 2016;71:750–759. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27940155>.

130. He J, Albertsen PC, Moore D, et al. Validation of a contemporary five-tiered Gleason grade grouping using population-based data. *Eur Urol* 2017;71:760–763. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27939073>.

131. Pompe RS, Davis-Bondarenko H, Zaffuto E, et al. Population-based validation of the 2014 ISUP Gleason grade groups in patients treated with radical prostatectomy, brachytherapy, external beam radiation, or no local treatment. *Prostate* 2017;77:686–693. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28156003>.

132. Kirmiz S, Qi J, Babitz SK, et al. Grade Groups provide improved predictions of pathological and early oncologic outcomes compared with Gleason score risk groups. *J Urol* 2019;201:278–283. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/30195846>.

133. Reese AC, Pierorazio PM, Han M, Partin AW. Contemporary evaluation of the National Comprehensive Cancer Network prostate cancer risk classification system. *Urology* 2012;80:1075–1079. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22995570>.

134. Muralidhar V, Chen MH, Reznor G, et al. Definition and validation of "favorable high-risk prostate cancer": implications for personalizing treatment of radiation-managed patients. *Int J Radiat Oncol Biol Phys* 2015;93:828–835. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/26530751>.

135. Gandaglia G, Karnes RJ, Sivaraman A, et al. Are all grade group 4 prostate cancers created equal? Implications for the applicability of the novel grade grouping. *Urol Oncol* 2017;35:461 e467–461 e414. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28359746>.

136. Dinh KT, Muralidhar V, Mahal BA, et al. Occult high-risk disease in clinically low-risk prostate cancer with $\geq 50\%$ positive biopsy cores: should national guidelines stop calling them low-risk? *Urology* 2015;87:125–132. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/26391387>.

137. Dinh KT, Mahal BA, Ziehr DR, et al. Incidence and predictors of upgrading and up staging among 10,000 contemporary patients with low risk prostate cancer. *J Urol* 2015;194:343–349. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/25681290>.

138. Zumsteg ZS, Spratt DE, Pei I, et al. A new risk classification system for therapeutic decision making with intermediate-risk prostate cancer patients undergoing dose-escalated external-beam radiation therapy. *Eur Urol* 2013;64:895–902. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23541457>.

139. Johns Hopkins Medicine. The Partin Tables. Available at: <https://www.hopkinsmedicine.org/brady-urology-institute/conditions-and-treatments/prostate-cancer/risk-assessment-tools/partin-tables>. Accessed October 24, 2025.



NCCN Guidelines Version 5.2026

Prostate Cancer

140. Makarov DV, Trock BJ, Humphreys EB, et al. Updated nomogram to predict pathologic stage of prostate cancer given prostate-specific antigen level, clinical stage, and biopsy Gleason score (Partin tables) based on cases from 2000 to 2005. *Urology* 2007;69:1095–1101. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17572194>.

141. Borque A, Rubio-Briones J, Esteban LM, et al. Implementing the use of nomograms by choosing threshold points in predictive models: 2012 updated Partin Tables vs a European predictive nomogram for organ-confined disease in prostate cancer. *BJU Int* 2014;113:878–886. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24529282>.

142. Tosoian JJ, Chappidi M, Feng Z, et al. Prediction of pathological stage based on clinical stage, serum prostate-specific antigen, and biopsy Gleason score: Partin Tables in the contemporary era. *BJU Int* 2017;119:676–683. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27367645>.

143. Kattan MW, Eastham JA, Wheeler TM, et al. Counseling men with prostate cancer: a nomogram for predicting the presence of small, moderately differentiated, confined tumors. *J Urol* 2003;170:1792–1797. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14532778>.

144. Leyh-Bannurah SR, Dell'Oglio P, Tian Z, et al. A proposal of a new nomogram for predicting upstaging in contemporary D'Amico low-risk prostate cancer patients. *World J Urol* 2017;35:189–197. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27289238>.

145. Wong LM, Neal DE, Finelli A, et al. Evaluation of models predicting insignificant prostate cancer to select men for active surveillance of prostate cancer. *Prostate Cancer Prostatic Dis* 2015;18:137–143. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25667108>.

146. Memorial Sloan-Kettering Cancer Center. Prostate Cancer Nomograms. Available at: <http://www.mskcc.org/mskcc/html/10088.cfm>. Accessed October 24, 2025.

147. Punnen S, Freedland SJ, Presti JC, Jr., et al. Multi-institutional validation of the CAPRA-S score to predict disease recurrence and

mortality after radical prostatectomy. *Eur Urol* 2014;65:1171–1177. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23587869>.

148. Stephenson AJ, Scardino PT, Eastham JA, et al. Preoperative nomogram predicting the 10-year probability of prostate cancer recurrence after radical prostatectomy. *J Natl Cancer Inst* 2006;98:715–717. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16705126>.

149. Stephenson AJ, Kattan MW, Eastham JA, et al. Prostate cancer-specific mortality after radical prostatectomy for patients treated in the prostate-specific antigen era. *J Clin Oncol* 2009;27:4300–4305. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19636023>.

150. Graefen M, Haese A, Pichlmeier U, et al. A validated strategy for side specific prediction of organ confined prostate cancer: a tool to select for nerve sparing radical prostatectomy. *J Urol* 2001;165:857–863. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11176486>.

151. Ohori M, Kattan MW, Koh H, et al. Predicting the presence and side of extracapsular extension: a nomogram for staging prostate cancer. *J Urol* 2004;171:1844–1849; discussion 1849. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15076291>.

152. Steuber T, Graefen M, Haese A, et al. Validation of a nomogram for prediction of side specific extracapsular extension at radical prostatectomy. *J Urol* 2006;175:939–944; discussion 944. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16469587>.

153. Briganti A, Chun FK, Salonia A, et al. A nomogram for staging of exclusive nonobturator lymph node metastases in men with localized prostate cancer. *Eur Urol* 2007;51:112–119; discussion 119–120. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16806662>.

154. Cagiannos I, Karakiewicz P, Eastham JA, et al. A preoperative nomogram identifying decreased risk of positive pelvic lymph nodes in patients with prostate cancer. *J Urol* 2003;170:1798–1803. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14532779>.



NCCN Guidelines Version 5.2026 Prostate Cancer

155. Gandaglia G, Fossati N, Zaffuto E, et al. Development and internal validation of a novel model to identify the candidates for extended pelvic lymph node dissection in prostate cancer. *Eur Urol* 2017;72:632–640. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28412062>.

156. Gandaglia G, Ploussard G, Valerio M, et al. A novel nomogram to identify candidates for extended pelvic lymph node dissection among patients with clinically localized prostate cancer diagnosed with magnetic resonance imaging-targeted and systematic biopsies. *Eur Urol* 2019;75:506–514. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30342844>.

157. Kattan MW, Potters L, Blasko JC, et al. Pretreatment nomogram for predicting freedom from recurrence after permanent prostate brachytherapy in prostate cancer. *Urology* 2001;58:393–399. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11549487>.

158. Potters L, Morgenstern C, Calugaru E, et al. 12-year outcomes following permanent prostate brachytherapy in patients with clinically localized prostate cancer. *J Urol* 2008;179:S20–24. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18405743>.

159. Potters L, Roach M, 3rd, Davis BJ, et al. Postoperative nomogram predicting the 9-year probability of prostate cancer recurrence after permanent prostate brachytherapy using radiation dose as a prognostic variable. *Int J Radiat Oncol Biol Phys* 2010;76:1061–1065. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19540064>.

160. Zelefsky MJ, Kattan MW, Fearn P, et al. Pretreatment nomogram predicting ten-year biochemical outcome of three-dimensional conformal radiotherapy and intensity-modulated radiotherapy for prostate cancer. *Urology* 2007;70:283–287. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17826490>.

161. Lee SJ, Lindquist K, Segal MR, Covinsky KE. Development and validation of a prognostic index for 4-year mortality in older adults. *JAMA* 2006;295:801–808. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16478903>.

162. Kattan MW, Wheeler TM, Scardino PT. Postoperative nomogram for disease recurrence after radical prostatectomy for prostate cancer. *J Clin Oncol* 1999;17:1499–1507. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10334537>.

163. Ondracek RP, Kattan MW, Murekeyisoni C, et al. Validation of the Kattan nomogram for prostate cancer recurrence after radical prostatectomy. *J Natl Compr Canc Netw* 2016;14:1395–1401. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27799510>.

164. Tendulkar RD, Agrawal S, Gao T, et al. Contemporary update of a multi-institutional predictive nomogram for salvage radiotherapy after radical prostatectomy. *J Clin Oncol* 2016;34:3648–3654. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27528718>.

165. Dearnaley DP, Khoo VS, Norman AR, et al. Comparison of radiation side-effects of conformal and conventional radiotherapy in prostate cancer: a randomised trial. *Lancet* 1999;353:267–272. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9929018>.

166. Khoo VS. Radiotherapeutic techniques for prostate cancer, dose escalation and brachytherapy. *Clin Oncol (R Coll Radiol)* 2005;17:560–571. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16238144>.

167. D'Amico AV, Cote K, Loffredo M, et al. Determinants of prostate cancer-specific survival after radiation therapy for patients with clinically localized prostate cancer. *J Clin Oncol* 2002;20:4567–4573. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12454114>.

168. Dell'Oglio P, Suardi N, Boorjian SA, et al. Predicting survival of men with recurrent prostate cancer after radical prostatectomy. *Eur J Cancer* 2016;54:27–34. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26707594>.

169. Abdollah F, Karnes RJ, Suardi N, et al. Predicting survival of patients with node-positive prostate cancer following multimodal treatment. *Eur Urol* 2014;65:554–562. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24094576>.



NCCN Guidelines Version 5.2026 Prostate Cancer

170. D'Amico AV, Moul JW, Carroll PR, et al. Surrogate end point for prostate cancer-specific mortality after radical prostatectomy or radiation therapy. *J Natl Cancer Inst* 2003;95:1376–1383. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/13130113>.

171. Murphy AB, Abern MR, Liu L, et al. Impact of a genomic test on treatment decision in a predominantly African American population with favorable-risk prostate cancer: A randomized trial. *J Clin Oncol* 2021;39:1660–1670. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33835822>.

172. Hu JC, Tosoian JJ, Qi J, et al. Clinical utility of gene expression classifiers in men with newly diagnosed prostate cancer JCO Precis Oncol 2018;published online, October 19, 2018 Available at: <http://ascopubs.org/doi/abs/10.1200/PO.18.00163>.

173. Marascio J, Spratt DE, Zhang J, et al. Prospective study to define the clinical utility and benefit of Decipher testing in men following prostatectomy. *Prostate Cancer Prostatic Dis* 2020;23:295–302. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31719663>.

174. Vince RA, Jr., Jiang R, Qi J, et al. Impact of Decipher Biopsy testing on clinical outcomes in localized prostate cancer in a prospective statewide collaborative. *Prostate Cancer Prostatic Dis* 2022;25:677–683. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34285350>.

175. Morgan TM, Daignault-Newton S, Spratt DE, et al. Impact of gene expression classifier testing on adjuvant treatment following radical prostatectomy: The G-MINOR prospective randomized cluster-crossover trial. *Eur Urol* 2025;87:228–237. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39379238>.

176. Jia AY, Sun Y, Patel M, et al. Cross-comparison individual patient-level analysis of three gene expression signatures in localized prostate in over 50,000 men. *JCO Precis Oncol* 2025;9:e2400705. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/40479622>.

177. Lin DW, Zheng Y, McKenney JK, et al. 17-gene genomic prostate score test results in the canary prostate active surveillance study (PASS)

cohort. *Journal of Clinical Oncology* 2020;38:1549–1557. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32130059>.

178. Spratt DE, Liu VYT, Michalski J, et al. Genomic classifier performance in intermediate-risk prostate cancer: Results from NRG Oncology/RTOG 0126 randomized phase 3 trial. *Int J Radiat Oncol Biol Phys* 2023;117:370–377. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37137444>.

179. Feng FY, Huang HC, Spratt DE, et al. Validation of a 22-Gene Genomic Classifier in Patients With Recurrent Prostate Cancer: An Ancillary Study of the NRG/RTOG 9601 Randomized Clinical Trial. *JAMA Oncol* 2021;7:544–552. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33570548>.

180. Nguyen PL, Huang HR, Spratt DE, et al. Analysis of a biopsy-based genomic classifier in high-risk prostate cancer: Meta-analysis of the NRG Oncology/Radiation Therapy Oncology Group 9202, 9413, and 9902 phase 3 randomized trials. *Int J Radiat Oncol Biol Phys* 2023;116:521–529. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36596347>.

181. Dal Pra A, Ghadjar P, Hayoz S, et al. Validation of the Decipher genomic classifier in patients receiving salvage radiotherapy without hormone therapy after radical prostatectomy - an ancillary study of the SAKK 09/10 randomized clinical trial. *Ann Oncol* 2022;33:950–958. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35636621>.

182. Cullen J, Rosner IL, Brand TC, et al. A biopsy-based 17-gene genomic prostate score predicts recurrence after radical prostatectomy and adverse surgical pathology in a racially diverse population of men with clinically low- and intermediate-risk prostate cancer. *Eur Urol* 2015;68:123–131. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25465337>.

183. Klein EA, Cooperberg MR, Magi-Galluzzi C, et al. A 17-gene assay to predict prostate cancer aggressiveness in the context of Gleason grade heterogeneity, tumor multifocality, and biopsy undersampling. *Eur Urol* 2014;66:550–560. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24836057>.



NCCN Guidelines Version 5.2026 Prostate Cancer

184. Cooperberg MR, Simko JP, Cowan JE, et al. Validation of a cell-cycle progression gene panel to improve risk stratification in a contemporary prostatectomy cohort. *J Clin Oncol* 2013;31:1428–1434. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23460710>.
185. Lin DW, Crawford ED, Keane T, et al. Identification of men with low-risk biopsy-confirmed prostate cancer as candidates for active surveillance. *Urol Oncol* 2018;36:310 e317–310 e313. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29655620>.
186. Tosoian JJ, Chappidi MR, Bishoff JT, et al. Prognostic utility of biopsy-derived cell cycle progression score in patients with National Comprehensive Cancer Network low-risk prostate cancer undergoing radical prostatectomy: implications for treatment guidance. *BJU Int* 2017;120:808–814. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28481440>.
187. Esteva A, Feng J, van der Wal D, et al. Prostate cancer therapy personalization via multi-modal deep learning on randomized phase III clinical trials. *NPJ Digit Med* 2022;5:71. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35676445>.
188. Spratt DE, Liu VYT, Jia AY, et al. Meta-analysis of individual patient-level data for a multimodal artificial intelligence biomarker in high-risk prostate cancer: Results from six NRG/RTOG phase 3 randomized trials. *Eur Urol* 2024;86:369–371. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39025748>.
189. Tward JD, Huang HC, Esteva A, et al. Prostate cancer risk stratification in NRG Oncology phase III randomized trials using multimodal deep learning with digital histopathology. *JCO Precis Oncol* 2024;8:e2400145. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39447096>.
190. Spratt DE, Tang S, Sun Y, et al. Artificial intelligence predictive model for hormone therapy use in prostate cancer. *NEJM Evid* 2023;2:EVID0a2300023. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/38320143>.
191. Armstrong AJ, Liu VYT, Selvaraju RR, et al. Development and validation of an artificial intelligence digital pathology biomarker to predict benefit of long-term hormonal therapy and radiotherapy in men with high-risk prostate cancer across multiple phase III trials. *J Clin Oncol* 2025;JCO2400365. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/40239134>.
192. Mason BR, Eastham JA, Davis BJ, et al. Current status of MRI and PET in the NCCN Guidelines for Prostate Cancer. *J Natl Compr Canc Netw* 2019;17:506–513. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31085758>.
193. Schoots IG, Barentsz JO, Bittencourt LK, et al. PI-RADS Committee position on MRI without contrast medium in biopsy-naïve men with suspected prostate cancer: Narrative review. *AJR Am J Roentgenol* 2021;216:3–19. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32812795>.
194. de Rooij M, Hamoen EH, Witjes JA, et al. Accuracy of Magnetic Resonance Imaging for Local Staging of Prostate Cancer: A Diagnostic Meta-analysis. *Eur Urol* 2016;70:233–245. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26215604>.
195. Turkbey B, Mani H, Shah V, et al. Multiparametric 3T prostate magnetic resonance imaging to detect cancer: histopathological correlation using prostatectomy specimens processed in customized magnetic resonance imaging based molds. *J Urol* 2011;186:1818–1824. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21944089>.
196. Siddiqui MM, Rais-Bahrami S, Truong H, et al. Magnetic resonance imaging/ultrasound-fusion biopsy significantly upgrades prostate cancer versus systematic 12-core transrectal ultrasound biopsy. *Eur Urol* 2013;64:713–719. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23787357>.
197. Rastinehad AR, Turkbey B, Salami SS, et al. Improving detection of clinically significant prostate cancer: magnetic resonance imaging/transrectal ultrasound fusion guided prostate biopsy. *J Urol*



NCCN Guidelines Version 5.2026

Prostate Cancer

2013;191:1749–1754. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24333515>.

198. Wysock JS, Rosenkrantz AB, Huang WC, et al. A prospective, blinded comparison of magnetic resonance (MR) imaging-ultrasound fusion and visual estimation in the performance of MR-targeted prostate biopsy: the PROFUS trial. *Eur Urol* 2014;66:343–351. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24262102>.

199. Ahdoot M, Wilbur AR, Reese SE, et al. MRI-targeted, systematic, and combined biopsy for prostate cancer diagnosis. *N Engl J Med* 2020;382:917–928. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/32130814>.

200. Somford DM, Hamoen EH, Futterer JJ, et al. The predictive value of endorectal 3 Tesla multiparametric magnetic resonance imaging for extraprostatic extension in patients with low, intermediate and high risk prostate cancer. *J Urol* 2013;190:1728–1734. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23680307>.

201. Park BH, Jeon HG, Jeong BC, et al. Influence of magnetic resonance imaging in the decision to preserve or resect neurovascular bundles at robotic assisted laparoscopic radical prostatectomy. *J Urol* 2014;192:82–88. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24440235>.

202. Pasoglou V, Larbi A, Collette L, et al. One-step TNM staging of high-risk prostate cancer using magnetic resonance imaging (MRI): toward an upfront simplified "all-in-one" imaging approach? *Prostate* 2014;74:469–477. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24375774>.

203. Heck MM, Souvatzoglou M, Retz M, et al. Prospective comparison of computed tomography, diffusion-weighted magnetic resonance imaging and [11C]choline positron emission tomography/computed tomography for preoperative lymph node staging in prostate cancer patients. *Eur J Nucl Med Mol Imaging* 2014;41:694–701. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24297503>.

204. Lecouvet FE, El Mouedden J, Collette L, et al. Can whole-body magnetic resonance imaging with diffusion-weighted imaging replace Tc

99m bone scanning and computed tomography for single-step detection of metastases in patients with high-risk prostate cancer? *Eur Urol* 2012;62:68–75. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22366187>.

205. Rosar F, Dewes S, Ries M, et al. New insights in the paradigm of upregulation of tumoral PSMA expression by androgen receptor blockade: Enzalutamide induces PSMA upregulation in castration-resistant prostate cancer even in patients having previously progressed on enzalutamide. *Eur J Nucl Med Mol Imaging* 2020;47:687–694. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/31901103>.

206. Staniszewska M, Fragoso Costa P, Eiber M, et al. Enzalutamide enhances PSMA expression of PSMA-low prostate cancer. *Int J Mol Sci* 2021;22. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34299051>.

207. Emmett L, Yin C, Crumbaker M, et al. Rapid modulation of PSMA expression by androgen deprivation: serial (68)Ga-PSMA-11 PET in men with hormone-sensitive and castrate-resistant prostate cancer commencing androgen blockade. *J Nucl Med* 2019;60:950–954. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30552200>.

208. Calais J, Ceci F, Eiber M, et al. (18)F-fluciclovine PET-CT and (68)Ga-PSMA-11 PET-CT in patients with early biochemical recurrence after prostatectomy: a prospective, single-centre, single-arm, comparative imaging trial. *Lancet Oncol* 2019;20:1286–1294. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31375469>.

209. Afshar-Oromieh A, Avtzi E, Giesel FL, et al. The diagnostic value of PET/CT imaging with the (68)Ga-labelled PSMA ligand HBED-CC in the diagnosis of recurrent prostate cancer. *Eur J Nucl Med Mol Imaging* 2015;42:197–209. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/25411132>.

210. Eiber M, Maurer T, Souvatzoglou M, et al. Evaluation of hybrid (68)Ga-PSMA ligand PET/CT in 248 patients with biochemical recurrence after radical prostatectomy. *J Nucl Med* 2015;56:668–674. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25791990>.



NCCN Guidelines Version 5.2026 Prostate Cancer

211. Hoffmann MA, Buchholz HG, Wieler HJ, et al. PSA and PSA kinetics thresholds for the presence of (68)Ga-PSMA-11 PET/CT-detectable lesions in patients with biochemical recurrent prostate cancer. *Cancers* (Basel) 2020;12. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32046318>.

212. Dietlein F, Kobe C, Neubauer S, et al. PSA-stratified performance of (18)F- and (68)Ga-PSMA PET in patients with biochemical recurrence of prostate cancer. *J Nucl Med* 2017;58:947–952. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27908968>.

213. Tan N, Oyoyo U, Bavadian N, et al. PSMA-targeted radiotracers versus (18)F-fluciclovine for the detection of prostate cancer biochemical recurrence after definitive therapy: A systematic review and meta-analysis. *Radiology* 2020;296:44–55. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32396045>.

214. Jani AB, Ravizzini GC, Gartrell BA, et al. Diagnostic Performance and Safety of (18)F-rhPSMA-7.3 Positron Emission Tomography in Men With Suspected Prostate Cancer Recurrence: Results From a Phase 3, Prospective, Multicenter Study (SPOTLIGHT). *J Urol* 2023;210:299–311. Available at:

215. Surasi DS, Eiber M, Maurer T, et al. Diagnostic Performance and Safety of Positron Emission Tomography with (18)F-rhPSMA-7.3 in Patients with Newly Diagnosed Unfavourable Intermediate- to Very-high-risk Prostate Cancer: Results from a Phase 3, Prospective, Multicentre Study (LIGHTHOUSE). *Eur Urol* 2023;84:361–370. Available at:

216. Hope TA, Eiber M, Armstrong WR, et al. Diagnostic accuracy of 68Ga-PSMA-11 PET for pelvic nodal metastasis detection prior to radical prostatectomy and pelvic lymph node dissection: A multicenter prospective phase 3 imaging trial. *JAMA Oncol* 2021;7:1635–1642. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34529005>.

217. Fendler WP, Calais J, Eiber M, et al. Assessment of 68Ga-PSMA-11 PET accuracy in localizing recurrent prostate cancer: A prospective single-arm clinical trial. *JAMA Oncol* 2019;5:856–863. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30920593>.

218. Pienta KJ, Gorin MA, Rowe SP, et al. A phase 2/3 prospective multicenter study of the diagnostic accuracy of prostate specific membrane antigen PET/CT with (18)F-DCFPyL in prostate cancer patients (OSPReY). *J Urol* 2021;206:52–61. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33634707>.

219. Jansen BHE, Bodar YJL, Zwezerijnen GJC, et al. Pelvic lymph-node staging with (18)F-DCFPyL PET/CT prior to extended pelvic lymph-node dissection in primary prostate cancer - the SALT trial. *Eur J Nucl Med Mol Imaging* 2021;48:509–520. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32789599>.

220. Morris MJ, Rowe SP, Gorin MA, et al. Diagnostic Performance of (18)F-DCFPyL-PET/CT in Men with Biochemically Recurrent Prostate Cancer: Results from the CONDOR Phase III, Multicenter Study. *Clin Cancer Res* 2021;27:3674–3682. Available at:

221. Fuccio C, Castellucci P, Schiavina R, et al. Role of 11C-choline PET/CT in the re-staging of prostate cancer patients with biochemical relapse and negative results at bone scintigraphy. *Eur J Radiol* 2012;81:e893–896. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22621862>.

222. Nanni C, Schiavina R, Brunocilla E, et al. 18F-fluciclovine PET/CT for the detection of prostate cancer relapse: a comparison to 11C-choline PET/CT. *Clin Nucl Med* 2015;40:e386–391. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26053708>.

223. Evangelista L, Zattoni F, Guttilla A, et al. Choline PET or PET/CT and biochemical relapse of prostate cancer: a systematic review and meta-analysis. *Clin Nucl Med* 2013;38:305–314. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23486334>.

224. Fanti S, Minozzi S, Castellucci P, et al. PET/CT with C-choline for evaluation of prostate cancer patients with biochemical recurrence: meta-analysis and critical review of available data. *Eur J Nucl Med Mol Imaging* 2015;43:55–69. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26450693>.



NCCN Guidelines Version 5.2026 Prostate Cancer

225. Fanti S, Minozzi S, Castellucci P, et al. PET/CT with (11)C-choline for evaluation of prostate cancer patients with biochemical recurrence: meta-analysis and critical review of available data. *Eur J Nucl Med Mol Imaging* 2016;43:55–69. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26450693>.

226. Giovacchini G, Picchio M, Coradeschi E, et al. Predictive factors of [(11)C]choline PET/CT in patients with biochemical failure after radical prostatectomy. *Eur J Nucl Med Mol Imaging* 2010;37:301–309. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/19756592>.

227. Kitajima K, Murphy RC, Nathan MA, et al. Detection of recurrent prostate cancer after radical prostatectomy: comparison of 11C-choline PET/CT with pelvic multiparametric MR imaging with endorectal coil. *J Nucl Med* 2014;55:223–232. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24434294>.

228. Mitchell CR, Lowe VJ, Rangel LJ, et al. Operational characteristics of (11)c-choline positron emission tomography/computerized tomography for prostate cancer with biochemical recurrence after initial treatment. *J Urol* 2013;189:1308–1313. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/23123372>.

229. Nanni C, Zanoni L, Pultrone C, et al. (18)F-FACBC (anti1-amino-3-(18)F-fluorocyclobutane-1-carboxylic acid) versus (11)C-choline PET/CT in prostate cancer relapse: results of a prospective trial. *Eur J Nucl Med Mol Imaging* 2016;43:1601–1610. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26960562>.

230. Reske SN, Blumstein NM, Glatting G. [11C]choline PET/CT imaging in occult local relapse of prostate cancer after radical prostatectomy. *Eur J Nucl Med Mol Imaging* 2008;35:9–17. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/17828534>.

231. Scattoni V, Picchio M, Suardi N, et al. Detection of lymph-node metastases with integrated [11C]choline PET/CT in patients with PSA failure after radical retropubic prostatectomy: results confirmed by open pelvic-retroperitoneal lymphadenectomy. *Eur Urol* 2007;52:423–429. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/17397992>.

232. Umbehr MH, Muntener M, Hany T, et al. The role of 11C-choline and 18F-fluorocholine positron emission tomography (PET) and PET/CT in prostate cancer: a systematic review and meta-analysis. *Eur Urol* 2013;64:106–117. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23628493>.

233. Odewole OA, Tade FI, Nieh PT, et al. Recurrent prostate cancer detection with anti-3-[(18)F]FACBC PET/CT: comparison with CT. *Eur J Nucl Med Mol Imaging* 2016;43:1773–1783. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27091135>.

234. Schuster DM, Nieh PT, Jani AB, et al. Anti-3-[(18)F]FACBC positron emission tomography-computerized tomography and (111)In-capromab pendetide single photon emission computerized tomography-computerized tomography for recurrent prostate carcinoma: results of a prospective clinical trial. *J Urol* 2014;191:1446–1453. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24144687>.

235. Scarsbrook AF, Bottomley D, Teoh EJ, et al. Effect of (18)F-fluciclovine positron emission tomography on the management of patients with recurrence of prostate cancer: Results from the FALCON trial. *Int J Radiat Oncol Biol Phys* 2020;107:316–324. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/32068113>.

236. Andriole GL, Kostakoglu L, Chau A, et al. The impact of positron emission tomography with (18)F-fluciclovine on the management of patients with biochemical recurrence of prostate cancer: Results from the LOCATE trial. *J Urol* 2018;201:322–331. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/30179618>.

237. Wu SY, Boreta L, Shinohara K, et al. Impact of staging (68)Ga-PSMA-11 PET scans on radiation treatment plans in patients with prostate cancer. *Urology* 2019;125:154–162. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/30580002>.

238. Fendler WP, Ferdinandus J, Czernin J, et al. Impact of (68)Ga-PSMA-11 PET on the management of recurrent prostate cancer in a prospective single-arm clinical trial. *J Nucl Med* 2020;61:1793–1799. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/32358094>.



NCCN Guidelines Version 5.2026 Prostate Cancer

239. Zacho HD, Fonager RF, Nielsen JB, et al. Observer agreement and accuracy of (18)F-sodium fluoride PET/CT in the diagnosis of bone metastases in prostate cancer. *J Nucl Med* 2020;61:344–349. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31481577>.

240. Rowe SP, Li X, Trock BJ, et al. Prospective comparison of PET imaging with PSMA-targeted (18)F-DCFPyL versus Na(18)F for bone lesion detection in patients with metastatic prostate cancer. *J Nucl Med* 2020;61:183–188. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31451492>.

241. Rowe SP, Pienta KJ, Pomper MG, Gorin MA. PSMA-RADS version 1.0: A step towards standardizing the interpretation and reporting of PSMA-targeted PET imaging studies. *Eur Urol* 2018;73:485–487. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29132714>.

242. Toriihara A, Nobashi T, Baratto L, et al. Comparison of 3 interpretation criteria for (68)Ga-PSMA11 PET based on inter- and intrareader agreement. *J Nucl Med* 2020;61:533–539. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31562226>.

243. American College of Radiology. ACR Appropriateness Criteria. Available at: <https://www.acr.org/Clinical-Resources/Clinical-Tools-and-Reference/Appropriateness-Criteria>. Accessed October 24, 2025.

244. Walsh L, Shore R, Auvinen A, et al. Risks from CT scans--what do recent studies tell us? *J Radiol Prot* 2014;34:E1–5. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24594968>.

245. American College of Radiology. ACR Manual on Contrast Media. Available at: <https://www.acr.org/Clinical-Resources/Clinical-Tools-and-Reference/Contrast-Manual>. Accessed October 24, 2025.

246. Johansson JE, Holmberg L, Johansson S, et al. Fifteen-year survival in prostate cancer. A prospective, population-based study in Sweden. *JAMA* 1997;277:467–471. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9020270>.

247. Bill-Axelson A, Holmberg L, Garmo H, et al. Radical prostatectomy or watchful waiting in early prostate cancer. *N Engl J Med* 2014;370:932–942. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24597866>.

248. Bill-Axelson A, Holmberg L, Garmo H, et al. Radical prostatectomy or watchful waiting in prostate cancer - 29-year follow-up. *N Engl J Med* 2018;379:2319–2329. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30575473>.

249. Wilt TJ, Jones KM, Barry MJ, et al. Follow-up of prostatectomy versus observation for early prostate cancer. *N Engl J Med* 2017;377:132–142. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28700844>.

250. Sakr WA, Grignon DJ, Crissman JD, et al. High grade prostatic intraepithelial neoplasia (HGPIN) and prostatic adenocarcinoma between the ages of 20-69: an autopsy study of 249 cases. *In Vivo* 1994;8:439–443. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7803731>.

251. Thompson IM, Pauler DK, Goodman PJ, et al. Prevalence of prostate cancer among men with a prostate-specific antigen level < or =4.0 ng per milliliter. *N Engl J Med* 2004;350:2239–2246. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15163773>.

252. Schroder FH, Hugosson J, Roobol MJ, et al. Prostate-cancer mortality at 11 years of follow-up. *N Engl J Med* 2012;366:981–990. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22417251>.

253. Schroder FH, Hugosson J, Roobol MJ, et al. Screening and prostate-cancer mortality in a randomized European study. *N Engl J Med* 2009;360:1320–1328. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19297566>.

254. Klotz L. Active surveillance for prostate cancer: for whom? *J Clin Oncol* 2005;23:8165–8169. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16278468>.

255. Draisma G, Etzioni R, Tsodikov A, et al. Lead time and overdiagnosis in prostate-specific antigen screening: importance of methods and context.



NCCN Guidelines Version 5.2026

Prostate Cancer

J Natl Cancer Inst 2009;101:374–383. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/19276453>.

256. Draisma G, Boer R, Otto SJ, et al. Lead times and overdetection due to prostate-specific antigen screening: estimates from the European Randomized Study of Screening for Prostate Cancer. J Natl Cancer Inst 2003;95:868–878. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/12813170>.

257. Vince RA, Jr., Sun Y, Mahal B, et al. The impact of a statewide active surveillance initiative: A roadmap for increasing active surveillance utilization nationwide. Eur Urol 2023;83:307–310. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/35688664>.

258. Cooperberg MR, Meeks W, Fang R, et al. Time trends and variation in the use of active surveillance for management of low-risk prostate cancer in the US. JAMA Netw Open 2023;6:e231439. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/36862409>.

259. Al Hussein Al Awamlh B, Barocas DA, Zhu A, et al. Use of active surveillance vs definitive treatment among men with low- and favorable intermediate-risk prostate cancer in the US between 2010 and 2018. JAMA Intern Med 2023;183:608–611. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/37010836>.

260. Loeb S, Byrne N, Makarov DV, et al. Use of conservative management for low-risk prostate cancer in the Veterans Affairs Integrated Health Care System from 2005-2015. JAMA 2018;319:2231–2233. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29800017>.

261. Hamdy FC, Donovan JL, Lane JA, et al. 10-Year outcomes after monitoring, surgery, or radiotherapy for localized prostate cancer. N Engl J Med 2016;375:1415–1424. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/27626136>.

262. Neal DE, Metcalfe C, Donovan JL, et al. Ten-year mortality, disease progression, and treatment-related side effects in men with localised prostate cancer from the ProtecT randomised controlled trial according to

treatment received. Eur Urol 2020;77:320–330. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/31771797>.

263. Hamdy FC, Donovan JL, Lane JA, et al. Fifteen-year outcomes after monitoring, surgery, or radiotherapy for prostate cancer. N Engl J Med 2023;388:1547–1558. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/36912538>.

264. Sushentsev N, Warren AY, Colling R, et al. Active monitoring, surgery, and radiotherapy for cribriform-positive and cribriform-negative prostate cancer: A secondary analysis of the PROTECT randomized clinical trial. JAMA Oncol 2025. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/41100113>.

265. Donovan JL, Hamdy FC, Lane JA, et al. Patient-reported outcomes 12 years after localized prostate cancer treatment. NEJM Evid 2023;2:EVIDoa2300018. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/38320051>.

266. Spratt DE. To ProtecT our patients with prostate cancer. JAMA Oncol 2017;3:1461–1462. Available at:
<https://www.ncbi.nlm.nih.gov/pubmed/28301652>.

267. Carter G, Clover K, Britton B, et al. Wellbeing during Active Surveillance for localised prostate cancer: a systematic review of psychological morbidity and quality of life. Cancer Treat Rev 2015;41:46–60. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25467109>.

268. Jeldres C, Cullen J, Hurwitz LM, et al. Prospective quality-of-life outcomes for low-risk prostate cancer: Active surveillance versus radical prostatectomy. Cancer 2015;121:2465–2473. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/25845467>.

269. Parker PA, Davis JW, Latini DM, et al. Relationship between illness uncertainty, anxiety, fear of progression and quality of life in men with favourable-risk prostate cancer undergoing active surveillance. BJU Int 2015;117:469–477. Available at:
<http://www.ncbi.nlm.nih.gov/pubmed/25714186>.



NCCN Guidelines Version 5.2026 Prostate Cancer

270. van den Bergh RC, Essink-Bot ML, Roobol MJ, et al. Anxiety and distress during active surveillance for early prostate cancer. *Cancer* 2009;115:3868–3878. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19637245>.

271. Pham KN, Cullen J, Hurwitz LM, et al. Prospective quality of life in men choosing active surveillance compared to those biopsied but not diagnosed with prostate cancer. *J Urol* 2016;196:392–398. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26976206>.

272. Klotz L, Vesprini D, Sethukavalan P, et al. Long-term follow-up of a large active surveillance cohort of patients with prostate cancer. *J Clin Oncol* 2015;33:272–277. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/25512465>.

273. Loeb S, Folkvaljon Y, Makarov DV, et al. Five-year nationwide follow-up study of active surveillance for prostate cancer. *Eur Urol* 2015;67:233–238. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24993868>.

274. Roemeling S, Roobol MJ, de Vries SH, et al. Active surveillance for prostate cancers detected in three subsequent rounds of a screening trial: characteristics, PSA doubling times, and outcome. *Eur Urol* 2007;51:1244–1250; discussion 1251. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17161520>.

275. Tosoian JJ, Mamawala M, Epstein JI, et al. Intermediate and longer-term outcomes from a prospective active-surveillance program for favorable-risk prostate cancer. *J Clin Oncol* 2015;33:3379–3385. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26324359>.

276. van As NJ, Norman AR, Thomas K, et al. Predicting the probability of deferred radical treatment for localised prostate cancer managed by active surveillance. *Eur Urol* 2008;54:1297–1305. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18342430>.

277. Cooley LF, Emeka AA, Meyers TJ, et al. Factors Associated with Time to Conversion from Active Surveillance to Treatment for Prostate Cancer in a Multi-Institutional Cohort. *J Urol* 2021;206:1147–1156. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34503355>.

278. Bokhorst LP, Valdagni R, Rannikko A, et al. A decade of active surveillance in the PRIAS study: An update and evaluation of the criteria used to recommend a switch to active treatment. *Eur Urol* 2016;70:954–960. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27329565>.

279. Newcomb LF, Thompson IM, Jr., Boyer HD, et al. Outcomes of active surveillance for the management of clinically localized prostate cancer in the prospective, multi-institutional Canary PASS cohort. *J Urol* 2015;195:313–320. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/26327354>.

280. Welty CJ, Cowan JE, Nguyen H, et al. Extended followup and risk factors for disease reclassification in a large active surveillance cohort for localized prostate cancer. *J Urol* 2015;193:807–811. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/25261803>.

281. Dall'Era MA, Konety BR, Cowan JE, et al. Active surveillance for the management of prostate cancer in a contemporary cohort. *Cancer* 2008;112:2664–2670. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18433013>.

282. Maggi M, Cowan JE, Fasulo V, et al. The long-term risks of metastases in men on active surveillance for early stage prostate cCancer. *J Urol* 2020;204:1222–1228. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/33157570>.

283. Simpkin AJ, Tilling K, Martin RM, et al. Systematic review and meta-analysis of factors determining change to radical treatment in active surveillance for localized prostate cancer. *Eur Urol* 2015;67:993–1005. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25616709>.

284. Spratt DE, Schaeffer EM. Revisiting prostate cancer active surveillance candidacy. *Lancet Oncol* 2025;26:1273–1275. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/41038181>.

285. Ng SP, Duchesne G, Tai KH, et al. Support for the use of objective comorbidity indices in the assessment of noncancer death risk in prostate cancer patients. *Prostate Int* 2017;5:8–12. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28352617>.



NCCN Guidelines Version 5.2026

Prostate Cancer

286. Cooperberg MR, Zheng Y, Faino AV, et al. Tailoring Intensity of Active Surveillance for Low-Risk Prostate Cancer Based on Individualized Prediction of Risk Stability. *JAMA Oncol* 2020;6:e203187. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32852532>.

287. Loneragan PE, Washington SL, 3rd, Cowan JE, et al. Risk factors for biopsy reclassification over time in men on active surveillance for early stage prostate cancer. *J Urol* 2020;204:1216–1221. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32519915>.

288. Musunuru HB, Yamamoto T, Klotz L, et al. Active surveillance for intermediate risk prostate cancer: Survival outcomes in the Sunnybrook experience. *J Urol* 2016;196:1651–1658. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27569437>.

289. Newcomb LF, Schenk JM, Zheng Y, et al. Long-term outcomes in patients using protocol-directed active surveillance for prostate cancer. *JAMA* 2024;331:2084–2093. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/38814624>.

290. Patel HD, Tosoian JJ, Carter HB, Epstein JI. Adverse pathologic findings for men electing immediate radical prostatectomy: Defining a favorable intermediate-risk group. *JAMA Oncol* 2018;4:89–92. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28715578>.

291. Gearman DJ, Morlacco A, Cheville JC, et al. Comparison of pathological and oncologic outcomes of favorable risk Gleason score 3 + 4 and low risk Gleason score 6 prostate cancer: Considerations for active surveillance. *J Urol* 2018;199:1188–1195. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29225057>.

292. Aghazadeh MA, Frankel J, Belanger M, et al. National Comprehensive Cancer Network(R) favorable intermediate risk prostate cancer-Is active surveillance appropriate? *J Urol* 2018;199:1196–1201. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29288120>.

293. Loeb S, Folkvaljon Y, Bratt O, et al. Defining intermediate-risk prostate cancer suitable for active surveillance. *J Urol* 2018;201:292–299. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30240688>.

294. Ginsburg KB, Jacobs JC, Qi J, et al. Impact of early confirmatory tests on upgrading and conversion to treatment in prostate cancer patients on active surveillance. *Urology* 2020;147:213–222. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32946908>.

295. Kornberg Z, Cowan JE, Westphalen AC, et al. Genomic prostate score, PI-RADS version 2 and progression in men with prostate cancer on active surveillance. *J Urol* 2019;201:300–307. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30179620>.

296. Dickinson L, Ahmed HU, Allen C, et al. Magnetic resonance imaging for the detection, localisation, and characterisation of prostate cancer: recommendations from a European consensus meeting. *Eur Urol* 2011;59:477–494. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21195536>.

297. Gallagher KM, Christopher E, Cameron AJ, et al. Four-year outcomes from a multiparametric magnetic resonance imaging (MRI)-based active surveillance programme: PSA dynamics and serial MRI scans allow omission of protocol biopsies. *BJU Int* 2018;123:429–438. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30113755>.

298. Cantiello F, Russo GI, Kaufmann S, et al. Role of multiparametric magnetic resonance imaging for patients under active surveillance for prostate cancer: a systematic review with diagnostic meta-analysis. *Prostate Cancer Prostatic Dis* 2018;22:206–220. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30487646>.

299. Liss MA, Newcomb LF, Zheng Y, et al. Magnetic resonance imaging for the detection of high grade cancer in the Canary Prostate Active Surveillance Study. *J Urol* 2020;204:701–706. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32343189>.

300. Chu CE, Loneragan PE, Washington SL, et al. Multiparametric magnetic resonance imaging alone is insufficient to detect grade reclassification in active surveillance for prostate cancer. *Eur Urol* 2020;78:515–517. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32631744>.



NCCN Guidelines Version 5.2026

Prostate Cancer

301. Bonekamp D, Bonekamp S, Mullins JK, et al. Multiparametric magnetic resonance imaging characterization of prostate lesions in the active surveillance population: incremental value of magnetic resonance imaging for prediction of disease reclassification. *J Comput Assist Tomogr* 2013;37:948–956. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24270118>.

302. Mullins JK, Bonekamp D, Landis P, et al. Multiparametric magnetic resonance imaging findings in men with low-risk prostate cancer followed using active surveillance. *BJU Int* 2013;111:1037–1045. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23464904>.

303. Nassiri N, Margolis DJ, Natarajan S, et al. Targeted biopsy to detect Gleason score upgrading during active surveillance for men with low versus intermediate risk prostate cancer. *J Urol* 2016;197:632–639. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27639713>.

304. Ma TM, Tosoian JJ, Schaeffer EM, et al. The role of multiparametric magnetic resonance imaging/ultrasound fusion biopsy in active surveillance. *Eur Urol* 2017;71:174–180. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27236496>.

305. Recabal P, Assel M, Sjöberg DD, et al. The efficacy of multiparametric magnetic resonance imaging and magnetic resonance imaging targeted biopsy in risk classification for patients with prostate cancer on active surveillance. *J Urol* 2016;196:374–381. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26920465>.

306. Tran GN, Leapman MS, Nguyen HG, et al. Magnetic resonance imaging-ultrasound fusion biopsy during prostate cancer active surveillance. *Eur Urol* 2016;72:275–281. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27595378>.

307. Klotz L. Point: active surveillance for favorable risk prostate cancer. *J Natl Compr Canc Netw* 2007;5:693–698. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17692173>.

308. Nguyen CT, Lew A, Pettit NN, et al. Microbiology of infection-related complications after transrectal ultrasound-guided prostate biopsy. *Can*

Urol Assoc J 2024;18:E204–E211. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/38466864>.

309. Bryant RJ, Marian IR, Williams R, et al. Local anaesthetic transperineal biopsy versus transrectal prostate biopsy in prostate cancer detection (TRANSLATE): a multicentre, randomised, controlled trial. *Lancet Oncol* 2025;26:583–595. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/40139210>.

310. Mian BM, Feustel PJ, Aziz A, et al. Complications following transrectal and transperineal prostate biopsy: Results of the ProBE-PC randomized clinical trial. *J Urol* 2024;211:205–213. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37976319>.

311. Dall'Era MA, Albertsen PC, Bangma C, et al. Active surveillance for prostate cancer: a systematic review of the literature. *Eur Urol* 2012;62:976–983. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22698574>.

312. Carter HB, Kettermann A, Warlick C, et al. Expectant management of prostate cancer with curative intent: an update of the Johns Hopkins experience. *J Urol* 2007;178:2359–2364; discussion 2364–2355. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17936806>.

313. Klotz L, Zhang L, Lam A, et al. Clinical results of long-term follow-up of a large, active surveillance cohort with localized prostate cancer. *J Clin Oncol* 2010;28:126–131. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19917860>.

314. Sheridan TB, Carter HB, Wang W, et al. Change in prostate cancer grade over time in men followed expectantly for stage T1c disease. *J Urol* 2008;179:901–904; discussion 904–905. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18207195>.

315. Tosoian JJ, Trock BJ, Landis P, et al. Active surveillance program for prostate cancer: an update of the Johns Hopkins experience. *J Clin Oncol* 2011;29:2185–2190. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21464416>.



NCCN Guidelines Version 5.2026 Prostate Cancer

316. Loblaw A, Zhang L, Lam A, et al. Comparing prostate specific antigen triggers for intervention in men with stable prostate cancer on active surveillance. J Urol 2010;184:1942–1946. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20846681>.

317. Ross AE, Loeb S, Landis P, et al. Prostate-specific antigen kinetics during follow-up are an unreliable trigger for intervention in a prostate cancer surveillance program. J Clin Oncol 2010;28:2810–2816. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20439642>.

318. Jain S, Loblaw A, Vesprini D, et al. Gleason upgrading with time in a large prostate cancer active surveillance cohort. J Urol 2015;194:79–84. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25660208>.

319. Yamamoto T, Musunuru B, Vesprini D, et al. Metastatic prostate cancer in men initially treated with active surveillance. J Urol 2016;195:1409–1414. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26707510>.

320. Tosoian JJ, Sundi D, Trock BJ, et al. Pathologic outcomes in favorable-risk prostate cancer: comparative analysis of men electing active surveillance and immediate surgery. Eur Urol 2015;69:576–581. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26456680>.

321. Dall'Era MA, Cowan JE, Simko J, et al. Surgical management after active surveillance for low-risk prostate cancer: pathological outcomes compared with men undergoing immediate treatment. BJU Int 2011;107:1232–1237. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20804478>.

322. Filippou P, Welty CJ, Cowan JE, et al. Immediate versus delayed radical prostatectomy: updated outcomes following active surveillance of prostate cancer. Eur Urol 2015;68:458–463. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26138041>.

323. Bill-Axelson A, Holmberg L, Filen F, et al. Radical prostatectomy versus watchful waiting in localized prostate cancer: the Scandinavian prostate cancer group-4 randomized trial. J Natl Cancer Inst

2008;100:1144–1154. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18695132>.

324. Holmberg L, Garmo H, Andersson SO, et al. Radical prostatectomy or watchful waiting in early prostate cancer. N Engl J Med 2024;391:1362–1364. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39383464>.

325. Eastham JA, Heller G, Halabi S, et al. Cancer and Leukemia Group B 90203 (Alliance): Radical prostatectomy with or without neoadjuvant chemohormonal therapy in localized, high-risk prostate cancer. J Clin Oncol 2020;38:3042–3050. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32706639>.

326. Gongora M, Stranne J, Johansson E, et al. Characteristics of patients in SPCG-15-a randomized trial comparing radical prostatectomy with primary radiotherapy plus androgen deprivation therapy in men with locally advanced prostate cancer. Eur Urol Open Sci 2022;41:63–73. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35813256>.

327. Chade DC, Eastham J, Graefen M, et al. Cancer control and functional outcomes of salvage radical prostatectomy for radiation-recurrent prostate cancer: a systematic review of the literature. Eur Urol 2012;61:961–971. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22280856>.

328. Shekarri B, Upadhyay J, Pontes JE. Salvage radical prostatectomy. Urol Clin North Am 2001;28:545–553. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11590813>.

329. Klein EA, Bianco FJ, Serio AM, et al. Surgeon experience is strongly associated with biochemical recurrence after radical prostatectomy for all preoperative risk categories. J Urol 2008;179:2212–2216; discussion 2216–2217. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18423716>.

330. Begg CB, Riedel ER, Bach PB, et al. Variations in morbidity after radical prostatectomy. N Engl J Med 2002;346:1138–1144. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11948274>.



NCCN Guidelines Version 5.2026 Prostate Cancer

331. Herrell SD, Smith JA, Jr. Robotic-assisted laparoscopic prostatectomy: what is the learning curve? *Urology* 2005;66:105–107. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16194715>.

332. Smith JA, Jr., Herrell SD. Robotic-assisted laparoscopic prostatectomy: do minimally invasive approaches offer significant advantages? *J Clin Oncol* 2005;23:8170–8175. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16278469>.

333. Ilic D, Evans SM, Allan CA, et al. Laparoscopic and robotic-assisted versus open radical prostatectomy for the treatment of localised prostate cancer. *Cochrane Database Syst Rev* 2017;9:CD009625. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28895658>.

334. Hu JC, Gu X, Lipsitz SR, et al. Comparative effectiveness of minimally invasive vs open radical prostatectomy. *JAMA* 2009;302:1557–1564. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19826025>.

335. Gandaglia G, Sammon JD, Chang SL, et al. Comparative effectiveness of robot-assisted and open radical prostatectomy in the postdissemination era. *J Clin Oncol* 2014;32:1419–1426. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24733797>.

336. Parsons JK, Bennett JL. Outcomes of retropubic, laparoscopic, and robotic-assisted prostatectomy. *Urology* 2008;72:412–416. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18267330>.

337. Ficarra V, Novara G, Rosen RC, et al. Systematic review and meta-analysis of studies reporting urinary continence recovery after robot-assisted radical prostatectomy. *Eur Urol* 2012;62:405–417. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22749852>.

338. Ficarra V, Novara G, Ahlering TE, et al. Systematic review and meta-analysis of studies reporting potency rates after robot-assisted radical prostatectomy. *Eur Urol* 2012;62:418–430. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22749850>.

339. Coughlin GD, Yaxley JW, Chambers SK, et al. Robot-assisted laparoscopic prostatectomy versus open radical retropubic prostatectomy:

24-month outcomes from a randomised controlled study. *Lancet Oncol* 2018;19:1051–1060. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30017351>.

340. Yaxley JW, Coughlin GD, Chambers SK, et al. Robot-assisted laparoscopic prostatectomy versus open radical retropubic prostatectomy: early outcomes from a randomised controlled phase 3 study. *Lancet* 2016;388:1057–1066. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27474375>.

341. Freire MP, Weinberg AC, Lei Y, et al. Anatomic bladder neck preservation during robotic-assisted laparoscopic radical prostatectomy: description of technique and outcomes. *Eur Urol* 2009;56:972–980. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19781848>.

342. Abel EJ, Masterson TA, Warner JN, et al. Nerve-sparing prostatectomy and urinary function: a prospective analysis using validated quality-of-life measures. *Urology* 2009;73:1336–1340. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19362347>.

343. Avulova S, Zhao Z, Lee D, et al. The effect of nerve sparing status on sexual and urinary function: 3-year results from the CEASAR study. *J Urol* 2018;199:1202–1209. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29253578>.

344. Davis JW, Chang DW, Chevray P, et al. Randomized phase II trial evaluation of erectile function after attempted unilateral cavernous nerve-sparing retropubic radical prostatectomy with versus without unilateral sural nerve grafting for clinically localized prostate cancer. *Eur Urol* 2009;55:1135–1143. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18783876>.

345. van As N, Yasar B, Griffin C, et al. Radical prostatectomy versus stereotactic radiotherapy for clinically localised prostate cancer: Results of the PACE-A randomised trial. *Eur Urol* 2024;86:566–576. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39266383>.



NCCN Guidelines Version 5.2026

Prostate Cancer

346. Briganti A, Blute ML, Eastham JH, et al. Pelvic lymph node dissection in prostate cancer. *Eur Urol* 2009;55:1251–1265. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19297079>.

347. Heidenreich A, Ohlmann CH, Polyakov S. Anatomical extent of pelvic lymphadenectomy in patients undergoing radical prostatectomy. *Eur Urol* 2007;52:29–37. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17448592>.

348. Masterson TA, Bianco FJ, Jr., Vickers AJ, et al. The association between total and positive lymph node counts, and disease progression in clinically localized prostate cancer. *J Urol* 2006;175:1320–1324; discussion 1324–1325. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16515989>.

349. Joslyn SA, Konety BR. Impact of extent of lymphadenectomy on survival after radical prostatectomy for prostate cancer. *Urology* 2006;68:121–125. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16806432>.

350. Allaf ME, Palapattu GS, Trock BJ, et al. Anatomical extent of lymph node dissection: impact on men with clinically localized prostate cancer. *J Urol* 2004;172:1840–1844. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15540734>.

351. Bader P, Burkhard FC, Markwalder R, Studer UE. Disease progression and survival of patients with positive lymph nodes after radical prostatectomy. Is there a chance of cure? *J Urol* 2003;169:849–854. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12576797>.

352. Daneshmand S, Quek ML, Stein JP, et al. Prognosis of patients with lymph node positive prostate cancer following radical prostatectomy: long-term results. *J Urol* 2004;172:2252–2255. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15538242>.

353. Wagner M, Sokoloff M, Daneshmand S. The role of pelvic lymphadenectomy for prostate cancer--therapeutic? *J Urol* 2008;179:408–413. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18076938>.

354. Fossati N, Willemse PM, van den Bergh RC, et al. The benefits and harms of different extents of lymph node dissection during radical prostatectomy for prostate cancer: a systematic review. *Eur Urol* 2017;72:84–109. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28126351>.

355. Efstathiou JA, Yeap BY, Michalski JM, et al. Prostate advanced radiation technologies investigating quality of life (PARTIQoL): Phase III randomized clinical trial of proton therapy vs. IMRT for localized prostate cancer [abstract]. *International Journal of Radiation Oncology, Biology, Physics* 2024;120:S1. Available at: <https://doi.org/10.1016/j.ijrobp.2024.08.012>.

356. Yu JB, Soulos PR, Herrin J, et al. Proton versus intensity-modulated radiotherapy for prostate cancer: patterns of care and early toxicity. *J Natl Cancer Inst* 2013;105:25–32. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23243199>.

357. Zelefsky MJ, Kollmeier M, Cox B, et al. Improved clinical outcomes with high-dose image guided radiotherapy compared with non-IGRT for the treatment of clinically localized prostate cancer. *Int J Radiat Oncol Biol Phys* 2012;84:125–129. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22330997>.

358. Kishan AU, Ma TM, Lamb JM, et al. Magnetic resonance imaging-guided vs computed tomography-guided stereotactic body radiotherapy for prostate cancer: The MIRAGE randomized clinical trial. *JAMA Oncol* 2023;9:365–373. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36633877>.

359. Pollack A, Walker G, Horwitz EM, et al. Randomized trial of hypofractionated external-beam radiotherapy for prostate cancer. *J Clin Oncol* 2013;31:3860–3868. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24101042>.

360. Arcangeli S, Strigari L, Gomellini S, et al. Updated results and patterns of failure in a randomized hypofractionation trial for high-risk prostate cancer. *Int J Radiat Oncol Biol Phys* 2012;84:1172–1178. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22537541>.



NCCN Guidelines Version 5.2026 Prostate Cancer

361. Arcangeli G, Saracino B, Arcangeli S, et al. Moderate Hypofractionation In High-Risk, Organ-Confined Prostate Cancer: Final Results Of A Phase III randomized trial. J Clin Oncol 2017;35:1891–1897. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28355113>.

362. Incrocci L, Wortel RC, Alemayehu WG, et al. Hypofractionated versus conventionally fractionated radiotherapy for patients with localised prostate cancer (HYPRO): final efficacy results from a randomised, multicentre, open-label, phase 3 trial. Lancet Oncol 2016;17:1061–1069. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27339116>.

363. Dearnaley D, Syndikus I, Mossop H, et al. Conventional versus hypofractionated high-dose intensity-modulated radiotherapy for prostate cancer: 5-year outcomes of the randomised, non-inferiority, phase 3 CHHiP trial. Lancet Oncol 2016;17:1047–1060. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27339115>.

364. Aluwini S, Pos F, Schimmel E, et al. Hypofractionated versus conventionally fractionated radiotherapy for patients with prostate cancer (HYPRO): acute toxicity results from a randomised non-inferiority phase 3 trial. Lancet Oncol 2015;16:274–283. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25656287>.

365. Lee WR, Dignam JJ, Amin MB, et al. Randomized phase III noninferiority study comparing two radiotherapy fractionation schedules in patients with low-risk prostate cancer. J Clin Oncol 2016;34:2325–2332. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27044935>.

366. Catton CN, Lukka H, Gu CS, et al. Randomized trial of a hypofractionated radiation regimen for the treatment of localized prostate cancer. J Clin Oncol 2017;35:1884–1890. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28296582>.

367. Hoffman KE, Voong KR, Levy LB, et al. Randomized trial of hypofractionated, dose-escalated, intensity-modulated radiation therapy (IMRT) versus conventionally fractionated IMRT for localized prostate cancer. J Clin Oncol 2018;36:2943–2949. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30106637>.

368. Bruner DW, Pugh SL, Lee WR, et al. Quality of life in patients with low-risk prostate cancer treated with hypofractionated vs conventional radiotherapy: A phase 3 randomized clinical trial. JAMA Oncol 2019;5:664–670. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30763425>.

369. Kishan AU, Sun Y, Tree AC, et al. Hypofractionated radiotherapy for prostate cancer (HYDRA): an individual patient data meta-analysis of randomised trials in the MARCAP consortium. Lancet Oncol 2025;26:459–469. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/40112848>.

370. Dasu A. Is the alpha/beta value for prostate tumours low enough to be safely used in clinical trials? Clin Oncol (R Coll Radiol) 2007;19:289–301. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17517328>.

371. Buyyounouski MK, Price RA, Jr., Harris EE, et al. Stereotactic body radiotherapy for primary management of early-stage, low- to intermediate-risk prostate cancer: report of the American Society for Therapeutic Radiology and Oncology Emerging Technology Committee. Int J Radiat Oncol Biol Phys 2010;76:1297–1304. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20338473>.

372. Jackson WC, Silva J, Hartman HE, et al. Stereotactic Body Radiation Therapy for Localized Prostate Cancer: A Systematic Review and Meta-Analysis of Over 6,000 Patients Treated On Prospective Studies. Int J Radiat Oncol Biol Phys 2019;104:778–789. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30959121>.

373. Widmark A, Gunnlaugsson A, Beckman L, et al. Ultra-hypofractionated versus conventionally fractionated radiotherapy for prostate cancer: 5-year outcomes of the HYPO-RT-PC randomised, non-inferiority, phase 3 trial. Lancet 2019;394:385–395. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31227373>.

374. van As N, Griffin C, Tree A, et al. Phase 3 trial of stereotactic body radiotherapy in localized prostate cancer. N Engl J Med 2024;391:1413–1425. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39413377>.



NCCN Guidelines Version 5.2026 Prostate Cancer

375. Mariados N, Sylvester J, Shah D, et al. Hydrogel spacer prospective multicenter randomized controlled pivotal trial: dosimetric and clinical effects of perirectal spacer application in men undergoing prostate image guided intensity modulated radiation therapy. *Int J Radiat Oncol Biol Phys* 2015;92:971–977. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/26054865>.

376. Miller LE, Efsthathiou JA, Bhattacharyya SK, et al. Association of the placement of a perirectal hydrogel spacer with the clinical outcomes of men receiving radiotherapy for prostate cancer: A systematic review and meta-analysis. *JAMA Netw Open* 2020;3:e208221. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/32585020>.

377. Hamstra DA, Mariados N, Sylvester J, et al. Continued benefit to rectal separation for prostate radiation therapy: final results of a phase III trial. *Int J Radiat Oncol Biol Phys* 2017;97:976–985. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28209443>.

378. Hamstra DA, Mariados N, Sylvester J, et al. Sexual quality of life following prostate intensity modulated radiation therapy (IMRT) with a rectal/prostate spacer: Secondary analysis of a phase 3 trial. *Pract Radiat Oncol* 2018;8:e7–e15. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28951089>.

379. Schorghofer A, Drerup M, Kunit T, et al. Rectum-spacer related acute toxicity - endoscopy results of 403 prostate cancer patients after implantation of gel or balloon spacers. *Radiat Oncol* 2019;14:47. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30876433>.

380. Levy JF, Khairnar R, Louie AV, et al. Evaluating the cost-effectiveness of hydrogel rectal spacer in prostate cancer radiation therapy. *Pract Radiat Oncol* 2019;9:e172–e179. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/30342180>.

381. Zumsteg ZS, Spratt DE, Daskivich TJ, et al. Effect of androgen deprivation on long-term outcomes of intermediate-risk prostate cancer stratified as favorable or unfavorable: A secondary analysis of the RTOG 9408 randomized clinical trial. *JAMA Netw Open* 2020;3:e2015083. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32902647>.

382. Kishan AU, Sun Y, Hartman H, et al. Androgen deprivation therapy use and duration with definitive radiotherapy for localised prostate cancer: an individual patient data meta-analysis. *Lancet Oncol* 2022;23:304–316. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35051385>.

383. Krauss DJ, Karrison T, Martinez AA, et al. Dose-escalated radiotherapy alone or in combination with short-term androgen deprivation for intermediate-risk prostate cancer: Results of a phase III multi-institutional trial. *J Clin Oncol* 2023;41:3203–3216. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/37104748>.

384. Kerkmeijer LGW, Groen VH, Pos FJ, et al. Focal boost to the intraprostatic tumor in external beam radiotherapy for patients with localized prostate cancer: Results from the FLAME randomized phase III trial. *J Clin Oncol* 2021;39:787–796. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/33471548>.

385. Morris WJ, Tyldesley S, Rodda S, et al. Androgen suppression combined with elective nodal and dose escalated radiation therapy (the ASCENDE-RT trial): An analysis of survival endpoints for a randomized trial comparing a low-dose-rate brachytherapy boost to a dose-escalated external beam boost for high- and intermediate-risk prostate cancer. *Int J Radiat Oncol Biol Phys* 2017;98:275–285. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28262473>.

386. Roach M, Moughan J, Lawton CAF, et al. Sequence of hormonal therapy and radiotherapy field size in unfavourable, localised prostate cancer (NRG/RTOG 9413): long-term results of a randomised, phase 3 trial. *Lancet Oncol* 2018;19:1504–1515. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/30316827>.

387. Mason MD, Parulekar WR, Sydes MR, et al. Final report of the Intergroup randomized study of combined androgen-deprivation therapy plus radiotherapy versus androgen-deprivation therapy alone in locally advanced prostate cancer. *J Clin Oncol* 2015;33:2143–2150. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25691677>.

388. Warde P, Mason M, Ding K, et al. Combined androgen deprivation therapy and radiation therapy for locally advanced prostate cancer: a



NCCN Guidelines Version 5.2026

Prostate Cancer

randomised, phase 3 trial. Lancet 2011;378:2104–2111. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22056152>.

389. Widmark A, Klepp O, Solberg A, et al. Endocrine treatment, with or without radiotherapy, in locally advanced prostate cancer (SPCG-7/SFUO-3): an open randomised phase III trial. Lancet 2009;373:301–308. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19091394>.

390. Fossa SD, Wiklund F, Klepp O, et al. Ten- and 15-yr prostate cancer-specific mortality in patients with nonmetastatic locally advanced or aggressive intermediate prostate cancer, randomized to lifelong endocrine treatment alone or combined with radiotherapy: final results of the Scandinavian Prostate Cancer Group-7. Eur Urol 2016;70:684–691. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27025586>.

391. Pommier P, Chabaud S, Lagrange JL, et al. Is there a role for pelvic irradiation in localized prostate adenocarcinoma? Preliminary results of GETUG-01. J Clin Oncol 2007;25:5366–5373. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18048817>.

392. Roach M, T.G. Karrison, H.T. Chung, et al. Androgen-deprivation therapy and high-dose definitive radiotherapy ± whole-pelvic RT in patients with unfavorable intermediate or favorable high-risk prostate cancer: early results of a phase III randomized controlled trial [abstract]. 2025:ASTRO LBA 05. Available at: https://www.astro.org/ASTRO/media/ASTRO/Meetings%20and%20Education/PDFs/AM25/AM25_LBAs.pdf.

393. Murthy V, Maitre P, Kannan S, et al. Prostate-Only Versus Whole-Pelvic Radiation Therapy in High-Risk and Very High-Risk Prostate Cancer (POP-RT): Outcomes From Phase III Randomized Controlled Trial. J Clin Oncol 2021;39:1234–1242. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33497252>.

394. Attard G, Murphy L, Clarke NW, et al. Abiraterone acetate and prednisolone with or without enzalutamide for high-risk non-metastatic prostate cancer: a meta-analysis of primary results from two randomised controlled phase 3 trials of the STAMPEDE platform protocol. Lancet

2022;399:447–460. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34953525>.

395. Nguyen PL, Sweeney C, Stockler MR, et al. Randomised phase III trial of androgen deprivation therapy (ADT) with radiation therapy with or without enzalutamide for high risk, clinically localised prostate cancer: ENZARAD (ANZUP 1303) [abstract]. ESMA 2025:LBA86. Available at: <https://cslide.ctimeetingtech.com/esmo2025/attendee/confcal/show/session/119>.

396. Brachman DG, Thomas T, Hilbe J, Beyer DC. Failure-free survival following brachytherapy alone or external beam irradiation alone for T1-2 prostate tumors in 2222 patients: results from a single practice. Int J Radiat Oncol Biol Phys 2000;48:111–117. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10924979>.

397. Masson S, Persad R, Bahl A. HDR brachytherapy in the management of high-risk prostate cancer. Adv Urol 2012;2012:980841. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22461791>.

398. Nag S, Bice W, DeWyngaert K, et al. The American Brachytherapy Society recommendations for permanent prostate brachytherapy postimplant dosimetric analysis. Int J Radiat Oncol Biol Phys 2000;46:221–230. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10656396>.

399. Hoskin P. High dose rate brachytherapy for prostate cancer. Cancer Radiother 2008;12:512–514. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18755623>.

400. Grills IS, Martinez AA, Hollander M, et al. High dose rate brachytherapy as prostate cancer monotherapy reduces toxicity compared to low dose rate palladium seeds. J Urol 2004;171:1098–1104. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14767279>.

401. Vargas C, Ghilezan M, Hollander M, et al. A new model using number of needles and androgen deprivation to predict chronic urinary toxicity for high or low dose rate prostate brachytherapy. J Urol



NCCN Guidelines Version 5.2026

Prostate Cancer

2005;174:882–887. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16093980>.

402. Hudson JM, Loblaw A, McGuffin M, et al. Prostate high dose-rate brachytherapy as monotherapy for low and intermediate-risk prostate cancer: Efficacy results from a randomized phase II clinical trial of one fraction of 19 Gy or two fractions of 13.5 Gy: A 9-year update.

Radiotherapy and Oncology 2024;198:110381. Available at:

<https://pubmed.ncbi.nlm.nih.gov/38879130/>.

403. Loblaw A, Glicksman R, Helou JA, et al. Androgen suppression combined with elective nodal irradiation and dose escalated prostate treatment: A non-inferiority, phase III randomized controlled trial of stereotactic body radiation therapy versus brachytherapy boost in patients with unfavourable risk localized prostate cancer (ASCENDE-SBRT; CCTG PR24; NCT06235697) [abstract]. J Clin Oncol 2025;43:TPS5132–TPS5132. Available at:

https://doi.org/10.1200/JCO.2025.43.16_suppl.TPS5132.

404. Sathya JR, Davis IR, Julian JA, et al. Randomized trial comparing iridium implant plus external-beam radiation therapy with external-beam radiation therapy alone in node-negative locally advanced cancer of the prostate. J Clin Oncol 2005;23:1192–1199. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15718316>.

405. Hoskin PJ, Motohashi K, Bownes P, et al. High dose rate brachytherapy in combination with external beam radiotherapy in the radical treatment of prostate cancer: initial results of a randomised phase three trial. Radiother Oncol 2007;84:114–120. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17531335>.

406. Hoskin PJ, Rojas AM, Bownes PJ, et al. Randomised trial of external beam radiotherapy alone or combined with high-dose-rate brachytherapy boost for localised prostate cancer. Radiother Oncol 2012;103:217–222.

Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22341794>.

407. Rodda S, Tyldesley S, Morris WJ, et al. Ascende-rt: An analysis of treatment-related morbidity for a randomized trial comparing a low-dose-rate brachytherapy boost with a dose-escalated external beam boost for

high- and intermediate-risk prostate cancer. Int J Radiat Oncol Biol Phys 2017;98:286–295. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28433432>.

408. Rodda S, Morris WJ, Hamm J, Duncan G. ASCENDE-RT: An analysis of health-related quality of life for a randomized trial comparing low-dose-rate brachytherapy boost with dose-escalated external beam boost for high- and intermediate-risk prostate cancer. Int J Radiat Oncol Biol Phys 2017;98:581–589. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28581398>.

409. Spratt DE, Carroll PR. Optimal radical therapy for localized prostate cancer: Recreation of the self-fulfilling prophecy with combination brachytherapy? J Clin Oncol 2018;36:2914–2917. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29782208>.

410. Joseph D, Denham JW, Steigler A, et al. Radiation dose escalation or longer androgen suppression to prevent distant progression in men with locally advanced prostate cancer: 10-year data from the TROG 03.04 RADAR trial. Int J Radiat Oncol Biol Phys 2020;106:693–702. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/32092343>.

411. Yorozu A, Namiki M, Saito S, et al. Trimodality therapy with iodine-125 brachytherapy, external beam radiation therapy, and short- or long-term androgen deprivation therapy for high-risk localized prostate cancer: Results of a multicenter, randomized phase 3 trial (TRIP/TRIGU0907). Int J Radiat Oncol Biol Phys 2024;118:390–401. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/37802225>.

412. Yamada Y, Kollmeier MA, Pei X, et al. A Phase II study of salvage high-dose-rate brachytherapy for the treatment of locally recurrent prostate cancer after definitive external beam radiotherapy. Brachytherapy 2014;13:111–116. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24373762>.

413. Crook JM, Zhang P, Pisansky TM, et al. A prospective phase II trial of trans-perineal ultrasound-guided brachytherapy for locally recurrent prostate cancer after external beam radiotherapy (NRG Oncology/RTOG -



NCCN Guidelines Version 5.2026 Prostate Cancer

0526). *Int J Radiat Oncol Biol Phys* 2018;103:335–343. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30312717>.

414. Konski A, James J, Hartsell W, et al. Economic analysis of radiation therapy oncology group 97-14: multiple versus single fraction radiation treatment of patients with bone metastases. *Am J Clin Oncol* 2009;32:423–428. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19546803>.

415. Hartsell WF, Scott CB, Bruner DW, et al. Randomized trial of short-versus long-course radiotherapy for palliation of painful bone metastases. *J Natl Cancer Inst* 2005;97:798–804. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15928300>.

416. Chow E, van der Linden YM, Roos D, et al. Single versus multiple fractions of repeat radiation for painful bone metastases: a randomised, controlled, non-inferiority trial. *Lancet Oncol* 2014;15:164–171. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24369114>.

417. Hoskin PJ, Hopkins K, Misra V, et al. Effect of single-fraction vs multifraction radiotherapy on ambulatory status among patients with spinal canal compression from metastatic cancer: The SCORAD randomized clinical trial. *JAMA* 2019;322:2084–2094. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31794625>.

418. Roy S, Sun Y, Eastham JA, et al. Radiotherapy- versus surgery-based treatment strategy in high-risk prostate cancer. *Eur Urol Oncol* 2025. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/40683825>.

419. Barocas DA, Alvarez J, Resnick MJ, et al. Association between radiation therapy, surgery, or observation for localized prostate cancer and patient-reported outcomes after 3 years. *JAMA* 2017;317:1126–1140. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28324093>.

420. Lardas M, Liew M, van den Bergh RC, et al. Quality of life outcomes after primary treatment for clinically localised prostate cancer: A systematic review. *Eur Urol* 2017;72:869–885. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28757301>.

421. Donnelly BJ, Saliken JC, Brasher PM, et al. A randomized trial of external beam radiotherapy versus cryoablation in patients with localized prostate cancer. *Cancer* 2010;116:323–330. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19937954>.

422. Robinson JW, Donnelly BJ, Siever JE, et al. A randomized trial of external beam radiotherapy versus cryoablation in patients with localized prostate cancer: quality of life outcomes. *Cancer* 2009;115:4695–4704. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19691092>.

423. Chin JL, Al-Zahrani AA, Autran-Gomez AM, et al. Extended followup oncologic outcome of randomized trial between cryoablation and external beam therapy for locally advanced prostate cancer (T2c-T3b). *J Urol* 2012;188:1170–1175. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22901586>.

424. Ehdaie B, Tempany CM, Holland F, et al. MRI-guided focused ultrasound focal therapy for patients with intermediate-risk prostate cancer: a phase 2b, multicentre study. *Lancet Oncol* 2022;23:910–918. Available at:

425. Duwe G, Boehm K, Haack M, et al. Single-center, prospective phase 2 trial of high-intensity focused ultrasound (HIFU) in patients with unilateral localized prostate cancer: good functional results but oncologically not as safe as expected. *World J Urol* 2023;41:1293–1299. Available at:

426. Ganzer R, Hadaschik B, Pahernik S, et al. Prospective multicenter phase II study on focal therapy (hemiblation) of the prostate with high intensity focused ultrasound. *J Urol* 2018;199:983–989. Available at:

427. Ghai S, Finelli A, Corr K, et al. Mri-guided focused ultrasound focal therapy for intermediate-risk prostate cancer: Final results from a 2-year phase II clinical trial. *Radiology* 2024;310:e231473. Available at:

428. Klotz L, Pavlovich CP, Chin J, et al. Magnetic resonance imaging-guided transurethral ultrasound ablation of prostate cancer. *J Urol* 2021;205:769–779. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33021440>.



NCCN Guidelines Version 5.2026 Prostate Cancer

429. Mehralivand S, George AK, Hoang AN, et al. MRI-guided focal laser ablation of prostate cancer: a prospective single-arm, single-center trial with 3 years of follow-up. *Diagn Interv Radiol* 2021;27:394–400. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34003127>.

430. Eggener SE, Yousuf A, Watson S, et al. Phase II evaluation of magnetic resonance imaging guided focal laser ablation of prostate cancer. *J Urol* 2016;196:1670–1675. Available at:

431. Pound CR, Partin AW, Eisenberger MA, et al. Natural history of progression after PSA elevation following radical prostatectomy. *JAMA* 1999;281:1591–1597. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10235151>.

432. Smith MR, Saad F, Oudard S, et al. Denosumab and bone metastasis-free survival in men with nonmetastatic castration-resistant prostate cancer: exploratory analyses by baseline prostate-specific antigen doubling time. *J Clin Oncol* 2013;31:3800–3806. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24043751>.

433. Kane CJ, Amling CL, Johnstone PA, et al. Limited value of bone scintigraphy and computed tomography in assessing biochemical failure after radical prostatectomy. *Urology* 2003;61:607–611. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12639656>.

434. Martino P, Scattoni V, Galosi AB, et al. Role of imaging and biopsy to assess local recurrence after definitive treatment for prostate carcinoma (surgery, radiotherapy, cryotherapy, HIFU). *World J Urol* 2011;29:595–605. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21553276>.

435. Dotan ZA, Bianco FJ, Jr., Rabbani F, et al. Pattern of prostate-specific antigen (PSA) failure dictates the probability of a positive bone scan in patients with an increasing PSA after radical prostatectomy. *J Clin Oncol* 2005;23:1962–1968. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15774789>.

436. Cher ML, Bianco FJ, Jr., Lam JS, et al. Limited role of radionuclide bone scintigraphy in patients with prostate specific antigen elevations after

radical prostatectomy. *J Urol* 1998;160:1387–1391. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9751361>.

437. Vale CL, Fisher D, Kneebone A, et al. Adjuvant or early salvage radiotherapy for the treatment of localised and locally advanced prostate cancer: a prospectively planned systematic review and meta-analysis of aggregate data. *Lancet* 2020;396:1422–1431. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33002431>.

438. Parker CC, Petersen PM, Cook AD, et al. Timing of radiotherapy (RT) after radical prostatectomy (RP): long-term outcomes in the RADICALS-RT trial (NCT00541047). *Ann Oncol* 2024;35:656–666. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/38583574>.

439. Parker CC, Clarke NW, Cook AD, et al. Timing of radiotherapy after radical prostatectomy (RADICALS-RT): a randomised, controlled phase 3 trial. *Lancet* 2020;396:1413–1421. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33002429>.

440. Sargos P, Chabaud S, Latorzeff I, et al. Adjuvant radiotherapy versus early salvage radiotherapy plus short-term androgen deprivation therapy in men with localised prostate cancer after radical prostatectomy (GETUG-AFU 17): a randomised, phase 3 trial. *Lancet Oncol* 2020;21:1341–1352. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33002438>.

441. Kneebone A, Fraser-Browne C, Duchesne GM, et al. Adjuvant radiotherapy versus early salvage radiotherapy following radical prostatectomy (TROG 08.03/ANZUP RAVES): a randomised, controlled, phase 3, non-inferiority trial. *Lancet Oncol* 2020;21:1331–1340. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33002437>.

442. Hackman G, Taari K, Tammela TL, et al. Randomised trial of adjuvant radiotherapy following radical prostatectomy versus radical prostatectomy alone in prostate cancer patients with positive margins or extracapsular extension. *Eur Urol* 2019;76:586–595. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31375279>.

443. Sachdev S, Carroll P, Sandler H, et al. Assessment of Postprostatectomy Radiotherapy as Adjuvant or Salvage Therapy in



NCCN Guidelines Version 5.2026

Prostate Cancer

Patients With Prostate Cancer: A Systematic Review. JAMA Oncol 2020;6:1793–1800. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32852528>.

444. Bhindi B, Lokeshwar SD, Klaassen Z, et al. Systematic review and meta-analysis of trials evaluating the role of adjuvant radiation after radical prostatectomy for prostate cancer: Implications for early salvage. Can Urol Assoc J 2020;14:330–336. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32432529>.

445. Renzulli JF, 2nd, Brito J, 3rd, Kim IY, Broccoli I. A meta-analysis on the use of radiotherapy after prostatectomy: adjuvant versus early salvage radiation. Prostate Int 2022;10:80–84. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35510080>.

446. Wenzel M, Burdinski K, Tselis N, et al. Real world comparison of adjuvant vs. salvage radiation therapy on cancer-control outcomes after radical prostatectomy. Strahlenther Onkol 2025. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/40328962>.

447. Kwon YS, Wang W, Srivastava A, et al. Observation with or without late radiotherapy is equivalent to early radiotherapy in high-risk prostate cancer after radical prostatectomy: A SEER-Medicare analysis on trends, survival outcomes, and complications. Prostate Int 2021;9:82–89. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34386450>.

448. Messing EM, Manola J, Yao J, et al. Immediate versus deferred androgen deprivation treatment in patients with node-positive prostate cancer after radical prostatectomy and pelvic lymphadenectomy. Lancet Oncol 2006;7:472–479. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16750497>.

449. Touijer KA, Mazzola CR, Sjoberg DD, et al. Long-term outcomes of patients with lymph node metastasis treated with radical prostatectomy without adjuvant androgen-deprivation therapy. Eur Urol 2014;65:20–25. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23619390>.

450. Gupta M, Patel HD, Schwen ZR, et al. Adjuvant radiation with androgen-deprivation therapy for men with lymph node metastases after

radical prostatectomy: identifying men who benefit. BJU Int 2019;123:252–260. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29626845>.

451. Cotter SE, Chen MH, Moul JW, et al. Salvage radiation in men after prostate-specific antigen failure and the risk of death. Cancer 2011;117:3925–3932. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21437885>.

452. Trock BJ, Han M, Freedland SJ, et al. Prostate cancer-specific survival following salvage radiotherapy vs observation in men with biochemical recurrence after radical prostatectomy. JAMA 2008;299:2760–2769. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18560003>.

453. Pollack A, Karrison TG, Balogh AG, et al. The addition of androgen deprivation therapy and pelvic lymph node treatment to prostate bed salvage radiotherapy (NRG Oncology/RTOG 0534 SPPORT): an international, multicentre, randomised phase 3 trial. Lancet 2022;399:1886–1901. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35569466>.

454. Parker CC, Kynaston H, Cook AD, et al. Duration of androgen deprivation therapy with postoperative radiotherapy for prostate cancer: a comparison of long-course versus short-course androgen deprivation therapy in the RADICALS-HD randomised trial. Lancet 2024;403:2416–2425. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/38763153>.

455. Carrie C, Hasbini A, de Laroche G, et al. Salvage radiotherapy with or without short-term hormone therapy for rising prostate-specific antigen concentration after radical prostatectomy (GETUG-AFU 16): a randomised, multicentre, open-label phase 3 trial. Lancet Oncol 2016;17:747–756. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27160475>.

456. Carrie C, Magne N, Burban-Provost P, et al. Short-term androgen deprivation therapy combined with radiotherapy as salvage treatment after radical prostatectomy for prostate cancer (GETUG-AFU 16): a 112-month follow-up of a phase 3, randomised trial. Lancet Oncol 2019;20:1740–1749. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31629656>.



NCCN Guidelines Version 5.2026 Prostate Cancer

457. Shipley WU, Seiferheld W, Lukka HR, et al. Radiation with or without antiandrogen therapy in recurrent prostate cancer. *N Engl J Med* 2017;376:417–428. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28146658>.

458. Dess RT, Sun Y, Jackson WC, et al. Association of presalvage radiotherapy PSA levels after prostatectomy with outcomes of long-term antiandrogen therapy in men with prostate cancer. *JAMA Oncol* 2020;6:735–743. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/32215583>.

459. Burdett S, Fisher DJ, Tierney JF, et al. Duration of androgen suppression with postoperative radiotherapy (DADSPORT) for nonmetastatic prostate cancer: a collaborative systematic review and meta-analysis of aggregate data. *Eur Urol* 2025;88:277–290. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/40571441>.

460. Koulikov D, Mohler MC, Mehedint DC, et al. Low detectable prostate specific antigen after radical prostatectomy--treat or watch? *J Urol* 2014;192:1390–1396. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24859441>.

461. Shinghal R, Yemoto C, McNeal JE, Brooks JD. Biochemical recurrence without PSA progression characterizes a subset of patients after radical prostatectomy. Prostate-specific antigen. *Urology* 2003;61:380–385. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12597952>.

462. Roach M, 3rd, Hanks G, Thames H, Jr., et al. Defining biochemical failure following radiotherapy with or without hormonal therapy in men with clinically localized prostate cancer: recommendations of the RTOG-ASTRO Phoenix Consensus Conference. *Int J Radiat Oncol Biol Phys* 2006;65:965–974. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16798415>.

463. Ismail M, Ahmed S, Kastner C, Davies J. Salvage cryotherapy for recurrent prostate cancer after radiation failure: a prospective case series of the first 100 patients. *BJU Int* 2007;100:760–764. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17662081>.

464. Ahmed HU, Cathcart P, McCartan N, et al. Focal salvage therapy for localized prostate cancer recurrence after external beam radiotherapy: a pilot study. *Cancer* 2012;118:4148–4155. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/22907704>.

465. Baco E, Gelet A, Crouzet S, et al. Hemi salvage high-intensity focused ultrasound (HIFU) in unilateral radiorecurrent prostate cancer: a prospective two-centre study. *BJU Int* 2014;114:532–540. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/24930692>.

466. Crouzet S, Murat FJ, Pommier P, et al. Locally recurrent prostate cancer after initial radiation therapy: early salvage high-intensity focused ultrasound improves oncologic outcomes. *Radiother Oncol* 2012;105:198–202. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23068708>.

467. Shah TT, Peters M, Kanthabalan A, et al. PSA nadir as a predictive factor for biochemical disease-free survival and overall survival following whole-gland salvage HIFU following radiotherapy failure. *Prostate Cancer Prostatic Dis* 2016;19:311–316. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27431499>.

468. Siddiqui KM, Billia M, Arifin A, et al. Pathological, oncologic and functional outcomes of a prospective registry of salvage high intensity focused ultrasound ablation for radiorecurrent prostate cancer. *J Urol* 2016;197:97–102. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/27422297>.

469. Uddin Ahmed H, Cathcart P, Chalasani V, et al. Whole-gland salvage high-intensity focused ultrasound therapy for localized prostate cancer recurrence after external beam radiation therapy. *Cancer* 2012;118:3071–3078. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22071795>.

470. Blazevski A, Geboers B, Scheltema MJ, et al. Salvage irreversible electroporation for radio-recurrent prostate cancer - the prospective FIRE trial. *BJU Int* 2023;131 Suppl 4:23–31. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/36495482>.

471. Geboers B, Scheltema MJ, Blazevski A, et al. Median 4-year outcomes of salvage irreversible electroporation for localized radio-



NCCN Guidelines Version 5.2026 Prostate Cancer

recurrent prostate cancer. *BJU Int* 2023;131 Suppl 4:14–22. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36594205>.

472. Ingrosso G, Becherini C, Lancia A, et al. Nonsurgical salvage local therapies for radiorecurrent prostate cancer: A systematic review and meta-analysis. *Eur Urol Oncol* 2020;3:183–197. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31411996>.

473. Valle LF, Lehrer EJ, Markovic D, et al. A systematic review and meta-analysis of local salvage therapies after radiotherapy for prostate cancer (MASTER). *Eur Urol* 2021;80:280–292. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33309278>.

474. Crook J, Rodgers JP, Pisansky TM, et al. Salvage low-dose-rate prostate brachytherapy: Clinical outcomes of a phase 2 trial for local recurrence after external beam radiation therapy (NRG Oncology/RTOG 0526). *Int J Radiat Oncol Biol Phys* 2022;112:1115–1122. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34740768>.

475. Bergamin S, Eade T, Kneebone A, et al. Interim results of a prospective prostate-specific membrane antigen-directed focal stereotactic reirradiation trial for locally recurrent prostate cancer. *Int J Radiat Oncol Biol Phys* 2020;108:1172–1178. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32659332>.

476. Pasquier D, Martinage G, Janoray G, et al. Salvage stereotactic body radiation therapy for local prostate cancer recurrence after radiation therapy: A retrospective multicenter study of the GETUG. *Int J Radiat Oncol Biol Phys* 2019;105:727–734. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31344433>.

477. Mohler JL, Halabi S, Ryan ST, et al. Management of recurrent prostate cancer after radiotherapy: long-term results from CALGB 9687 (Alliance), a prospective multi-institutional salvage prostatectomy series. *Prostate Cancer Prostatic Dis* 2018;22:309–316. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30385835>.

478. Freedland SJ, de Almeida Luz M, De Giorgi U, et al. Improved Outcomes with Enzalutamide in Biochemically Recurrent Prostate Cancer.

N Engl J Med 2023;389:1453–1465. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37851874>.

479. Aggarwal R, Heller G, Hillman DW, et al. PRESTO: A Phase III, Open-Label Study of Intensification of Androgen Blockade in Patients With High-Risk Biochemically Relapsed Castration-Sensitive Prostate Cancer (AFT-19). *J Clin Oncol* 2024;42:1114–1123. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/38261983>.

480. Potosky AL, Haque R, Cassidy-Bushrow AE, et al. Effectiveness of primary androgen-deprivation therapy for clinically localized prostate cancer. *J Clin Oncol* 2014;32:1324–1330. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24638009>.

481. Lu-Yao GL, Albertsen PC, Moore DF, et al. Fifteen-year survival outcomes following primary androgen-deprivation therapy for localized prostate cancer. *JAMA Intern Med* 2014;174:1460–1467. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25023796>.

482. Klotz L, O'Callaghan C, Ding K, et al. Nadir testosterone within first year of androgen-deprivation therapy (ADT) predicts for time to castration-resistant progression: a secondary analysis of the PR-7 trial of intermittent versus continuous ADT. *J Clin Oncol* 2015;33:1151–1156. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25732157>.

483. Shore ND, Saad F, Cookson MS, et al. Oral relugolix for androgen-deprivation therapy in advanced prostate cancer. *N Engl J Med* 2020;382:2187–2196. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32469183>.

484. Dearnaley DP, Saltzstein DR, Sylvester JE, et al. The oral gonadotropin-releasing hormone receptor antagonist relugolix as neoadjuvant/adjuvant androgen deprivation therapy to external beam radiotherapy in patients with localised intermediate-risk prostate cancer: A randomised, open-label, parallel-group phase 2 trial. *Eur Urol* 2020;78:184–192. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32273183>.



NCCN Guidelines Version 5.2026 Prostate Cancer

485. Spratt DE, Malone S, Roy S, et al. Prostate radiotherapy with adjuvant androgen deprivation therapy (ADT) improves metastasis-free survival compared to neoadjuvant ADT: An individual patient meta-analysis. *J Clin Oncol* 2021;39:136–144. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33275486>.

486. Bolla M, Van Tienhoven G, Warde P, et al. External irradiation with or without long-term androgen suppression for prostate cancer with high metastatic risk: 10-year results of an EORTC randomised study. *Lancet Oncol* 2010;11:1066–1073. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20933466>.

487. Pilepich MV, Winter K, Lawton CA, et al. Androgen suppression adjuvant to definitive radiotherapy in prostate carcinoma--long-term results of phase III RTOG 85-31. *Int J Radiat Oncol Biol Phys* 2005;61:1285–1290. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15817329>.

488. Jackson WC, Hartman HE, Dess RT, et al. Addition of Androgen-Deprivation Therapy or Brachytherapy Boost to External Beam Radiotherapy for Localized Prostate Cancer: A Network Meta-Analysis of Randomized Trials. *J Clin Oncol* 2020;38:3024–3031. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32396488>.

489. James ND, Spears MR, Clarke NW, et al. Failure-free survival and radiotherapy in patients with newly diagnosed nonmetastatic prostate cancer: data from patients in the control arm of the STAMPEDE trial. *JAMA Oncol* 2016;2:348–357. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26606329>.

490. Labrie F, Dupont A, Belanger A, Lachance R. Flutamide eliminates the risk of disease flare in prostatic cancer patients treated with a luteinizing hormone-releasing hormone agonist. *J Urol* 1987;138:804–806. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/3309363>.

491. Schulze H, Senge T. Influence of different types of antiandrogens on luteinizing hormone-releasing hormone analogue-induced testosterone surge in patients with metastatic carcinoma of the prostate. *J Urol* 1990;144:934–941. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/2144596>.

492. Shaw GL, Wilson P, Cuzick J, et al. International study into the use of intermittent hormone therapy in the treatment of carcinoma of the prostate: a meta-analysis of 1446 patients. *BJU Int* 2007;99:1056–1065. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17346277>.

493. Akakura K, Bruchovsky N, Goldenberg SL, et al. Effects of intermittent androgen suppression on androgen-dependent tumors. Apoptosis and serum prostate-specific antigen. *Cancer* 1993;71:2782–2790. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7682149>.

494. Crook JM, O'Callaghan CJ, Duncan G, et al. Intermittent androgen suppression for rising PSA level after radiotherapy. *N Engl J Med* 2012;367:895–903. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22931259>.

495. Schulman C, Cornel E, Matveev V, et al. Intermittent Versus Continuous Androgen Deprivation Therapy in Patients with Relapsing or Locally Advanced Prostate Cancer: A Phase 3b Randomised Study (ICELAND). *Eur Urol* 2016;69:720–727. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26520703>.

496. Dong Z, Wang H, Xu M, et al. Intermittent hormone therapy versus continuous hormone therapy for locally advanced prostate cancer: a meta-analysis. *Aging Male* 2015;18:233–237. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26225795>.

497. Hussain M, Tangen CM, Berry DL, et al. Intermittent versus continuous androgen deprivation in prostate cancer. *N Engl J Med* 2013;368:1314–1325. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23550669>.

498. Hershman DL, Unger JM, Wright JD, et al. Adverse health events following intermittent and continuous androgen deprivation in patients with metastatic prostate cancer. *JAMA Oncol* 2016;2:453–461. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26720308>.

499. Tsai HT, Pfeiffer RM, Philips GK, et al. Risks of serious toxicities from intermittent versus continuous androgen deprivation therapy for advanced



NCCN Guidelines Version 5.2026 Prostate Cancer

prostate cancer: a population based study. J Urol 2017;197:1251–1257. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27993663>.

500. Botrel TE, Clark O, dos Reis RB, et al. Intermittent versus continuous androgen deprivation for locally advanced, recurrent or metastatic prostate cancer: a systematic review and meta-analysis. BMC Urol 2014;14:9. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24460605>.

501. Magnan S, Zarychanski R, Pilote L, et al. Intermittent vs continuous androgen deprivation therapy for prostate cancer: a systematic review and meta-analysis. JAMA Oncol 2015;1:1–10. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26378418>.

502. Niraula S, Le LW, Tannock IF. Treatment of prostate cancer with intermittent versus continuous androgen deprivation: a systematic review of randomized trials. J Clin Oncol 2013;31:2029–2036. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23630216>.

503. Hussain M, Tangen C, Higano C, et al. Evaluating intermittent androgen-deprivation therapy phase III clinical trials: the devil is in the details. J Clin Oncol 2015;34:280–285. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26552421>.

504. Ahmadi H, Daneshmand S. Androgen deprivation therapy: evidence-based management of side effects. BJU Int 2013;111:543–548. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23351025>.

505. Gaztanaga M, Crook J. Androgen deprivation therapy: minimizing exposure and mitigating side effects. J Natl Compr Canc Netw 2012;10:1088–1095; quiz 1088, 1096. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22956808>.

506. Lapi F, Azoulay L, Niazi MT, et al. Androgen deprivation therapy and risk of acute kidney injury in patients with prostate cancer. JAMA 2013;310:289–296. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23860987>.

507. Albertsen PC, Klotz L, Tombal B, et al. Cardiovascular morbidity associated with gonadotropin releasing hormone agonists and an

antagonist. Eur Urol 2014;65:565–573. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24210090>.

508. Lopes RD, Higano CS, Slovin SF, et al. Cardiovascular safety of degarelix versus leuprolide in patients with prostate cancer: The primary results of the PRONOUNCE randomized trial. Circulation 2021;144:1295–1307. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34459214>.

509. Sun M, Choueiri TK, Hamnvik OP, et al. Comparison of gonadotropin-releasing hormone agonists and orchiectomy: effects of androgen-deprivation therapy. JAMA Oncol 2016;2:500–507. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26720632>.

510. Kolinsky M, de Bono JS. The ongoing challenges of targeting the androgen receptor. Eur Urol 2016;69:841–843. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26585581>.

511. Gonzalez BD, Jim HS, Booth-Jones M, et al. Course and predictors of cognitive function in patients with prostate cancer receiving androgen-deprivation therapy: a controlled comparison. J Clin Oncol 2015;33:2021–2027. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25964245>.

512. Nead KT, Gaskin G, Chester C, et al. Androgen deprivation therapy and future Alzheimer's Disease risk. J Clin Oncol 2015;34:566–571. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26644522>.

513. Khosrow-Khavar F, Rej S, Yin H, et al. Androgen deprivation therapy and the risk of dementia in patients with prostate cancer. J Clin Oncol 2017;35:201–207. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27870566>.

514. Baik SH, Kury FSP, McDonald CJ. Risk of Alzheimer's disease among senior medicare beneficiaries treated with androgen deprivation therapy for prostate cancer. J Clin Oncol 2017;35:3401–3409. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28841388>.

515. Deka R, Simpson DR, Bryant AK, et al. Association of androgen deprivation therapy with dementia in men with prostate cancer who



NCCN Guidelines Version 5.2026

Prostate Cancer

receive definitive radiation therapy. JAMA Oncol 2018;4:1616–1617.
Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30325986>.

516. Jayadevappa R, Chhatre S, Malkowicz SB, et al. Association between androgen deprivation therapy use and diagnosis of dementia in men with prostate cancer. JAMA Netw Open 2019;2:e196562. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31268539>.

517. Ospina-Romero M, Glymour MM, Hayes-Larson E, et al. Association Between Alzheimer Disease and Cancer With Evaluation of Study Biases: A Systematic Review and Meta-analysis. JAMA Netw Open 2020;3:e2025515. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33185677>.

518. Sari Motlagh R, Quhal F, Mori K, et al. The Risk of New Onset Dementia and/or Alzheimer Disease among Patients with Prostate Cancer Treated with Androgen Deprivation Therapy: A Systematic Review and Meta-Analysis. J Urol 2021;205:60–67. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32856962>.

519. Nowakowska MK, Ortega RM, Wehner MR, Nead KT. Association of second-generation antiandrogens with cognitive and functional toxic effects in randomized clinical trials: A systematic review and meta-analysis. JAMA Oncol 2023;9:930–937. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37227736>.

520. Shahinian VB, Kuo YF, Freeman JL, Goodwin JS. Risk of fracture after androgen deprivation for prostate cancer. N Engl J Med 2005;352:154–164. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15647578>.

521. Smith MR, Boyce SP, Moyneur E, et al. Risk of clinical fractures after gonadotropin-releasing hormone agonist therapy for prostate cancer. J Urol 2006;175:136–139; discussion 139. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16406890>.

522. Smith MR, Lee WC, Brandman J, et al. Gonadotropin-releasing hormone agonists and fracture risk: a claims-based cohort study of men

with nonmetastatic prostate cancer. J Clin Oncol 2005;23:7897–7903. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16258089>.

523. Daniell HW, Dunn SR, Ferguson DW, et al. Progressive osteoporosis during androgen deprivation therapy for prostate cancer. J Urol 2000;163:181–186. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10604342>.

524. Diamond T, Campbell J, Bryant C, Lynch W. The effect of combined androgen blockade on bone turnover and bone mineral densities in men treated for prostate carcinoma: longitudinal evaluation and response to intermittent cyclic etidronate therapy. Cancer 1998;83:1561–1566. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9781950>.

525. Maillefert JF, Sibilia J, Michel F, et al. Bone mineral density in men treated with synthetic gonadotropin-releasing hormone agonists for prostatic carcinoma. J Urol 1999;161:1219–1222. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10081873>.

526. Smith MR, McGovern FJ, Zietman AL, et al. Pamidronate to prevent bone loss during androgen-deprivation therapy for prostate cancer. N Engl J Med 2001;345:948–955. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11575286>.

527. Smith MR, Finkelstein JS, McGovern FJ, et al. Changes in body composition during androgen deprivation therapy for prostate cancer. J Clin Endocrinol Metab 2002;87:599–603. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11836291>.

528. LeBoff MS, Greenspan SL, Insogna KL, et al. The clinician's guide to prevention and treatment of osteoporosis. Osteoporos Int 2022;33:2049–2102. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35478046>.

529. Smith MR, Eastham J, Gleason DM, et al. Randomized controlled trial of zoledronic acid to prevent bone loss in men receiving androgen deprivation therapy for nonmetastatic prostate cancer. J Urol 2003;169:2008–2012. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12771706>.



NCCN Guidelines Version 5.2026 Prostate Cancer

530. Michaelson MD, Kaufman DS, Lee H, et al. Randomized controlled trial of annual zoledronic acid to prevent gonadotropin-releasing hormone agonist-induced bone loss in men with prostate cancer. *J Clin Oncol* 2007;25:1038–1042. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17369566>.

531. Greenspan SL, Nelson JB, Trump DL, Resnick NM. Effect of once-weekly oral alendronate on bone loss in men receiving androgen deprivation therapy for prostate cancer: a randomized trial. *Ann Intern Med* 2007;146:416–424. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17371886>.

532. Smith MR, Egerdie B, Hernandez Toriz N, et al. Denosumab in men receiving androgen-deprivation therapy for prostate cancer. *N Engl J Med* 2009;361:745–755. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19671656>.

533. Sturgeon KM, Deng L, Bluethmann SM, et al. A population-based study of cardiovascular disease mortality risk in US cancer patients. *Eur Heart J* 2019;40:3889–3897. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/31761945>.

534. Keating NL, O'Malley AJ, Smith MR. Diabetes and cardiovascular disease during androgen deprivation therapy for prostate cancer. *J Clin Oncol* 2006;24:4448–4456. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16983113>.

535. Okwuosa TM, Morgans A, Rhee JW, et al. Impact of hormonal therapies for treatment of hormone-dependent cancers (breast and prostate) on the cardiovascular system: Effects and modifications: A scientific statement from the American Heart Association. *Circ Genom Precis Med* 2021;14:e000082. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/33896190>.

536. El-Taji O, Taktak S, Jones C, et al. Cardiovascular events and androgen receptor signaling inhibitors in advanced prostate cancer: A systematic review and meta-analysis. *JAMA Oncol* 2024;10:874–884. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/38842801>.

537. Gardner JR, Livingston PM, Fraser SF. Effects of exercise on treatment-related adverse effects for patients with prostate cancer receiving androgen-deprivation therapy: a systematic review. *J Clin Oncol* 2014;32:335–346. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24344218>.

538. Fizazi K, Tran N, Fein L, et al. Abiraterone plus prednisone in metastatic, castration-sensitive prostate cancer. *N Engl J Med* 2017;377:352–360. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28578607>.

539. Fizazi K, Tran N, Fein L, et al. Abiraterone acetate plus prednisone in patients with newly diagnosed high-risk metastatic castration-sensitive prostate cancer (LATITUDE): final overall survival analysis of a randomised, double-blind, phase 3 trial. *Lancet Oncol* 2019;20:686–700. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/30987939>.

540. Chi KN, Protheroe A, Rodriguez-Antolin A, et al. Patient-reported outcomes following abiraterone acetate plus prednisone added to androgen deprivation therapy in patients with newly diagnosed metastatic castration-naïve prostate cancer (LATITUDE): an international, randomised phase 3 trial. *Lancet Oncol* 2018;19:194–206. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29326030>.

541. James ND, de Bono JS, Spears MR, et al. Abiraterone for prostate cancer not previously treated with hormone therapy. *N Engl J Med* 2017;377:338–351. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/28578639>.

542. Szmulewitz RZ, Peer CJ, Ibraheem A, et al. Prospective international randomized phase II study of low-dose abiraterone with food versus standard dose abiraterone in castration-resistant prostate cancer. *J Clin Oncol* 2018;36:1389–1395. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29590007>.

543. Chi KN, Agarwal N, Bjartell A, et al. Apalutamide for metastatic, castration-sensitive prostate cancer. *N Engl J Med* 2019;381:13–24. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/31150574>.



NCCN Guidelines Version 5.2026 Prostate Cancer

544. Agarwal N, McQuarrie K, Bjartell A, et al. Health-related quality of life after apalutamide treatment in patients with metastatic castration-sensitive prostate cancer (TITAN): a randomised, placebo-controlled, phase 3 study. *Lancet Oncol* 2019;20:1518–1530. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31578173>.

545. Chi KN, Chowdhury S, Bjartell A, et al. Apalutamide in Patients With Metastatic Castration-Sensitive Prostate Cancer: Final Survival Analysis of the Randomized, Double-Blind, Phase III TITAN Study. *J Clin Oncol* 2021;39:2294–2303. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33914595>.

546. Davis ID, Martin AJ, Stockler MR, et al. Enzalutamide with standard first-line therapy in metastatic prostate cancer. *N Engl J Med* 2019;381:121–131. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31157964>.

547. Sweeney CJ, Martin AJ, Stockler MR, et al. Testosterone suppression plus enzalutamide versus testosterone suppression plus standard antiandrogen therapy for metastatic hormone-sensitive prostate cancer (ENZAMET): an international, open-label, randomised, phase 3 trial. *Lancet Oncol* 2023;24:323–334. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36990608>.

548. Armstrong AJ, Szmulewitz RZ, Petrylak DP, et al. ARCHES: A randomized, phase iii study of androgen deprivation therapy with enzalutamide or placebo in men with metastatic hormone-sensitive prostate cancer. *J Clin Oncol* 2019;37:2974–2986. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31329516>.

549. Armstrong AJ, Azad AA, Iguchi T, et al. Improved survival with enzalutamide in patients with metastatic hormone-sensitive prostate cancer. *J Clin Oncol* 2022;40:1616–1622. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35420921>.

550. Saad F, Vjaters E, Shore N, et al. Darolutamide in combination with androgen-deprivation therapy in patients with metastatic hormone-sensitive prostate cancer from the phase III ARANOTE trial. *J Clin Oncol*

2024;42:4271–4281. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39279580>.

551. James ND, Sydes MR, Clarke NW, et al. Addition of docetaxel, zoledronic acid, or both to first-line long-term hormone therapy in prostate cancer (STAMPEDE): survival results from an adaptive, multiarm, multistage, platform randomised controlled trial. *Lancet* 2016;387:1163–1177. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26719232>.

552. Sweeney CJ, Chen YH, Carducci M, et al. Chemohormonal therapy in metastatic hormone-sensitive prostate cancer. *N Engl J Med* 2015;373:737–746. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26244877>.

553. Kyriakopoulos CE, Chen YH, Carducci MA, et al. Chemohormonal therapy in metastatic hormone-sensitive prostate cancer: Long-term survival analysis of the randomized phase III E3805 CHAARTED trial. *J Clin Oncol* 2018;36:1080–1087. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29384722>.

554. Gravis G, Fizazi K, Joly F, et al. Androgen-deprivation therapy alone or with docetaxel in non-castrate metastatic prostate cancer (GETUG-AFU 15): a randomised, open-label, phase 3 trial. *Lancet Oncol* 2013;14:149–158. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23306100>.

555. Gravis G, Boher JM, Joly F, et al. Androgen deprivation therapy (ADT) plus docetaxel versus ADT alone in metastatic non castrate prostate cancer: impact of metastatic burden and long-term survival analysis of the randomized phase 3 GETUG-AFU15 trial. *Eur Urol* 2015;70:256–262. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/26610858>.

556. Fizazi K, Foulon S, Carles J, et al. Abiraterone plus prednisone added to androgen deprivation therapy and docetaxel in de novo metastatic castration-sensitive prostate cancer (PEACE-1): a multicentre, open-label, randomised, phase 3 study with a 2 x 2 factorial design. *Lancet* 2022;399:1695–1707. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/35405085>.



NCCN Guidelines Version 5.2026

Prostate Cancer

557. Smith MR, Hussain M, Saad F, et al. Darolutamide and survival in metastatic, hormone-sensitive prostate cancer. *N Engl J Med* 2022;386:1132–1142. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/35179323>.

558. Vale CL, Fisher DJ, Godolphin PJ, et al. Which patients with metastatic hormone-sensitive prostate cancer benefit from docetaxel: a systematic review and meta-analysis of individual participant data from randomised trials. *Lancet Oncol* 2023;24:783–797. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/37414011>.

559. Hussain M, Tombal B, Saad F, et al. Darolutamide plus androgen-deprivation therapy and docetaxel in metastatic hormone-sensitive prostate cancer by disease volume and risk subgroups in the phase III ARASENS trial. *J Clin Oncol* 2023;41:3595–3607. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/36795843>.

560. Attard G, Agarwal N, Graff JN, et al. Niraparib and abiraterone acetate plus prednisone for HRR-deficient metastatic castration-sensitive prostate cancer: a randomized phase 3 trial. *Nat Med* 2025. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/41057655>.

561. Parker CC, James ND, Brawley CD, et al. Radiotherapy to the primary tumour for newly diagnosed, metastatic prostate cancer (STAMPEDE): a randomised controlled phase 3 trial. *Lancet* 2018;392:2353–2366. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/30355464>.

562. Parker CC, James ND, Brawley CD, et al. Radiotherapy to the prostate for men with metastatic prostate cancer in the UK and Switzerland: Long-term results from the STAMPEDE randomised controlled trial. *PLoS Med* 2022;19:e1003998. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/35671327>.

563. Bossi A, Foulon S, Maldonado X, et al. Efficacy and safety of prostate radiotherapy in de novo metastatic castration-sensitive prostate cancer (PEACE-1): a multicentre, open-label, randomised, phase 3 study with a 2 x 2 factorial design. *Lancet* 2024;404:2065–2076. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/39580202>.

564. Ali A, Hoyle A, Haran AM, et al. Association of bone metastatic burden with survival benefit from prostate radiotherapy in patients with newly diagnosed metastatic prostate cancer: A secondary analysis of a randomized clinical trial. *JAMA Oncol* 2021;7:555–563. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/33599706>.

565. Deek MP, Van der Eecken K, Sutera P, et al. Long-term outcomes and genetic predictors of response to metastasis-directed therapy versus observation in oligometastatic prostate cancer: Analysis of STOMP and ORIOLE trials. *J Clin Oncol* 2022;40:3377–3382. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/36001857>.

566. Francolini G, Allegra AG, Detti B, et al. Stereotactic body radiation therapy and abiraterone acetate for patients affected by oligometastatic castrate-resistant prostate cancer: A randomized phase II trial (ARTO). *J Clin Oncol* 2023;41:5561–5568. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/37733977>.

567. Marvaso G, Corrao G, Zaffaroni M, et al. ADT with SBRT versus SBRT alone for hormone-sensitive oligorecurrent prostate cancer (RADIOA): a randomised, open-label, phase 2 clinical trial. *Lancet Oncol* 2025;26:300–311. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/40049196>.

568. Tang C, Sherry AD, Haymaker C, et al. Addition of metastasis-directed therapy to intermittent hormone therapy for oligometastatic prostate cancer: The EXTEND phase 2 randomized clinical trial. *JAMA Oncol* 2023;9:825–834. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/37022702>.

569. Ost P, Reynders D, Decaestecker K, et al. Surveillance or metastasis-directed therapy for oligometastatic prostate cancer recurrence: A prospective, randomized, multicenter phase II trial. *J Clin Oncol* 2018;36:446–453. Available at:

<https://www.ncbi.nlm.nih.gov/pubmed/29240541>.

570. Phillips R, Shi WY, Deek M, et al. Outcomes of observation vs stereotactic ablative radiation for oligometastatic prostate cancer: The



NCCN Guidelines Version 5.2026 Prostate Cancer

ORIOLE phase 2 randomized clinical trial. JAMA Oncol 2020;6:650–659. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32215577>.

571. Sherry AD, Siddiqui BA, Haymaker C, et al. Continuous androgen deprivation therapy with or without metastasis-directed therapy for oligometastatic prostate cancer: The multicenter phase 2 randomized EXTEND trial. Eur Urol 2025. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/40750497>.

572. Palma DA, Olson R, Harrow S, et al. Stereotactic Ablative Radiotherapy for the Comprehensive Treatment of Oligometastatic Cancers: Long-Term Results of the SABR-COMET Phase II Randomized Trial. J Clin Oncol 2020;38:2830–2838. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32484754>.

573. Scher HI, Halabi S, Tannock I, et al. Design and end points of clinical trials for patients with progressive prostate cancer and castrate levels of testosterone: recommendations of the Prostate Cancer Clinical Trials Working Group. J Clin Oncol 2008;26:1148–1159. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/18309951>.

574. Smith MR, Kabbavar F, Saad F, et al. Natural history of rising serum prostate-specific antigen in men with castrate nonmetastatic prostate cancer. J Clin Oncol 2005;23:2918–2925. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15860850>.

575. Abida W, Armenia J, Gopalan A, et al. Prospective genomic profiling of prostate cancer across disease states reveals germline and somatic alterations that may affect clinical decision making. JCO Precis Oncol 2017;2017. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28825054>.

576. Ryan CJ, Shah S, Efsthathiou E, et al. Phase II study of abiraterone acetate in chemotherapy-naïve metastatic castration-resistant prostate cancer displaying bone flare discordant with serologic response. Clin Cancer Res 2011;17:4854–4861. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21632851>.

577. Scher HI, Morris MJ, Stadler WM, et al. Trial design and objectives for castration-resistant prostate cancer: updated recommendations from

the Prostate Cancer Clinical Trials Working Group 3. J Clin Oncol 2016;34:1402–1418. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26903579>.

578. Holzbeierlein J, Lal P, LaTulippe E, et al. Gene expression analysis of human prostate carcinoma during hormonal therapy identifies androgen-responsive genes and mechanisms of therapy resistance. Am J Pathol 2004;164:217–227. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14695335>.

579. Mohler JL, Gregory CW, Ford OH, 3rd, et al. The androgen axis in recurrent prostate cancer. Clin Cancer Res 2004;10:440–448. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14760063>.

580. de Bono JS, Logothetis CJ, Molina A, et al. Abiraterone and increased survival in metastatic prostate cancer. N Engl J Med 2011;364:1995–2005. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21612468>.

581. Fizazi K, Scher HI, Molina A, et al. Abiraterone acetate for treatment of metastatic castration-resistant prostate cancer: final overall survival analysis of the COU-AA-301 randomised, double-blind, placebo-controlled phase 3 study. Lancet Oncol 2012;13:983–992. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22995653>.

582. Logothetis CJ, Basch E, Molina A, et al. Effect of abiraterone acetate and prednisone compared with placebo and prednisone on pain control and skeletal-related events in patients with metastatic castration-resistant prostate cancer: exploratory analysis of data from the COU-AA-301 randomised trial. Lancet Oncol 2012;13:1210–1217. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23142059>.

583. Ryan CJ, Smith MR, de Bono JS, et al. Abiraterone in metastatic prostate cancer without previous chemotherapy. N Engl J Med 2013;368:138–148. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23228172>.

584. Ryan CJ, Smith MR, Fizazi K, et al. Abiraterone acetate plus prednisone versus placebo plus prednisone in chemotherapy-naïve men



NCCN Guidelines Version 5.2026 Prostate Cancer

with metastatic castration-resistant prostate cancer (COU-AA-302): final overall survival analysis of a randomised, double-blind, placebo-controlled phase 3 study. *Lancet Oncol* 2015;16:152–160. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25601341>.

585. Hussaini A, Olszanski AJ, Stein CA, et al. Impact of an alternative steroid on the relative bioavailability and bioequivalence of a novel versus the originator formulation of abiraterone acetate. *Cancer Chemother Pharmacol* 2017;80:479–486. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28695267>.

586. Goldwater R, Hussaini A, Bosch B, Nemeth P. Comparison of a novel formulation of abiraterone acetate vs. The originator formulation in healthy male subjects: Two randomized, open-label, crossover studies. *Clin Pharmacokinet* 2017;56:803–813. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28425029>.

587. Stein CA, Levin R, Given R, et al. Randomized phase 2 therapeutic equivalence study of abiraterone acetate fine particle formulation vs. originator abiraterone acetate in patients with metastatic castration-resistant prostate cancer: The STAAR study. *Urol Oncol* 2018;36:81 e89–81 e16. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29150328>.

588. Romero-Laorden N, Lozano R, Jayaram A, et al. Phase II pilot study of the prednisone to dexamethasone switch in metastatic castration-resistant prostate cancer (mCRPC) patients with limited progression on abiraterone plus prednisone (SWITCH study). *Br J Cancer* 2018;119:1052–1059. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30131546>.

589. Fenioux C, Louvet C, Charton E, et al. Switch from abiraterone plus prednisone to abiraterone plus dexamethasone at asymptomatic PSA progression in patients with metastatic castration-resistant prostate cancer. *BJU Int* 2019;123:300–306. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30099821>.

590. Scher HI, Fizazi K, Saad F, et al. Increased survival with enzalutamide in prostate cancer after chemotherapy. *N Engl J Med*

2012;367:1187–1197. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22894553>.

591. Fizazi K, Scher HI, Miller K, et al. Effect of enzalutamide on time to first skeletal-related event, pain, and quality of life in men with castration-resistant prostate cancer: results from the randomised, phase 3 AFFIRM trial. *Lancet Oncol* 2014;15:1147–1156. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25104109>.

592. Drugs@FDA: FDA-Approved Drugs. U.S. Food & Drug Administration; Available at: <https://www.accessdata.fda.gov/scripts/cder/daf/index.cfm>. Accessed August 15, 2025.

593. Beer TM, Armstrong AJ, Rathkopf DE, et al. Enzalutamide in metastatic prostate cancer before chemotherapy. *N Engl J Med* 2014;371:424–433. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24881730>.

594. Beer TM, Armstrong AJ, Rathkopf D, et al. Enzalutamide in men with chemotherapy-naïve metastatic castration-resistant prostate cancer: extended analysis of the phase 3 PREVAIL study. *Eur Urol* 2017;71:151–154. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27477525>.

595. Hussain M, Fizazi K, Saad F, et al. Enzalutamide in men with nonmetastatic, castration-resistant prostate cancer. *N Engl J Med* 2018;378:2465–2474. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29949494>.

596. Sternberg CN, Fizazi K, Saad F, et al. Enzalutamide and survival in nonmetastatic, castration-resistant prostate cancer. *N Engl J Med* 2020;382:2197–2206. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32469184>.

597. Tombal B, Saad F, Penson D, et al. Patient-reported outcomes following enzalutamide or placebo in men with non-metastatic, castration-resistant prostate cancer (PROSPER): a multicentre, randomised, double-blind, phase 3 trial. *Lancet Oncol* 2019;20:556–569. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30770294>.



NCCN Guidelines Version 5.2026 Prostate Cancer

598. Smith MR, Saad F, Chowdhury S, et al. Apalutamide treatment and metastasis-free survival in prostate cancer. *N Engl J Med* 2018;378:1408–1418. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29420164>.
599. Saad F, Cella D, Basch E, et al. Effect of apalutamide on health-related quality of life in patients with non-metastatic castration-resistant prostate cancer: an analysis of the SPARTAN randomised, placebo-controlled, phase 3 trial. *Lancet Oncol* 2018;19:1404–1416. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30213449>.
600. Smith MR, Saad F, Chowdhury S, et al. Apalutamide and overall survival in prostate cancer. *Eur Urol* 2020;79:150–158. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32907777>.
601. Fizazi K, Shore N, Tammela TL, et al. Darolutamide in nonmetastatic, castration-resistant prostate cancer. *N Engl J Med* 2019;380:1235–1246. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30763142>.
602. Fizazi K, Shore N, Tammela TL, et al. Nonmetastatic, castration-resistant prostate cancer and survival with darolutamide. *N Engl J Med* 2020;383:1040–1049. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32905676>.
603. Small EJ, Halabi S, Dawson NA, et al. Antiandrogen withdrawal alone or in combination with ketoconazole in androgen-independent prostate cancer patients: a phase III trial (CALGB 9583). *J Clin Oncol* 2004;22:1025–1033. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15020604>.
604. Dupont A, Gomez JL, Cusan L, et al. Response to flutamide withdrawal in advanced prostate cancer in progression under combination therapy. *J Urol* 1993;150:908–913. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/7688437>.
605. Sartor AO, Tangen CM, Hussain MH, et al. Antiandrogen withdrawal in castrate-refractory prostate cancer: a Southwest Oncology Group trial (SWOG 9426). *Cancer* 2008;112:2393–2400. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18383517>.
606. Denmeade SR, Wang H, Agarwal N, et al. TRANSFORMER: A Randomized Phase II Study Comparing Bipolar Androgen Therapy Versus Enzalutamide in Asymptomatic Men With Castration-Resistant Metastatic Prostate Cancer. *J Clin Oncol* 2021;39:1371–1382. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33617303>.
607. Petrylak DP, Tangen CM, Hussain MH, et al. Docetaxel and estramustine compared with mitoxantrone and prednisone for advanced refractory prostate cancer. *N Engl J Med* 2004;351:1513–1520. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15470214>.
608. Tannock IF, de Wit R, Berry WR, et al. Docetaxel plus prednisone or mitoxantrone plus prednisone for advanced prostate cancer. *N Engl J Med* 2004;351:1502–1512. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15470213>.
609. Berthold DR, Pond GR, Soban F, et al. Docetaxel plus prednisone or mitoxantrone plus prednisone for advanced prostate cancer: updated survival in the TAX 327 study. *J Clin Oncol* 2008;26:242–245. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18182665>.
610. Kellokumpu-Lehtinen PL, Harmenberg U, Joensuu T, et al. 2-Weekly versus 3-weekly docetaxel to treat castration-resistant advanced prostate cancer: a randomised, phase 3 trial. *Lancet Oncol* 2013;14:117–124. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23294853>.
611. de Morree ES, Vogelzang NJ, Petrylak DP, et al. Association of survival benefit with docetaxel in prostate cancer and total number of cycles administered: A post hoc analysis of the mainsail study. *JAMA Oncol* 2017;3:68–75. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27560549>.
612. Lavaud P, Gravis G, Foulon S, et al. Anticancer activity and tolerance of treatments received beyond progression in men treated upfront with androgen deprivation therapy with or without docetaxel for metastatic castration-naïve prostate cancer in the GETUG-AFU 15 phase 3 trial. *Eur Urol* 2018;73:696–703. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29074061>.



NCCN Guidelines Version 5.2026

Prostate Cancer

613. de Bono JS, Oudard S, Ozguroglu M, et al. Prednisone plus cabazitaxel or mitoxantrone for metastatic castration-resistant prostate cancer progressing after docetaxel treatment: a randomised open-label trial. *Lancet* 2010;376:1147–1154. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20888992>.

614. Bahl A, Oudard S, Tombal B, et al. Impact of cabazitaxel on 2-year survival and palliation of tumour-related pain in men with metastatic castration-resistant prostate cancer treated in the TROPIC trial. *Ann Oncol* 2013;24:2402–2408. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23723295>.

615. Meisel A, von Felten S, Vogt DR, et al. Severe neutropenia during cabazitaxel treatment is associated with survival benefit in men with metastatic castration-resistant prostate cancer (mCRPC): A post-hoc analysis of the TROPIC phase III trial. *Eur J Cancer* 2016;56:93–100. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26829012>.

616. de Wit R, de Bono J, Sternberg CN, et al. Cabazitaxel versus abiraterone or enzalutamide in metastatic prostate cancer. *N Engl J Med* 2019;381:2506–2518. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31566937>.

617. Fizazi K, Kramer G, Eymard JC, et al. Quality of life in patients with metastatic prostate cancer following treatment with cabazitaxel versus abiraterone or enzalutamide (CARD): an analysis of a randomised, multicentre, open-label, phase 4 study. *Lancet Oncol* 2020;21:1513–1525. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32926841>.

618. Eisenberger M, Hardy-Bessard AC, Kim CS, et al. Phase III study comparing a reduced dose of cabazitaxel (20 mg/m²) and the currently approved dose (25 mg/m²) in postdocetaxel patients with metastatic castration-resistant prostate cancer-PROSELICA. *J Clin Oncol* 2017;35:3198–3206. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28809610>.

619. Oudard S, Fizazi K, Sengelov L, et al. Cabazitaxel versus docetaxel as first-line therapy for patients with metastatic castration-resistant prostate cancer: a randomized phase III trial-FIRSTANA. *J Clin Oncol*

2017;35:JCO2016721068. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28753384>.

620. Oudard S, Ratta R, Voog E, et al. Biweekly vs triweekly cabazitaxel in older patients with metastatic castration-resistant prostate cancer: The CABASTY phase 3 randomized clinical trial. *JAMA Oncol* 2023;9:1629–1638. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37883073>.

621. Sarantopoulos J, Mita AC, He A, et al. Safety and pharmacokinetics of cabazitaxel in patients with hepatic impairment: a phase I dose-escalation study. *Cancer Chemother Pharmacol* 2017;79:339–351. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28058445>.

622. Corn PG, Heath EI, Zurita A, et al. Cabazitaxel plus carboplatin for the treatment of men with metastatic castration-resistant prostate cancers: a randomised, open-label, phase 1-2 trial. *Lancet Oncol* 2019;20:1432–1443. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31515154>.

623. Kantoff PW, Higano CS, Shore ND, et al. Sipuleucel-T immunotherapy for castration-resistant prostate cancer. *N Engl J Med* 2010;363:411–422. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20818862>.

624. Higano CS, Armstrong AJ, Sartor AO, et al. Real-world outcomes of sipuleucel-T treatment in PROCEED, a prospective registry of men with metastatic castration-resistant prostate cancer. *Cancer* 2019;125:4172–4180. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31483485>.

625. Graff JN, Alumkal JJ, Drake CG, et al. Early evidence of anti-PD-1 activity in enzalutamide-resistant prostate cancer. *Oncotarget* 2016;7:52810–52817. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27429197>.

626. Le DT, Durham JN, Smith KN, et al. Mismatch repair deficiency predicts response of solid tumors to PD-1 blockade. *Science* 2017;357:409–413. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28596308>.



NCCN Guidelines Version 5.2026 Prostate Cancer

627. Hansen AR, Massard C, Ott PA, et al. Pembrolizumab for advanced prostate adenocarcinoma: findings of the KEYNOTE-028 study. *Ann Oncol* 2018;29:1807–1813. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29992241>.

628. Tucker MD, Zhu J, Marin D, et al. Pembrolizumab in men with heavily treated metastatic castrate-resistant prostate cancer. *Cancer Med* 2019;8:4644–4655. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31270961>.

629. Marabelle A, Le DT, Ascierto PA, et al. Efficacy of pembrolizumab in patients with noncolorectal high microsatellite instability/mismatch repair-deficient cancer: Results from the phase II KEYNOTE-158 study. *J Clin Oncol* 2020;38:1–10. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31682550>.

630. Antonarakis ES, Piulats JM, Gross-Goupil M, et al. Pembrolizumab for treatment-refractory metastatic castration-resistant prostate cancer: Multicohort, open-label phase II KEYNOTE-199 study. *J Clin Oncol* 2020;38:395–405. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31774688>.

631. Lenis AT, Ravichandran V, Brown S, et al. Microsatellite instability, tumor mutational burden, and response to immune checkpoint blockade in patients with prostate cancer. *Clin Cancer Res* 2024;30:3894–3903. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/38949888>.

632. Marabelle A, Fakih M, Lopez J, et al. Association of tumour mutational burden with outcomes in patients with advanced solid tumours treated with pembrolizumab: prospective biomarker analysis of the multicohort, open-label, phase 2 KEYNOTE-158 study. *Lancet Oncol* 2020;21:1353–1365. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32919526>.

633. Tannock IF, Osoba D, Stockler MR, et al. Chemotherapy with mitoxantrone plus prednisone or prednisone alone for symptomatic hormone-resistant prostate cancer: a Canadian randomized trial with palliative end points. *J Clin Oncol* 1996;14:1756–1764. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8656243>.

634. Kantoff PW, Halabi S, Conaway M, et al. Hydrocortisone with or without mitoxantrone in men with hormone-refractory prostate cancer: results of the cancer and leukemia group B 9182 study. *J Clin Oncol* 1999;17:2506–2513. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10561316>.

635. Kaufman B, Shapira-Frommer R, Schmutzler RK, et al. Olaparib monotherapy in patients with advanced cancer and a germline BRCA1/2 mutation. *J Clin Oncol* 2015;33:244–250. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/25366685>.

636. Mateo J, Carreira S, Sandhu S, et al. DNA-repair defects and olaparib in metastatic prostate cancer. *N Engl J Med* 2015;373:1697–1708. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26510020>.

637. Clarke N, Wiechno P, Alekseev B, et al. Olaparib combined with abiraterone in patients with metastatic castration-resistant prostate cancer: a randomised, double-blind, placebo-controlled, phase 2 trial. *Lancet Oncol* 2018;19:975–986. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29880291>.

638. Farmer H, McCabe N, Lord CJ, et al. Targeting the DNA repair defect in BRCA mutant cells as a therapeutic strategy. *Nature* 2005;434:917–921. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/15829967>.

639. Imyanitov EN, Moiseyenko VM. Drug therapy for hereditary cancers. *Hered Cancer Clin Pract* 2011;9:5. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/21819606>.

640. Cheng HH, Pritchard CC, Boyd T, et al. Biallelic inactivation of BRCA2 in platinum-sensitive metastatic castration-resistant prostate cancer. *Eur Urol* 2016;69:992–995. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26724258>.

641. Pomerantz MM, Spisak S, Jia L, et al. The association between germline BRCA2 variants and sensitivity to platinum-based chemotherapy among men with metastatic prostate cancer. *Cancer* 2017;123:3532–3539. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28608931>.



NCCN Guidelines Version 5.2026 Prostate Cancer

642. Mota JM, Barnett E, Nauseef JT, et al. Platinum-based chemotherapy in metastatic prostate cancer with DNA repair gene alterations. *JCO Precis Oncol* 2020;4:355–366. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32856010>.

643. Schmid S, Omlin A, Higano C, et al. Activity of Platinum-Based Chemotherapy in Patients With Advanced Prostate Cancer With and Without DNA Repair Gene Aberrations. *JAMA Netw Open* 2020;3:e2021692. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33112397>.

644. Hager S, Ackermann CJ, Joerger M, et al. Anti-tumour activity of platinum compounds in advanced prostate cancer—a systematic literature review. *Ann Oncol* 2016;27:975–984. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/27052650>.

645. Antonarakis ES, Lu C, Luber B, et al. Germline DNA-repair gene mutations and outcomes in men with metastatic castration-resistant prostate cancer receiving first-line abiraterone and enzalutamide. *Eur Urol* 2018;74:218–225. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29439820>.

646. Mateo J, Cheng HH, Beltran H, et al. Clinical outcome of prostate cancer patients with germline DNA repair mutations: Retrospective analysis from an international study. *Eur Urol* 2018;73:687–693. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29429804>.

647. Antonarakis ES, Isaacsson Velho P, Fu W, et al. CDK12-Altered Prostate Cancer: Clinical Features and Therapeutic Outcomes to Standard Systemic Therapies, Poly (ADP-Ribose) Polymerase Inhibitors, and PD-1 Inhibitors. *JCO Precis Oncol* 2020;4:370–381. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32462107>.

648. Schweizer MT, Ha G, Gulati R, et al. CDK12-Mutated Prostate Cancer: Clinical Outcomes With Standard Therapies and Immune Checkpoint Blockade. *JCO Precis Oncol* 2020;4:382–392. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32671317>.

649. Nguyen CB, Reimers MA, Perera C, et al. Evaluating immune checkpoint blockade in metastatic castration-resistant prostate cancers with deleterious CDK12 alterations in the phase 2 IMPACT trial. *Clin Cancer Res* 2024;30:3200–3210. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/38787530>.

650. van Wilpe S, Kloots ISH, Slootbeek PHJ, et al. Ipilimumab with nivolumab in molecularly selected patients with castration-resistant prostate cancer: primary analysis of the phase II INSPIRE trial. *Ann Oncol* 2024;35:1126–1137. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39293514>.

651. Mateo J, Porta N, Bianchini D, et al. Olaparib in patients with metastatic castration-resistant prostate cancer with DNA repair gene aberrations (TOPARP-B): a multicentre, open-label, randomised, phase 2 trial. *Lancet Oncol* 2020;21:162–174. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31806540>.

652. de Bono J, Mateo J, Fizazi K, et al. Olaparib for metastatic castration-resistant prostate cancer. *N Engl J Med* 2020;382:2091–2102. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32343890>.

653. Hussain M, Mateo J, Fizazi K, et al. Survival with olaparib in metastatic castration-resistant prostate cancer. *N Engl J Med* 2020;383:2345–2357. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32955174>.

654. Abida W, Patnaik A, Campbell D, et al. Rucaparib in men with metastatic castration-resistant prostate cancer harboring a BRCA1 or BRCA2 gene alteration. *J Clin Oncol* 2020;38:3763–3772. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32795228>.

655. Abida W, Campbell D, Patnaik A, et al. Rucaparib for the treatment of metastatic castration-resistant prostate cancer associated with a DNA damage repair gene alteration: Final results from the phase 2 TRITON2 study. *Eur Urol* 2023. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37277275>.



NCCN Guidelines Version 5.2026 Prostate Cancer

656. Fizazi K, Piulats JM, Reaume MN, et al. Rucaparib or physician's choice in metastatic prostate cancer. *N Engl J Med* 2023;388:719–732. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36795891>.

657. Abida W, Campbell D, Patnaik A, et al. Non-BRCA DNA Damage Repair Gene Alterations and Response to the PARP Inhibitor Rucaparib in Metastatic Castration-Resistant Prostate Cancer: Analysis From the Phase II TRITON2 Study. *Clin Cancer Res* 2020;26:2487–2496. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/32086346>.

658. Schiewer MJ, Goodwin JF, Han S, et al. Dual roles of PARP-1 promote cancer growth and progression. *Cancer Discov* 2012;2:1134–1149. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/22993403>.

659. Polkinghorn WR, Parker JS, Lee MX, et al. Androgen receptor signaling regulates DNA repair in prostate cancers. *Cancer Discov* 2013;3:1245–1253. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24027196>.

660. Li L, Karanika S, Yang G, et al. Androgen receptor inhibitor-induced "BRCAness" and PARP inhibition are synthetically lethal for castration-resistant prostate cancer. *Sci Signal* 2017;10. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28536297>.

661. Clarke NW, Armstrong AJ, Thiery-Vuillemin A, et al. Abiraterone and olaparib for metastatic castration-resistant prostate cancer. *NEJM Evidence* 2022;1:EVIDoa2200043. Available at: <https://evidence.nejm.org/doi/abs/10.1056/EVIDoa2200043>

662. Saad F, Clarke NW, Oya M, et al. Olaparib plus abiraterone versus placebo plus abiraterone in metastatic castration-resistant prostate cancer (PROpel): final prespecified overall survival results of a randomised, double-blind, phase 3 trial. *Lancet Oncol* 2023;24:1094–1108. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37714168>.

663. de Bono JS, Mehra N, Scagliotti GV, et al. Talazoparib monotherapy in metastatic castration-resistant prostate cancer with DNA repair alterations (TALAPRO-1): an open-label, phase 2 trial. *Lancet Oncol*

2021;22:1250–1264. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34388386>.

664. Agarwal N, Azad AA, Carles J, et al. Talazoparib plus enzalutamide in men with first-line metastatic castration-resistant prostate cancer (TALAPRO-2): a randomised, placebo-controlled, phase 3 trial. *Lancet* 2023;402:291–303. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37285865>.

665. Agarwal N, Azad AA, Carles J, et al. Talazoparib plus enzalutamide in men with metastatic castration-resistant prostate cancer: final overall survival results from the randomised, placebo-controlled, phase 3 TALAPRO-2 trial. *Lancet* 2025;406:447–460. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/40683290>.

666. Chi KN, Rathkopf D, Smith MR, et al. Niraparib and abiraterone acetate for metastatic castration-resistant prostate cancer. *J Clin Oncol* 2023;41:3339–3351. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/36952634>.

667. Chi KN, Sandhu S, Smith MR, et al. Niraparib plus abiraterone acetate with prednisone in patients with metastatic castration-resistant prostate cancer and homologous recombination repair gene alterations: second interim analysis of the randomized phase III MAGNITUDE trial. *Ann Oncol* 2023. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/37399894>.

668. Sartor O, de Bono J, Chi KN, et al. Lutetium-177-PSMA-617 for metastatic castration-resistant prostate cancer. *N Engl J Med* 2021;385:1091–1103. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34161051>.

669. Morris MJ, Castellano D, Herrmann K, et al. (177)Lu-PSMA-617 versus a change of androgen receptor pathway inhibitor therapy for taxane-naïve patients with progressive metastatic castration-resistant prostate cancer (PSMAfore): a phase 3, randomised, controlled trial. *Lancet* 2024;404:1227–1239. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39293462>.



NCCN Guidelines Version 5.2026 Prostate Cancer

670. Parker C, Nilsson S, Heinrich D, et al. Alpha emitter radium-223 and survival in metastatic prostate cancer. *N Engl J Med* 2013;369:213–223. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23863050>.

671. Hoskin P, Sartor O, O'Sullivan JM, et al. Efficacy and safety of radium-223 dichloride in patients with castration-resistant prostate cancer and symptomatic bone metastases, with or without previous docetaxel use: a prespecified subgroup analysis from the randomised, double-blind, phase 3 ALSYMPCA trial. *Lancet Oncol* 2014;15:1397–1406. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/25439694>.

672. Sartor O, Coleman R, Nilsson S, et al. Effect of radium-223 dichloride on symptomatic skeletal events in patients with castration-resistant prostate cancer and bone metastases: results from a phase 3, double-blind, randomised trial. *Lancet Oncol* 2014;15:738–746. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24836273>.

673. Nilsson S, Cisko P, Sartor O, et al. Patient-reported quality-of-life analysis of radium-223 dichloride from the phase III ALSYMPCA study. *Ann Oncol* 2016;27:868–874. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26912557>.

674. Smith M, Parker C, Saad F, et al. Addition of radium-223 to abiraterone acetate and prednisone or prednisolone in patients with castration-resistant prostate cancer and bone metastases (ERA 223): a randomised, double-blind, placebo-controlled, phase 3 trial. *Lancet Oncol* 2019;20:408–419. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/30738780>.

675. Tombal B, Choudhury A, Saad F, et al. Enzalutamide plus radium-223 in metastatic castration-resistant prostate cancer: results of the EORTC 1333/PEACE-3 trial. *Ann Oncol* 2025. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/40450503>.

676. Gillesen S, Choudhury A, Rodriguez-Vida A, et al. Decreased fracture rate by mandating bone protecting agents in the EORTC 1333/PEACEIII trial combining Ra223 with enzalutamide versus enzalutamide alone: An updated safety analysis [abstract]. *Journal of*

Clinical Oncology 2021;39:5002–5002. Available at: https://ascopubs.org/doi/abs/10.1200/JCO.2021.39.15_suppl.5002.

677. Francolini G, Bertini N, Di Cataldo V, et al. Impact of stereotactic body radiotherapy after progression in castrate resistant prostate cancer patients undergoing first line abiraterone treatment. A subgroup analysis from ARTO trial (NCT03449719). *Prostate Cancer Prostatic Dis* 2025. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/39972049>.

678. Niazi T, Saad F, Koul R, et al. Metastases-directed therapy in addition to standard systemic therapy in oligometastatic castration resistant prostate cancer: A randomized phase II trial (GROUQ-PCS 9) [abstract]. *Journal of Clinical Oncology* 2025;43:22–22. Available at: https://ascopubs.org/doi/abs/10.1200/JCO.2025.43.5_suppl.22.

679. Beltran H, Tagawa ST, Park K, et al. Challenges in recognizing treatment-related neuroendocrine prostate cancer. *J Clin Oncol* 2012;30:e386–389. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/23169519>.

680. Aggarwal R, Huang J, Alumkal JJ, et al. Clinical and genomic characterization of treatment-emergent small-cell neuroendocrine prostate cancer: A multi-institutional prospective study. *J Clin Oncol* 2018;36:2492–2503. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/29985747>.

681. Brennan SM, Gregory DL, Stillie A, et al. Should extrapulmonary small cell cancer be managed like small cell lung cancer? *Cancer* 2010;116:888–895. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20052730>.

682. Sella A, Konichezky M, Flex D, et al. Low PSA metastatic androgen-independent prostate cancer. *Eur Urol* 2000;38:250–254. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10940696>.

683. Spiess PE, Pettaway CA, Vakar-Lopez F, et al. Treatment outcomes of small cell carcinoma of the prostate: a single-center study. *Cancer* 2007;110:1729–1737. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17786954>.



NCCN Guidelines Version 5.2026 Prostate Cancer

684. Saad F, Gleason DM, Murray R, et al. A randomized, placebo-controlled trial of zoledronic acid in patients with hormone-refractory metastatic prostate carcinoma. *J Natl Cancer Inst* 2002;94:1458–1468. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12359855>.

685. Saad F, Gleason DM, Murray R, et al. Long-term efficacy of zoledronic acid for the prevention of skeletal complications in patients with metastatic hormone-refractory prostate cancer. *J Natl Cancer Inst* 2004;96:879–882. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15173273>.

686. James ND, Pirrie SJ, Pope AM, et al. Clinical outcomes and survival following treatment of metastatic castrate-refractory prostate cancer with docetaxel alone or with strontium-89, zoledronic acid, or both: The trapeze randomized clinical trial. *JAMA Oncol* 2016;2:493–499. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/26794729>.

687. Fizazi K, Carducci M, Smith M, et al. Denosumab versus zoledronic acid for treatment of bone metastases in men with castration-resistant prostate cancer: a randomised, double-blind study. *Lancet* 2011;377:813–822. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21353695>.

688. Himelstein AL, Foster JC, Khatcheressian JL, et al. Effect of longer-interval vs standard dosing of zoledronic acid on skeletal events in patients with bone metastases: A randomized clinical trial. *JAMA* 2017;317:48–58. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28030702>.

689. Smith MR, Halabi S, Ryan CJ, et al. Randomized controlled trial of early zoledronic acid in men with castration-sensitive prostate cancer and bone metastases: results of CALGB 90202 (alliance). *J Clin Oncol* 2014;32:1143–1150. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/24590644>.

690. Coleman RE. Risks and benefits of bisphosphonates. *Br J Cancer* 2008;98:1736–1740. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18506174>.

691. Tarassoff P, Csermak K. Avascular necrosis of the jaws: risk factors in metastatic cancer patients. *J Oral Maxillofac Surg* 2003;61:1238–1239. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14586868>.

692. Smith MR, Saad F, Coleman R, et al. Denosumab and bone-metastasis-free survival in men with castration-resistant prostate cancer: results of a phase 3, randomised, placebo-controlled trial. *Lancet* 2012;379:39–46. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22093187>.

693. Noonan KL, North S, Bitting RL, et al. Clinical activity of abiraterone acetate in patients with metastatic castration-resistant prostate cancer progressing after enzalutamide. *Ann Oncol* 2013;24:1802–1807. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23585511>.

694. Lortot Y, Bianchini D, Ileana E, et al. Antitumour activity of abiraterone acetate against metastatic castration-resistant prostate cancer progressing after docetaxel and enzalutamide (MDV3100). *Ann Oncol* 2013;24:1807–1812. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23576708>.

695. Bianchini D, Lorente D, Rodriguez-Vida A, et al. Antitumour activity of enzalutamide (MDV3100) in patients with metastatic castration-resistant prostate cancer (CRPC) pre-treated with docetaxel and abiraterone. *Eur J Cancer* 2014;50:78–84. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24074764>.

696. Smith MR, Saad F, Rathkopf DE, et al. Clinical outcomes from androgen signaling-directed therapy after treatment with abiraterone acetate and prednisone in patients with metastatic castration-resistant prostate cancer: Post hoc analysis of COU-AA-302. *Eur Urol* 2017;72:10–13. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/28314611>.

697. Khalaf DJ, Annala M, Taavitsainen S, et al. Optimal sequencing of enzalutamide and abiraterone acetate plus prednisone in metastatic castration-resistant prostate cancer: a multicentre, randomised, open-label, phase 2, crossover trial. *Lancet Oncol* 2019;20:1730–1739. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31727538>.



NCCN Guidelines Version 5.2026 Prostate Cancer

698. Hofman MS, Emmett L, Sandhu S, et al. [(177)Lu]Lu-PSMA-617 versus cabazitaxel in patients with metastatic castration-resistant prostate cancer (TheraP): a randomised, open-label, phase 2 trial. *Lancet* 2021;397:797–804. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/33581798>.

699. Wallis CJD, Klaassen Z, Jackson WC, et al. Olaparib vs cabazitaxel in metastatic castration-resistant prostate cancer. *JAMA Netw Open* 2021;4:e2110950. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/34028551>.

700. Abratt RP, Brune D, Dimopoulos MA, et al. Randomised phase III study of intravenous vinorelbine plus hormone therapy versus hormone therapy alone in hormone-refractory prostate cancer. *Ann Oncol* 2004;15:1613–1621. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15520061>.

701. Aparicio AM, Harzstark AL, Corn PG, et al. Platinum-based chemotherapy for variant castrate-resistant prostate cancer. *Clin Cancer Res* 2013;19:3621–3630. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23649003>.

702. Beer TM, Garzotto M, Katovic NM. High-dose calcitriol and carboplatin in metastatic androgen-independent prostate cancer. *Am J Clin Oncol* 2004;27:535–541. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15596926>.

703. Cabrespine A, Guy L, Khenifar E, et al. Randomized Phase II study comparing paclitaxel and carboplatin versus mitoxantrone in patients with hormone-refractory prostate cancer. *Urology* 2006;67:354–359. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16442593>.

704. Harris KA, Harney E, Small EJ. Liposomal doxorubicin for the treatment of hormone-refractory prostate cancer. *Clin Prostate Cancer* 2002;1:37–41. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15046711>.

705. Ladoire S, Eymard JC, Zanetta S, et al. Metronomic oral cyclophosphamide prednisolone chemotherapy is an effective treatment

for metastatic hormone-refractory prostate cancer after docetaxel failure. *Anticancer Res* 2010;30:4317–4323. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21036758>.

706. Lee JL, Ahn JH, Choi MK, et al. Gemcitabine-oxaliplatin plus prednisolone is active in patients with castration-resistant prostate cancer for whom docetaxel-based chemotherapy failed. *Br J Cancer* 2014;110:2472–2478. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24736579>.

707. Loriot Y, Massard C, Gross-Goupil M, et al. Combining carboplatin and etoposide in docetaxel-pretreated patients with castration-resistant prostate cancer: a prospective study evaluating also neuroendocrine features. *Ann Oncol* 2009;20:703–708. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19179557>.

708. Nakabayashi M, Sartor O, Jacobus S, et al. Response to docetaxel/carboplatin-based chemotherapy as first- and second-line therapy in patients with metastatic hormone-refractory prostate cancer. *BJU Int* 2008;101:308–312. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18184327>.

709. Torti FM, Aston D, Lum BL, et al. Weekly doxorubicin in endocrine-refractory carcinoma of the prostate. *J Clin Oncol* 1983;1:477–482. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/6668511>.

710. Shamash J, Powles T, Sarker SJ, et al. A multi-centre randomised phase III trial of dexamethasone vs dexamethasone and diethylstilbestrol in castration-resistant prostate cancer: immediate vs deferred diethylstilbestrol. *Br J Cancer* 2011;104:620–628. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21285990>.

711. Tosoian JJ, Mamawala M, Epstein JI, et al. Active surveillance of grade group 1 prostate cancer: Long-term outcomes from a large prospective cohort. *Eur Urol* 2020;77:675–682. Available at: <https://www.ncbi.nlm.nih.gov/pubmed/31918957>.