KENYA MEDICAL TRAINING COLLEGE
FACULTY OF NURSING
DEPARTMENT OF NURSING

YEAR 2, END OF SEMESTER 1 EXAMINATION
FOR
DIPLOMA IN COMMUNITY HEALTH NURSING

MODULE: PHARMACOLOGY II

DATE: 16TH JULY 2024 **TIME: 2 HOURS (9.00-11.00PM)**

INSTRUCTIONS

1. This paper consists of:
   - Section 1 (20 Multiple Choice Questions)
   - Section 2 (8 Short Answer Questions)
   - Section 3 (2 Long Answer Questions)
2. Attempt **ALL** Questions
3. ALL Multiple choice questions MUST be answered in **CAPITAL LETTERS**
4. Write your **COLLEGE NUMBER** on all the answer sheets provided and on the question paper.
5. No other writing should be done on the question paper except your COLLEGE NUMBER.
6. Ensure that all examination answer scripts are handed in at the end of the examination
7. Ensure you sign the examination register provided

COLLEGE NUMBER .....

PHARMACOLOGY II SET 1

SECTION I: MULTIPLE CHOICE QUESTIONS (20 MARKS)

1. Flumazenil blocks the actions of
   A. Phenobarbital
   B. Morphine
   C. Zolpidem
   D. Ethanol
2. Which of the following antiseizure drugs is also effective in treating trigeminal neuralgia?
   A. Primidone
   B. Topiramate
   C. Carbamazepine
   D. Lamotrigine
3. Barbiturates are used in the emergency treatment of status epilepticus in infants and children because of
   A. They decrease oxygen utilization by the brain
   B. Short onset and duration of action
   C. They do not have effect on sleep architecture
   D. They induce liver microsomal enzymes
4. The preferred treatment of status epilepticus is intravenous administration of
   A. Chlorpromazine
   B. Diazepam
   C. Succinylcholine
   D. Ethosuximide.
5. Which of the following inhalation anesthetics is most likely to produce diffusionhypoxia?
   A. Isoflurane
   B. Enflurance
   C. Halothane
   D. Nitrous oxide
6. Mood elevator drugs are known as;
   A. Analgesics
   B. Soporifics
   C. Tranquilizers
   D. Barbiturates
7. A 35 year old lady who is on anticancer therapy is now suffering from severe nausea and vomiting; which of the following drugs is having most effective antiemetic actions in this case?
   A. Dexamethasone.
   B. Levodopa.
   C. Apomorphine.
   D. Sucrulfate
8. While prescribing a treatment to a young female patient suffering from peptic ulcer associated with H. Pylori the gastroenterologist will neglect:
   A. Tetracycline.
   B. Vancomycin.
   C. Amoxicillin.
   D. Metronidazole.

9. The drug of choice for treatment of absence seizures is
   A. Ethosuximide
   B. Phenytoin
   C. Carbamezapine
   D. Imipramine
10. The drugs associated with enlargement of breasts (gynaecomastia) in males include
   A. Thiopental
   B. Cimetidine
   C. Atropine
   D. Metoclopromide
11. A mixture of gases for inhalational anaesthesia containing nitrous oxide and oxygen in 50:50 proportions is referred as
   A. Isoflurane
   B. Halothane
   C. Entonox
   D. Ether
12. Disulfuram effects are experienced when a patient takes alcohol when on which drug
   A. Fluphenazine
   B. Atracurium
   C. Metronidazole
   D. Imipramine
13. Hypnotic drugs are mainly used in management of
   A. Psychotic disorders
   B. Parkinsons disease
   C. Insomnia
   D. Bulimia
14. Intravenous anaesthetics agents are preferred for induction than inhalation because
   A. Act faster
   B. Are more safer
   C. Do not cross blood brain barrier
   D. Are easy to reverse
15. One of the following statements is correct concerning local anaesthesia:
   A. Its mechanism of action is through blockade of the membrane effects of the excitatory neurotransmitter glutamic acid at the NMDA (N-methyl-D-aspartate) receptor subtype
   B. Its mechanism of action is blockade of voltage-gated sodium channels
   C. It is used in management of peptic ulcer disease
   D. It is a highly flammable medication
16. Which of the following classes of drugs of Purgatives work by retaining water in the lumen of the gut thereby increasing the bulk of faeces;
   A. Stimulant purgatives
   B. Bulk Forming laxatives
   C. Osmotic laxatives
   D. Stool softeners

17. The role of the nurse prior to administration of digoxin is
    A. Starve the patient for six hours
    B. Administer I.V fluids
    C. Monitor the heart rate
    D. Monitor the respiratory rate
18. Reduction of gastric secretions can be achieved by use of:
    A. Antidiuretics
    B. Antihistamine
    C. Antiemetics
    D. Antispasmodic
19. An emetic drug of central action is
    A. Ipecacuanha derivatives
    B. Promethazine
    C. Tropisetron
    D. Apomorphine hydrochloride
20. Cocaine exerts its central action by
    A. Inhibiting phosphodiesterase
    B. Increasing a release of catecholaminergic neurotransmitters, including dopamine
    C. Inhibiting dopamine and norepinephrine reuptake
    D. Altering serotonin turnover

SECTION II: SHORT ANSWER QUESTIONS (40 MARKS)

21. Outline five (5) classes of a drugs for treating peptic ulcer disease. **5 marks**
22. State five (5) indications of insulin therapy. **5 marks**
23. Outline five (5) clinical uses of diazepam. **5 marks**
24. State five (5) goals of treatment in management of Diabetes Mellitus. **5 marks**
25. Explain the mechanism of action of loperamide. **6 marks**
26. State five (5) side effects of glucocorticoids. **5 marks**
27. Outline five (5) specific health messages you will share with a client on phenytoin. **5 marks**
28. Explain mechanism of action of local anesthetic drugs. **4 marks**

SECTION III: LONG ANSWER QUESTIONS (40 MARKS)

29. To manage hypertension, the patient and the nurse must maintain a good therapeutic relationship. Explain five (5) classes of antihypertensive drugs and describe five (5) reasons why the patient and the nurse must maintain a therapeutic relationship in pharmacological management of hypertension. **20 marks**
30. Ms. Y has been diagnosed with depression and has been initiated on anti-depressants. Explain five (5) classes of anti-depressants and describe five (5) health messages for Ms. Y on antidepressants therapy. **20 marks**