

B R A I N P I L L

User Guide

Please carefully read this guide before using our product
and keep it for any future reference or issues.

Table Of Contents

1	Introduction.....	3
2	Quick Start	3
3	Uses for <i>Brain Pill™</i>	3
4	Before Using <i>Brain Pill™</i>	4
	4.1 Allergies	4
	4.2 Pediatric and Breast Feeding	4
	4.3 Storage	4
5	Precautions While Using <i>Brain Pill™</i>	4
6	<i>Brain Pill™</i> Side Effects	4

1 Introduction

Brain Pill™ is based on SSRIs to inject a micro-controller to one's hippocampus and take over the body remotely.

Brain Pill™ is endowed with a *plug and play* technology allowing you to start controlling *Brain Pill™* within seconds¹ by simply connecting a game controller to your device.

2 Quick Start

1. Grab our software at <https://brain-pill.vanille.bid/> or flash our QRCode;
2. Swallow a pill;
3. Wait 2 minutes;
4. Plug in a game controller and enjoy!




Figure 1: controls for a PS4 game controller.

3 Uses for *Brain Pill™*

Brain Pill™ belongs to the *Norwegian Psychiatrist Research Institute*, which researched and developed this product for the sole objective to allow people to overcome their fears and memories.

Brain Pill™ is internally and actively tested to treat:

- Acute Stress Disorders (ASR),
- Agoraphobia,
- Claustrophobia,
- Obsessive-Compulsive Disorders (OCD),
- self-harm,
- pyromania,
- social phobias,
- some forms of schizoaffective disorders.

¹The pill requires two to three minutes to be fully active after swallowing.

4 Before Using *Brain Pill™*

Before anything, the risks must be weighed against the good the pill will do to one's body. Taking this pill is a decision only for you and your doctor to make.

To lower the risk of memory loss, you only should take this pill when you have had a full night of sleep (at least 6 or 7 hours).

Always put at least a 5 hour gap between each *Brain Pill™* usage.

4.1 Allergies

Tell your doctor if you encounter any unusual allergic reaction during or after the pill's effects.

4.2 Pediatric and Breast Feeding

Studies on *Brain Pill™* were only done on adult psychiatric patients. Thus, no information or studies are available on the usage risks for children or infants.

4.3 Storage

The pills must be kept in a closed container at room temperature away from children, liquids, moisture, freezing and direct light.

5 Precautions While Using *Brain Pill™*

Make sure you are in a safe environment, where you cannot easily be hurt. We also recommend that you stay seated on a chair or in bed when waiting for the pill to become active.

Brain Pill™ may cause some people to feel drowsy, dizzy, lightheaded, clumsy or unsteady. Make sure, when getting up, to carefully take your time and do not hesitate to abort if you are feeling heavy or at risk.

As the pill is flagged as an unknown entity, thus, as a normal reaction, will be slowly eliminated. The effects will slowly withdraw. From our experiments, the pills are lasting at least an hour and up to two hours.

6 *Brain Pill™* Side Effects

Along with its desired effects, *Brain Pill™* can come with some unwanted effects. If anything uncommon occurs, make sure to check with your doctor as soon as possible.

Most common:

- Confusion;
- Clumsiness or unsteadiness;
- Difficulty with coordination;
- Mood or mental changes.

Less common:

- Severe drowsiness;
- Shortness of breath and difficult or labored breathing;
- Tightness in the chest;
- Memory loss.

Rare:

- Loss of consciousness;
- Increased appetite;
- Weight loss;
- Vomiting.