Patient ID	: QLD24/3922	Sample ID	: 24081284
Name	**: MRS K ARUNA**	Registration Date	: 24/08/2024 07:58 AM
Age/Gender	: 51 Yrs/Female	Report Date	: 24/08/2024 03:50 PM
Ref. Dr.	: SELF	QLD24/3922	
Center	: Q LABS DIAGNOSTICS		

BIOCHEMISTRY REPORT

Test Description	Result	Unit	Biological Reference Ranges
Fasting Plasma Glucose *Method: Hexokinase*	108	mg/dl	Normal : 74 - 106 Impaired GT : 107 - 125 Diabetic : >or=126

> or =126	Diabetic	Its a group of metabolic disorders characterized by a high blood sugar level over a prolonged period of time
101-125	Impaired GT	Impaired glucose tolerance means that blood glucose is raised beyond normal levels, but not high enough to warrant a diabetes diagnosis. With impaired glucose tolerance one can face a much greater risk of developing diabetes and cardiovascular disease.
70-100	Normal	The American Diabetes Association recommends a routine screening for type 2 diabetes. If the results are normal, the screening should be repeated every 3 years.

Serum Creatinine *Method: Modified Jaffe's*	0.70	mg/dL	0.60 - 1.30

Dr.SYEDA SALEHA FAREED MD PATHOLOGY
Authorised Signature

Patient ID	: QLD24/3922	Sample ID	: 24081284
Name	**: MRS K ARUNA**	Registration Date	: 24/08/2024 07:58 AM
Age/Gender	: 51 Yrs/Female	Report Date	: 24/08/2024 03:50 PM
Ref. Dr.	: SELF	QLD24/3922	
Center	: Q LABS DIAGNOSTICS		

LIPID PROFILE

Cholesterol-Total	**235.24**	mg/dL	< 200 - Desirable
Method: Spectrophotometry			200 - 239 -Boderline High
			> 240 - High
Triglycerides level	**344.14**	mg/dL	< 150 - Normal
Method: Serum, Enzymatic, endpoint			150 - 199 -Boderline High
			200 - 499 -High
			>500 Very -High
HDL Cholesterol	**47.0**	mg/dL	< 40 - Low
Method: Serum, Direct measure-PEG			> 40 - Normal
LDL Cholesterol	119.41	mg/dL	< 100 - Optimal
Method: Enzymatic selective protection			100 - 129 - Near/Above Optimal
			130 - 159 - Borderline high
			160 - 189 - High
			> 190 - Very High
VLDL Cholesterol	**68.83**	mg/dL	6 - 38
Method: Serum, Enzymatic			
CHOL/HDL RATIO	**5.01**		0.27 - 5.0
Method: Serum, Enzymatic			
LDL/HDL RATIO	2.54		2.5 - 3.5
Method: Serum, Enzymatic			

NOTE

8-10 hours fasting sample is required

**** End of the report. ****
Please correlate clinically.

Dr.SYEDA SALEHA FAREED MD PATHOLOGY
Authorised Signature