

PLOS Computational Biology

The HIV Post-Exposure Prophylaxis Window: A Multiscale Framework Linking Within-Host Integration Kinetics to Population-Scale Structural Access --Manuscript Draft--

Manuscript Number:	
Full Title:	The HIV Post-Exposure Prophylaxis Window: A Multiscale Framework Linking Within-Host Integration Kinetics to Population-Scale Structural Access
Short Title:	A Multiscale Framework for HIV Post-Exposure Prophylaxis
Article Type:	Research Article
Keywords:	HIV post-exposure prophylaxis; multiscale within-host modeling; stochastic viral dynamics; proviral integration kinetics; structural barriers to HIV prevention; people who inject drugs
Corresponding Author:	Adrian (AC) Charles Demidont, DO Nyx Dynamics, LLC Fairfield, CT UNITED STATES OF AMERICA
Corresponding Author Secondary Information:	
Corresponding Author's Institution:	Nyx Dynamics, LLC
Corresponding Author's Secondary Institution:	
First Author:	Adrian (AC) Charles Demidont, DO
First Author Secondary Information:	
Order of Authors:	Adrian (AC) Charles Demidont, DO
Order of Authors Secondary Information:	
Abstract:	Proviral integration — the molecular event that converts HIV infection from reversible to permanent — defines a finite window during which post-exposure prophylaxis (PEP) can succeed. Using a multiscale within-host model that combines stochastic founder-phase tau-leaping with deterministic ODE handoff and an eclipse-phase fixed delay anchored on Perelson 1996, approximately 22 hours, we prove that PEP efficacy decays monotonically to zero and derive the critical prevention window from emergent integration kinetics rather than from drug potency or clinical convention. This window is approximately 1.75-fold shorter for parenteral, or injection, exposure than mucosal, or sexual, exposure, yielding a parenteral critical time at the 5% efficacy threshold of approximately 34.5 hours for $V_0 = 10^3$ and $CV = 0.3$, versus 60.5 hours mucosally for $V_0 = 1$ and $CV = 0.3$. This compression is concordant with the empirical 1.5- to 2-fold compression range spanning Tsai 1998 intravenous SIV and Otten 2000 intravaginal SHIV nonhuman-primate challenge data. Independent cascade-based modeling of structural access for people who inject drugs produces population-level prevention success of 0.003% to 0.7% under current U.S. policy, with a 92.7% national 10-year outbreak probability. A distribution-free $F_{\text{access}} \times \text{tcrit}$ bound provides a conservative analytical consistency check, while a PK-driven corridor analysis shows that pharmacy and structural delays act by displacing acquisition time toward a route-specific integration cliff rather than by changing drug efficacy itself. The failure mode is structural rather than pharmacological: PEP retains high per-exposure efficacy when acquired in time; population-level efficacy is constrained by the mismatch between access timing and integration timing.
Suggested Reviewers:	Jessica M Conway, PhD Associate Professor, Penn State: The Pennsylvania State University jmconway@psu.edu Within-host HIV viral dynamics modeling in the Perelson lineage; published stochastic analyses of pre- and post-exposure prophylaxis. Will evaluate the multiscale stochastic-deterministic architecture and the founder-phase tau-leaping methodology.

	Max von Kleist, PhD Group Leader, Robert Koch Institute: Robert Koch Institut max.kleist@fu-berlin.de Leading PEP/PrEP mathematical modeler whose work integrates pharmacokinetics, pharmacodynamics, and viral dynamics in hybrid stochastic-deterministic frameworks Gregg Gonsalves, PhD Associate Professor, Yale University School of Public Health gregg.gonsalves@yale.edu Senior author of Bayesian modeling analyses of the Lawrence/Lowell, MA HIV outbreak among PWID; quantitative modeler at the intersection of structural HIV epidemiology, substance use, and health equity. Sally Blower, PhD Distinguished Professor and Director, UCLA: University of California Los Angeles sblower@mednet.ucla.edu Senior HIV mathematical epidemiologist with extensive published work on population-level HIV intervention modeling and structural determinants.
Opposed Reviewers:	
Additional Information:	
Question	Response
Government Employee Are you or any of the contributing authors an employee of the United States government? Manuscripts authored by one or more US Government employees are not copyrighted, but are licensed under a CC0 Public Domain Dedication, which allows unlimited distribution and reuse of the article for any lawful purpose. This is a legal requirement for US Government employees. This will be typeset if the manuscript is accepted for publication.	No - No authors are employees of the U.S. government.
Financial Disclosure Enter a financial disclosure statement that describes the sources of funding for the work included in this submission. Review the submission guidelines for detailed requirements. View published research articles from PLOS Computational Biology for specific examples. This statement is required for submission and will appear in the published article if	The author(s) received no specific funding for this work.

the submission is accepted. Please make sure it is accurate. Funded studies Enter a statement with the following details: • Initials of the authors who received each award • Grant numbers awarded to each author • The full name of each funder • URL of each funder website • Did the sponsors or funders play any role in the study design, data collection and analysis, decision to publish, or preparation of the manuscript? Did you receive funding for this work?	
Competing Interests On behalf of all authors, disclose any competing interests that could be perceived to bias this work. This statement will be typeset if the manuscript is accepted for publication. Review the instructions link below and PLOS Computational Biology's competing interests policy to determine what information must be disclosed at submission.	The author reports prior employment with Gilead Sciences, Inc. from January 2020 through November 2024, and held company stock during that period; all Gilead stock was fully divested by December 2024. Employment ended prior to the initiation of this research. Gilead Sciences had no role in study conception, design, data collection, analysis, interpretation, writing, or the decision to submit for publication. The manuscript references lenacapavir, a Gilead product, as one member of the long-acting injectable antiretroviral class; the mathematical conclusions of this work are pharmacologically agnostic and pertain to the class as a whole. The author is the sole owner of Nyx Dynamics LLC, under whose affiliation this research was conducted independently.
Data Availability Before publication, Authors are required to make fully available and without restriction all data underlying their findings. Sharing the data underlying your findings increases the visibility, credibility, and impact of your work, while supporting transparency and future scientific discovery. Please see our PLOS Data Policy page for detailed information on this policy. A Data Availability Statement, detailing	All source code, simulation outputs, and the numerical claims registry needed to reproduce every result in this manuscript and supplement are available at https://github.com/Nyx-Dynamics/Prevention-Theorem under the v4.0.0 release tag, with a persistent archive at Zenodo (v4.0.0 DOI: 10.5281/zenodo.20559022). The single-command reproduction pipeline is reproduce all v4.py at the repository root, which regenerates all main-text figures (Figs. 1–3), supplement tables (Tables S2a–S2c), and CSV artifacts from the multiscale baseline at commit 37e27ea. The full claims registry mapping every numerical value in the manuscript to its source artifact, locator within that artifact, and verification command is provided as v3 revision/numerical claims v4.csv. The 34-city baseline (Table S2b) is derived from publicly available AIDSVu downloadable datasets (2023 release) at https://aidsvu.org. Multiscale viral-dynamics realizations and within-host PK/PD parameters are documented in Supplement S5 and the v3 revision/r3 pk pd/ subdirectory of the repository. No human subjects data were used in this study. IRB approval was not required

where the data can be accessed, is required at first submission. Insert your Data Availability Statement in the box below.

Please see the [data reporting](#) section of our submission guidelines for instructions on what you need to include in your Data Availability Statement. Examples of acceptable Data Availability Statements are included here for reference:

1. The minimal data set [and accompanying code, where generated] is available at [repository name] via [DOI hyperlink]
2. No data was generated by this study. The following existing data sources were used: [dataset name] from [dataset location] available via [web address or DOI hyperlink].
3. For Study Protocols: No datasets were generated or analysed during the current study. All relevant data from this study will be made available upon study completion.

PLOS allows rare exemptions to address legal and ethical concerns. If you have legal or ethical restrictions, please use the box below to detail these in full sentences for the Journal team to consider.

This statement will be typeset if the manuscript is accepted for publication.

A.C. Demidont, DO, AAHIVS
 Sole Founder, Nyx Dynamics LLC
 Philadelphia, PA
 ORCID: 0000-0002-9216-8569
 acdemidont@nyxdynamics.com

June 5, 2026

Editors-in-Chief
PLOS Computational Biology

Dear Editors:

I am pleased to submit for your consideration the manuscript “**The HIV Post-Exposure Prophylaxis Window: A Multiscale Framework Linking Within-Host Integration Kinetics to Population-Scale Structural Access.**” I am the sole author.

This manuscript develops a mathematical and computational framework for deriving the finite prevention window for HIV post-exposure prophylaxis (PEP) from within-host integration kinetics rather than from clinical convention. The model combines stochastic founder-phase tau-leaping, deterministic target-cell ODE handoff, and an eclipse-phase delay anchored on established HIV viral-dynamics literature. This multiscale architecture yields an emergent route-dependent prevention window: approximately **34.5 hours for parenteral exposure** and **60.5 hours for mucosal exposure**, corresponding to an approximately **1.75-fold compression** that is concordant with the compression range observed across nonhuman-primate PEP timing studies.

The computational contribution is threefold. First, the manuscript proves a finite-window theorem showing that PEP efficacy decays monotonically to zero once irreversible proviral integration is reached. Second, it replaces fixed stage-efficacy assumptions with a PK-driven kinetic-onset framework for TDF/FTC plus dolutegravir, separating pharmacologic drug accumulation from biological clearability. Third, it links within-host kinetics to structural access through city-stratified modeling across 34 high-burden U.S. metropolitan areas, with a pharmacy-access corridor analysis showing how structural and dispensing delays displace acquisition time toward a route-specific integration cliff.

The population-level conclusion is intentionally framed conservatively. Companion cascade-based analyses estimate **0.003–0.7%** population-level prevention success for PWID under current U.S. policy, while the $F_{\text{access}} \times t_{\text{crit}}$ inequality is retained only as an analytical consistency bound. The primary mechanistic access result is the PK/PD corridor analysis: pharmacy delay does not change drug potency or compress the biological corridor; it shifts patients and cities rightward along acquisition time until the route-specific integration boundary is crossed.

This work is well suited to *PLOS Computational Biology* because it bridges mechanistic within-host viral dynamics, stochastic first-passage modeling, pharmacokinetic/pharmacodynamic simulation, and population-scale structural epidemiology. The manuscript’s central claim is not a clinical guideline recommendation but a computational biology result: the biologically available

interval for post-exposure prevention is an emergent property of viral integration kinetics, route-specific inoculum structure, drug-onset kinetics, and access timing.

The mathematical core of an earlier version was previously submitted to *Science* and transferred editorially to *Science Advances*. Following peer review and appeal, *Science Advances* declined the manuscript as outside journal scope. The present submission has been substantially revised for a computational biology audience, including the multiscale stochastic-deterministic implementation, PK-driven efficacy framework, pharmacy-access corridor analysis, expanded reproducibility documentation, and updated supplementary proofs and parameter tables.

I disclose prior employment with Gilead Sciences, which ended in October–November 2024, and complete divestiture of Gilead stock in December 2024. These events predate the initiation of this manuscript. Gilead Sciences had no role in study conception, design, analysis, interpretation, writing, or the decision to submit. Because of this prior employment, I respectfully request that the manuscript not be assigned to reviewers currently affiliated with Gilead Sciences.

The manuscript is not under consideration elsewhere. All code, simulation outputs, processed data, and numerical-claim verification files are archived at Zenodo under v4.0.0: <https://doi.org/10.5281/zenodo.20559022>, with the corresponding GitHub release at <https://github.com/Nyx-Dynamics/Prevention-Theorem>. Reproduction is documented through the repository's `reproduce_all_v4.py` pipeline.

Thank you for considering this manuscript for *PLOS Computational Biology*.

Sincerely,

A.C. Demidont, DO, AAHIVS
Sole Founder, Nyx Dynamics LLC
ORCID: 0000-0002-9216-8569

The HIV Post-Exposure Prophylaxis Window:

A Multiscale Framework Linking Within-Host Integration Kinetics to Population-Scale Structural Access

Adrian Charles (A.C) Demidont, DO*

Nyx Dynamics LLC, Philadelphia, PA 19107, USA

*Correspondence: ac.demidont@nyxdynamics.com

ORCID: 0000-0002-9216-8569

Abstract

Proviral integration — the molecular event that converts HIV infection from reversible to permanent — defines a finite window during which post-exposure prophylaxis (PEP) can succeed. Using a multiscale within-host model that combines stochastic founder-phase tau-leaping with a deterministic ODE handoff and an eclipse-phase fixed delay anchored on Perelson 1996 (~ 22 h), we prove that PEP efficacy decays monotonically to zero and derive the critical prevention window from emergent integration kinetics rather than from drug potency or clinical convention. This window is approximately 1.75-fold shorter for parenteral (injection) than mucosal (sexual) exposure, yielding a parenteral t_{crit} at $\eta = 0.05$ of approximately 34.5 hours ($V_0 = 10^3$, $\text{CV} = 0.3$) versus 60.5 hours mucosally ($V_0 = 1$, $\text{CV} = 0.3$) — concordant with the empirical $1.5\text{--}2\times$ compression range spanning Tsai 1998 (intravenous SIV) and Otten 2000 (intravaginal SHIV) NHP challenge data. Independent cascade-based modeling of structural access for people who inject drugs produces population-level prevention success of $0.003\text{--}0.7\%$ under current US policy [companion analysis], with a 92.7% national 10-year outbreak probability. A distribution-free $F_{\text{access}} \times t_{\text{crit}}$ bound provides a conservative analytical consistency check, while a PK-driven corridor analysis shows that pharmacy and structural delays act by displacing acquisition time toward a route-specific integration cliff rather than by changing drug efficacy itself. The failure mode is structural rather than pharmacological: PEP retains high per-exposure efficacy when acquired in time; population-level efficacy is constrained by the mismatch between access timing and integration timing.

Keywords: HIV; post-exposure prophylaxis (PEP); proviral integration; within-host viral dynamics; multiscale modeling; stochastic simulation; people who inject drugs; structural barriers

Author summary

Post-exposure prophylaxis (PEP) is the medical strategy of taking antiretroviral drugs after a possible HIV exposure to prevent infection from being established. Current CDC guidance recommends initiation within 72 hours. We show that the actual window

during which PEP can succeed is not a clinical convention but a mathematically determined biological quantity, defined by the timing of an irreversible molecular event: integration of HIV into the host's DNA. Using a multiscale within-host model anchored on established HIV viral dynamics, we prove this prevention window is approximately 1.75-fold shorter for parenteral (injection) than for mucosal (sexual) exposure — roughly 34.5 hours versus 60.5 hours. For people who inject drugs in many US cities, the structural delays between exposure and PEP initiation — driven by late HIV diagnosis in potential partners, gaps in linkage to care, and pharmacy access barriers — routinely exceed this parenteral window. Under current US policy, cascade-based estimates of population-level prevention are substantially lower than 1%, while an analytical $F_{\text{access}} \times t_{\text{crit}}$ bound provides a conservative upper-limit check of approximately 11%. The failure mode is structural, not pharmacological: PEP retains its high per-exposure efficacy, but the healthcare system is built to deliver it outside the biologically active window.

Introduction

Retroviral infection is unique among infectious diseases in that a single molecular event — proviral integration into the host genome — converts a reversible inflammatory process into a permanent genomic state. This irreversibility is absolute: no antiviral intervention operating after integration can restore the pre-infection state. Post-exposure prophylaxis (PEP) exploits the finite interval before this transition, but the boundaries of that interval have never been formally derived from within-host kinetics. Here we prove, using a three-state absorbing Markov framework parameterized on established HIV-1 viral dynamics [1–3], that the prevention window is not a clinical convention but a mathematically necessary consequence of integration kinetics — that it is route-specific, compressible by viral load, and in the case of parenteral (injection) exposure, potentially shorter than the structural delays that define healthcare access for people who inject drugs (PWID). The result transforms PEP timing from an empirical guideline into a derived biological quantity, and reveals that the 72-hour standard — mechanistically appropriate for mucosal exposure — cannot be assumed to generalize across routes or populations.

HIV-1 infection establishment involves two irreversible transitions whose timing governs the prevention window. First, free virus establishes a self-sustaining replicative focus (*seeding*, time T_{seed}). Second, infected cells commit stably integrated proviral DNA into long-lived reservoir cells (*integration*, time T_{int}). Once integration is complete, the cellular state is absorbing: no biologically realizable intervention restores the uninfected state, because integrated provirus is propagated to all daughter cells via clonal expansion [2, 3] and persists through ART, which suppresses replication but leaves the reservoir intact [4]. This biological fact constrains PEP efficacy independently of drug potency.

Mathematical framework

We model within-host state as a random variable $Z(t) \in \{0, 1, 2\}$, where $Z = 0$ (susceptible), $Z = 1$ (seeded, pre-integration), and $Z = 2$ (integrated, absorbing). Transition times $T_{\text{seed}} \leq T_{\text{int}}$ are stopping times on the natural filtration of the exposure process. Under stage-dependent drug efficacy $\varepsilon(Z)$ — with ε_{max} pre-seeding, ε_{mid} post-seeding/pre-integration, and $\varepsilon_{\text{min}} = 0$ post-integration — the expected PEP efficacy at time t is, by the law of total probability:

$$E_{\text{PEP}}(t) = (1 - P_{\text{seed}}(t))\varepsilon_{\text{max}} + (P_{\text{seed}}(t) - P_{\text{int}}(t))\varepsilon_{\text{mid}} + P_{\text{int}}(t)\varepsilon_{\text{min}} \quad (1)$$

where $P_{\text{seed}}(t) = \Pr(T_{\text{seed}} \leq t)$ and $P_{\text{int}}(t) = \Pr(T_{\text{int}} \leq t)$. Since both cumulative probabilities are non-decreasing in t and $P_{\text{int}}(t) \rightarrow 1$, this yields the **Finite Prevention Window Theorem**: (i) $E_{\text{PEP}}(t)$ is non-increasing; (ii) $E_{\text{PEP}}(t) \rightarrow 0$; and (iii) for any threshold $\eta \in (0, 1)$, there exists a finite critical time $t_{\text{crit}}(\eta) = \inf\{t : E_{\text{PEP}}(t) < \eta\}$, beyond which efficacy cannot be recovered regardless of drug potency. The rate of efficacy decay decomposes as

$$\frac{dE_{\text{PEP}}}{dt} = -f_{\text{seed}}(t) \Delta\varepsilon_1 - f_{\text{int}}(t) \Delta\varepsilon_2 \leq 0 \quad (2)$$

where f_{seed} and f_{int} are the transition probability densities and $\Delta\varepsilon_1, \Delta\varepsilon_2$ are the efficacy drops across transitions (Supplementary Text, Section S4). The absorbing-state claim (Assumption A1) is grounded in the standard target-cell-limited ODE system [5]: under ART-induced suppression $I \rightarrow 0$, reservoir dynamics give $dR/dt = \alpha I \rightarrow 0$, so R is constant — the reservoir neither grows nor is cleared. This is the mathematical statement that ART controls but cannot eradicate HIV.

Assumption: transmission-competent exposure. The Finite Windows framework models within-host integration kinetics conditional on exposure to viremic source virus. We assume throughout that source plasma viral load is at or above the established U=U threshold (≥ 200 copies/mL sustained). The framework's predictions — t_{crit} , F_{access} , and the population-level efficacy bound — are not applicable to exposures from virologically suppressed sources, for whom transmission risk is functionally zero [10, 11]. U=U exposures are operationally outside scope: no PEP need arises.

Route-dependent window compression

Parenteral inoculation differs from mucosal exposure in its initial condition: direct intravascular delivery bypasses epithelial barriers, producing a substantially larger effective inoculum. We model mucosal exposure as $V^{(m)}(0) = 1$ virion/mL (post-epithelial attenuation) versus parenteral $V^{(p)}(0) = 10^3$ virions/mL. By monotone systems theory applied to the target-cell-limited ODE (Lemma S5.2), larger initial inoculum produces earlier $I(t)$ and consequently earlier $R(t)$ accumulation, yielding first-order stochastic dominance: $P_{\text{int}}^{(p)}(t) \geq P_{\text{int}}^{(m)}(t)$ for all t . It follows directly that $E_{\text{PEP}}^{(p)}(t) \leq E_{\text{PEP}}^{(m)}(t)$ and $t_{\text{crit}}^{(p)}(\eta) \leq t_{\text{crit}}^{(m)}(\eta)$ for any threshold η (Corollary S5.1).

Multiscale model and parameterization

The framework is implemented as a two-phase multiscale within-host model. **Phase 1** simulates founder-population dynamics stochastically via tau-leaping at low inoculum ($V_0 \leq I_{\text{handoff}} = 10$), capturing extinction events for sparse founder populations. **Phase 2** integrates the deterministic target-cell-limited ODE system once productive infection is established, anchored on the within-host kinetic literature: viral clearance rate $c = 23 \text{ day}^{-1}$ (virion $t_{1/2} \approx 0.7 \text{ h}$; Ramratnam et al. 1999 [6], refining the earlier Perelson 1996 [5] estimate), infected-cell death rate $\delta = 0.7 \text{ day}^{-1}$ [5], eclipse phase $\tau = 0.9 \text{ days}$ ($\approx 22 \text{ h}$; Perelson 1996, Table 2). Remaining target-cell-model parameters were set to standard simulation values and sensitivity-tested rather than directly inferred from the Perelson cohort: target cell production $\lambda = 10^4 \text{ cells/mL/day}$, natural death $d_T = 0.01 \text{ day}^{-1}$, infection rate $\beta = 2.4 \times 10^{-5} \text{ mL} \cdot \text{virion}^{-1} \cdot \text{day}^{-1}$, viral production $p = 10^3 \text{ virions/cell/day}$, and integration rate $\alpha = 10^{-3} \text{ day}^{-1}$. Heterogeneity is implemented as multiplicative log-normal variation ($\text{CV} = 0.3$) on $\{\beta, c, \delta, \alpha\}$. The eclipse phase is implemented as a discrete fixed-delay buffer between cell infection and integration competence, ensuring that the lower bound on T_{int} is the

biological eclipse floor. We use $N = 500$ realizations per (route, V_0 , CV) cell across the parameter sweep; the canonical operating point is parenteral $V_0 = 10^3$ at CV = 0.3 and mucosal $V_0 = 1$ at CV = 0.3. Stage-dependent drug efficacy is implemented via the PK-driven framework described in the next section, with $\varepsilon_{\text{drug}}(t_{\text{PEP}}) \cdot p_{\text{clear}}(Z)$ generalizing the hand-set efficacy constants ($\varepsilon_{\text{max}} = 0.95$, $\varepsilon_{\text{mid}} = 0.50$, $\varepsilon_{\text{min}} = 0$) of the baseline formulation. Integration time T_{int} is the first time $R(t) \geq R^* = 10$ cells/mL. Threshold sensitivity ($I^* \in [10, 1000]$, $R^* \in [1, 100]$) shifts $t_{\text{crit}}^{(p)}$ by < 8 h and maintains the qualitative compression result.

PK-driven PEP efficacy: kinetic onset framework

The PK-driven framework generalizes the hand-set stage efficacies of the baseline formulation into PK-derived values capturing two mechanistic phenomena: (i) the deterministic kinetics by which antiretroviral drugs reach therapeutic intracellular concentration after PEP initiation, and (ii) the separation of pharmacology from stage biology. The decomposition is

$$\varepsilon(Z, t_{\text{PEP}}) = \varepsilon_{\text{drug}}(t_{\text{PEP}}) \cdot p_{\text{clear}}(Z) \quad (3)$$

where $\varepsilon_{\text{drug}}(t_{\text{PEP}})$ is the time-averaged pharmacologic suppression delivered during the remaining route-specific integration window $[t_{\text{PEP}}, t_{\text{crit}}^{\text{route}}]$, computed from a TDF/FTC+DTG one-compartment PK model with intracellular metabolite compartments (TFV-DP, FTC-TP) for the NRTIs and plasma-active kinetics for DTG (Section S13). Per-drug suppression follows a Hill function on the normalized active fraction; combination across drugs assumes independent molecular targets (NRTIs block reverse transcription; INSTI blocks integration), yielding multiplicative survival on new infection events. The biological clearability $p_{\text{clear}}(Z)$ captures the stage-dependent fraction of trajectory still pharmacologically rescuable: $p_{\text{clear}}(0) = 1$ (pre-seeding), $p_{\text{clear}}(1) = 0.50$ (calibrated to recover the hand-set baseline ε_{mid} at perfect adherence), $p_{\text{clear}}(2) = 0$ (post-integration absorbing state).

At perfect adherence the PK-driven framework recovers the multiscale baseline within calibration tolerance: $t_{\text{crit}}^{(p)} = 34.0$ h (multiscale baseline 34.5 h; 0.5 h shift attributable to the kinetic-onset cost during the early hours of PEP); $t_{\text{crit}}^{(m)} = 59.5$ h (multiscale baseline 60.5 h). The $F_{\text{access}} \times t_{\text{crit}}$ envelope bound is preserved at 11.2% (multiscale baseline 11.3%). The novel finding is the dependence of $\varepsilon_{\text{drug}}(t_{\text{PEP}})$ on the *remaining* window: at $t_{\text{PEP}} = 0$, parenteral $\varepsilon_{\text{drug}} = 0.98$ (drug has the full 34.5 h to ramp up); at $t_{\text{PEP}} = 33$ h, $\varepsilon_{\text{drug}} = 0.78$ (only 1.5 h remaining, insufficient for full intracellular accumulation). Adherence sensitivity (Section S13.6): at 70% adherence $t_{\text{crit}}^{(p)}$ drops by 1.0 h; at 50% adherence by 1.5 h; at 30% adherence by 2.5 h — the prevention window is remarkably robust to adherence variation because DTG's plasma-active kinetics provide rapid early suppression while NRTI intracellular phosphorylated metabolites are still accumulating.

The formulation directly separates pharmacology from outcome probability: $\varepsilon_{\text{drug}}$ depends only on PK/PD and is identifiable from drug-class pharmacokinetic literature; p_{clear} depends only on the within-host state at PEP initiation and is identifiable from viral integration dynamics. The two terms multiply but do not co-mingle, addressing the methodological concern that fixed stage efficacies conflate distinct quantities.

Results: route-specific windows

Figure 1 shows the predicted $E_{\text{PEP}}(t)$ curves for both routes derived from the multiscale model at the cited submission SHA. The mucosal window $t_{\text{crit}}^{(m)}(0.05) \approx 60.5$ hours at

$V_0 = 1$ (CV = 0.3) emerges from within-host kinetics and falls within the operational 72-hour CDC mucosal-PEP guideline [1], providing mechanistic context for the clinical threshold. The parenteral window compresses to $t_{\text{crit}}^{(p)}(0.05) \approx 34.5$ hours at $V_0 = 10^3$ (CV = 0.3) — a ≈ 1.75 -fold reduction. The route compression ratio is concordant with the empirical NHP range of $1.5\text{--}2\times$ spanning Tsai et al. [8] (intravenous SIV: complete protection at 24 h, 50% at 48 h) and Otten et al. [9] (intravaginal SHIV: complete protection through 36 h, 50% at 72 h). At the within-host kinetic level, the model captures full protection at early delays (Otten 12 h, 36 h: model $E_{\text{PEP}} = 0.95$ vs. NHP 1.00) but under-predicts protection at later timepoints (Tsai 24 h, 48 h; Otten 72 h). We interpret this gap as evidence of NHP-specific protective contributions not captured by the within-host kinetic framework alone — an empirical limit of the model’s domain rather than a refutation of the compression result. Table 1 summarizes per-timepoint concordance.

Stochastic efficacy analysis

Monte Carlo integration of Equation 1 over source viral load (VL) distributions (Fig. 2A–B) reveals three empirically distinct temporal regimes defined by the 95% credible-interval (CI) width of the stochastic efficacy distribution. *Regime 1* (0–22 h): CI is narrow (<15 percentage points) and efficacy is high (>90% mean for PWID-relevant VL distributions); VL knowledge adds limited decision-relevant precision. *Regime 2* (22–78 h for PWID/acute; 22–88 h for mixed/general distributions): CI width is maximal, peaking at 39.7 percentage points at 49 h for untreated PWID VL distributions (mean $\log_{10} = 4.5$, SD = 1.1, parameterized from NHBS/NHAS surveillance data), reflecting the phase in which integration is complete for some VL draws but not others. *Regime 3* (>78 h): CI collapses as efficacy converges to zero across all VL draws. The regime transition at ≈ 22 h is mechanistically interpretable: it corresponds to the eclipse phase lower bound from Perelson et al. [5] (≈ 0.9 days ≈ 22 h), below which integration cannot yet be complete for any draw. Threshold probability analysis shows that $P(\text{efficacy} < 50\%) = 100\%$ across all distributions at 48 h and that $P(\text{futile})$ — defined as $P(\text{efficacy} < 5\%)$ — reaches 91.8% (PWID untreated) and 98.8% (acute-infection-enriched network) by 72 h (Table 2).

City-stratified analysis

To extend the stochastic model to an empirically grounded, policy-actionable instrument, we parameterized the VL distributions and structural access delays for 34 high-burden US metropolitan areas using AIDSVu 2023 surveillance data. City-level VL distributions were derived via a two-component mixture model (suppressed: mean $\log_{10} = 1.5$, SD = 0.4; unsuppressed: mean $\log_{10} = 4.5$, SD = 1.1), weighted by city-specific viral suppression fraction with an IDU-specific adjustment (0.12 percentage points per percent IDU prevalence, capped at 12 percentage points).

City-level structural delay (population-level access friction)

Structural access delay was estimated from AIDSVu late-diagnosis and linkage-to-care percentages as

$$\Delta t_{\text{structural}} = \beta_1 (\text{LateDx}\% - 10\%) + \beta_2 \max(90\% - \text{Linkage}\%, 0) + \Delta t_{\text{SDOH}} \quad (4)$$

with coefficients $\beta_1 = 1.2$ h/% and $\beta_2 = 0.3$ h/% calibrated to yield ≈ 6 hours at the median city. City-level structural delay is operationalized as a composite indicator

combining AIDSvu late-diagnosis percentage and linkage-to-care percentage according to the functional form specified in Equation 4. The composite is a constructed measure of healthcare access friction; it is not a proxy under proximal causal inference [14], and the analysis does not invoke the identification conditions required for proxy-based estimation [15]. The distinction matters because related work in this research program does invoke proxy-based identification under those conditions.

Expected PEP efficacy at city-specific structural delay ranged from 78.8% (Hartford CT; structural delay 24.4 h; IDU prevalence 23.4%) to 98.3% (Milwaukee WI; 1.9 h), a 19.5 percentage-point spread correlated almost entirely with structural delay rather than community VL (Pearson $r = -0.935$ vs. -0.084 , respectively; Fig. 3). Cities with high IDU prevalence exhibit consistently longer structural delays and lower viral suppression simultaneously, narrowing the PEP window from both directions at once.

Pharmacy-access barrier: envelope-corridor analysis

The structural delay formula (Eq. 4) captures system-level friction up to clinical decision-making but treats prescription writing and first-dose ingestion as effectively coincident. In real-world ED practice this assumption is often false. The CDC 2025 nPEP recommendations explicitly document that pharmacy dispensing practices have sometimes been a barrier to timely nPEP access [1]; a recent quality-improvement initiative at Alameda Health System reported that initial PEP compliance is materially impeded by limited access to pharmacies and cost [23]; national survey data from 587 U.S. EDs show that only 65.3% of EDs offer both HIV testing and PEP to sexual assault survivors [23, 24].

We treat the pharmacy-access barrier $\Delta t_{\text{pharmacy}} \in \{0, 2, 4, 6, 8, 10, 12\}$ hours as an upstream gating event: patients who fail to acquire medication before t_{crit} never accumulate drug, and PK-driven efficacy is defined only conditional on successful acquisition. Under this causal structure, the achievable E_{PEP} envelope is a two-dimensional corridor in $(t_{\text{acq}}, E_{\text{PEP}})$ space, bounded above by perfect-adherence kinetics ($\rho = 1.0$, cliff at $t_{\text{crit}}(\rho=1.0) = 34.0$ h) and below by low-adherence kinetics ($\rho = 0.30$, cliff at 32.0 h). Pharmacy delay does not compress this corridor; it slides each city rightward along $t_{\text{acq}} = \Delta t_{\text{structural}} + \Delta t_{\text{pharmacy}}$ toward the cliff (Section S14).

The effect is concentrated in already-high-barrier cities. Hartford (structural delay 24.4 h) enters at $t_{\text{acq}} = 24.4$ h with $E_{\text{PEP}}^{\text{upper}} = 0.479$ and $E_{\text{PEP}}^{\text{lower}} = 0.098$ — the corridor is already narrow at this position. A pharmacy delay of $\Delta t_{\text{pharmacy}} = 8$ h slides Hartford to $t_{\text{acq}} = 32.4$ h with $E_{\text{PEP}}^{\text{upper}} = 0.175$, still on the corridor. At $\Delta t_{\text{pharmacy}} = 10$ h, Hartford reaches $t_{\text{acq}} = 34.4$ h, exceeding $t_{\text{crit}}(\rho=1.0) = 34.0$ h, and is displaced *off* the corridor; E_{PEP} collapses to zero — not as a probabilistic floor effect, but because the drug never accumulates before viral integration is complete in that regime. The other five high-vulnerability cities in Fig. S1 (San Juan, Jackson, Phoenix, Denver, Milwaukee) remain on the corridor with $E_{\text{PEP}}^{\text{upper}} > 0.96$ even at $\Delta t_{\text{pharmacy}} = 12$ h because their baseline structural delays are 1.3–8.1 h, leaving substantial margin to $t_{\text{crit}}(\rho=1.0)$. Across the 34-city panel, the cohort mean of $E_{\text{PEP}}^{\text{upper}}$ declines from 0.962 at $\Delta t_{\text{pharmacy}} = 0$ to 0.865 at $\Delta t_{\text{pharmacy}} = 12$ h. The substantively interpretable summary is the off-corridor count: at $\Delta t_{\text{pharmacy}} \in \{0, 2, 4, 6, 8\}$ h, zero of 34 cities are displaced past t_{crit} ; at $\Delta t_{\text{pharmacy}} \in \{10, 12\}$ h, one city (Hartford) is displaced. The analytical $F_{\text{access}} \times t_{\text{crit}}$ bound (Eq. 5) therefore remains a separate population-level consistency check; the pharmacy analysis answers a different mechanistic question: which cities are displaced along the PK/PD–integration corridor toward or beyond the parenteral cliff.

The geographic and structural patterning of pharmacy access is non-random: Qato and colleagues demonstrated that pharmacy accessibility and cost-related underuse of prescription medications cluster in low-income Black and Hispanic urban communities [25] — the same populations bearing disproportionate HIV transmission

risk and represented in Hartford’s IDU-dominated parenteral exposure profile (23.4% IDU prevalence). The pharmacy-access barrier therefore compounds rather than averages across the existing structural inequities the city-stratified analysis already documents. Granular data on stocked-on-shelf inventory of HIV PEP regimens at 24-hour retail pharmacies during overnight and weekend hours — particularly at hospital-adjacent locations — is a research gap that limits more precise per-city quantification; systematic ED-pharmacy linkage data, formulary-inclusion audits, and time-stamped dispensing records would allow this barrier layer to be incorporated as a quantified term in the structural-delay decomposition rather than as a sensitivity range. We identify this as a high-priority research direction for HIV prevention implementation science.

The pharmacy-corridor analysis and the $F_{\text{access}} \times t_{\text{crit}}$ inequality answer complementary questions. The corridor analysis is individual- and city-position specific: it asks whether acquisition time remains inside the route-specific PK/PD-integration window. The $F_{\text{access}} \times t_{\text{crit}}$ inequality is population-distributional: it asks how large expected prevention could be under a conservative upper-limit timing assumption. We therefore use the corridor for mechanistic interpretation of access barriers and the F_{access} bound only as an analytical consistency check on the scale of population-level failure.

Population-level implications: cascade estimates, outbreak forecasts, and analytical consistency bound

(a) Cascade-based point estimates

Companion modeling [22] applies an architectural barrier framework ($n = 100,000$ PWID, 5-year horizon) and an independent PWID Monte Carlo simulation to compute the probability that an individual PWID achieves the full prevention cascade ($P(R_0 = 0)$) under structural conditions. Under current US policy: the architectural-barrier model produces $P(R_0 = 0) = 0.003\%$ [95% CI 0.000–0.006%]; the independent PWID simulation produces 0.007%. Decriminalization-only intervention raises this to 0.198–0.814%; full harm reduction raises it to 9.55%. The same architectural model applied to MSM under identical structural assumptions produces 16.3% prevention success — a $\approx 5,400$ -fold population-level disparity between routes.

(b) Outbreak probability forecasting

Independent stochastic outbreak modeling [22] forecasts a 73.8% probability of major HIV outbreak among PWID nationally within 5 years under current policy (95% PSA CI 63.5–82.0%) and 92.7% within 10 years; regional probabilities reach 86.3% (Pacific Northwest) and 78.4% (Appalachia) within 5 years. Median time-to-outbreak is 3 years.

(c) Analytical $F_{\text{access}} \times t_{\text{crit}}$ consistency bound

The timing mismatch can be formalized as a distribution-free inequality. For access delay distribution F_{access} , the expected population-level PEP efficacy satisfies

$$\bar{E}_{\text{PEP}} \leq F_{\text{access}}(t_{\text{crit}}^{(p)}) \varepsilon_{\text{max}} + (1 - F_{\text{access}}(t_{\text{crit}}^{(p)})) \eta \quad (5)$$

(Proposition S6.1). The bound requires only the fraction of the population accessing PEP within the parenteral window — not the full parametric form of F_{access} . It should be interpreted as a conservative analytical ceiling, not as a full population preventability estimator, because it collapses access timing into a scalar distributional term and does

not model the per-patient PK/PD corridor described above. Modeling access delay as log-normal (median 96 h, geometric SD 2.0; modeling choice informed by qualitative discussion in the PWID PEP-access literature [12,13]; the specific median is a defensible parameterization rather than an empirical anchor lifted from primary timing data) yields $F_{\text{access}}(34.5 \text{ h}) \approx 0.070$. With $\eta = 0.05$ and $\varepsilon_{\text{max}} = 0.95$, Equation 5 yields $\bar{E}_{\text{PEP}} \leq 0.070 \times 0.95 + 0.930 \times 0.05 \approx 11.3\%$. Sensitivity analysis across plausible median values $F_{\text{median}} \in \{72, 96, 110, 120\}$ hours yields analytical ceiling estimates of 8.2%–18.0% (Table 3); in all cases, the ceiling remains one to four orders of magnitude above the cascade point estimates in (a), supporting the conclusion that the cascade estimates are not artifacts of an overly pessimistic access assumption.

Convergent finding

These analyses converge at different inferential levels. The cascade model supplies central estimates of realized prevention success; the outbreak model supplies an epidemiologic consequence of those barriers; the PK/PD corridor identifies the mechanism by which structural and pharmacy delays move acquisition time toward the parenteral integration cliff; and the $F_{\text{access}} \times t_{\text{crit}}$ inequality provides a conservative upper-limit check. Together, they support the conclusion that PEP cannot serve as a population-level safety net for PWID under current structural conditions. The analytical bound is deliberately least pessimistic; the cascade-based central estimates are 100–3,000 $\times$ lower and remain qualitatively consistent.

Limitation: F_{access} denominator

F_{access} here is derived from population-wide PEP-access surveillance, which includes exposures from both transmission-competent and virologically suppressed sources. Under U=U [10,11], exposures from suppressed sources have functionally zero transmission risk and are outside the framework’s domain. Restricting F_{access} to transmission-competent exposures only would reduce the relevant denominator and yield a tighter (smaller) F_{access} and a tighter bound. This refinement strengthens, rather than weakens, the conclusion.

Stochastic VL uncertainty and the critical window

The source VL knowledge premium — the gain from knowing source VL precisely versus marginalizing over the distribution — peaks at 2.9 percentage points at 21 h for PWID untreated and 3.8 percentage points for acute-infection-enriched networks, both within Regime 2. This establishes the clinical decision value of rapid source VL determination: it is most actionable precisely in the window where the biological outcome remains uncertain.

Implications

Two conclusions follow. First, the 72-hour PEP guideline is mechanistically grounded but route-specific: it accurately characterizes the mucosal window and should not be generalized to parenteral exposure. The parenteral window is shorter by a factor determined by within-host kinetics, and that factor is concordant with NHP empirical data. Second, for PWID under current structural conditions, PEP cannot function as a population-level safety net. Since PEP and pre-exposure prophylaxis (PrEP) are the only two biomedical prophylaxis modalities for HIV [16,17], and PrEP uptake among PWID remains below 1% of those with indication [13], this population effectively lacks

both options. Long-acting injectable PrEP, demonstrated to provide effective pre-exposure protection against parenteral challenge in NHP studies [18,19], is the modality most capable of bridging this gap — if structural delivery barriers are addressed [20]. The mathematical framework presented here provides the quantitative grounding for that conclusion: the prevention gap for PWID is not primarily a pharmacological problem but a structural one, and its boundary is defined by biology.

All efficacy and access estimates herein are conditional on a transmission-competent exposure; the framework does not apply to U=U exposures (source viral load < 200 copies/mL sustained), for whom transmission risk is functionally zero, and the policy implications drawn here pertain only to populations with non-suppressed source viremia.

Conclusion

PEP is not the safety net for parenteral HIV exposure that it is for sexual transmission. We prove that proviral integration establishes a finite, route-specific prevention window; that this window is approximately 1.75-fold shorter for parenteral than for mucosal exposure; and that structural access delays for PWID systematically consume this window. Across the cascade model, system-level outbreak forecasting, PK/PD corridor analysis, and analytical $F_{\text{access}} \times t_{\text{crit}}$ consistency bound, population-level PEP efficacy for PWID under current structural conditions is estimated at approximately 0.003–0.7% in cascade models and remains below an intentionally conservative analytical ceiling of approximately 11%, with corresponding 92.7% 10-year outbreak probability under current policy. The conclusion is not a statement against PEP for individuals who present in time. It is a quantitative argument that pre-exposure prevention is the primary biomedical tool available for this population — and that current deployment levels are insufficient to close the gap.

Materials and Methods

Mathematical proofs

All formal proofs (Theorem 1, Corollaries 1–2, Lemmas 1–2, Propositions 1–3) are provided in the Supplementary Text. The absorbing-state ODE system, hazard-rate decomposition, monotone systems comparison argument, and population-efficacy bound are each derived in full.

Software environment and reproducibility

All computational analyses were performed in Python 3.11 with NumPy, SciPy, matplotlib, and pandas. Complete environment specifications, including pinned package versions, are provided in `requirements.txt` in the project repository. Monte Carlo simulations used fixed random seeds (`np.random.default_rng(base_seed*1,000,000+k)`) to ensure deterministic reproducibility from the cited commit (SHA 37e27ea). Independent re-execution from the cloned repository at the cited SHA reproduces all numerical claims to full floating-point precision; this was verified prior to submission.

Numerical simulation

The multiscale model (Phase 1 stochastic tau-leaping + Phase 2 deterministic ODE handoff + eclipse-phase fixed-delay buffer) was implemented in Python. Monte Carlo

replicate count ($N = 500$ per (route, V_0 , CV) cell; $\sim 27,000$ realizations across the full parameter sweep) was selected to achieve standard error below 1 percentage point on cumulative integration probabilities at clinical timepoints (24 h, 48 h, 72 h). Sensitivity to N at the canonical operating point ($V_0 = 10^3$ parenteral, CV = 0.3) was tested at $N \in \{200, 500, 1000, 5000\}$; t_{crit} estimates stable to within ± 0.5 h across this range. Integration time T_{int} was computed as the first time $R(t) \geq R^* = 10$ cells/mL. Efficacy curves were computed as in Equation 1 using the PK-driven framework $\varepsilon(Z, t_{\text{PEP}}) = \varepsilon_{\text{drug}}(t_{\text{PEP}}) \cdot p_{\text{clear}}(Z)$ documented in the next subsection, with $p_{\text{clear}}(Z)$ calibrated to recover the hand-set baseline efficacies ($\varepsilon_{\text{max}} = 0.95$, $\varepsilon_{\text{mid}} = 0.50$, $\varepsilon_{\text{min}} = 0$) at perfect adherence. Threshold sensitivity analysis varied $I^* \in [10, 1000]$ cells/mL and $R^* \in [1, 100]$ cells/mL across a 20×20 grid. Stochastic VL integration used 50 time points, 0–120 h, with 10,000 VL draws per time point.

PK-driven PEP efficacy framework

The kinetic-onset framework couples a one-compartment Bateman PK model (with intracellular phosphorylated-metabolite compartments for TFV-DP and FTC-TP, plasma-active kinetics for DTG) to Hill-function pharmacodynamics on the normalized active fraction, combined across drug classes via multiplicative survival. The pharmacologic component $\varepsilon_{\text{drug}}(t_{\text{PEP}})$ is the time-averaged combined suppression over the remaining route-specific window $[t_{\text{PEP}}, t_{\text{crit}}^{\text{route}}]$; the biological component $p_{\text{clear}}(Z)$ is the stage-dependent fraction of trajectory still pharmacologically rescuable, calibrated to recover the hand-set baseline mid-stage efficacy at perfect adherence. Complete model specification, parameter values, validation against the multiscale Monte Carlo baseline (commit 37e27ea), and adherence sensitivity ($\rho \in \{1.0, 0.7, 0.5, 0.3\}$) are documented in Section S13. Implementation: `r3_pk_pd/` in the project repository.

Sex as a biological variable

The within-host kinetic framework operates on cellular and viral dynamics that have not been established to differ by sex at the resolution of the available data; the foundational Perelson 1996 cohort [5] was small (5 patients) and did not establish sex-specific kinetic parameters. The city-stratified analysis pools across sex within AIDSVu surveillance categories, which provide sex-stratified late-diagnosis and viral suppression data; sex-stratified replication of the city analysis is a defensible extension but is not pursued here as the structural-barrier conclusions are robust to within-stratum variation.

City-stratified parameterization

City-level viral suppression, late-diagnosis, and linkage-to-care data were obtained from AIDSVu downloadable datasets for 34 high-burden US metropolitan areas (2023 reporting year). VL mixture parameters were derived using the law of total variance with suppressed component (mean $\log_{10} = 1.5$, SD = 0.4) and unsuppressed component (mean $\log_{10} = 4.5$, SD = 1.1). IDU suppression adjustment: 0.12 percentage points per percent IDU prevalence, capped at 12 percentage points. Structural delay coefficients calibrated to median city yield ≈ 6 hours. Counterfactual analysis assigned all cities a uniform 24-hour delay. Full dataset and derivation scripts are in the code repository.

Data Availability

All source code, simulation outputs, and the numerical claims registry needed to reproduce every result in this manuscript and supplement are available at

<https://github.com/Nyx-Dynamics/Prevention-Theorem> under the v4.0.0 release tag, with a persistent archive at Zenodo (v4.0.0 DOI: 10.5281/zenodo.20559022).

The single-command reproduction pipeline is `reproduce_all_v4.py` at the repository root, which regenerates all main-text figures (Figs. 1–3), supplement tables (Tables S2a–S2c), and CSV artifacts from the multiscale baseline at commit `37e27ea`. The full claims registry mapping every numerical value in the manuscript to its source artifact, locator within that artifact, and verification command is provided as `v3_revision/numerical_claims_v4.csv`.

The 34-city baseline (Table S2b) is derived from publicly available AIDSVu downloadable datasets (2023 release) at <https://aidsvu.org>. Multiscale viral-dynamics realizations and within-host PK/PD parameters are documented in Supplement S5 and the `v3_revision/r3_pk_pd/` subdirectory of the repository.

No human subjects data were used in this study. IRB approval was not required.

References and Notes

1. M.R. Tanner *et al.*, Antiretroviral postexposure prophylaxis after sexual, injection drug use, or other nonoccupational exposure to HIV — CDC recommendations, United States, 2025. *MMWR Recomm. Rep.* **74**(1), 1–56 (2025).
2. T.W. Chun, D. Finzi, J. Margolick, K. Chadwick, D. Schwartz, R.F. Siliciano, In vivo fate of HIV-1-infected T cells: quantitative analysis of the transition to stable latency. *Nat. Med.* **1**, 1284–1290 (1995).
3. J.D. Siliciano, R.F. Siliciano, Enhanced culture assay for detection and quantitation of latently infected, resting CD4⁺ T-cells carrying replication-competent virus in HIV-1-infected individuals, in *Human Retrovirus Protocols* (Humana Press, New Jersey, 2005), vol. 304, pp. 3–16.
4. A.S. Perelson *et al.*, Decay characteristics of HIV-1-infected compartments during combination therapy. *Nature* **387**, 188–191 (1997).
5. A.S. Perelson, A.U. Neumann, M. Markowitz, J.M. Leonard, D.D. Ho, HIV-1 dynamics *in vivo*: virion clearance rate, infected cell life-span, and viral generation time. *Science* **271**, 1582–1586 (1996).
6. B. Ramratnam *et al.*, Rapid production and clearance of HIV-1 and hepatitis C virus assessed by large volume plasma apheresis. *Lancet* **354**, 1782–1785 (1999).
7. A. J. McMichael, P. Borrow, G. D. Tomaras, N. Goonetilleke, B. F. Haynes, The immune response during acute HIV-1 infection: clues for vaccine development. *Nat Rev Immunol* **10**, 11–23 (2010).
8. C.C. Tsai *et al.*, Prevention of SIV infection in macaques by (R)-9-(2-phosphonylmethoxypropyl)adenine. *Science* **270**, 1197–1199 (1995). [Timing data: *J. Virol.* **72**, 4265–4273 (1998)]
9. R. A. Otten, D. K. Smith, D. R. Adams, J. K. Pullium, E. Jackson, C. N. Kim, H. Jaffe, R. Janssen, S. Butera, T. M. Folks, Efficacy of postexposure prophylaxis after intravaginal exposure of pig-tailed macaques to a human-derived retrovirus (human immunodeficiency virus type 2). *J Virol* **74**, 9771–9775 (2000).
10. A.J. Rodger *et al.*, Risk of HIV transmission through condomless sex in serodifferent gay couples with the HIV-positive partner taking suppressive antiretroviral therapy (PARTNER): final results of a multicentre, prospective, observational study. *Lancet* **393**, 2428–2438 (2019).

11. B.R. Bavinton *et al.*, Viral suppression and HIV transmission in serodiscordant male couples: an international, prospective, observational, cohort study (Opposites Attract). *Lancet HIV* **5**, e438–e447 (2018). 486
487
488
12. J.L. Taylor, A.Y. Walley, A.R. Bazzi, Stuck in the window with you: HIV exposure prophylaxis in the highest risk people who inject drugs. *Subst. Abus.* **40**, 441–443 (2019). [Commentary; cited for qualitative discussion of structural barriers, not primary timing data.] 489
490
491
492
13. A.R. Baugher *et al.*, Are we ending the HIV epidemic among persons who inject drugs? Key findings from 19 US cities. *AIDS* **39**(12), 1813–1819 (2025). 493
494
14. E.J. Tchetgen Tchetgen, A. Ying, Y. Cui, X. Shi, W. Miao, An introduction to proximal causal inference. *Statistical Science* **39**, 375–390 (2024). 495
496
15. J. Pearl, *Causality: Models, Reasoning, and Inference*, 2nd ed. (Cambridge University Press, 2009). 497
498
16. R.M. Grant *et al.*, Preexposure chemoprophylaxis for HIV prevention in men who have sex with men. *N. Engl. J. Med.* **363**, 2587–2599 (2010). 499
500
17. P.L. Anderson *et al.*, Emtricitabine–tenofovir concentrations and pre-exposure prophylaxis efficacy. *Sci. Transl. Med.* **4**, 151ra125 (2012). 501
502
18. A. E. Swanstrom, R. J. Gorelick, J. L. Welker, F. Schmidt, B. Lu, K. Wang, W. Rowe, M. W. Breed, K. E. Killoran, J. A. Kramer, D. Donohue, J. D. Roser, P. D. Bieniasz, T. Hatzioannou, C. Pyle, J. A. Thomas, C. M. Trubey, J. Zheng, W. Blair, S. R. Yant, J. D. Lifson, G. Q. Del Prete, Long-acting lenacapavir protects macaques against intravenous challenge with simian-tropic HIV. *eBioMedicine* **95**, 104764 (2023). 503
504
505
506
507
508
19. S. J. Vidal, E. Bekerman, D. Hansen, B. Lu, K. Wang, J. Mwangi, W. Rowe, F. Campigotto, J. Zheng, D. Kato, A. Chandrashekar, J. Barrett, S. Patel, H. Wan, T. Anioke, N. B. Mercado, J. P. Nkolola, M. J. Ferguson, W. J. Rinaldi, C. Callebaut, W. Blair, T. Cihlar, R. Geleziunas, S. R. Yant, D. H. Barouch, Long-acting capsid inhibitor protects macaques from repeat SHIV challenges. *Nature* **601**, 612–616 (2022). 509
510
511
512
513
514
20. A.C. Demidont, Clinical decision support algorithm for LAI-PrEP bridge period navigation at UNAIDS global target scale. *Viruses* (2026; accepted). 515
516
21. P. Emau *et al.*, Post-exposure prophylaxis for SIV revisited: animal model for HIV prevention. *AIDS Res. Ther.* **3**, 8 (2006). 517
518
22. A.C. Demidont, Structural barriers, stochastic avoidance, and outbreak risk in HIV prevention for people who inject drugs. [Companion manuscript; in preparation / preprint DOI to be assigned]. 519
520
521
23. D.A.E. White, M. Jewett, M. Burns, C. Rodriguez, C.M. Pinto, G. Regalado, K. Del Angel, H.J. Larkin, E.S. Anderson, A quality improvement initiative to increase HIV post-exposure prophylaxis treatment for emergency department patients after sexual assault: A pre-post study. *PLoS ONE* **20**(3), e0320690 (2025). doi:10.1371/journal.pone.0320690. 522
523
524
525
526
24. H. Lopez Castillo, D.J. Sheridan, E.E. Vasilevskis *et al.*, HIV post-exposure prophylaxis in the emergency department: An updated assessment and opportunities for HIV prevention identified. *Am. J. Emerg. Med.* **38**(7), 1373–1377 (2020). PMID: 33069548. 527
528
529
530

25. D. Qato, J. Wilder, S. Zenk, A. Davis, J. Makelarski, S. Lindau, Pharmacy accessibility and cost-related underuse of prescription medications in low-income Black and Hispanic urban communities. *J. Am. Pharm. Assoc.* **57**(2), 162–169.e1 (2017). doi:10.1016/j.japh.2016.12.065.

531
532
533
534

Acknowledgments

535

The author thanks the HIV prevention research community whose published work informed model parameterization, and PWID community advocates whose testimony informed characterization of structural barriers.

536
537
538

Funding: No external funding was received.

539

Author contributions: A.C.D.: conceptualization, formal analysis, mathematical proofs, simulation, writing.

540
541

Competing interests: The author reports prior employment with Gilead Sciences, Inc. from January 2020 through November 2024, and held company stock during that period; all Gilead stock was fully divested by December 2024. Employment ended prior to the initiation of this research. Gilead Sciences had no role in study conception, design, data collection, analysis, interpretation, writing, or the decision to submit for publication. The manuscript references lenacapavir, a Gilead product, as one member of the long-acting injectable antiretroviral class; the mathematical conclusions of this work are pharmacologically agnostic and pertain to the class as a whole. The author is the sole owner of Nyx Dynamics LLC, under whose affiliation this research was conducted independently.

542
543
544
545
546
547
548
549
550
551

AI tool disclosure: Computational analyses were conducted in Python (NumPy, SciPy, matplotlib). Large language model assistants (Anthropic Claude, OpenAI ChatGPT) were used to support literature search, code review, manuscript readability, and verification of numerical claims against source code. JetBrains Junie was used for code correction; Zotero AI for reference management. All AI tools were used as assistive technologies; the author retains full responsibility for design, analysis, interpretation, and conclusions. Independent reproducibility of all numerical claims from committed code at the cited SHA was verified prior to submission.

552
553
554
555
556
557
558
559

Supplementary Materials: Supplementary Text S1–S14 with complete proofs, parameter tables, extended methods, NHP concordance details, PK/PD implementation, pharmacy-access corridor analysis, and figure inventory.

560
561
562




Fig 1. Route-dependent emergent t_{crit} from multiscale dynamics, with the three-state within-host framework. (A) Expected PEP efficacy $E_{\text{PEP}}(t)$ as a function of hours from exposure to PEP initiation, comparing mucosal exposure (blue solid; $V_0 = 1$ virion/mL, post-epithelial attenuation, $\text{CV} = 0.3$) and parenteral exposure (red solid; $V_0 = 10^3$ virions/mL, direct intravascular delivery, $\text{CV} = 0.3$). Both curves are computed from $N = 500$ stochastic realizations of the multiscale viral dynamics model. Emergent critical times at the $\eta = 5\%$ efficacy threshold are $t_{\text{crit}}^{(m)} \approx 60.5$ h (mucosal) and $t_{\text{crit}}^{(p)} \approx 34.5$ h (parenteral), yielding a route-dependent compression ratio of $1.75\times$, concordant with the NHP empirical range of $1.5\text{--}2\times$. Markers indicate study-level NHP concordance ($\blacktriangle$ concordant; $\blacktriangledown$ not concordant) at the corresponding timepoints (Methods, §NHP Concordance). Horizontal reference at $\eta = 5\%$; vertical dashed at the CDC 72-hour guideline. The framework applies only to transmission-competent exposures with source $\text{VL} \geq 200$ c/mL (Rodger 2019; Bavinton 2018); U=U exposures are outside scope. (B) Three-state Markov framework for within-host viral dynamics: the susceptible compartment $Z = 0$ transitions to seeded $Z = 1$ at hitting time T_{seed} , then to integrated $Z = 2$ at T_{int} . PEP efficacy is $\varepsilon = \varepsilon_{\text{max}} = 0.95$ in $Z = 0$, $\varepsilon_{\text{mid}} = 0.50$ in $Z = 1$, and $\varepsilon_{\text{min}} = 0$ in the absorbing $Z = 2$ state. Route-dependent inoculum scaling ($V_0 = 1$ for mucosal post-attenuation; $V_0 = 10^3$ for parenteral) propagates through the hitting-time distributions, yielding $T_{\text{int}}^{(p)} \leq T_{\text{int}}^{(m)}$ pathwise (Lemma S5.2) and hence $t_{\text{crit}}^{(p)} \leq t_{\text{crit}}^{(m)}$ at any threshold η (Corollary S5.1). Panel A is the empirical consequence of the inoculum-monotonicity proven in Panel B. Values reproducible from Nyx-Dynamics/Prevention-Theorem at commit 37e27ea via v2_revision/make_figure_1.v2.py.




Fig 2. Stochastic efficacy analysis under source VL uncertainty (Monte Carlo extension, $N = 10,000$). **(A)** Expected PEP efficacy (mean $\pm$ 50% / 95% CI bands) for four source VL distributions (PWID untreated: mean $\log_{10} = 4.5$; PWID mixed treatment: 3.2; general HIV+ community: 3.8; acute-infection-enriched: 5.0) over 0–120 h. Vertical reference lines: 24-h optimal (purple dashed); 72-h CDC guideline (gray dashed). Horizontal reference at 50% efficacy threshold. The framework applies to transmission-competent exposures only (source VL ≥ 200 c/mL); U=U exposures are outside scope (Methods). **(B)** Source VL probability densities on a $\log_{10}$ axis from LLoD (~ 20 c/mL) to 10^7 c/mL. **Vertical green dotted line at LLoQ = 200 c/mL** marks the U=U operational threshold (per PARTNER2; Rodger 2019; Bavinton 2018); below this VL, transmission risk is functionally zero and the framework is not applicable. Vertical orange dotted line at 10^4 c/mL and red dotted line at 10^5 c/mL mark inflection points used in the route-compression analysis. Panel B is the natural location for the U=U threshold marker; Panel A's hours-from-exposure axis precludes a literal VL-axis band. **(C)** Probability of sub-therapeutic PEP, $P(\text{efficacy} < 50\%)$, by hours-from-exposure for each VL distribution. By 48 h, $P(< 50\%) = 100\%$ across all distributions regardless of VL profile. **(D)** VL knowledge premium — the absolute gain in expected efficacy from knowing the source VL at PEP initiation versus marginalizing over the VL distribution — for the two most clinically actionable distributions (PWID untreated; acute-infection-enriched network). Premium peaks within Regime 2 (~ 18 – 24 h) at 2.85–3.65 pp, where source VL testing is most decision-relevant.




Fig 3. City-stratified PEP efficacy across 34 US metropolitan areas. (A) Expected PEP efficacy at city-specific structural delay for 34 cities (AIDSvu 2023), sorted by efficacy. Color: barrier category (green: low <8 h; orange: moderate 8–30 h; red: high >30 h). (B) Structural delay vs. expected efficacy (Pearson $r = -0.935$); bubble size proportional to IDU prevalence. Labeled outliers: Hartford CT (24.4 h, highest structural delay, 23.4% IDU); San Juan PR (13.3 h, highest community VL burden, 22.2% IDU); Milwaukee WI (1.9 h, lowest barrier). (C) Joint VL burden × structural delay risk surface; color encodes expected efficacy. (D) Actual vs. 24-hour counterfactual efficacy by city. For 33/34 cities, actual efficacy exceeds counterfactual because structural delays are currently shorter than 24 h; Hartford is the exception. The 19.5 pp range across cities is almost entirely attributable to structural delay ($r = -0.935$) rather than community VL ($r = -0.084$). *Scope notes:* (i) AIDSvu surveillance data are aggregated across sex; the city-stratified analysis pools across sex within AIDSvu categories (see Methods §Sex as a biological variable). (ii) AIDSvu late-diagnosis and viral suppression data are aggregated across mode of HIV acquisition and cannot be reliably subset to U=U status; this figure pools all source-VL strata in the surveillance denominator, consistent with the Methods specification of F_{access} .

Tables

Table 1. NHP PEP timing data by route and per-timepoint model concordance. Multiscale model output computed at the canonical operating point ($CV = 0.3$); see numerical claims summary CSV for reproducibility provenance.

Route	Study	Delay	NHP protection	Model E_{PEP}	Per-timepoint
Parenteral (IV)	Tsai 1998 [8]	24 h	1.00	0.50	not concordant
	Tsai 1998	48 h	0.50	0.00	not concordant
Mucosal (vaginal)	Otten 2000 [9]	12 h	1.00	0.95	concordant
	Otten 2000	36 h	1.00	0.95	concordant
	Otten 2000	72 h	0.50	0.016	not concordant

Compression-ratio concordance. The model’s mucosal/parenteral t_{crit} ratio at $\eta = 0.05$ is $60.5/34.5 = 1.75\times$, concordant with the empirical NHP range of $\sim 1.5\text{--}2\times$ spanned by Tsai 1998 (parenteral half-protection at 48 h) and Otten 2000 (mucosal half-protection at 72 h: $72/48 = 1.50\times$). **Per-timepoint discordance.** The within-host kinetic model captures full protection at early Otten timepoints but under-predicts protection at the late Otten and both Tsai timepoints; we interpret this as evidence of NHP-specific protective contributions (innate immunity, eclipse-phase containment) not captured by the within-host kinetic framework alone. The compression result and the absolute-timepoint result are independent: the compression result is robust to the model’s domain limitations.

Table 2. Stochastic efficacy metrics by VL distribution at key time points.


Distribution	24 h mean	36 h mean	48 h $P(< 50\%)$	48 h $P(\text{futile})$	72 h $P(\text{futile})$	Peak CI (time)
PWID untreated	65.5%	44.6%	100%	9.1%	91.8%	39.7 pp (49 h)
PWID mixed Tx	75.4%	52.8%	100%	3.1%	56.5%	40.1 pp (54 h)
General community	71.2%	49.2%	100%	3.5%	76.0%	39.8 pp (54 h)
Acute-enriched	61.2%	41.3%	100%	13.8%	98.8%	37.4 pp (49 h)

$P(\text{futile}) = P(\text{efficacy} < 5\%)$. CI = peak 95% credible interval width in percentage points (pp).
 $P(< 50\%)$ = probability that efficacy falls below majority protection. At 48 h, $P(< 50\%) = 100\%$ across all distributions regardless of VL profile.

Table 3. Analytical $F_{\text{access}} \times t_{\text{crit}}$ consistency-bound sensitivity (Equation 5, $\eta = 0.05$, $\varepsilon_{\text{max}} = 0.95$, $t_{\text{crit}}^{(p)} = 34.5$ h).

F_{access} parameterization	$F_{\text{access}}(34.5 \text{ h})$	Analytical ceiling for $\bar{E}_{PEP}$
Log-normal median 72 h, GSD 2.0 (JID-consistent)	0.144	$\leq 18.0\%$
Log-normal median 96 h, GSD 2.0 (canonical)	0.070	$\leq 11.3\%$
Log-normal median 110 h, GSD 2.0	0.047	$\leq 9.2\%$
Log-normal median 120 h, GSD 2.0	0.036	$\leq 8.2\%$

Reading the bound. The $F_{\text{access}} \times t_{\text{crit}}$ inequality is a conservative analytical ceiling, not a full population preventability estimator. Cascade-based point estimates (Section: Population-level implications) yield 0.003–0.7% under current US policy, one to four orders of magnitude below this ceiling. The qualitative conclusion is supported primarily by the cascade estimates and the PK/PD corridor analysis; the F_{access} value functions as a consistency check and depends on the modeled access-delay median.



B. Three-state within-host model & route-dependent inoculum




Figure 2

[Click here to access/download;Figure;Figure_2_Stochastic_Efficacy_VL_Uncertainty.png](#)

PEP Efficacy Under Source Viral Load Uncertainty






Figure 3

[Click here to access/download;Figure;Figure_3_City-Stratified_PEP_Efficacy.png](#)



Selected City Comparison: Structural Determinants of PEP Window Access







[Click here to access/download](#)

Supporting Information


[finite_windows_v4_supplement_plos_comp_bio.pdf](#)





Click here to access/download

Other
references.bib







[Click here to access/download](#)

LaTeX Source File (TEX file)

`finite_windows_v4_main_plos_comp_bio.tex`





[Click here to access/download](#)

LaTeX Source File (TEX file)

[finite_windows_v4_supplement_plos_comp_bio.tex](#)