

Guidance and surveillance capability for bystander *Staphylococcus aureus* tetracycline resistance under doxy-PEP: a three-stream evidence audit

Adrian C. DEMIDONT*

Nyx Dynamics, LLC, Fairfield, Connecticut, USA

Running title: Surveillance and doxy-PEP bystander *S. aureus* resistance

*Corresponding author. Nyx Dynamics, LLC, Fairfield, Connecticut, USA. Tel: +1-215-901-2366. E-mail: acdemidont@nyxdynamics.org

ORCID: 0000-0002-9216-8569

Synopsis

Background. Doxycycline post-exposure prophylaxis (doxy-PEP) prevents bacterial sexually transmitted infections and has scaled into guidance internationally. Doxycycline also selects tetracycline resistance beyond its target, and *Staphylococcus aureus*—a commensal and pathogen for which doxycycline is one of a limited number of oral options—is the natural bystander.

Objectives. To determine whether the guidance and surveillance deployed around doxy-PEP could follow the bystander-resistance signal the pivotal trial's final analysis resolved, rather than to estimate a population effect.

Methods. A reproducible, public-data audit of three evidence streams—clinical trials, governmental guidance and deployed US surveillance—each scored against one yardstick: the within-exposed relative risk it could detect, benchmarked to an empirical colonisation estimate.

Results. The final randomised DoxyPEP analysis reported a 3.89-fold higher hazard of incident doxycycline-resistant *S. aureus* under doxy-PEP (95% CI 1.42–10.68; 68 of 393 versus 5 of 163 incident events), with no effect on colonisation clearance. Yet 0 of 6 outbreak-matched governmental guidelines (0 of 10 coded) required *S. aureus* monitoring, and 0 of 12 established

US surveillance systems linked doxy-PEP exposure to an *S. aureus* tetracycline phenotype at a common population denominator. Under realistic exposure density and isolate volumes, an ordinary population system would need an implausible within-exposed effect ($RR \approx 6\text{--}28$) to move a rate.

Conclusions. The upstream trial generated information the downstream architecture is not built to receive. Following the signal requires sentinel surveillance linking doxy-PEP exposure to the *S. aureus* tetracycline phenotype at a common denominator. No population-level causal claim is made.

Introduction

Doxycycline post-exposure prophylaxis (doxy-PEP) substantially reduces bacterial sexually transmitted infections (STIs) and has moved quickly from trial result into guidance at every level, from the World Health Organization through national and municipal health departments.^{1,2} Every dose is selective pressure, and not only on the organisms it targets. *Staphylococcus aureus* is the consequential bystander: it colonises the anatomical sites these trials swabbed and carries mobile tetracycline resistance determinants.³ For two decades, community MRSA in these networks has been documented not as steady-state carriage but as clustered, outbreak-pattern transmission—the USA300 lineage moving through sexual networks in San Francisco, Tokyo and Chicago^{4,5}; the organism colonising these networks is the same one carried on skin community-wide, so resistance selected within the network could, if transmitted, extend beyond it. Tetracycline-resistant USA300 was in fact documented in an MSM-predominant community health population two decades ago, although by the doxycycline-sparing *tet(K)* mechanism, with those isolates remaining doxycycline-susceptible.⁶ Doxycycline is one of a limited number of oral options for outpatient skin and soft-tissue infection, a choice often made empirically before any methicillin result returns,^{7,8} so tetracycline susceptibility can matter clinically before methicillin status is known.

Clinicians who prescribe doxy-PEP have registered the concern directly: in a survey of Italian STI-clinic dermatologists, 91.7% named methicillin-resistant *S. aureus* (MRSA) selection

as a worry, tied with gonococcal resistance as the highest-rated,⁹ and a systematic review of 1840 healthcare professionals across four countries likewise placed antimicrobial resistance among the leading concerns about doxy-PEP.¹⁰ The trial literature has now answered a version of it. In the final DoxyPEP analysis, among participants free of doxycycline-resistant *S. aureus* at baseline, doxy-PEP carried a 3.89-fold higher hazard of incident doxycycline-resistant *S. aureus* (95% CI 1.42–10.68; 68/393 versus 5/163; $P = 0.0044$), while the hazard of clearing colonisation did not differ between arms (HR 1.01, 0.69–1.46).¹¹ The trial’s authors called for public-health surveillance of antimicrobial resistance (AMR) in *S. aureus* and other commensals. One national programme has since taken up that call: the UK Health Security Agency’s 2026 doxy-PEP monitoring plan for England proposes tetracycline-resistance surveillance in bystander pathogens, naming *S. aureus*.¹²

That call defines the present question. We ask not whether doxy-PEP drives population-level resistance—a claim we do not make—but whether the guidance and surveillance deployed around doxy-PEP could follow the signal the trial resolved, at the population level where the clinical stake lies. The clinically decisive axis is tetracycline (doxycycline) susceptibility used in empirical oral therapy, not the methicillin axis around which staphylococcal surveillance is organised. We audit the three places an answer would have to come from—the trials, the guidance and the deployed surveillance—each against a single detectability yardstick.

Materials and methods

Ethics

This study analysed publicly available guidance, surveillance documentation and aggregate published data; it did not involve individual-level human-participant data, and institutional review board approval was therefore not required.

Design

The study is a documentary and quasi-computational audit assembled in three streams, collected and analysed in sequence: clinical trials (Stream A), the governmental guidance built

on them (Stream B) and the population surveillance meant to monitor them (Stream C). No individual-level or restricted data were used; every input is public. Sources were located, read, coded with page/section/table locators and SHA-256-pinned before analysis; analysis decisions that could go either way were date-logged before results were known; and every figure and table regenerates from a single command over a test-gated pipeline archived with a citable DOI.¹³ A large language model assisted with writing and analysis engineering; the author verified every numerical value and quotation against primary sources and takes full responsibility (see Acknowledgements).

Detectability yardstick

Each stream is scored on one commensurable quantity: the within-exposed relative risk (RR) it could detect. The empirical benchmark is Soge *et al.*¹⁴ (tetracycline-resistant *S. aureus* colonisation, 18% versus 8% in doxy-PEP users with >3 doses/month versus non-users; $RR \approx 2.25$); a cross-organism gonococcal $RR \approx 1.42$ is carried as a flagged, more conservative secondary. Because the trial's within-exposed hazard ratio (3.89) sits well above this band, the detectability verdict below is robust to which anchor is used: the relative risk a population system would need (RR 6–28, Results) exceeds the cross-sectional benchmark across the realistic parameter range, and exceeds the trial's own hazard ratio except at the limiting dose assumption that every user takes >3 doses/month (dose-sensitivity in the Supplementary data). Hazard ratio and risk ratio are in any case different estimands, so this is a magnitude reference, not an equivalence.

Sources and coding

The author identified, retrieved and screened all source documents for eligibility; each eligible document was then coded from full text against a prespecified schema, every value carrying a page, section or table locator. For the guidance corpus (Stream B), a first LLM pass coded all 10 documents against the schema; a second, blinded LLM pass—a different interface (chat rather than the command-line agent) but the same model family—independently re-coded a 5-document subsample (25 field-codings) for reliability, and disagreements were resolved by the author against the source. Per-field agreement is reported as raw concordance, which measures

the reproducibility of the coding protocol rather than its accuracy; because the passes share a
model family their errors are not independent, so a Cohen's κ is given for context only (Lim-
itations). The surveillance enumeration (Stream C) was coded by a single pass; its inclusion
and exclusion list, with a stated reason per system, is given in the Supplementary data and re-
flects systems as deployed as of August 2026. All coding prompts and outputs are in the public
repository.

Stream A—trials

For each primary-trial *S. aureus* arm comparison (first-party published data only), we com-
puted, from arm sizes and observed control rate, the exact (Fisher) power to detect the exter-
nally fixed benchmark effect and the minimum detectable RR at 80% power; fixing the effect
externally makes this a design-based sensitivity analysis rather than observed power. The *S. au-*
reus assay is doxycycline throughout (E-test, MIC $\geq$ 16 mg/L). We tabulated how the resistance
result was reported across trial venues and secondary syntheses (numerator, denominator ba-
sis, conditioning set and terminology). The final trial performed the appropriate participant-
level time-to-event test itself; we cite that randomised result directly and did not re-estimate
it. Cross-sectional carriage cannot resolve the clustered transmission structure of *S. aureus*
in these networks; the supporting dispersion, trend-test degradation and outbreak-shape anal-
yses are MRSA-scoped (MRSA being a subset, not the analytic unit) and are reported in the
Supplementary data (available as Supplementary data at JAC Online).

Stream B—guidance

To make the denominator principled, we anchored it to the same clustering evidence used in
Stream A: US jurisdictions with documented community-associated MRSA/USA300 transmis-
sion in men who have sex with men.^{4,5} We took governmental public-health guidance only
(clinic and provider-organisation documents excluded, with reasons). Each document was
coded from full text, one record per document, every finding carrying a locator, for whether
it requires, suggests or omits *S. aureus* monitoring and whether it names, counsels on or mea-
sures the organism (coding provenance above). An independence check separated genuinely

independent declines from documents that defer to, or are adapted from, one another. 100

Stream C—surveillance 101

We examined a prespecified set of 12 established US surveillance and data systems (federal, state or metropolitan) plausibly capable of capturing, at the required level, either doxy-102
PEP/doxycycline exposure or *S. aureus* AMR phenotype, and coded each on three capabil-103
ities: exposure-side capture, phenotype-side capture (of the *S. aureus* tetracycline unit, not104
the MRSA subset) and the join, recording the geographic resolution of each side. A sup-105
porting analysis asked whether, even at a common denominator, the exposed subgroup is106
dense enough to move a population rate. The exposed population is indexed on the com-107
bined PrEP and people-living-with-HIV male count (primary specification; the trial enrolled108
both, AIDSVu 2024 state PrEP and prevalence), so the dilution fraction is $f = ((\text{male-PrEP} +109
(\text{male-PLWH}) \text{ density} / 10^5) \text{ uptake } d \kappa$, where d is the share of users taking >3 doses/month (the110
subgroup carrying the benchmark effect) and κ the isolate-enrichment factor; f is population-111
independent because it tracks exposure density, not headcount, and the minimum detectable112
change in a proportion near baseline R_0 inverts to $RR_{\text{needed}} = 1 + \text{MDE} / (f R_0)$. Parame-113
ters were not uniformly generous: isolate volume was set generously high (favouring detec-114
tion), while the realistic under-sampling ($\kappa = 0.5$) and dose-threshold ($d = 0.5$) factors bound115
against it. Both ranges and the resulting RR_{needed} are reported; full grids and the metropolitan116
break-even are in the Supplementary data. 117
118

Scope 119

The corpus and the surveillance enumeration were restricted by design: Stream C is a prespec-120
ified set of US systems, and Stream B is US governmental guidance with the World Health121
Organization included only as the international apex reference. The resulting counts describe122
US architecture, not a global census. 123

Results

124

Stream A: trial measurement and reporting of the *S. aureus* endpoint

125

When the analysis matched the question, the signal appeared. The final DoxyPEP analysis ran the participant-level test the interim had not—time to first detection of doxycycline-resistant *S. aureus* by arm, standard-of-care censored at crossover, by Cox proportional hazards—and returned HR 3.89 (95% CI 1.42–10.68; $P = 0.0044$), with colonisation clearance unchanged (HR 1.01).¹¹ The 2023 interim, by contrast, was presented as reassuring on a cross-sectional resistance fraction and was underpowered for any externally specified effect (0 of 4 comparisons powered; minimum detectable RR 3.8–5.1 against the benchmark 1.42–2.25). The hazard ratio is not comparable to that cross-sectional minimum: the estimand, not the sample size alone, became adequate to the question.

126

127

128

129

130

131

132

133

134

Two measurement features of the interim record persist and matter downstream. First, a unit mismatch: the trial’s data table is headed “phenotypic resistance to the tetracycline antibiotic class” while its methods define the *S. aureus* assay as doxycycline, and the federal guideline then describes the doxycycline result as “tetracycline resistance in *S. aureus*”.² Because efflux and ribosomal-protection mechanisms differ in whether doxycycline still works (below), an endpoint scored as “the tetracycline class” does not answer the treatment-relevant question. Second, the signal was reported over differing, incompletely documented denominators and then propagated into secondary reporting under further contrasts (Table 1): the resistant numerator changes from 5 to 16 to 28 and the denominator basis shifts from colonised carriers to all-swabbed, while the percentage travels more reliably than the denominator, contrast and terminology needed to interpret it. The individual-level data that could reconcile these first-party reports are not publicly available.¹ Even the intensively surveilled gonococcus was phenotyped in fewer than one diagnosis in five (resistance testing unavailable for 83%, 212/256).¹⁵

135

136

137

138

139

140

141

142

143

144

145

146

147

Across trials the phenotype was measured on incompatible terms: DoxyPEP measured doxycycline resistance by E-test over two denominators; DuDHS measured it by disc diffusion in single-digit carriers; DOXYVAC measured only MRSA carriage—a methicillin, not a tetracycline, quantity, and one that itself rose within the doxy-PEP arm (1.8% to 9.9%) while being

148

149

150

151

reported as a between-arms null.¹⁶ A pooled estimate is not defensible, and the heterogeneity is itself a finding.

The estimand shift did not propagate downstream: no guidance document in the corpus cites the final time-to-event analysis or any hazard ratio, and the two documents revised after its online publication in March 2025—WHO (2026) and San Francisco (May 2026)—retained a cross-sectional or generic frame. Non-citation is telling here because this corpus demonstrably transmits numbers when it has them: CDC carried the cross-sectional 5%→13% figures and WHO cites Soge’s colonisation estimate.¹⁴

Stream B: *S. aureus* monitoring requirements in governmental guidance

The governmental guidance shows a consistent within-document asymmetry: the efficacy question received a graded, systematic evidence review, while the AMR-harm question—explicitly including resistant-pathogen development—received an ungraded review in the same document. Of the six documented-outbreak jurisdictions with governmental doxy-PEP guidance (San Francisco, Chicago, New York City, Boston/Massachusetts, Los Angeles County and San Diego), 0 require or suggest *S. aureus* monitoring. The decline is not an artefact of copying: four jurisdictions are independently authored, one (Los Angeles County) is adapted from San Francisco and one (Massachusetts) from an external template. Nor is it for want of naming the organism—three of the six name *S. aureus* specifically, two invoke only a generic “microbiome” risk and one is silent; even where named, none measure it. Across the full coded corpus (10 documents: CDC, WHO and eight US jurisdictions) the result is identical: 0 instrument the bystander. Double-coding of a 5-of-10-document subsample (25 field-codings) gave 92% raw agreement (23/25); the one field with meaningful disagreement (bystander treatment; $\kappa = 0.58$) is reported unrounded. Providers are nonetheless required to counsel patients about resistance in commensal organisms—a harm no system measures.

Stream C: surveillance capability and exposure–phenotype linkage

Following the signal requires one system to carry, at a common population denominator, both who is exposed to doxy-PEP and what tetracycline phenotype *S. aureus* carries. Across this

prespecified set of twelve systems, none does (Figure 1). Two systems capture exposure at a population denominator (AIDSVu and CDC AtlasPlus, as a pre-exposure prophylaxis proxy, not doxy-PEP itself),¹⁹ and zero capture *S. aureus* tetracycline resistance at a population denominator: no join is possible because one half does not exist.

The absence is structural, not incidental. Wherever *S. aureus* is surveilled at a population denominator the reported axis is methicillin: the one population-based platform, Active Bacterial Core surveillance/Emerging Infections Program (ABCs/EIP), reports invasive MRSA; the National Healthcare Safety Network reports MRSA events; and the only routine source of *S. aureus* tetracycline susceptibility, the facility cumulative antibiogram, is non-standardised and carries no population denominator, because CLSI M39 mandates the MRSA/MSSA split and makes tetracycline a selectively suppressible supplemental agent. The National Antimicrobial Resistance Monitoring System excludes *S. aureus* altogether, and tetracycline-resistant *S. aureus* is nationally notifiable nowhere. A system could be built—most directly by adding *S. aureus* tetracycline susceptibility to ABCs/EIP, which already holds the denominator—but none is deployed.

Nor could an ordinary population system recover the missing side by collecting more isolates. Indexing the exposed population on the combined PrEP and people-living-with-HIV male count—the primary specification, because the trial enrolled both (Materials and methods)—the realistic-cell within-exposed effect required to move a population rate runs to $RR \approx 6\text{--}28$ across the design-effect range (Figure 2); the more restrictive PrEP-only indexing gives $RR \approx 14\text{--}64$, which overstates infeasibility. Either way it is far above any plausible effect. Detection appears in a single grid cell that maxes every generous assumption at once (RR_{needed} 1.03, or 1.79 under single-comparison inference; parameters in the Supplementary data). The verdict turns on quantities no deployed system reports—exposure density, isolate enrichment and dosing frequency. Against the more conservative cross-organism benchmark (RR 1.42), the best metropolitan cell needs about twice the combined male exposure density of the densest US geography (Washington, D.C.), and realistic cells far more. The only configuration that restores sensitivity is to sample *S. aureus* directly from doxy-PEP-exposed persons, converting surveillance into a sentinel cohort.




Figure 1: Schematic summary of the 12-system surveillance audit: no deployed US system links doxy-PEP exposure to *S. aureus* tetracycline resistance at a common population denominator. Exposure is captured only as a PrEP proxy (AIDSVu, CDC AtlasPlus); the *S. aureus* tetracycline phenotype is captured at a population denominator nowhere, because the one population-based platform reports the methicillin (MRSA) subset. Of the 12 systems assessed, 0 join exposure to phenotype.

Alt text: A schematic summarising the 12-system surveillance audit: exposure is captured only as a PrEP proxy and the *S. aureus* tetracycline phenotype at a population denominator nowhere, so no system links exposure to phenotype.

Alt text: A logarithmic relative-risk axis with a shaded benchmark band at 1.42 to 2.25; the trial hazard ratio 3.89 sits above the band, surveillance detectability thresholds of about 6 to 28 (primary combined-denominator specification; 14 to 64 under the more restrictive PrEP-only specification) sit far to the right, and guidance has no threshold.

The three streams are summarised against the yardstick in Table 2; a PubMed literature-coverage analysis of the bystander organism (361 papers, of which few name or measure it) is reported in the Supplementary data (S5).




Figure 2: One axis—within-exposed relative risk (log scale)—carries all three streams against the plausible benchmark band (RR 1.42–2.25). The trial’s resolved effect (HR 3.89) clears the band; guidance requires no *S. aureus* measurement so no detection threshold exists; and deployed surveillance would need RR $\approx$ 6–28 to move a population rate (combined PrEP+PLWH exposed denominator; the more restrictive PrEP-only indexing gives 14–64). The trial HR is shown as an empirical magnitude reference only; it is not numerically interchangeable with the surveillance RR threshold because the estimands differ. The signal is above the line while the systems built to follow it are not.

Discussion

218

The pivotal trial has now resolved the bystander question it was long thought unable to reach: 219
a significant increase in incident doxycycline-resistant *S. aureus* under randomised exposure, 220
with no matching effect on carriage, and a call from its authors for surveillance of exactly this.¹¹ 221
Across the two systems that would have to receive it—guidance and deployed surveillance— 222
they cannot: guidance instruments nothing (0/6; 0/10) and no deployed system joins exposure 223
to the *S. aureus* tetracycline phenotype at a common denominator (0/12). 224

The sharper reading is that the trial’s measurement system evolved to meet its own question 225
while the implementation system did not: the same endpoint moved from a cross-sectional 226
fraction read as reassurance to a participant-level HR of 3.89, yet guidance and surveillance did 227
not follow²⁰—a demonstrated signal handed to systems built to measure something else. 228

Tetracycline resistance mechanisms and the phenotype unit

229

Staphylococcal surveillance is standardised around methicillin, yet the treatment-relevant axis
here is tetracycline susceptibility, used empirically before methicillin status is known; the two
are orthogonal. So is the molecular basis of tetracycline resistance: *tet(K)* efflux leaves doxy-
cycline largely usable whereas *tet(M)* ribosomal protection can compromise it, and the trial
resolved incident doxycycline resistance without resolving which.^{3,8} Of 823 invasive USA300
isolates in national surveillance, 72 (9%) were tetracycline-resistant, yet 69 were doxycycline-
susceptible *tet(K)* and only 3 doxycycline-resistant *tet(M)*²¹—a single breakpoint reporting
one number for two clinically opposite situations. An exposure cohort shows the same blind-
ness: doxycycline prophylaxis in 168 military personnel left overall tetracycline-class resis-
tance in *S. aureus* unmoved while *tet(M)* genes were significantly more common in the exposed
($p=0.031$)²²—a military antimalarial cohort, not doxy-PEP. An inherited surveillance variable
can stay valid for its original question while being insufficient for the one now imposed on it;
we do not forecast which mechanism is spreading, only that the inherited variable cannot say.

230

231

232

233

234

235

236

237

238

239

240

241

242

Instrument requirements for population monitoring

243

Following the signal to the population requires an instrument the deployed architecture does
not provide: sentinel sampling of *S. aureus* from doxy-PEP-exposed individuals, which fixes
the exposed fraction and the denominator; sampling of the oropharyngeal and peri-anal sites,
since nares-only screening under-detects colonisation by about 20% in this population²³; re-
sistance reported per carrier and by mechanism (*tet(M)* versus *tet(K)*) rather than as a pooled
tetracycline-class rate; dosing frequency rather than prescription status; a population-denominator
feed of *S. aureus* tetracycline susceptibility, most directly by adding it to the platform that al-
ready tracks invasive MRSA and joining it to exposure on geography; and access to individual-
level trial data. None is exotic; none is in place at scale.

244

245

246

247

248

249

250

251

252

England's national doxy-PEP monitoring plan is an instructive out-of-frame case: postdat-
ing our audit and outside a US corpus, it nonetheless proposes the phenotype half we find
nowhere—tetracycline resistance in bystander pathogens including *S. aureus*—through routine
laboratory AMR surveillance.¹² Yet its specimens come mainly from hospitalised patients; it

253

254

255

256

records neither doxy-PEP use nor sexual orientation; and its exposure contrast is therefore a
male-to-female resistance ratio in an 18–45 age band, an STI-surveillance conditioning set
carried onto an organism that does not respect it. The same programme already runs exposure-
linked sentinel surveillance that records recent doxy-PEP use—but only for the gonococcus:
the instrument exists, and inheritance points it at the in-category organism.

These failures share a structure. A guideline or a surveillance system carries in its sam-
pling frame, cadence, coding, denominator and unit the organism and question it was built
for. The doxy-PEP apparatus, built to measure gonococcal efficacy, inherits for the bystander
commensal a denominator the intervention itself moves, a phenotype label spanning two mech-
anisms and a surveillance architecture standardised around methicillin. We call this *measure-
ment inheritance*: a downstream system retains the dimensions—estimand, denominator, unit,
cadence, linkage, measurement responsibility—appropriate to the upstream question but not
those the downstream question requires, so the question becomes unanswerable not for want
of data but for want of an inherited dimension. Built from familiar parts but concerned with
how they couple *across* stages of an evidence chain,^{24,25} the downstream gap is not closable by
a better analysis of the trial’s data—that analysis has been done, and it is significant—but is a
property of the guidance and surveillance instruments.

Doxy-PEP’s benefit accrues to the recipient; the tetracycline selection cost falls on peo-
ple outside the prescribing decision—outpatients with skin infections and, at the sharp end,
dialysis, surgical and oncology patients for whom an oral anti-staphylococcal agent is no con-
venience. A negative externality that no deployed instrument can price is not thereby small; it
is unweighable. The wider distributive argument—that the two sides are observed on unequal
terms, a property we term *distributive observability*—is developed in the Supplementary data
(S6).²⁶

Limitations

We make no population-level causal claim and no transmission claim: whether the individual-
level signal propagates into a population externality is precisely what no deployed system can
determine. The trial’s own randomised result is significant and we accept rather than dispute

it; our earlier aggregate reconstruction of the interim is retired in its favour. Colonisation is not
infection, and the colonisation–infection link is itself unmeasured here. Much of the evidence
for how *S. aureus* clusters and moves through these networks exists only for its MRSA subset,
so where our clustering analysis relies on MRSA we mark it as such—an instance of the in-
herited limitation, not an exception to it. Finally, the trial reaches only a roughly twelve-month
horizon; persistence, dose–response and mechanism remain open. Our coding carries two ac-
knowledgeed asymmetries. The guidance corpus was double-coded on a 5-document subsample
by two passes run through different interfaces—a command-line agent and a chat interface—
but the same model family, so their errors are not independent; the 92% concordance measures
the reproducibility of the coding protocol, not its accuracy. The surveillance enumeration was
coded by a single pass. Consistent with a falsification standard set in advance, the inheritance
claim fails if a doxy-PEP guideline comes to require *S. aureus* monitoring, or if a deployed
system comes to link doxy-PEP exposure to an *S. aureus* tetracycline phenotype at a common
population denominator; a proposed national programme does not yet meet this standard.¹²

Conclusions

The randomised instrument generated a bystander-resistance signal that the guidance and surveil-
lance built around doxy-PEP are not instrumented to follow. The gap is one of measurement
architecture, not of missing data, and it is closable—by sentinel surveillance that links doxy-
PEP exposure to the *S. aureus* tetracycline phenotype at a common denominator. Until it is
built, the population question the trial’s authors posed cannot be run.

Acknowledgements

A large language model was used for writing and analysis-engineering assistance beyond copy-
editing. The scientific argument, the reading of primary sources, the model specification and all
conclusions are the author’s own; the author reviewed all generated content, independently ver-
ified every numerical value and quotation against primary sources and the regenerable analysis
outputs, and takes full responsibility. No artificial-intelligence tool is credited as an author.

Funding

311

This work received no specific grant from any funding agency in the public, commercial or not-for-profit sectors. It was conducted as an unfunded investigator-initiated study.

312

313

Transparency declarations

314

A. C. D. reports former employment at Gilead Sciences (2020–2024) with prior ownership of corporate stock, all divested by 2024; Gilead played no role in this work. A. C. D. is sole owner of Nyx Dynamics, LLC. No other conflicts to declare. All data are public (AIDS Vu; published trials and guidelines) and all analysis code is openly available (<https://github.com/Nyx-Dynamics/doxyep-bystander>) and archived at Zenodo (doi:10.5281/zenodo.22725070),²⁷ regenerating every figure and table by a single command; every coded source is SHA-256-pinned. No restricted-access data were used.

315

316

317

318

319

320

321

References

322

[1] Luetkemeyer AF, Donnell D, Dombrowski JC, et al. Postexposure doxycycline to prevent bacterial sexually transmitted infections. *N Engl J Med*. 2023;388:1296-306.

323

324

[2] Bachmann LH, Barbee LA, Chan P, Reno H, Workowski KA, Hoover K, et al. CDC clinical guidelines on the use of doxycycline postexposure prophylaxis for bacterial sexually transmitted infection prevention, United States, 2024. *MMWR Recomm Rep*. 2024;73(2):1-8.

325

326

327

328

[3] Grossman TH. Tetracycline antibiotics and resistance. *Cold Spring Harb Perspect Med*. 2016;6(4):a025387.

329

330

[4] Diep BA, Chambers HF, Graber CJ, Szumowski JD, Miller LG, Han LL, et al. Emergence of multidrug-resistant, community-associated, methicillin-resistant *Staphylococcus aureus* clone USA300 in men who have sex with men. *Ann Intern Med*. 2008;148(4):249-57.

331

332

333

334

- [5] de Jong GM, van der Boor SC, van Bokhoven C, et al. Methicillin-resistant *Staphylococcus aureus* (MRSA) in men having sex with men (MSM): a systematic review. *BMC Infect Dis.* 2025;25:299. 335 336 337
- [6] Han LL, McDougal LK, Gorwitz RJ, Mayer KH, Patel JB, Sennott JM, et al. High frequencies of clindamycin and tetracycline resistance in methicillin-resistant *Staphylococcus aureus* pulsed-field type USA300 isolates collected at a Boston ambulatory health center. *J Clin Microbiol.* 2007;45(4):1350-2. 338 339 340 341
- [7] Stevens DL, Bisno AL, Chambers HF, Dellinger EP, Goldstein EJC, Gorbach SL, et al. Practice guidelines for the diagnosis and management of skin and soft tissue infections: 2014 update by the Infectious Diseases Society of America. *Clin Infect Dis.* 2014;59(2):e10-52. 342 343 344 345
- [8] Liu C, Bayer A, Cosgrove SE, Daum RS, Fridkin SK, Gorwitz RJ, et al. Clinical practice guidelines by the Infectious Diseases Society of America for the treatment of methicillin-resistant *Staphylococcus aureus* infections in adults and children. *Clin Infect Dis.* 2011;52(3):e18-55. 346 347 348 349
- [9] Donà MG, Zaccarelli M, Giuliani E, et al. Doxycycline postexposure prophylaxis: a cross-sectional survey on knowledge, attitudes and prescribing practices among Italian dermatologists of sexually transmitted infection centers. *Ital J Dermatol Venereol.* 2026;161(2):120-5. 350 351 352 353
- [10] Spence M, Fowler C, Absalom S, Roper T, Williams D, Richardson D. Healthcare professionals' beliefs and concerns about the use of doxycycline post-exposure prophylaxis (doxyPEP): a systematic review. *Venereology.* 2026;5(1):9. 354 355 356
- [11] Luetkemeyer AF, Donnell D, Cohen SE, et al. Doxycycline to prevent bacterial sexually transmitted infections in the USA: final results from the DoxyPEP multicentre, open-label, randomised controlled trial and open-label extension. *Lancet Infect Dis.* 2025;25:873-83. 357 358 359 360

- [12] UK Health Security Agency. Monitoring and evaluation plan of doxycycline post-exposure prophylaxis (doxyPEP); 2026. Guidance; published 30 April 2026; applies to England. <https://www.gov.uk/government/publications/doxyypep-monitoring-and-evaluation/monitoring-and-evaluation-plan-of-doxycycline-post-exposure-prophylaxis>
- [13] Armond ACV, Cobey KD, Moher D. Research Integrity definitions and challenges. *Journal of Clinical Epidemiology*. 2024;171:111367.
- [14] Soge OO, Thibault CS, Cannon CA, et al. Potential impact of doxycycline post-exposure prophylaxis on tetracycline resistance in *Neisseria gonorrhoeae* and colonization with tetracycline-resistant *Staphylococcus aureus* and Group A *Streptococcus*. *Clin Infect Dis*. 2025;80:1188-96.
- [15] Luetkemeyer AF, DoxyPEP Study Team. DoxyPEP and antimicrobial resistance in *N. gonorrhoeae*, commensal *Neisseria* and *S. aureus*. Seattle, WA; 2023. Conference on Retroviruses and Opportunistic Infections (CROI), Oral Abstract OA-3.
- [16] Vanbaelen T, Manoharan-Basil SS, Kenyon C. Studies of postexposure prophylaxis with doxycycline should consider population-level selection for antimicrobial resistance. *Lancet Infect Dis*. 2024;24:e606-7.
- [17] Mayer KH, Traeger M, Marcus JL. Doxycycline Postexposure Prophylaxis and Sexually Transmitted Infections. *JAMA*. 2023;330(14):1381-2.
- [18] Flores J, Davis AM, Hazra A. Doxycycline Postexposure Prophylaxis to Prevent Bacterial Sexually Transmitted Infection. *JAMA*. 2025;333(3):248-9.
- [19] AIDSVu. State and county PrEP use and PrEP-to-Need Ratio, 2012–2025; 2026. Public data; accessed 2026. Emory University, Rollins School of Public Health. <https://aidsvu.org>.
- [20] Renson A, Montoya L, Goin DE, Díaz I, Ross RK. Pulling Back the Curtain: The Road from Statistical Estimand to Machine-Learning-Based Estimator for Epidemiologists (No Wizard Required). *American Journal of Epidemiology*. 2025;194(12):3566-71.

- [21] McDougal LK, Fosheim GE, Nicholson A, Bulens SN, Limbago BM, Shearer JES, et al. 388
Emergence of resistance among USA300 methicillin-resistant *Staphylococcus aureus* 389
isolates causing invasive disease in the United States. *Antimicrob Agents Chemother.* 390
2010;54(9):3804-11. 391
- [22] Mende K, Beckius ML, Zera WC, Yu X, Li P, Tribble DR, et al. Lack of doxycycline 392
antimalarial prophylaxis impact on *Staphylococcus aureus* tetracycline resistance. *Diagn* 393
Microbiol Infect Dis. 2016;86(2):211-20. 394
- [23] Kyaw WM, Lee LK, Siong WC, Ping ACL, Ang B, Leo YS. Prevalence of and risk 395
factors for MRSA colonization in HIV-positive outpatients in Singapore. *AIDS Res Ther.* 396
2012;9:33. 397
- [24] Tancredi S, Cullati S, Chiolerio A. Surveillance bias in the assessment of the size of 398
COVID-19 epidemic waves: a case study. *Public Health.* 2024;234:98-104. 399
- [25] Treem JW, Barley WC, Weber MS, Barbour JB. Signaling and Meaning in Organiza- 400
tional Analytics: Coping with Goodhart's Law in an Era of Digitization and Datafication. 401
Journal of Computer-Mediated Communication. 2023;28(4):zmad023. 402
- [26] Montoya LM, Geng EH, Valancius M, Kosorok MR, Petersen ML. Effects Among the 403
Affected. *Statistics in Medicine.* 2025;44(28-30):e70353. 404
- [27] Demidont AC. Guidance and surveillance capability for bystander *Staphylococcus au-* 405
reus tetracycline resistance under doxy-PEP: a three-stream evidence audit — research 406
compendium. Zenodo; 2026. Software and data compendium. 407

Table 1: Propagation of the DoxyPEP *S. aureus* resistance signal across reporting venues. Rows are not assumed to be identical estimands; differences in contrast, conditioning set, denominator and terminology are the object of comparison.

Venue	Reported result	Frame / conditioning set
NEJM appendix ¹	5/31 = 16%	Month-12 doxy-PEP arm; resistant among carriers
CROI abstract ¹⁵	16/137 = 12%	Doxy-PEP arm; all-swabbed denominator
CDC guideline ²	28/222 = 13%; summarised 5%→13%	Doxy-PEP arm; guideline frame
JAMA Insights ¹⁷	5% versus 4%	Doxy-PEP versus control; “infected with”
JAMA Synopsis ¹⁸	5%→16%	Within arm, among <i>S. aureus</i> -isolated

Table 2: Headline audit of the doxy-PEP evidence base against one detectability yardstick. Each cell is licensed by the Results above; no composite score is implied.

Stream	Question	Result
A. Trials	Was the bystander signal resolved?	Yes—HR 3.89 (1.42–10.68), incident doxy-R <i>S. aureus</i>
B. Guidance	Is <i>S. aureus</i> monitoring required?	No—0/6 outbreak-matched; 0/10 coded
C. Surveillance	Is exposure linked to phenotype?	No—0/12 systems; $RR_{\text{needed}} \approx 6\text{--}28$ to detect