


# Total Synthesis of Violaceoid A and (–)- and (+)-Violaceoid B

Takatsugu Murata,<sup>†</sup> Teppei Kuboki,<sup>†</sup> Ryo Ishikawa,<sup>†</sup> Takahiro Saito,<sup>†</sup> Shotaro Taguchi,<sup>†</sup> Kazuma Takeuchi,<sup>†</sup> Emiko Hatano,<sup>†</sup> Motoyuki Shimonaka,<sup>‡</sup> and Isamu Shiina<sup>\*,†,§</sup>

<sup>†</sup>Department of Applied Chemistry, Faculty of Science, Tokyo University of Science, 1-3 Kagurazaka, Shinjuku-ku, Tokyo 162-8601, Japan

<sup>‡</sup>Department of Chemistry, Faculty of Science, Tokyo University of Science, 1-3 Kagurazaka, Shinjuku-ku, Tokyo 162-8601, Japan

§ Supporting Information




**ABSTRACT:** The first total synthesis of violaceoid A, a cytotoxic agent, and the asymmetric total synthesis of (–)- and (+)-violaceoid B are reported. The precursor was accessed by desymmetrization of a substituted quinol moiety, and the racemic secondary alcohol was kinetically resolved using a chiral nucleophilic catalyst. The asymmetric synthesis of (–)- and (+)-violaceoid B elucidated the absolute configuration of the naturally occurring violaceoid B. Synthetic violaceoid A inhibited the growth of human breast cancer cell lines MCF-7 and Hs 578T at concentrations of less than 100  $\mu$ M, while (S)- and (R)-violaceoid B were inactive.

In 2014, Sugawara and co-workers reported a series of unique alkylated hydroquinones, violaceoids A–F (compounds 1–6), which were isolated from a culture broth of *Aspergillus violaceofuscus* Gasperini coexisting with moss.<sup>1</sup> Violaceoids B and D–F are chiral compounds, and the absolute configurations of violaceoids B, D, and E have not yet been determined. Sugawara's group also reported that violaceoids exhibit cytotoxicity. Among them, violaceoid A (1) inhibited the growth of several human cancer cell lines. In addition, neither the total synthesis nor the synthetic approach for violaceoids A–E has ever been reported to the best of our knowledge.<sup>2</sup> Considering this, we initiated a program to synthesize 1 and 2, and the details are presented in this paper.

## RESULTS AND DISCUSSION

The retrosynthetic analysis of 1 and 2 is shown in Scheme 1.

Optically active (S)- or (R)-violaceoid B (2) can be obtained by the deprotection of the acetonide group from (S)- or (R)-7, respectively. Diols (S)- and (R)-7 can be separated by using a kinetic resolution method of a racemic compound (*rac*-7) that our group has already reported.<sup>3</sup> The substrate for kinetic resolution can be obtained from the racemic secondary alcohol 8, which can be prepared from 3,6-dihydroxyphthalonitrile (11) via direct protection of 11.




Following this strategy, we first tried to protect two hydroxy groups of 3,6-dihydroxyphthalonitrile as *tert*-butyldiphenylsilyl (TBDPS) ether depicted in Scheme 2. However, we could not obtain the desired phthalonitrile 10. Then we conducted the protection transforming it into methoxymethyl (MOM) ether groups, but we could get the desired compound 12 in low yield. On the basis of the above results, we decided to change the synthetic plan for providing violaceoids A and B as shown in Scheme 3.


Received: March 14, 2018

Published: October 30, 2018

Scheme 1. Retrosynthetic Analysis of 1 and 2




Scheme 2. Protection of 3,6-Dihydroxyphthalonitrile (11)





In the revised synthetic plan, *rac*-7, the substrate for kinetic resolution, can be obtained from the racemic secondary alcohol 14 by several functional transformations. Violaceoid A (1) can also be derived from 14. The key intermediate 14 can be prepared from 11 via reduction of a derivative of 11.

The preparation of 14 was carried out as depicted in Scheme 4. The hydrolysis of 11 followed by esterification gave the dimethyl ester 17.<sup>4</sup> The two hydroxy groups of 17 were protected with TBDPS groups to afford 18. The symmetric diol 16 was obtained by the reduction of diester 18. Monotetrahydropyranylation gave the unilateral protected alcohol 19. After the oxidation of alcohol 19 to aldehyde 15 followed by alkylation with an alkyl lithium reagent, we obtained the key intermediate 14.

Scheme 3. Revised Retrosynthetic Analysis of 1 and 2



## Scheme 4. Synthesis of Key Intermediate 14

Scheme 5. Synthesis of 1 and *rac*-2

We next attempted to synthesize **1** and *rac*-**2** (Scheme 5).

The double bond in **20** was generated by mesylation of **14**. Then, the THP group was removed, and we obtained the alcohol **21**. Finally, by deprotection of the two TBDPS groups, we accomplished the total synthesis of violaceoid A (**1**). In addition, by sequential deprotection of the THP and TBDPS groups of **14**, we achieved the synthesis of *rac*-violaceoid B (*rac*-**2**). We compared the  $^1\text{H}$  and  $^{13}\text{C}$  NMR data of the synthetic **1** and **2** with those of naturally occurring violaceoids A and B reported in the literature to determine the true structure. The results are shown in Tables 1 and 2. As a result, the  $^1\text{H}$  and  $^{13}\text{C}$  NMR data of synthetic **1** and **2** were shown to be in accordance with those reported for the natural compounds.

Subsequently, we attempted to determine the absolute configuration of naturally occurring violaceoid B (**2**), as depicted in Scheme 6.

We obtained the alcohol **24** by acetylation of **14** followed by deprotection of the THP group. Next, deprotection of the TBDPS group was conducted to afford the 1,3-diol **13**. The acetyl and TBDPS groups were reductively removed together to yield *rac*-**7**, which is the substrate for kinetic resolution.<sup>3</sup> The kinetic resolution was conducted to afford the enantiomer (R)-**26**<sup>5</sup> and enantiomer (S)-**7**. Further, the same reaction was repeated five times to obtain the enantiomer (S)-**2** in 95% ee. Finally, by deprotection of the acetonide group, we accomplished the asymmetric total synthesis of (S)-violaceoid B ((S)-**2**). However, the optical rotation of (S)-**2** was not consistent with the natural product, which motivated us to synthesize the enantiomer, (R)-violaceoid B ((R)-**2**) (Scheme 7).

As for enantiopure (S)-**7**, we conducted the kinetic resolution procedure. Using the recovered enantiomer (S)-**7**, the

**Table 1.** Comparison of the  $^1\text{H}$  and  $^{13}\text{C}$  NMR Data of Naturally Occurring Violaceoid A with Those of Synthetic **1** in  $\text{CD}_3\text{OD}$


natural violaceoid A <sup>a</sup>			synthetic violaceoid A		
position	$\delta_{\text{C}}$	$\delta_{\text{H}}$ (J in Hz)	position	$\delta_{\text{C}}$ <sup>b</sup>	$\delta_{\text{H}}$ (J in Hz) <sup>c</sup>
1	148.8		1	148.8	
2	127.6		2	127.6	
3	125.9		3	125.9	
4	150.6		4	150.6	
5	115.0	6.57, d (8.7)	5	114.9	6.54, d (8.7)
6	116.3	6.60, d (8.7)	6	116.2	6.59, d (8.7)
1'	124.6	6.45, dt (16.1, 6.9)	1'	124.6	6.50–6.42, m
2'	137.9	6.06, dt (16.1, 6.9)	2'	137.8	6.08, dt (15.6, 6.9)
3'	34.9	2.24, tdd (6.9, 6.9, 1.5)	3'	34.9	2.32–2.19, m
4'	30.3	1.51, m	4'	30.4	1.61–1.43, m
5'	32.7	1.37, m	5'	32.7	1.43–1.30, m
6'	23.6	1.37, m	6'	23.7	1.43–1.30, m
7'	14.4	0.93, t (7.1)	7'	14.5	0.94, t (7.2)
1''	58.5	4.72, s	1''	58.4	4.71, s

<sup>a</sup>300 MHz for  $^1\text{H}$  NMR and 75 MHz for  $^{13}\text{C}$  NMR. See ref 1. <sup>b</sup>125 MHz using JNM-LAS00. <sup>c</sup>300 MHz using JNM-AL300.

the same reaction was then repeated five times to obtain the enantiomer (R)-**2** in 93% ee. Finally, deprotection of the



Scheme 7. Synthesis of (R)-Violaceoid B ((R)-2)



atoms of all compounds are numbered according to IUPAC nomenclature.

**3,6-Bis(methoxymethoxy)phthalonitrile (12).** To a solution of 3,6-dihydroxyphthalonitrile (**11**) (400 mg, 2.56 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (25.6 mL) were added diisopropylethylamine (1.76 mL, 10.24 mmol) and methyloxymethyl chloride (0.77 mL, 10.24 mmol) at 0 °C. The reaction mixture was stirred for 1 h and warmed up to room temperature (rt) and stirred for 12 h. The solution was diluted with water and extracted with CH<sub>2</sub>Cl<sub>2</sub>. The organic layer was washed with H<sub>2</sub>O and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtrated, and concentrated. The residue was purified by silica gel column chromatography (*n*-hexane/EtOAc = 4:1). Compound **12** (127 mg, 20%) was obtained as a white solid.

**Dimethyl 3,6-Dihydroxyphthalate (17).** To a solution of KOH (32.0 g, 570 mmol) in water (32.0 mL) was added 3,6-dihydroxyphthalonitrile (**11**) (5.00 g, 31.2 mmol) at rt. The reaction mixture was refluxed for 1 h. Aqueous 20% H<sub>2</sub>SO<sub>4</sub> (100 mL) was slowly poured into the reaction mixture and extracted with Et<sub>2</sub>O and CHCl<sub>3</sub> sequentially. The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtrated, and concentrated in vacuo, giving 3,6-dihydroxyphthalic acid (6.20 g) as the crude product including starting material **11**. The mixture was used for the next reaction without further purification.

To a solution of the crude 3,6-dihydroxyphthalic acid (6.20 g) in MeOH (240 mL) was added boron trifluoride diethyl etherate (12.8 mL, 100 mmol) at room temperature, and reaction mixture was refluxed for 15 h. The solution was cooled to rt, concentrated in vacuo, quenched with saturated aqueous NaHCO<sub>3</sub>, and extracted with EtOAc. The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub>, filtrated, and concentrated in vacuo to yield dimethyl 3,6-dihydroxyphthalate (**17**) (3.69 g, 52% in two steps) as a white solid.

**Dimethyl 3,6-Bis((*tert*-butyldiphenylsilyl)oxy)phthalate (18).** 3,6-Dihydroxyphthalate (**17**) (500 mg, 2.21 mmol) was dissolved in CH<sub>2</sub>Cl<sub>2</sub> (22 mL) and cooled to 0 °C. Imidazole (1.20 g, 17.68 mmol) and TBDPSCI (2.27 mL, 8.84 mmol) were added to the solution, and the reaction mixture was stirred for 2 h at rt. The mixture was quenched by saturated aqueous NaHCO<sub>3</sub> at 0 °C, extracted with EtOAc, washed with brine, and concentrated. The residue was purified by silica gel column chromatography (gradient, *n*-hexane/EtOAc = 20:1–5:1). Compound **18** (1.61 g, 95%) was obtained as a white solid.

**(3,6-Bis((*tert*-butyldiphenylsilyl)oxy)-1,2-phenylene)-dimethanol (16).** To a solution of **18** in CH<sub>2</sub>Cl<sub>2</sub> was added diisobutylaluminum hydride (42.7 mL, 42.7 mmol) at –78 °C, and the reaction mixture was stirred at 0 °C for 2 h. The reaction mixture was quenched by MeOH and saturated aqueous Rochell's salt, extracted with CH<sub>2</sub>Cl<sub>2</sub>, washed with brine, and concentrated. The residue was purified by silica gel column chromatography (*n*-hexane/EtOAc = 10:1). Compound **16** (1.60 g, 93%) was obtained as a colorless solid.

**(3,6-Bis((*tert*-butyldiphenylsilyl)oxy)-2-(((tetrahydro-2H-pyran-2'-yl)oxy)methyl)phenyl)methanol (19).** To a solution of

**16** (1.16 g, 1.79 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (44.8 mL, 0.04 M) were added pyridinium *p*-toluenesulfonate (90.4 mg, 0.36 mmol) and 3,4-dihydro-2H-pyran (0.23 mL, 2.69 mmol), and the reaction mixture was stirred for 4 h at rt. The reaction mixture was quenched by saturated aqueous NaHCO<sub>3</sub>, extracted with CH<sub>2</sub>Cl<sub>2</sub>, and concentrated. The residue was purified by silica gel column chromatography (gradient, *n*-hexane/EtOAc = 20:1–1:1). Compound **19** (1.22 g, 93%) was obtained as a white solid.

**3,6-Bis((*tert*-butyldiphenylsilyl)oxy)-2'-(((tetrahydro-2H-pyran-2'-yl)oxy)methyl)benzaldehyde (15).** To a solution of **19** (198 mg, 0.271 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (5.42 mL, 0.05 M) were added tetrapropylammonium perruthenate (28.6 mg, 0.0813 mmol) and 4-methylmorpholine *N*-oxide (95.5 mg, 0.813 mmol), and the reaction mixture was stirred for 2 h at 0 °C. The reaction mixture was filtrated through a short pad of silica gel with EtOAc, and the filtrate was concentrated. Compound **15** (197 mg, quant.) was obtained as a white solid.

**1-(3',6'-Bis((*tert*-butyldiphenylsilyl)oxy)-2'-(((tetrahydro-2H-pyran-2'-yl)oxy)methyl)phenyl)heptan-1-ol (14).** To a solution of **15** (457 mg, 0.628 mmol) in THF (12.6 mL, 0.05 M) was added a 1.15 M hexyllithium solution in *n*-hexane at –78 °C. The reaction mixture was stirred for 5 min, quenched with saturated aqueous NaHCO<sub>3</sub>, extracted with EtOAc, and concentrated. The residue was purified by silica gel column chromatography (*n*-hexane/EtOAc = 5:1). Compound **14** (382 mg, 75%) was obtained as a white solid.

**(E)-((2-(Hept-1'-en-1'-yl)-3-(((tetrahydro-2H-pyran-2'-yl)oxy)methyl)-1,4-phenylene)bis(oxy))bis((*tert*-butyldiphenylsilyl)oxy)methyl)phenyl)methanol (20).** To a solution of **14** (180 mg, 0.220 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (4.35 mL, 0.05 M) were added methanesulfonyl chloride (0.047 mL, 0.660 mmol) and triethylamine (0.186 mL, 1.321 mmol) at 0 °C. The reaction mixture was stirred for 3 h at rt, quenched with saturated aqueous NaHCO<sub>3</sub>, extracted with CH<sub>2</sub>Cl<sub>2</sub>, and concentrated. The residue was purified by silica gel column chromatography (*n*-hexane/EtOAc = 5:1). Compound **20** (122 mg, 70%) was obtained as a white solid.

**(E)-(3,6-Bis((*tert*-butyldiphenylsilyl)oxy)-2-(hept-1'-en-1'-yl)phenyl)methanol (21).** To a solution of **20** (84.0 mg, 0.106 mmol) in MeOH (4.24 mL, 0.025 M) was added *p*-toluenesulfonic acid monohydrate (30.0 mg, 0.158 mmol) at 0 °C. The reaction mixture was stirred for 2 h at rt, diluted with water, extracted with chloroform, and concentrated. The residue was purified by silica gel column chromatography (*n*-hexane/EtOAc = 5:1). Compound **21** (53.8 mg, 72%) was obtained as a colorless oil.

**(E)-2-(Hept-1'-en-1'-yl)-3-(hydroxymethyl)benzene-1,4-diol (Violaceoid A (1)).** To a solution of **21** (18.5 mg, 0.026 mmol) in THF/pyridine (1.7 mL, *v/v* = 1/1, 0.015 M) was added hydrogen fluoride pyridine complex (0.60 mL) at 0 °C. The reaction mixture was stirred for 2 h, quenched with saturated aqueous NaHCO<sub>3</sub>, extracted with EtOAc, and washed with saturated aqueous copper sulfate, water, saturated aqueous NaHCO<sub>3</sub>, and brine. The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. The residue was



purified by silica gel chromatography (*n*-hexane/EtOAc = 2:1). Violaceoid A (1) (4.1 mg, 67%) was obtained as a white solid.

**1-(3',6'-Bis((*tert*-butyldiphenylsilyl)oxy)-2'-(hydroxymethyl)phenyl)heptan-1-ol (22).** To a solution of 14 (200 mg, 0.246 mmol) in MeOH (5.00 mL, 0.05 M) was added *p*-toluenesulfonic acid monohydrate (56.0 mg, 0.295 mmol) at 0 °C. The reaction mixture was stirred for 2 h at rt, diluted with water, extracted with chloroform, and concentrated. The residue was purified by silica gel column chromatography (*n*-hexane/EtOAc = 5:1). Compound 22 (119 mg, 66%) was obtained as a white solid.

**2-(1'-Hydroxyheptyl)-3-(hydroxymethyl)benzene-1,4-diol (rac-Violaceoid B (2)).** To a solution of 22 (77 mg, 0.105 mmol) in THF/pyridine (5.25 mL, *v/v* = 1:1, 0.02 M) was added hydrogen fluoride pyridine complex (2.00 mL) at 0 °C. The reaction mixture was stirred for 2 h, quenched with saturated aqueous NaHCO<sub>3</sub>, extracted with EtOAc, and washed with saturated aqueous copper sulfate, water, saturated aqueous NaHCO<sub>3</sub>, and brine. The organic layer was dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated. The residue was purified by silica gel chromatography (*n*-hexane/EtOAc = 2:1). *rac*-Violaceoid B (2) (19.2 mg, 72%) was obtained as a white solid.

**1-(3',6'-Bis((*tert*-butyldiphenylsilyl)oxy)-2'-(hydroxymethyl)phenyl)heptyl Acetate (24).** To a solution of 14 (32.4 mg, 0.040 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (2.00 mL, 0.02 M) were added acetic anhydride (7.5 μL, 0.0796 mmol), triethylamine (22.2 μL, 0.159 mmol), and *N,N*-dimethylpyridin-4-amine (1.0 mg, 0.00796 mmol) at rt. The reaction mixture was stirred for 12 h, quenched with saturated aqueous NaHCO<sub>3</sub>, extracted with CH<sub>2</sub>Cl<sub>2</sub> × 2 and EtOAc × 2, and dried over Na<sub>2</sub>SO<sub>4</sub>. The organic layer was filtrated and concentrated. The residue was purified by silica gel chromatography (*n*-hexane/EtOAc = 5:1). The crude product 23 was used for the following reaction without further purification.

To a solution of the crude product 23 in MeOH/THF (3.35 mL, *v/v* = 1:1, 0.01 M) was added *p*-toluenesulfonic acid monohydrate (9.6 mg, 0.0503 mmol) at 0 °C. The reaction mixture was stirred for 2 h at rt, quenched with saturated aqueous NaHCO<sub>3</sub>, extracted with CH<sub>2</sub>Cl<sub>2</sub> × 2 and EtOAc × 2, and dried over Na<sub>2</sub>SO<sub>4</sub>. The organic layer was filtrated and concentrated. The residue was purified by silica gel chromatography (*n*-hexane/EtOAc = 5:1). Compound 24 (20.4 mg, 66% in 2 steps) was obtained as a white solid.

**1-(6'-((*tert*-Butyldiphenylsilyl)oxy)-3'-hydroxy-2'-(hydroxymethyl)phenyl)heptyl Acetate (13).** To a solution of 24 (1.45 g, 1.70 mmol) in THF/pyridine (34.0 mL, *v/v* = 2:1, 0.050 M) was added hydrogen fluoride pyridine complex (1.0 mL) at 0 °C. The reaction mixture was stirred for 2 h, quenched with saturated aqueous NaHCO<sub>3</sub>, extracted with EtOAc, and concentrated. Pyridine was removed by using azeotrope with benzene, and the residue was purified by silica gel chromatography (*n*-hexane/EtOAc = 3:1). Compound 13 (851 mg, 94%) was obtained as a white solid.

**1-(6'-((*tert*-Butyldiphenylsilyl)oxy)-2',2'-dimethyl-4H-benzo[d][1',3']dioxin-5'-yl)heptyl Acetate (25).** To a solution of 13 (851 mg, 1.59 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (53.1 mL, 0.03 M) were added 2,2-dimethoxypropane (0.49 mL, 3.98 mmol) and *p*-toluenesulfonic acid monohydrate (1.5 mg, 0.0796 mmol). The reaction mixture was stirred for 30 min, quenched with saturated aqueous NaHCO<sub>3</sub>, extracted with EtOAc, and concentrated. The residue was purified by silica gel chromatography (*n*-hexane/EtOAc = 3:1). Compound 25 (838 mg, 92%) was obtained as a colorless oil.

**5-(1'-Hydroxyheptyl)-2,2-dimethyl-4H-benzo[d][1,3]dioxin-6-ol (rac-7).** To a solution of 25 (422 mg, 0.734 mmol) in THF (24.5 mL, 0.03 M) was added 1.0 M lithium aluminum hydride in THF (1.84 mL) at 0 °C. The reaction mixture was stirred for 5 min, quenched with MeOH and saturated aqueous Rochell's salt, extracted with EtOAc, and dried over Na<sub>2</sub>SO<sub>4</sub>. The organic layer was filtrated and concentrated. The residue was purified by silica gel chromatography (*n*-hexane/EtOAc = 3:1). Compound *rac*-7 (214 mg, 99%) was obtained as a white solid.

**(R)-1-(6'-Hydroxy-2',2'-dimethyl-4H-benzo[d][1',3']dioxin-5'-yl)heptyl 2,2-Diphenylacetate ((R)-26) and (S)-5-(1'-Hydroxyheptyl)-2,2-dimethyl-4H-benzo[d][1,3]dioxin-6-ol ((S)-7).** To a solution of *rac*-7 (69.3 mg, 0.235 mmol) in Et<sub>2</sub>O (2.3 mL, 0.1 M) were added *N,N*-diisopropylethylamine (0.030 mL, 0.169

mmol), (*R*)-benzotetramisole (3.0 mg, 0.0118 mmol), and diphenylacetic anhydride (57.0 mg, 0.141 mmol). The reaction mixture was stirred for 12 h, quenched with saturated aqueous NaHCO<sub>3</sub>, extracted with EtOAc, and dried over Na<sub>2</sub>SO<sub>4</sub>. The organic layer was filtrated and concentrated. The residue was purified by silica gel chromatography (*n*-hexane/EtOAc = 4:1, then CH<sub>2</sub>Cl<sub>2</sub>). Compound (*R*)-26 (33.1 mg, 29%, 12% ee) was obtained as a colorless oil, and compound (*S*)-7 (37.1 mg, 54%, 50% ee) was obtained as a white solid.

**(S)-5-(1'-Hydroxyheptyl)-2,2-dimethyl-4H-benzo[d][1,3]dioxin-6-ol ((S)-7) (95% ee).** To a solution of (*S*)-7 (37.1 mg, 0.126 mmol, 50% ee) in Et<sub>2</sub>O (1.3 mL, 0.1 M) were added *N,N*-diisopropylethylamine (0.014 mL, 0.0339 mmol), (*R*)-benzotetramisole (1.6 mg, 0.00628 mmol), and diphenylacetic anhydride (13.8 mg, 0.0339 mmol). The reaction mixture was stirred for 12 h, quenched with saturated aqueous NaHCO<sub>3</sub>, extracted with EtOAc, and dried over Na<sub>2</sub>SO<sub>4</sub>. The organic layer was filtrated and concentrated. The residue was purified by silica gel chromatography (*n*-hexane/EtOAc = 4:1, then CH<sub>2</sub>Cl<sub>2</sub>). Compound (*S*)-7 (35.9 mg, 97%, 54% ee) was obtained as a white solid. The chiral enriched (*S*)-7 (18.8 mg, 95% ee) was obtained by the same kinetic resolution after four repetitions from the above (*S*)-7 (35.9 mg, 0.126 mmol, 54% ee).

(*S*)-7 (95% ee): [ $\alpha$ ]<sub>D</sub><sup>25</sup> −46.3 (*c* 1.12, CHCl<sub>3</sub>). Other spectra are the same as those of *rac*-7.

**(S)-2-(1'-Hydroxyheptyl)-3-(hydroxymethyl)benzene-1,4-diol ((S)-Violaceoid B, (S)-2).** To a solution of (*S*)-7 (69.5 mg, 0.236 mmol, 95% ee) in THF/H<sub>2</sub>O (2.36 mL, *v/v* = 1:1, 0.1 M) was added *p*-toluenesulfonic acid monohydrate (22.4 mg, 0.118 mmol). The reaction mixture was stirred for 6 h, quenched with saturated aqueous NaHCO<sub>3</sub>, extracted with EtOAc, and concentrated. The residue was purified by silica gel chromatography (*n*-hexane/EtOAc = 3:1) to afford (*S*)-violaceoid B (9.2 mg, 15%) as a white solid, and (*S*)-7 (59.6 mg, 86%) was recovered.

Additionally, (*S*)-violaceoid B (25.7 mg) and (*S*)-7 (27.1 mg) were obtained by the same reaction after two repetitions from the recovered (*S*)-7 (59.6 mg, 0.202 mmol). Therefore, 34.9 mg (total amount) of (*S*)-violaceoid (58%) was prepared from the starting 69.5 mg of (*S*)-7.

(*S*)-Violaceoid B ((S)-2) (95% ee): [ $\alpha$ ]<sub>D</sub><sup>23</sup> −17.0 (*c* 0.093, CHCl<sub>3</sub>), [ $\alpha$ ]<sub>D</sub><sup>23</sup> −22.5 (*c* 0.667, MeOH). Other spectra are the same as those of *rac*-2.

**(S)-1-(6'-Hydroxy-2',2'-dimethyl-4H-benzo[d][1',3']dioxin-5'-yl)heptyl 2,2-diphenylacetate ((S)-26) and (R)-5-(1'-Hydroxyheptyl)-2,2-dimethyl-4H-benzo[d][1,3]dioxin-6-ol ((R)-7).** To a solution of *rac*-7 (123.7 mg, 0.420 mmol) in Et<sub>2</sub>O (4.2 mL, 0.1 M) were added *N,N*-diisopropylethylamine (0.050 mL, 0.303 mmol), (*S*)-benzotetramisole (5.0 mg, 0.0210 mmol), and diphenylacetic anhydride (102 mg, 0.252 mmol). The reaction mixture was stirred for 12 h, quenched with saturated aqueous NaHCO<sub>3</sub>, extracted with EtOAc, and dried over Na<sub>2</sub>SO<sub>4</sub>. The organic layer was filtrated and concentrated. The residue was purified by silica gel chromatography (*n*-hexane/EtOAc = 4:1, then CH<sub>2</sub>Cl<sub>2</sub>). Compound (*S*)-26 (65.9 mg, 32%, 19% ee) was obtained as a colorless oil, and compound (*R*)-7 (60.1 mg, 49%, 58% ee) was obtained as a white solid.

(*S*)-26 (19% ee): [ $\alpha$ ]<sub>D</sub><sup>26</sup> −0.20 (*c* 2.39, CHCl<sub>3</sub>). Other spectra are the same as those of (*R*)-26.

**(R)-5-(1'-Hydroxyheptyl)-2,2-dimethyl-4H-benzo[d][1,3]dioxin-6-ol ((R)-7) (93% ee).** To a solution of (*R*)-7 (60.1 mg, 0.204 mmol, 58% ee) in Et<sub>2</sub>O (2.0 mL, 0.10 M) were added *N,N*-diisopropylethylamine (9.6 μL, 0.0550 mmol), (*S*)-benzotetramisole (2.6 mg, 0.0118 mmol), and diphenylacetic anhydride (22.3 mg, 0.0550 mmol). The reaction mixture was stirred for 12 h, quenched with saturated aqueous NaHCO<sub>3</sub>, extracted with EtOAc, and dried over Na<sub>2</sub>SO<sub>4</sub>. The organic layer was filtrated and concentrated. The residue was purified by silica gel chromatography (*n*-hexane/EtOAc = 4:1, then CH<sub>2</sub>Cl<sub>2</sub>). Compound (*R*)-7 (56.8 mg, 95%, 66% ee) was obtained as a white solid. The chiral enriched (*R*)-7 (31.0 mg, 93% ee) was obtained by the same kinetic resolution after four repetitions from the above (*R*)-7 (56.8 mg, 0.193 mmol, 66% ee).

(*R*)-7 (93% ee): [ $\alpha$ ]<sub>D</sub><sup>25</sup> +44.6 (*c* 1.87, CHCl<sub>3</sub>). Other spectra are the same as those of *rac*-7.

(*R*)-2-(1'-Hydroxyheptyl)-3-(hydroxymethyl)benzene-1,4-diol ((*R*)-Violaceoid B, (*R*)-2)). To a solution of (*R*)-7 (25.1 mg, 0.0853 mmol, 93% ee) in THF/H<sub>2</sub>O (1.71 mL, v/v = 1:1, 0.05 M) was added *p*-toluenesulfonic acid monohydrate (8.1 mg, 0.0426 mmol). The reaction mixture was stirred for 6 h, quenched with saturated aqueous NaHCO<sub>3</sub>, extracted with EtOAc, and concentrated. The residue was purified by silica gel chromatography (*n*-hexane/EtOAc = 3:1) to afford (*R*)-violaceoid B (4.6 mg, 21%) as a white solid, and (*R*)-7 (19.6 mg, 78%) was recovered.

Additionally, (*R*)-violaceoid B (4.6 mg) and (*R*)-7 (12.2 mg) were obtained by the same reaction after two repetitions from the recovered (*R*)-7 (19.6 mg, 0.0618 mmol). Therefore, 9.2 mg (total amount) of (*R*)-violaceoid (54%) was prepared from the starting 25.1 mg of (*R*)-7.

(*R*)-Violaceoid B ((*R*)-2) (93% ee):  $[\alpha]_D^{23} +15.9$  (c 0.087, CHCl<sub>3</sub>),  $[\alpha]_D^{23} +22.9$  (c 0.667, MeOH). Other spectra are the same as those of *rac*-2.

**Biological Assay.** Human breast cancer cell lines MCF-7 and Hs 578T were obtained from Japanese Collection of Research Bioresources (JCRB) Cell Bank (Osaka, Japan) and American Type Culture Collection (Manassas, VA, USA), respectively. Cells were cultured with Dulbecco's modified Eagle's medium supplemented with antibiotic-antimycotic (100 U/mL penicillin, 100 µg/mL streptomycin, 0.25 µg/mL amphotericin B; Thermo Fisher Scientific, Inc., Waltham, MA, USA), gentamicin (10 µg/mL, Thermo Fisher Scientific, Inc.), and 10% heat-inactivated fetal bovine serum. MCF-7 cells or Hs 578T cells were seeded in 96-well plates (1000 cells/well) and incubated for 48 h at 37 °C. Cells were then incubated with violaceoid A or (*S*)- and (*R*)-violaceoid B (10–100 µM) for 48 h, and cell number was estimated by WST-8 reagent (Cell Counting Kit-8, Dojindo Laboratories, Japan).

## ■ ASSOCIATED CONTENT

### 📄 Supporting Information

The Supporting Information is available free of charge on the ACS Publications website at DOI: 10.1021/acs.jnatprod.8b00215.

Copies of <sup>1</sup>H and <sup>13</sup>C NMR spectra for all new compounds (1, 2, 7, and 12–26), HPLC data of compounds *rac*-7, (*S*)-7, and (*R*)-7, and additional data (procedure for the reduction of ester 26 to the corresponding alcohol 7 and for the determination of the absolute configuration of chiral alcohols (*S*)-7 and (*R*)-7) (PDF)

## ■ AUTHOR INFORMATION

### Corresponding Author

\*E-mail: shiina@rs.kagu.tus.ac.jp.

### ORCID

Isamu Shiina: 0000-0002-8749-2540

### Notes

The authors declare no competing financial interest.

## ■ ACKNOWLEDGMENTS

The authors wish to thank Profs. F. Sugawara and S. Kamisuki (Tokyo University of Science) for providing the spectroscopic data for violaceoids A and B. This study was partly supported by a Research Grant from the Center for Chirality and Grants-in-Aid for Scientific Research from the Ministry of Education, Science, Sports and Culture, Japan.

## ■ REFERENCES

(1) Myobatake, Y.; Takemoto, K.; Kamisuki, S.; Inoue, N.; Takasaki, A.; Takeuchi, T.; Mizushima, Y.; Sugawara, F. *J. Nat. Prod.* **2014**, *77*, 1236–1240.

(2) Violaceoid F had already been prepared as a synthetic intermediate in the total synthesis of the enantiomer of eupenoxide in 2004. See: Mehta, G.; Roy, S. *Org. Lett.* **2004**, *6*, 2389–2392.

(3) (a) Shiina, I.; Nakata, K. *Tetrahedron Lett.* **2007**, *48*, 8314–8317. (b) Shiina, I.; Nakata, K.; Sugimoto, M.; Onda, Y.; Iizumi, T.; Ono, K. *Heterocycles* **2009**, *77*, 801–810. (c) Nakata, K.; Shiina, I. *Heterocycles* **2010**, *80*, 169–176. (d) Shiina, I.; Ono, K.; Nakata, K. *Chem. Lett.* **2011**, *40*, 147–149. (e) Nakata, K.; Ono, K.; Shiina, I. *Heterocycles* **2011**, *82*, 1171–1180. (f) Nakata, K.; Shiina, I. *Org. Biomol. Chem.* **2011**, *9*, 7092–7096. (g) Shiina, I.; Nakata, K.; Ono, K.; Mukaiyama, T. *Helv. Chim. Acta* **2012**, *95*, 1891–1911.

(4) Kelly, T. R.; Bell, S. H.; Ohashi, N.; Armstrong-Chong, R. J. *J. Am. Chem. Soc.* **1988**, *110*, 6471–6480.

(5) The enantiomeric excess (ee) was determined after being reduced to the corresponding alcohol 7.