


Papua New Guinea — ERA5 rainfall × ENSO teleconnection

Pixelwise partial correlation of trimester rainfall with Niño3.4 — the other climate modes held constant — ERA5 0.25° grid, 1981–2025


Unique ENSO signal — strongest significant trimester per pixel




Papua New Guinea — current SEAS5 seasonal forecast

Issued August 2026 · drought/flood return-period categories vs the same-issuance SEAS5 hindcast (1981–2025) · skill-shaded · native 0.4° grid · ADM1 boundaries


SON 2026 — lead 1 mo — mean 3rd pct ← driest overall




OND 2026 — lead 2 mo — mean 13th pct



NDJ 2026/27 — lead 3 mo — mean 43rd pct



DJF 2026/27 — lead 4 mo — mean 59th pct




Forecast (high skill):

- Drought — ≥ 10 -yr return period
- Drought — 3-10-yr return period
- Roughly normal
- Flood — 3-10-yr return period
- Flood — ≥ 10 -yr return period

Other:

- Roughly normal (mod skill)
- Low skill
- Outside rainy season

Country-mean rainfall (mm/day)



Categories from the detrended skill cube's forecast percentile vs the same issued month + leads. RPs under the bars: country mean vs the 45-yr hindcast. Niño3.4: +1.7 °C in Jul 2026. Data: ECMWF SEAS5 · Boundaries: PNG ADM1 (COD)