


# Timor-Leste — ERA5 rainfall × ENSO teleconnection

Pixelwise partial correlation of trimester rainfall with Niño3.4 — the other climate modes held constant — ERA5 0.25° grid, 1981–2025

Unique ENSO signal — strongest significant trimester per pixel

## Partial correlation with Niño3.4




Method (the teleconnections page's partial pass):

- Trimester rainfall means per pixel, all 12 overlapping 3-month windows
- Partial  $r$  of rainfall vs Niño3.4, holding DMI (IOD), TNA, TSA and AMM constant (PDO kept out of the controls, as on the page)
- Each mode enters at its per-pixel best lag (0–3 mo) frozen from the total sweep
- Two-tailed  $p < 0.05$ , df adjusted for controls
- Unlike the global page, all seasons are kept (no rainy-season filter), so dry-season ENSO signals stay visible

Negative (brown) = El Niño → drier than normal.

Data: ERA5 monthly precip (0.25°), NOAA PSL Niño3.4. Bottom row: canonical trimesters.




DJF

MAM

JAS


OND




# Timor-Leste — current SEAS5 seasonal forecast

Issued August 2026 · drought/flood return-period categories vs the same-issuance SEAS5 hindcast (1981–2025) · skill-shaded · native 0.4° grid · ADM1 boundaries

SON 2026 — lead 1 mo — mean 0th pct



OND 2026 — lead 2 mo — mean 0th pct ← driest overall




### Forecast (high skill):

- Drought —  $\geq$  10-yr return period
- Drought — 3-10-yr return period
- Roughly normal
- Flood — 3-10-yr return period
- Flood —  $\geq$  10-yr return period


### Other:

- Roughly normal (mod skill)
- Low skill
- Outside rainy season


NDJ 2026/27 — lead 3 mo — mean 3rd pct



DJF 2026/27 — lead 4 mo — mean 9th pct



### Country-mean rainfall (mm/day)



Categories from the detrended skill cube's forecast percentile vs the same issued month + leads. RPs under the bars: country mean vs the 45-yr hindcast. Niño3.4: +1.7 °C in Jul 2026. Data: ECMWF SEAS5 · Boundaries: TLS ADM1 (COD)