


Niger — ERA5 rainfall × ENSO teleconnection

Pixelwise partial correlation of trimester rainfall with Niño3.4 — the other climate modes held constant — ERA5 0.25° grid, 1981–2025


Unique ENSO signal — strongest significant trimester per pixel




Niger — current SEAS5 seasonal forecast

Issued August 2026 · drought/flood return-period categories vs the same-issuance SEAS5 hindcast (1981–2025) · skill-shaded · native 0.4° grid · ADM1 boundaries


JJA 2026 — in season — 22nd pct ← driest overall




JAS 2026 — in season — 24th pct



ASO 2026 — in season — 29th pct



SON 2026 — lead 1 mo — 40th pct




Forecast (high skill):

- Drought — ≥ 10 -yr return period
- Drought — 3-10-yr return period
- Roughly normal
- Flood — 3-10-yr return period
- Flood — ≥ 10 -yr return period

Other:

- Roughly normal (mod skill)
- Low skill
- Outside rainy season

Country-mean rainfall (mm/day)



Categories from the detrended skill cube vs the same issued month + leads. Bar colours + RPs: the app's country-level forecast (* = off-season).
Niño3.4: +1.7 °C in Jul 2026. In season = obs + fcst.
Data: ECMWF SEAS5 · Boundaries: NER ADM1 (COD)