


Timor-Leste — ERA5 rainfall × ENSO teleconnection

Pixelwise partial correlation of trimester rainfall with Niño3.4 — the other climate modes held constant — ERA5 0.25° grid, 1981–2025

Unique ENSO signal — strongest significant trimester per pixel




Data: ERA5 monthly precip (0.25°), NOAA PSL Niño3.4. Bottom row: canonical trimesters.


Timor-Leste — current SEAS5 seasonal forecast

Issued August 2026 · drought/flood return-period categories vs the same-issuance SEAS5 hindcast (1981–2025) · skill-shaded · native 0.4° grid · ADM1 boundaries

SON 2026 — lead 1 mo — mean 0th pct



OND 2026 — lead 2 mo — mean 0th pct ← driest overall




Forecast (high skill):

-  Drought — $\geq$ 10-yr return period
-  Drought — 3-10-yr return period
-  Roughly normal
-  Flood — 3-10-yr return period
-  Flood — $\geq$ 10-yr return period


Other:

-  Roughly normal (mod skill)
-  Low skill
-  Outside rainy season


NDJ 2026/27 — lead 3 mo — mean 3rd pct



DJF 2026/27 — lead 4 mo — mean 9th pct



Country-mean rainfall (mm/day)



Categories from the detrended skill cube vs the same issued month + leads. Bar colours + RPs: the app's country-level forecast (* = off-season).

Niño3.4: +1.7 °C in Jul 2026.

Data: ECMWF SEAS5 · Boundaries: TLS ADM1 (COD)