


Russia: how does El Niño generally affect seasonal rainfall?

Pixelwise partial correlation of rainfall with Niño3.4 (ENSO), other climate modes held constant — ERA5 0.25° grid, 1981-2025



Brown = El Niño years tend to be drier than normal in that season; blue = wetter.

Method (the teleconnections page's partial pass):


- Trimester rainfall means per pixel, all 12 overlapping 3-month windows, 1981–2025
- Partial r of rainfall vs Niño3.4, holding DMI (IOD), TNA, TSA and AMM constant
- Each mode at its per-pixel best lag (0–3 mo)
- Two-tailed $p < 0.05$, df adjusted for controls
- All seasons kept (no rainy-season filter)



Data: ERA5 monthly precip (0.25°), NOAA PSL Niño3.4. Bottom row: the four quarters of the year. Distant territories beyond the map edge not shown.

Russia: what seasonal rainfall is predicted this year (August-issued forecast)?

SEAS5 (issued August 2026) accounts for El Niño and every other driver · return-period categories vs the 1981–2025 hindcast · skill-shaded · ADM1



Forecast (high skill):


- Drought — ≥ 10 -yr return period
- Drought — 3–10-yr return period
- Roughly normal
- Flood — 3–10-yr return period
- Flood — ≥ 10 -yr return period

Other:

- Mod. skill
- Low skill
- Outside rainy season



Country-mean rainfall (mm/day)
black = normal · coloured = forecast



White hatch on a colour = moderate skill.
Tile subtitles: the country-level return period and percentile from the alerts app.
In season = observed months + forecast.
Niño3.4: +1.7 °C in Jul 2026.

Data: ECMWF SEAS5 · Boundaries: RUS ADM1 (COD)
Distant territories beyond the map edge not shown.