



# Guinea: how does El Niño generally affect seasonal rainfall?

Pixelwise partial correlation of rainfall with Niño3.4 (ENSO), other climate modes held constant — ERA5 0.25° grid, 1981–2025

Unique ENSO signal — strongest significant trimester per pixel



## Partial correlation with Niño3.4



Brown = El Niño years tend to be drier than normal in that season; blue = wetter.


Method (the teleconnections page's partial pass):

- Trimester rainfall means per pixel, all 12 overlapping 3-month windows, 1981–2025
- Partial  $r$  of rainfall vs Niño3.4, holding DMI (IOD), TNA, TSA and AMM constant
- Each mode at its per-pixel best lag (0–3 mo)
- Two-tailed  $p < 0.05$ , df adjusted for controls
- All seasons kept (no rainy-season filter)


JFM



AMJ



JAS



OND




Data: ERA5 monthly precip (0.25°), NOAA PSL Niño3.4. Bottom row: the four quarters of the year.


# Guinea: what seasonal rainfall is predicted this year (August-issued forecast)?

SEAS5 (issued August 2026) accounts for El Niño and every other driver · return-period categories vs the 1981–2025 hindcast · skill-shaded · ADM1

**JJA 2026 — in season**  
1-in-46 dry · 0th pct



**JAS 2026 — in season**  
1-in-9 dry · 9th pct



**ASO 2026 — in season**  
1-in-3 wet · 71st pct



**SON 2026 — lead 1 mo**  
1-in-12 wet · 93rd pct



## Forecast (high skill):

- Drought —  $\geq 10$ -yr return period
- Drought — 3–10-yr return period
- Roughly normal
- Flood — 3–10-yr return period
- Flood —  $\geq 10$ -yr return period

## Other:

- Roughly normal (mod skill)
- Low skill
- Outside rainy season

**OND 2026 — lead 2 mo**  
off-season




**NDJ 2026/27 — lead 3 mo**  
off-season



**DJF 2026/27 — lead 4 mo**  
off-season



Country-mean rainfall (mm/day)  
black = normal · coloured = forecast



White hatch on a colour = moderate skill.  
Tile subtitles: the country-level return period and percentile from the alerts app.  
Driest outlook: JJA 2026 (1-in-46 dry).  
In season = observed months + forecast.  
Niño3.4: +1.7 °C in Jul 2026.