


St. Kitts and Nevis: how does El Niño generally affect seasonal rainfall?

Pixelwise partial correlation of rainfall with Niño3.4 (ENSO), other climate modes held constant — ERA5 0.25° grid, 1981–2025


Unique ENSO signal — strongest significant trimester per pixel



Data: ERA5 monthly precip (0.25°), NOAA PSL Niño3.4. Bottom row: the four quarters of the year.

St. Kitts and Nevis: what seasonal rainfall is predicted this year (August-issued forecast)?

SEAS5 (issued August 2026) accounts for El Niño and every other driver · return-period categories vs the 1981-2025 hindcast · skill-shaded · ADM1



Forecast (high skill):

- Drought — ≥ 10 -yr return period
- Drought — 3-10-yr return period
- Roughly normal
- Flood — 3-10-yr return period
- Flood — ≥ 10 -yr return period

Other:

- Mod. skill
- Low skill
- Outside rainy season

Country-mean rainfall (mm/day)
black = normal · coloured = forecast

White hatch on a colour = moderate skill.
Tile subtitles: the country-level return period and percentile from the alerts app.
Driest outlook: NDJ 2026/27 (1-in-46 dry).
In season = observed months + forecast.
Niño3.4: +1.7 °C in Jul 2026.

Data: ECMWF SEAS5 · Boundaries: KNA ADM1 (COD)