


# Martinique: how does El Niño generally affect seasonal rainfall?

Pixelwise partial correlation of rainfall with Niño3.4 (ENSO), other climate modes held constant — ERA5 0.25° grid, 1981–2025

Unique ENSO signal — strongest significant trimester per pixel



Data: ERA5 monthly precip (0.25°), NOAA PSL Niño3.4. Bottom row: the four quarters of the year.

# Martinique: what seasonal rainfall is predicted this year (August-issued forecast)?

SEAS5 (issued August 2026) accounts for El Niño and every other driver · return-period categories vs the 1981–2025 hindcast · skill-shaded · ADM1

JJA 2026 — in season  
1-in-23 dry · 2nd pct



**JAS 2026 — in season**  
**1-in-46 dry · 0th pct**



ASO 2026 — in season  
1-in-46 dry · 0th pct



SON 2026 — lead 1 mo  
1-in-46 dry · 0th pct



### Forecast (high skill):

- Drought —  $\geq 10$ -yr return period
- Drought — 3–10-yr return period
- Roughly normal
- Flood — 3–10-yr return period
- Flood —  $\geq 10$ -yr return period

### Other:

- Mod. skill
- Low skill
- Outside rainy season

OND 2026 — lead 2 mo  
1-in-46 dry · 0th pct




NDJ 2026/27 — lead 3 mo  
1-in-46 dry · 0th pct



DJF 2026/27 — lead 4 mo  
1-in-46 dry · 0th pct



Country-mean rainfall (mm/day)  
black = normal · coloured = forecast



White hatch on a colour = moderate skill.  
Tile subtitles: the country-level return period and percentile from the alerts app.  
Driest outlook: JAS 2026 (1-in-46 dry).  
In season = observed months + forecast.  
Niño3.4: +1.7 °C in Jul 2026.