



# Guatemala: how does El Niño generally affect seasonal rainfall?

Pixelwise partial correlation of rainfall with Niño3.4 (ENSO), other climate modes held constant — ERA5 0.25° grid, 1981–2025

Unique ENSO signal — strongest significant trimester per pixel



## Partial correlation with Niño3.4



Brown = El Niño years tend to be drier than normal in that season; blue = wetter.

Method (the teleconnections page's partial pass):


- Trimester rainfall means per pixel, all 12 overlapping 3-month windows, 1981–2025
- Partial  $r$  of rainfall vs Niño3.4, holding DMI (IOD), TNA, TSA and AMM constant
- Each mode at its per-pixel best lag (0–3 mo)
- Two-tailed  $p < 0.05$ , df adjusted for controls
- All seasons kept (no rainy-season filter)

Data: ERA5 monthly precip (0.25°), NOAA PSL Niño3.4. Bottom rows: all 12 rolling trimesters.

JFM




FMA




MAM




AMJ




MJJ




JJA




JAS



ASO



SON



OND



NDJ





DJF



# Guatemala: what seasonal rainfall is predicted this year (August-issued forecast)?

SEAS5 (issued August 2026) accounts for El Niño and every other driver · return-period categories vs the 1981–2025 hindcast · skill-shaded · ADM1



White hatch on a colour = moderate skill.  
Tile subtitles: the country-level return period and percentile from the alerts app.  
Driest outlook: JAS 2026 (1-in-46 dry).  
In season = observed months + forecast.  
Niño3.4: +1.7 °C in Jul 2026.