


Micronesia: how does El Niño generally affect seasonal rainfall?

Pixelwise partial correlation of rainfall with Niño3.4 (ENSO), other climate modes held constant — ERA5 0.25° grid, 1981–2025

Unique ENSO signal — strongest significant trimester per pixel



Partial correlation with Niño3.4

- Positive, strong ($r \geq 0.5$)
- Positive, moderate ($0.3 \leq r < 0.5$)
- Negative, moderate ($-0.5 < r \leq -0.3$)
- Negative, strong ($r \leq -0.5$)
- No significant signal ($p \geq 0.05$)

Brown = El Niño years tend to be drier than normal in that season; blue = wetter.

Method (the teleconnections page's partial pass):

- Trimester rainfall means per pixel, all 12 overlapping 3-month windows, 1981–2025
- Partial r of rainfall vs Niño3.4, holding DMI (IOD), TNA, TSA and AMM constant
- Each mode at its per-pixel best lag (0–3 mo)
- Two-tailed $p < 0.05$, df adjusted for controls
- All seasons kept (no rainy-season filter)

Data: ERA5 monthly precip (0.25°), NOAA PSL Niño3.4. Bottom row: the four quarters of the year. Distant territories beyond the map edge not shown.

Micronesia: what seasonal rainfall is predicted this year (August-issued forecast)?

SEAS5 (issued August 2026) accounts for El Niño and every other driver · return-period categories vs the 1981–2025 hindcast · skill-shaded · ADM1

