

COVID-19 Projections: Iraq

Report Date: 18 Jan 2021

This report summarizes the COVID-19 model results for Iraq, developed by the OCHA Centre for Humanitarian Data in partnership with the Johns Hopkins University Applied Physics Laboratory. These projections are based on COVID-19 cases and deaths data up to 18 January 2021. The data is sourced from World Health Organization (WHO) and the country's Ministry of Public Health (MOPH). For dynamic updates to this data and more, see the [HDX COVID-19 Map Explorer](#). For additional information, please contact Leonardo Milano at: leonardo.milano@un.org.

1. Key Messages

Current Situation (as of 18 Jan 2021)

- Per the WHO, a total of 608,232 cases and 12,944 deaths have been reported with the current number of daily new cases standing at 756.3 (7-day average).
- The current number of active severe cases requiring hospitalization is estimated at 1,584 - 2,697.
- Progress towards containment is insufficient: the entire country is seeing Community or Accelerated Spread (yellow or orange risk levels), while Al-Sulaymaniyah is at a Tipping Point (red risk level). More intense control efforts are critically needed (see section 2).
- We note that mitigation measures currently in place include an awareness campaign, face mask wearing (50% compliance) and limits on public gatherings (50% compliance). Please email us (leonardo.milano@un.org) if this information is inaccurate or incomplete as it affects the accuracy of projections.
- Please keep mind that the current assessment of the situation and all projections depend on complete, accurate data. The model might underestimate the crisis even after accounting for some underreporting.

National Projections (in the next 4 weeks or by 15 Feb 2021)

- In 4 weeks, we project that an additional 22,456 - 27,956 cases and 206 - 290 deaths will be reported if current NPIs are maintained.
- Currently implemented mitigation strategies are insufficient to curb the spread of the virus: cases are projected to increase even if NPIs are maintained. More intense control efforts are needed.
- Lifting the NPIs would lead to a markedly sharp increase in cases and deaths (up to 59,440 more cases and up to 219 more deaths; see sections 2 and 3 for details). Due to the lag between cases and deaths, a larger number of cases will ultimately result in higher deaths 2 - 4 weeks later.
- Active severe cases requiring healthcare are projected to reach 1,897 - 2,541 if NPIs are maintained, and to jump to 6,405 - 8,827 if they are lifted (see section 3).

Subnational Projections (in the next 2 weeks or by 01 Feb 2021)

- In the next two weeks and assuming NPIs are maintained, the situation in the country is projected to significantly worsen in Baghdad, although most other regions will see slight decreases in incidence. Still, those decreases will not suffice to suppress community spread and regions will remain at a yellow or worse level of risk (see section 4).
- The projected incidence indicates a need for more intense control strategies.

2. Current Situation (as of 18 Jan 2021)


Containment Progress

This report leverages a framework to provide guidance in the interpretation of incidence rates so decision-makers can more readily understand how effective the response has been in containing the virus. The framework was devised by experts from the Harvard Global Health Institute, Harvard's Edmond J. Safra Center for Ethics, and a network of research and policy organizations (more about the collaborative framework [here](#)).

The framework defines risk levels that indicate whether a region is on track for containment and can help decision-makers know where they are at the moment. The levels do not in themselves provide information about how to respond but do communicate the intensity of effort needed for control of COVID at varying levels of community spread. In addition to paying attention to the levels, decision-makers should pay close attention to direction of trend and rate of change (see section 4 for those metrics at the subnational level).

The map below illustrates regional risk levels as defined by estimated total incidence rate (daily new cases per 100,000 people as of 18 Jan 2021). Total cases are estimated from case reporting rate to adjust for underreporting. The table details the cutoffs for each risk level along with strategies of disease response needed for containment.

Current Estimated Total New Daily Cases
Per 100,000 People



| Risk Level | Case Incidence* | Status | Intensity of Control Effort Needed |
|------------|-----------------|--------------------------|--|
| Red | 25+ | Tipping Point | Stay-at-home orders necessary |
| Orange | 10-25 | Accelerated Spread | Strategic choices must be made about which package of non-pharmaceutical interventions to use for control. Stay-at-home orders are advised, unless viral testing and contact tracing capacity are implementable at levels meeting surge indicator standards. |
| Yellow | 1-10 | Community Spread | Strategic choices must be made about which package of non-pharmaceutical interventions to use for control |
| Green | <1 | On Track for Containment | On track for containment, conditional on continuing use of viral testing and contact tracing for surveillance and to contain spikes and outbreaks. |

*Daily new cases per 100,000 people as reported by MOPH

See [Key Metrics for COVID Suppression](#) for additional guidance on control effort needed.

Key Figures: Current Cases and Deaths¹²³

| | Cases | | Deaths | |
|--------------------|-----------------|------------|-----------------|------------|
| | Daily New Cases | Cumulative | Daily New Cases | Cumulative |
| Based on MOPH data | NA | 608,231 | NA | 12,944 |
| Based on WHO data | 756.3 | 608,232 | 7 | 12,944 |

"Daily new cases" in this table is the average over the last 7 days.

Most recent data from MOPH: 2021-01-17

Key Figures: Current Severe Cases

| | Active |
|----------|--|
| | Severe Cases Requiring Hospitalization |
| Estimate | 1,584 - 2,697 |

Note on data reliability

The limitations of COVID-19 reported data should be taken into consideration when interpreting metrics and projections. Sources may diverge in the counts they report (see WHO vs MOPH figures above); data reports may lag by several days or be missing altogether on certain days (see date of latest data above); cases and deaths are almost certainly underreported and their numbers are affected by testing practices. Scenario modelling (NPI vs non-NPI projections) relies on the freshness and accuracy of the information provided in the ACAPS database (see footnote 3). We strongly encourage the reader to ensure the database is up to date and to contact the Centre for Humanitarian Data with any suggestions of additional data sources or improvements to existing ones.

These are common limitations. This report aims to help the reader understand the situation on the ground through comparing and contrasting multiple data sources and estimates. For instance, we present data reported by the WHO and the MOPH, projected cases and deaths, and incidence of total cases, an estimate of the true number of cases factoring in the case reporting rate (ie., how many cases are likely unreported.) The projections are best estimates based on available data.

¹**Reported cases** refers to the number of infections expected (current situation) or expected to be reported (projections). Projections take into account the **case reporting rate** which corresponds to the estimated number of COVID-19 infections that are actually tested, confirmed and reported. The case reporting rate is calculated based on the number of deaths and cases reported by the WHO in the last 30 days.

²**Severe cases** refers to the number of people which will have severe symptoms and may require healthcare support. Projections are calculated as a proportion of the reported cases, and are based on planning parameters for case severity and the vulnerability of a given region.



³**Case Fatality Rate** refers to the estimated proportion of deaths compared to the total number of people diagnosed with the disease.

3. National Projections⁴⁵ (for the next 4 weeks or by 15 Feb 2021)



Projected Cases and Deaths

| | Change | |
|------------------------------|-----------------|--------------|
| | Added Cases | Added Deaths |
| With current NPIs maintained | 22,456 - 27,956 | 206 - 290 |
| With no NPIs | 61,718 - 81,896 | 305 - 425 |

The figures below present the historical data on daily new cases and deaths, and their projected trends. Trends are represented by a green line for the “Current NPIs maintained” scenario and a red line for the “No NPIs in place” scenario. Note that deaths typically lag reported cases by 2-4 weeks.



The figures below show the comparison between two data sources: national level data from WHO in light blue and subnational data from the Ministry of Public Health in dark blue.




⁴The regional data provided by the Ministry of Public Health are used to generate projections at the subnational level, which are then aggregated to the national level.

⁵**Non-pharmaceutical interventions - NPIs** are all measures implemented by different actors with the aim of reducing the spread and the impact of COVID-19. The NPIs currently in place are extracted from the [ACAPS database](#) and complemented with additional contextual information provided by our partners in the country.

Projected Severe Cases

The figures below show the projected trends for active severe cases requiring hospitalizations. In green are the projections under the “Current NPIs maintained” scenario while in red are the projections under “No NPIs in place” scenario.

| | Active Severe Cases |
|------------------------------|--------------------------------|
| With current NPIs maintained | 1,897 - 2,541 |
| With no NPIs | 6,405 - 8,827 |




4. Subnational Projections (for the next 2 weeks or by 01 Feb 2021)

Projected Incidence

The map below displays the projected risk level of every ADMIN1 region, assuming the current NPIs are maintained. It represents the projected total cases as estimated from case reporting rate to adjust for underreporting. See Section 1 for detail on the relationship between risk level and progress towards virus containment.

Projected Estimated Total New Daily Cases
Per 100,000 People



| Risk Level | Status |
|------------|--------------------------|
| Red | Tipping Point |
| Orange | Accelerated Spread |
| Yellow | Community Spread |
| Green | On Track for Containment |

Projected Changes In Incidence

Below are the projected incidence rates (= projected daily new cases per 100k people on 01 Feb 2021) and projected absolute changes in incidence between then and today for every region. These metrics do not include adjustments for underreporting.

Regions are ordered in decreasing order of change magnitude, from largest increase to largest decrease. Note that Incidence Rates are rounded to the third decimal; therefore a 0.000 incidence rate may mean that no new daily cases are projected or that fewer than 1 cases per 100 million people are projected.

While the magnitude of change may signal how successful mitigation strategies are projected to be, it should be evaluated in conjunction with incidence when considering adding or lifting suppression strategies. For instance, a region may be projected to show a large decrease but remain at a concerning high level of incidence that require intense control efforts.

| Region | Projections in cases per 100k people | |
|------------------------------------|--------------------------------------|-----------|
| | Change | Incidence |
| <i>Increasing or Stable</i> | | |
| Baghdad | +10.77 | 26.860 |
| Al-Anbar | +0.5 | 3.945 |
| <i>Decreasing</i> | | |
| Ninewa | -0.02 | 8.593 |
| Maysan | -0.42 | 2.553 |
| Babil | -0.46 | 9.775 |
| Thi Qar | -0.46 | 2.861 |
| Salah Al-Din | -0.76 | 7.087 |
| Al-Muthanna | -0.86 | 4.071 |
| Al-Basrah | -0.88 | 5.740 |
| Al-Qadissiya | -0.89 | 5.588 |
| Erbil | -1.4 | 5.760 |
| Al-Najaf | -1.55 | 6.775 |
| Kerbala | -1.84 | 13.722 |
| Diyala | -2.41 | 16.238 |
| Duhok | -3.18 | 12.074 |
| Wassit | -3.86 | 9.832 |
| Kirkuk | -4.14 | 19.970 |
| Al-Sulaymaniyah | -14.53 | 18.041 |

Background on Model Methodology

The Centre established a partnership with the Johns Hopkins University Applied Physics Laboratory to develop a COVID-19 model which provides projections and insights related to the **scale** of the crisis, the **duration** of the crisis in a specific location, and how different response **interventions** are expected to impact the epidemic curve.

The team is using an **SEIR (Susceptible, Exposed, Infectious, Recovered)** model of infectious disease dynamics which is considered the simplest and most effective technique used in the literature. The model is based on a progression from susceptible to either recovered or dead. Inputs include the reproduction rate (R_0), case fatality rate (CFR), and estimated probabilities that an individual person may contract COVID-19. The model then simulates an outbreak and provides estimates for cases, severe cases/hospitalizations, and deaths.



The key features of the model include:

- **Tuning on reported data** The estimation of the main parameters (mainly the reproduction rate R_0 and the case reporting rate) is tuned according to the observed recent trends in reported COVID-19 cases.
- **Subnational** The model provides COVID-19 projections at the subnational level, matching the administrative level at which COVID-19 cases are reported.
- **Spatial spread** The density of roads is used to estimate the expected mobility patterns and to simulate the spread of COVID-19 between administrative units.
- **Population stratification** The model fidelity is increased by taking into consideration:
 - The age structure of the population at the subnational level
 - The expected probability of contact between populations of different age groups, including contacts expected to happen at work, school, home and everywhere else (social mixing)
 - Vulnerability factors such as food insecurity and household air pollution.
- **Non-pharmaceutical interventions (NPIs)** The model simulates the expected impact of NPIs at the subnational level, and also how the outbreaks is influenced by changing NPIs implemented over time. The NPIs currently implemented can be categorised in three main groups:
 - Mobility based NPIs, which would limit the spread of disease between administrative units (e.g. border closures)
 - Contact based NPIs, which reduce the probability of contact between specific groups (e.g. shielding of the elderly, closing schools)
 - R_0 based NPIs, which reduce the overall reproduction rate (e.g. awareness campaigns, curfews)