

Hippocampal and prefrontal cortical mechanisms for goal-directed and memory-guided behavior.

Michael E. Hasselmo  
Department of Psychology  
Center for Memory and Brain  
and Program in Neuroscience,  
Boston University

#### ABSTRACT:

Mechanisms of memory-guided behavior were analyzed in a simulation of the hippocampus and prefrontal cortex which guided movements of a virtual rat in a virtual delayed spatial alternation task, in which correct turning response depended upon the previous episode. At the choice point, both possible episodes were retrieved in entorhianl cortex and temporal context from dentate gyrus and region CA3 allowed selection of one episode in CA1. Episodic retrieval allowed selection of correct goal-directed movement by prefrontal cortex circuits. Network activity replicates current source density during theta rhythm, as well as theta phase precession and context sensitivity of place cells.

#### SUMMARY:

Lesions of the hippocampus cause impairments in memory guided tasks including delayed spatial alternation, delayed match to position and the 8-arm radial maze (Markowska et al., 1989; Ennaceur et al., 2002, see Eichenbaum and Cohen, 2001). Many models have addressed hippocampal memory function, but few have explicitly modeled the actions of a virtual rat in these types of tasks. Ongoing research in the Hasselmo laboratory uses the MATLAB program and the CATACOMB simulation package (Cannon et al., 2002) to analyze cortical mechanisms involved in memory guided behavior in these tasks.

A number of models have addressed memory-guided behavior, but do not simultaneously address neurophysiological phenomena such as theta rhythm (Sharp et al., 1996; Redish and Touretzky, 2000). Recent models from the Hasselmo lab have addressed the role of theta rhythm in memory guided behavior (Hasselmo et al., 2002b, 2002c). However, several new features have been addressed in the models to be presented at CNS 2004. In particular, previous models from this lab used an interaction of a reverse spread of activity from the goal and forward retrieval activity to select next desired destination in the form of region CA1 activity. However, general goal-directed behavior is spared by hippocampal lesions, and appears to involve prefrontal cortical circuits. Therefore, we now use a model of prefrontal cortical circuitry to address the interaction of goal and current state, to obtain selection of the next action. This network implementation of goal directed function has features resembling aspects of temporal difference learning and action selection in reinforcement learning (Sutton and Barto, 1998). The hippocampal simulation interacts with this representation of prefrontal cortex.

In addition, the new simulations address context-dependent retrieval of full sequences of activity, which is necessary for memory-guided behavior in spatial alternation. This requires both encoding of the sequential associations between elements of

an episode, as well as encoding of temporal context analogous to that described by Howard and Kahana (Howard and Kahana, 2002; Howard et al., 2004). Activity necessary for both aspects can be provided by sustained activity in entorhinal cortex due to cholinergic activation of intrinsic regenerating cation currents (Klink and Alonso, 1997; Fransen et al., 2002). As described in previous models (Jensen and Lisman, 1996; Hasselmo et al., 2002c), these currents allow slow behavioral transitions to cause sequential spiking which falls within the window of spike timing dependent synaptic plasticity (Levy and Steward, 1983; Bi and Poo, 1998). In addition, gradual decay of these representations allows formation of associations between a gradually changing temporal context (Howard and Kahana, 2002) and individual behavioral events.

Previous simulations of the hippocampus have demonstrated forward retrieval of individual sequences (Levy, 1996; Jensen and Lisman, 1996; Lisman, 1999). The simulation presented here addresses the additional problem of selective episodic retrieval without interference from other episodes. In the new simulations, sequential retrieval of all possible previous episodes occurs due to associations within the circuit. Selection of a single current episode without interference from other episodes involves the convergence of sequential retrieval with temporal context in region CA1 of the simulation. This utilizes dynamical interactions during theta rhythm which match features of the physiological data on theta rhythm obtained with current source density analysis (Brankack et al., 1993). The sequential read out of previous episodes resembles theta phase precession (Skaggs et al., 1996), but has the additional feature of context-dependent retrieval, which matches evidence for place cell responses specific to prior or future responses in spatial alternation (Wood et al., 2000). As before, the model requires separate phases of encoding and retrieval, such that synaptic modification for encoding occurs on one phase of theta rhythm (Holscher et al., 1997; Huerta and Lisman, 1993; Hyman et al., 2003), whereas retrieval occurs on a separate phase of theta rhythm (Hasselmo et al., 2002a).

Bi GQ, Poo MM (1998) Synaptic modifications in cultured hippocampal neurons: dependence on spike timing, synaptic strength, and postsynaptic cell type. *J. Neurosci.* 18(24):10464-72.

Brankack, J, Stewart, M, Fox, SE (1993). Current source density analysis of the hippocampal theta rhythm: associated sustained potentials and candidate synaptic generators. *Brain Research*, 615(2): 310-327

Cannon RC, Hasselmo ME, Koene RA (2002) From biophysics to behavior: Catacomb2 and the design of biologically plausible models for spatial navigation. *Neuroinformatics* 1:3-42.

Eichenbaum, H. & Cohen, N.J. (2001) *From Conditioning to Conscious Recollection: Memory Systems of the Brain*. Oxford University Press.

Ennaceur A., Neave N. , Aggleton J.P. (1996) Neurotoxic lesions of the perirhinal cortex do not mimic the behavioural effects of fornix transection in the rat. *Behav. Brain Res.* 80: 9-25.

Fransen, E., Alonso, A.A. and Hasselmo, M.E. (2002) Simulations of the role of the muscarinic-activated calcium-sensitive non-specific cation current I(NCM) in entorhinal neuronal activity during delayed matching tasks. *J. Neurosci.* 22(3):1081-1097.

Hasselmo, M.E., Bodelon, C. and Wyble, B.P. (2002a) A proposed function for

hippocampal theta rhythm: Separate phases of encoding and retrieval enhance reversal of prior learning. *Neural Computation* 14: 793-817.

Hasselmo, M.E., Hay, J., Ilyn, M. and Gorchetnikov, A. (2002b) Neuromodulation, theta rhythm and rat spatial navigation. *Neural Networks*, 15: 689-707.

Hasselmo, M.E., Cannon, R.C. and Koene, R.A. (2002c) A simulation of parahippocampal and hippocampal structures guiding spatial navigation of a virtual rat in a virtual environment: A functional framework for theta theory. In: Witter, M.P., Wouterlood, F.G. (eds.) *The parahippocampal region: organisation and role in cognitive functions*. Oxford University Press: Oxford. pp. 139-161.

Holscher, C, Anwyl, R, Rowan, MJ (1997). Stimulation on the positive phase of hippocampal theta rhythm induces long-term potentiation that can be depotentiated by stimulation on the negative phase in area CA1 in vivo. *J Neurosci*, 17(16): 6470-6477.

Howard, M.W., Fotedar, M.S., Datey, A.V., Hasselmo, M.E. (2003) The temporal context model in spatial navigation and relational learning: Toward a common explanation of medial temporal lobe function across domains. *Psychological Review*, submitted.

Howard, M.W. and Kahana, M.J. (2002) A distributed representation of temporal context. *J. Math. Psychol.* 46(3): 269-299.

Huerta PT, Lisman JE (1995). Bidirectional synaptic plasticity induced by a single burst during cholinergic theta oscillation in CA1 in vitro. *Neuron*, 15(5): 1053-1063.

Hyman, J.M., Wyble, B.P., Goyal, V., Rossi, C.A., Hasselmo, M.E. (2003) Stimulation in hippocampal region CA1 in behaving rats yields LTP when delivered to the peak of theta and LTD when delivered to the trough. *J. Neurosci.* 23(37):11725-31.

Jensen O, Lisman JE (1996) Novel lists of 7 +/- 2 known items can be reliably stored in an oscillatory short-term memory network: interaction with long-term memory. *Learn Mem.* 3(2-3):257-63.

Klink R, Alonso A. (1997) Muscarinic modulation of the oscillatory and repetitive firing properties of entorhinal cortex layer II neurons. *J. Neurophysiol.* 77(4): 1813-1828.

Levy WB (1996) A sequence predicting CA3 is a flexible associator that learns and uses context to solve hippocampal-like tasks. *Hippocampus*, 6, 579-590.

Levy WB, Steward O (1983) Temporal contiguity requirements for long-term associative potentiation/ depression in the hippocampus. *Neuroscience*. 8(4):791-7.

Lisman JE (1999) Relating hippocampal circuitry to function: recall of memory sequences by reciprocal dentate-CA3 interactions. *Neuron* 22:233-242.

Lisman JE, Idiart MA (1995) Storage of 7 +/- 2 short-term memories in oscillatory subcycles. *Science* 267:1512-1515.

Markowska AL, Olton D.S., Murray E.A. and Gaffan D. (1989) A comparative analysis of the role of fornix and cingulate cortex in memory: rats. *Exp. Brain Res.* 74(1): 187-201.

Redish AD, Touretzky DS (1998) The role of the hippocampus in solving the Morris water maze. *Neural Comput* 10:73-111.

Sharp PE, Blair HT, Brown M (1996) Neural network modeling of the hippocampal formation spatial signals and their possible role in navigation: a modular approach. *Hippocampus* 6:720-734.

Skaggs, W.E., McNaughton, B.L., Wilson, M.A. and Barnes, C.A. (1996) Theta phase precession in hippocampal neuronal populations and the compression of temporal sequences. *Hippocampus* 6, 149-172.

Sohal VS, Hasselmo ME (1998b) GABA(B) modulation improves sequence disambiguation in computational models of hippocampal region CA3. *Hippocampus* 8:171-193.

Sutton R.S. and Barto A.G. (1998) *Reinforcement Learning: An Introduction*. MIT Press: Cambridge, MA.

Wood ER, Dudchenko PA, Robitsek RJ, Eichenbaum H (2000) Hippocampal neurons encode information about different types of memory episodes occurring in the same location. *Neuron* 27:623-633.