

The Role of Glyclinergeric Interneurons in the Dorsal Column Nuclei

E. Sánchez¹, J. Aguilar², C. Rivadulla³ and A. Canedo²

¹Grupo de Sistemas Intelixentes (GSI)
Departamento de Electrónica e Ciencias da Computación, Facultade de Física.
Universidade de Santiago de Compostela
15782 Santiago de Compostela, Spain
eduardos@usc.es

²Departamento de Fisioloxía, Facultade de Medicina.
Universidade de Santiago de Compostela
15782 Santiago de Compostela, Spain
jreyes@usc.es, fsancala@usc.es

³Departamento de Medicina, Escola universitaria de Fisioterapia.
Universidade da Coruña
15006 A Coruña, Spain
casto@udc.es

Abstract

The aim of this paper is to provide new insights about the circuitry and the role of the Dorsal Column Nuclei (DCN) in the processing of somatosensory information. The presence of Glycinergeric cells, a second type of DCN interneurons in addition to well-known GABAergic interneurons, opens the door to more complex interactions between cuneate cells as well as to new hypothesis about the computational implications of such interactions. The research posed here fits in a broader context in the field of the sensory systems and regards with the general issue about the role of subcortical structures (i.e. the Thalamus) in the processing of sensory information.

Keywords: Dorsal Column Nuclei, Somatosensory System, Computational Models, Optimal Coding.

1. Introduction

The DCN are located within the brainstem, close to the most rostral area of the spinal chord, and are constituted by the Gracile (GN) and the Cuneate (CN) nuclei. They present a similar organization, but differ on the origin of their primary afferents. The DCN process somatosensory information coming from Primary Afferent Fibres (PAF) and send it to the Thalamus via the Medial Lemniscus (ML). The experimental data were obtained from the dorsal part of the middle region of the Cuneate Nucleus (CN), which is located from the obex to about 4 mm caudal to it in the cat. This region receives information from somatotopically organized primary afferents sensible to both light touching or hair displacement, and distal Receptive Fields (RFs). On top of that, the CN receives an important corticofugal projection from layer V of sensory motor cortex [8, 13, 14, 18], with a topographic organization similar to the Primary afferents [3, 4, 6, 9, 10, 11].

A key point to better understand the cuneate function would undoubtedly be the precise knowledge of the RF structure of the different cell types implicated in the CN circuitry: Relay or CuneoLemniscal (CL) cells, and GABAergic (GAB) interneurons or non-CuneoLemniscal cells. The current knowledge is limited mainly to CL cells. It has been recently shown that their RF has an excitatory center and an inhibitory surround [7]. This spatial arrangement is generated through somatotopically organized afferent organization consisting on direct excitatory input on CL cells and GABAergic mediated inhibition from surrounding areas [7]. It has been demonstrated that when this circuit is constructed with realistic models of cuneate cells [15], it can perform edge detection as well as motion discrimination [16].

Moreover, there exists evidence of additional GAB interneurons performing recurrent lateral inhibition [2]. GABA participation seems evident when ejection of Bicuculline (BiCu) after Lemniscal Stimulation uncovers recurrent excitation (figure 1-A1). The effect of recurrent inhibition is clarified when recurrent activation is blocked after increasing Lemniscal stimulation to 2 mA (figure 1-A2). GAB interneurons seem confirmed in this circuitry when BiCu ejection (figure 1-A2, right) imposes the activity initially found under Control conditions (0.2 mA stimulation).

At this point, the presence of a second interneuronal type, the Glycinergic (GLY) interneurons, which constitute about 30% of total neuronal population in the rat [12], introduces further complexity into the circuit. It is known that GLY neurons evoke facilitation of Relay cells by inhibiting GAB interneurons [1, 2]. Figure 1-B shows that application of Strychnine, a glycine antagonist, after ML stimulation, allowed the blocking of the second, recurrent, spike.

Unfortunately we do not have experimental evidences related to the RF structure of these interneurons that could help us to better understand its role. So, we have resorted to computational models in order to explore the role that different spatial arrangements of GLY cells could have on the processing of CL cells.




Fig 1. Lemniscal stimulation and recurrent effects. ML stimulation and BiCu shows GABAergic neurons participating in lateral recurrent effects (A). ML stimulation and Strychnine ejection uncovers inhibitory effects on CL cells (B).

2. Methods

The computational model assumes that CL cells show excitatory center-inhibitory surround RFs, in which inhibition is performed by GABAergic interneurons. With regard the recurrent organization, and considering the experimental data discussed so far, we also have lateral inhibition through GAB interneurons over CL cells as well as a disinhibition mechanism mediated by GLY interneurons (figure 2).

For the simulation, we have implemented a bidimensional version of the circuitry in which all neurons have a square-shaped RF. Under this configuration, CL neurons have an afferent RF, which is constituted by excitatory input coming from the receptors at the centre of the square and inhibitory input from the surrounding edges, and a recurrent inhibitory contribution from corresponding GAB interneurons. These ones receive excitatory input only from CL cells with non-overlapped RF. Finally, GLY interneurons show a fully excitatory RF coming from CL cells with overlapped RF.

As we focus on circuit behaviour, we have simplified the neuronal models. The neurons are McCulloch-Pitts units, in which the output of the j th neuron is $y_j = \psi(\sum w_{ji}x_i)$, with the activation function ψ being of the threshold type. Contribution of synapse between neuron i th and neuron j th is modelled through weight w_{ji} . Weights and RF configurations are the most relevant free parameters in order to investigate the processing role of each neuronal type.




Fig 2. A model of the CN circuitry. CuneoLemniscal neurons (CL) have both afferent and recurrent influences. GABAergic neurons (GAB) seem to be involved in afferent inhibition as well as recurrent lateral inhibition. Glycinergic interneurons (GLY) perform disinhibition over CL cells by inhibiting GAB interneurons.

3. Results

We have built and test the circuit step by step, starting with the afferent connections, and iteratively adding the recurrent GAB and GLY layers. Figure 3 shows how a layer of CL neurons can perform edge detection based on afferent centre-surround organization. CL cells having object edges at the centre of their RF show a higher




Fig 3. Edge detection through afferent connections. Inputs of different forms and sizes were presented (top). CL cells (bottom) respond most remarkably to edges (black) than other parts of the objects (grey).

activity (black) than those cells receiving input from inner parts (grey). We have found that edge detection is quite robust against size, form and also intensity of the stimulus. However, in order to achieve robustness against stimulus intensity, a gain control mechanism like, for instance, shunting inhibition, seems to be required. As this is a problem out of the scope of this paper, the mechanism was not shown in figure 2.

Recurrent lateral inhibition was then added to the circuitry (figure 4), with a RF size of 9x9. After the first iteration, in which regular edge detection is performed, the output shows an oscillatory activity consisting of two states: in the first one, object edges are detected; in the second one, it is the middle region of the object what is highlighted. This result could be understood as an optimal spatial coding as CL output encodes the most relevant information: (1) edge location, and (2) that the object is of a solid type.

The encoding is robust against lateral inhibition strength, i.e the value of synapse weights between CL neurons and GAB interneurons. However, the effect depends on GABAergic interneurons RF. When RF has a size bigger than 21x21, the two-state oscillation disappears and the output seems to enter a chaotic pattern.




Fig 4. Afferent connections and recurrent lateral inhibition. The output shows an oscillatory pattern between two states that encode the edges and the object middle region.

The most interesting result is obtained when GLY interneurons are introduced (figure 5). The size of GLY RF was initially set to 3x3. The output evolves from an initial state, where the edges are detected, into a fill-in process that finishes with the complete covering of the object (figure 5). The bigger the GLY RF, the higher the speed of the whole process. As it happened with the GAB layer and the spatial coding, this temporal coding effect is also robust against inhibition strength.

The fill-in process could be explained by considering the activity carried out in both GAB and GLY maps. The GAB map presents a final state in which the object edges are represented, while the GLY map finally represents the object size. So it seems that CL output is caused by the interplay between the lateral recurrent inhibition (GAB map) and the recurrent disinhibition (GLY map).




Fig 5. CL outputs with the complete circuitry. Iterations (1-6) indicate how CL output evolves with time.

4. Discussion

The CN processing shown in Fig. 5 can be understood as an optimal spatio-temporal coding. Spatial coding is clearly performed in Fig. 4, in which the most salient regions of the object (edges and middle zone) are detected. On top of that, an optimal temporal coding, introduced by GLY neurons, allows a progressive transmission of information up to the Vento-Postero-Lateral (VPL) nucleus in the Thalamus. It starts with high-level informative regions (edges, Fig. 5-2) and finishes with low-level informative regions (core middle zone, Fig. 5-6). This behaviour presents certain similarities with the popular JPEG compression algorithm, which is the responsible of the progressive fill-in of images in web browsers. On each domain, both VPL and the user might stop the transmission of information when the available information is sufficient enough to identify the incoming pattern.

There is an issue that concerns about why the whole object appears interleaved between the fill-in stages (Figs. 5-3 and 5-5). There is experimental evidence that VPL responds with much higher reliability to a number of consecutive spikes than to single spikes [17]. Considering this, we can claim that: (1) single spikes generated from CL cells related with the middle region of the object (Figs 5-1, 5-3 and 5-5) might not generate any activity in VPL, (2) the interleaved pattern ensures that VPL will start firing just when CL cells start to generate spike trains as a result of the fill-in process.

The CN processing discussed so far support the thesis that the nervous system is designed to detect changes in the environment, discard those signals that do not contribute with relevant information, and encode it in an optimal manner [5].

References:

- [1] J. Aguilar, C. Rivadulla, C. Soto, A. Canedo. New corticocuneate cellular mechanisms underlying the modulation of cutaneous ascending transmission in anesthetized cats. *J Neurophysiol.* 89 (2003) 3328-39.
- [2] J. Aguilar, C. Soto, C. Rivadulla, A. Canedo. The lemniscal-cuneate recurrent excitation is suppressed by strychnine and enhanced by GABA_A antagonists in the anaesthetized cat. *Eur J Neurosci* 16 (2002) 1697-1704.
- [3] V.E. Amassian, J.L. De Vito. La transmission dans le noyau de Burdach (nucleus cuneatus). Etude analytique par unités isolées d'un relais somatosensoriel primaire. *Coll Int Centre Nat Rech Sci* 67 (1957) 353-393.
- [4] P. Andersen, J.C. Eccles, R.F. Schmidt, T. Yokota. Depolarization of presynaptic fibres in the cuneate nucleus. *J Neurophysiol* 27 (1964) 92-106.
- [5] H.B. Barlow. The coding of sensory messages. In Thorpe WH and Zangwill OL (Eds.): Current problems in animal behaviour, (Cambridge University Press, 1961) 330-360.
- [6] K.J. Berkley, R.J. Budell, A. Blomqvist, M. Bull. Output systems of the dorsal column nuclei in the cat. *Brain Res Rev* 11 (1986) 199-225.
- [7] A. Canedo, J. Aguilar. Spatial and cortical influences exerted on cuneothalamic and thalamocortical neurons of the cat. *Eur J Neurosci* 12 (2000) 2515-2533.
- [8] W.W. Chambers, C.N. Liu. Cortico-spinal tract of the cat. An attempt to correlate the pattern of degeneration with deficits in reflex activity following neocortical lesions. *J. Comp Neurol* 108 (1957) 23-55.
- [9] S. Cheema, B.L. Whitsel, A. Rustioni. The corticocuneate pathway in the cat: relations among terminal distribution patterns, cytoarchitecture, and single neuron functional properties. *Somatosen Res* 1 (1983) 169-205.
- [10] R.W. Dykes, D.D. Rasmusson, D. Sretavan, N.B. Rehman. Submodality segregation and receptive-field sequences in cuneate, gracile, and external cuneate nuclei of the cat. *J Neurophysiol* 47 (1982) 389-416.
- [11] G. Gordon, W.A. Seed. An investigation of the nucleus gracilis of the cat by antidromic stimulation. *J Physiol (Lond)* 155 (1961) 589-601.
- [12] J.H. Lue, S.M. Luai, T.J. Wang, J.Y. Shieh, C.Y. Wen. Synaptic relationships between corticocuneate terminals and glycine-immunoreactive neurons in the rat cuneate nucleus. *Brain Res* 771 (1997) 167-171.
- [13] L. Martinez, J.A. Lamas, A. Canedo. Pyramidal tract and corticospinal neurons with branching axons to the dorsal column nuclei of the cat. *Neuroscience* 68 (1995) 195-206.
- [14] A. Rustioni, N.L. Hayes. Corticospinal tract collaterals to the dorsal column nuclei of cats. *Exp Brain Res* 43 (1981) 237-245.
- [15] E. Sánchez, S. Barro, J. Mariño, A. Canedo. A computational model of cuneothalamic projection neurons. *Network: Comput neural syst.* 14 (2003) 211-231.
- [16] E. Sánchez, S. Barro, A. Canedo. Edge Detection and Motion Discrimination in the Cuneate Nucleus. *Lect. Notes in Comp. Sci.* 2415 (2002) 198-203.
- [17] W.M. Usrey, J.B. Reppas, R.C. Reid. Paired-spike interactions and synaptic efficacy of retinal inputs to the thalamus. *Nature.* 395 (1998) 384-387.
- [18] F. Walberg. Corticofugal fibres to the nuclei of the dorsal columns. An experimental study in the cat. *Brain* 80 (1957) 273-287.