

Characterizing *in vitro* Hippocampal Ripples using Time-Frequency Analysis

J.A. Gillis^{1,2}; W.P. Luk^{1,4}; L. Zhang^{1,3}; F.K. Skinner^{1,2,3,5}

1. Toronto West. Res. Inst., UHN, Toronto, ON, Canada

2. Physiol., 3. Med. (Neurol.), 4. Inst. Med. Sci, 5. IBBME, Univ. Toronto, Toronto, ON, Canada

Abstract

We analyze *in vitro* hippocampal activity exhibiting drug induced ripple-like rhythms using time-frequency distributions. The different tradeoffs between time-frequency resolution, interference, and range are examined with reference to the suitability of each for use in analyzing ripple data. An unbiased approach is taken to characterizing ripple rhythms uniquely by determining the parameter choices in which its behaviour is most distinct from baseline activity. Mixture distribution analysis of the two components generated from correlating frequency behaviour allows a clear method for determining the distinctiveness of ripple-like rhythms given a particular parameter choice. These parameters thus serve to characterize hippocampal ripples.

Keywords: Hippocampus; Time-Frequency; Ripples; Mixture distributions; EEG

1. Introduction

Electroencephalogram (EEG) recordings of the hippocampus reveal the presence of specific frequency associations with distinct behavioural states. Sharp wave associated ripple-like activity occurring in the 150-200 Hz range is present during awake immobility, consummatory behaviour, and slow-wave sleep [2]. This frequency would be suitable for LTP induction, suggesting an important physiological role for ripple-like rhythms [1]. One question which arises from recordings of these EEG signals is whether the classes of rhythmic activity characterized by different mean frequencies share underlying similarities in their frequency behaviour outside their differences in mean: How are the mechanism and properties of ripples related to those of baseline data? One example of this might be a similarity in the distribution of activity in a given frequency range. If these similarities were present in a

continuum, they might suggest a single mechanism underlying otherwise different classes of activity; this is not true of mean frequency data for ripples [2].

Because the rhythms observed are biological, techniques to characterize frequency which assume stationarity, and specifically, a distribution of frequencies that does not change with time, are often inappropriate. Time-frequency analysis is one solution to the problem of stationarity, and it consists, essentially, of performing analysis in small enough signal segments that variation is negligible. To generate time-frequency plots, a common choice is to use Cohen's class of distributions, the most familiar of which are the spectrogram and the Wigner distribution.

Unfortunately, the Wigner distribution generates substantial interference patterns from the interaction of cross-terms, rendering it less suitable for multicomponent signals; this is a common problem. Outside of Cohen's class of distribution, the wavelet transform results in a time-frequency domain tiled so that the technique gives good frequency resolution at low frequencies, and good time resolution at high frequencies. Since signals which have high frequency might be thought to change frequency more rapidly with time, this is often an appropriate choice [5]. After comparing the suitability of these techniques, they are used to allow for a more complete characterization of ripple-like behaviour through an examination of its similarities with baseline activity. In this work we propose a simple method for characterizing ripple rhythms with respect to baseline rhythmic activity.

2. Method

Ripples were induced in an in vitro preparation of the hippocampi of mice by application of 1,3-dipropyl-8-cyclopentylxanthine. Eight CA1/3 field EEG recordings of two to three minutes, sampled at 20 kHz comprised the data set [4]. Time-

frequency plots of the data were generated using the spectrogram, the Wigner distribution, and the wavelet transform. The breakdown between time and frequency resolution was also altered in the plots generated, running from 0.01 seconds up to 4 seconds in time resolution by doubling for each power of 10 (e.g., 0.01, 0.02, 0.04, 0.08, 0.1, 0.2. . .). Of course, this also determines the correspondingly better frequency resolutions. Having generated the frequency distributions for each unit time at a given resolution, they were compared for similarities in their frequency distributions. Simply taking the correlation between the different frequency distributions obtained at each time serves to do this, for the most part. In order to reflect trends of different magnitude, a chirp was added to the signal, resulting in a default high correlation unless there were different trends. In this way, purely noisy parts of the signal correlated well with one another, as we would like (indicating similar behaviour). This done, each time-frequency representation, and over varying frequency ranges, the frequency distributions were correlated. This was performed for each plot generated, and also for subdivisions in frequency (e.g., 1-30 Hz, 150-200 Hz, etc). Histograms were then constructed from the correlation values obtained. Clearly, the usefulness of any characterization is context dependent, and our analysis of ripple rhythms is explicitly dependent on its context - the baseline rhythmic activity. Because the signal exhibits two clear modes of activity (ripple, baseline), the histograms will be a mixture distribution of roughly two normal components, reflecting times where frequency distributions are the same (ripple to ripple, baseline to baseline giving similarly high correlations, but the former is also vanishingly infrequent), or different (ripple to baseline giving a relatively low correlation). The EM algorithm was used to characterize the underlying components giving rise to the distribution, under the assumption that there were two primarily contributing normal

distributions ([3], [6]). The statistical significance of the separation was determined from the pseudo-Chi square values and was taken to indicate the ability of a given set of parameters to characterize ripples (Figure 1). That is, parameters have been chosen which well characterize ripples when their frequency distribution is distinct. The analysis was also performed on artificial rhythmic data. Errors were inspected both by examining log-likelihood slopes, and by using a range of starting parameters and observing resultant fits.

3. Results

Ripples exhibit highly distinctive frequency distributions, and are normally defined by those characteristics in the high frequency domain (Figure 2). Time-Frequency analysis did not show any stationary characteristics preceding ripples, and non-stationarity in ripples was similar across all frequency ranges, eliminating the particular advantage of the wavelet transform. The baseline rhythmic activity varied substantially in frequency and regularity even within individual datasets, but approximated a 3 Hz rhythm. While the baseline rhythm varied substantially, there was no detectable relationship indicating the onset of a ripple. Ripples did exhibit a characteristic frequency distribution in the low frequency range, even after normalization within that range. This could be seen in correlating the low-frequency signal contribution, and observing the clear presence of a mixture distribution in the resultant histograms, reflecting the substantial difference between ripples and baseline in this frequency range. Regardless of the specific frequency range, duration, or times chosen, the correlation of frequencies exhibited qualitatively similar features with a tartan like appearance. This is also reflective of the lack of stationarity associated with ripples at all frequency ranges.

The frequency distributions gave the greatest separation into two distinct components for time resolution of 0.2 second, in the range of the approximate duration of ripples (Figure 2). In general two component mixture distribution fits were statistically significant ($p < 0.01$), with variation in the degree to which they were significant being the important point in determining defining frequencies. There were three clusters of frequency ranges that most distinctly separated ripples from baseline activity, 1-40 Hz, 120-170 Hz, and 150-220 Hz, with only that latter corresponding to the rough range most often used to define ripple behaviour. Each of the eight datasets had similar qualitative features with respect to the characterization of ripples by frequency. In each, there were three peaks in ripple separability corresponding to the given frequency ranges, although the relative contribution of each varied from dataset to dataset. Using both of the two latter overlapping regions generated less distinct distributions, suggesting a qualitative change in behaviour during the overlap. Ripples (to ripple correlations) constitute a small enough fraction of the higher-valued distribution so as to be relatively insignificant, even if their frequency distributions do not constitute precisely the same distribution as the same-same values generated by baseline activity. Variation in frequency range chosen did not alter the identification of ripples as determined from the separation of the mixture distribution of correlations. The same analysis performed on artificial spiking did not yield the distinctive mixture distribution of correlation values, since the spikes were not distinctly different in the population of frequency characteristics they displayed.

Overlap between the two distributions depended upon the frequency range and resolutions used, but even with high overlap there was excellent statistical separation and the existence of two classes of activity was very clear. Varying starting values rarely altered the fits obtained for the mixture distribution. Attempting three

component fits was rarely successful. In general, the weighting of a third component was very low, or the mean was very close to a pre-existing distribution, most often the cross-term distribution. Pieces of the signal contributing to intermediate points in the correlation histogram sometimes exhibited ripple-like characteristics in the low frequency band (e.g., $r > 0.95$) but no ripple-like character in the high frequency band, and otherwise did not change. That is, while most activity categorized as ripple-like in one parameterization remained ripple-like in another, to the extent there was migration, it was one way: more activity sharing distinctly ripple-like character in lower frequencies.

4. Conclusions and Discussion

Ripples are defined largely by their high-frequency behaviour, to the point where other contributing frequencies are filtered out in analysis. We have attempted to determine a way to best characterize ripples with fewer *a priori* assumptions as to their behaviour. So, while the identification of ripples is trivial, the determination of which parameters *best* characterizes them is not. In an *in vitro* hippocampal preparation known to exhibit ripples and a baseline rhythm, we assumed that the distribution of frequencies would exist in broadly two classes, and checked this assumption. Many frequency distributions were generated over time and with different parameter choices, and tested for their statistical separation into two classes, under the reasoning that the parameter which best separated the data into two classes would also be those that best characterize ripples with respect to the baseline rhythm. The primary intent is to develop a single metric for reasonably characterizing ripples with a minimum of bias - distinctiveness from baseline. Of course, one possibility would be to determine the contribution of different measured factors (e.g., factor

analysis, component analysis) to our measure, but first it is necessary to have some characterization or output of interest.

Unsurprisingly, we found that the data separated well for temporal resolutions in the range of the duration of ripples. More interestingly, the frequency ranges determined included not only the roughly tradition range (150-220 Hz), but also a slightly lower range (120-170 Hz), and a much lower one (1-40 Hz), surprising for the duration of activity. This last frequency range may represent ongoing activity (baseline rhythms) altered due to the occurrence of the ripples. It is not simply due to change in total power since we are comparing frequency distributions. Further, the low frequency characterization being merely a side-effect of ripples is argued against by the occurrence of an intermediate population which demonstrates the low frequency aspect even where rippling is not present.


Acknowledgements

This work was supported by the Natural Sciences and Engineering Research Council of Canada.


References

- [1] G. Buzsaki, Hippocampal sharp waves: Their origin and significance, *Brain Research* (1986), 398:242–252.
- [2] J. Csicsvari, H. Hirase, A. Czurko, A. Mamiya, and G. Buzsaki, Oscillatory coupling of hippocampal pyramidal cells and interneurons in the behaving rat, *Journal of Neuroscience* (1999), 19:274-287.


- [3] A.P. Dempster, N.M. Laird, and D.B. Rubin, Maximum likelihood from incomplete data via the em-algorithm plus "discussions on the paper", Journal of the Royal Statistical Society (1977), 39(B):1-38.
- [4] W.P. Luk , T. Wong, C.P. Wu, F.K. Skinner, and L. Zhang, An in-vitro model of mouse hippocampal sharp waves and ripples, unpublished manuscript, 2004.
- [5] A. Teolis, Computational Signal Processing with Wavelets (Birkhauser, Boston, 1998).
- [6] J.H. Wolfe, Pattern Clustering by multivariate mixture analysis, Multivariate Behavioural Research (1970) 5:329-350.



1.



2.



3.

Figure 1. A flow chart of the methodology is shown; A: The raw signal; B: Time-frequency representations of the signal, and in different ranges; C: Correlation between each time's frequency activity; D: Histograms for each set of correlations; E: Measuring separability in those histograms.

Figure 2. A sample of the data is shown at three scales to observe different characteristics; A: One entire recording; B: The baseline rhythm and ripple; C: The ripple in close-up.

Figure 3. Two histograms constructed from correlating frequency distributions as described; A: A fairly good separation around 140-190 Hz; B: A poor separation around 40-90 Hz.