

Sharpening the spatial tuning curves of place cells in a feed-forward network: the possible role of theta oscillation-based dynamics in the hippocampus

Máté Lengyel^{a,b}, Péter Érdi^{c,a}

- a, Department of Biophysics, KFKI Research Institute of Particle and Nuclear Physics, Hungarian Academy of Sciences, 29-33 Konkoly-Thege M. út, Budapest H-1121, Hungary
- b, Department of Physiology and Neurobiology, Faculty of Sciences, Eötvös Loránd University, 1/C Pázmány Péter sétány, Budapest H-1117, Hungary
- c, Center for Complex Systems Studies, Kalamazoo College, 1200 Academy Street, Kalamazoo, MI 49006-3295

Activity of the principal cells of the hippocampus during exploratory behavior is known to encode for the spatial location of the rat both by their firing rates^{1,2} and by the precise timing of their discharges relative to the ongoing theta rhythm field potential-oscillation, provided by the phase precession effect^{3,4}. Such 'place cells' with doubly (rate and phase) coded firing patterns were found in all three subregions of the hippocampus: in the dentate gyrus (DG), in CA3 and in CA1^{4,5}. These subregions possess different anatomical layouts, and are mainly connected by excitatory feed-forward connections in the DG→CA3→CA1 loop⁶. Moreover, the cortical input of the hippocampus is only weakly correlated with the location of the animal⁷, and place cell activity was shown to persist in downstream areas (CA3 and CA1) even in the absence of upstream place specific activity (DG⁸ and CA3⁹, respectively). Thus, place cells in each hippocampal area appear to bear significant autonomy in generating spatially coded firing and to only a smaller extent rely on topographical input or on a specific arrangement of intra-regional connections. Although a huge body of experimental and model works has accumulated in this field over the past years^{10,11}, no theory has yet been able to explain this autonomous doubly coded place specific activity throughout different areas of the hippocampal loop.

In this work, a simple feed-forward network was simulated, where the layers of the network corresponded to different hippocampal subfields, and were connected with realistic convergence and divergence ratios. This network architecture was shown to have two distinct modes of operation. In the 'classical mode', only the rate code of place cell firing could be considered: postsynaptic neurons simply summed the activity of their presynaptic partners, and thus place-specific activity in each layer depended on correctly formed spatial tuning curves in its afferent layer, with tuning curves in downstream areas being usually more blurred than in upstream areas due to the convergence of feed-forward connections. In 'theta mode', two consequences of the hippocampal theta oscillation were

taken into account. First, firing probability of interneurons was modulated by the phase of theta oscillation^{4,12}, and thus synaptic transmission between layers could only be effective in a short time window within each theta cycle. Ensemble activity of phase precessing place cells in this time window was shown to be strongly correlated with the velocity of the animal. Second, somatic membrane potential of place cells oscillated in the subthreshold regime with theta frequency¹³, while the denritic membrane sustained intrinsic membrane potential oscillations, frequency modulated in the theta range by the level of depolarization¹⁴. Thus, as shown in our previous work¹⁵, neurons of each layer were capable to generate place specific firing patterns, simultaneously rate and phase coded, by relying primarily on the velocity-correlatedness and only weakly on the location-correlatedness of the activity in their afferent areas. This predicts that hippocampal subregions show considerable autonomy in maintaining properly tuned firing rate curves in their place cells, and perform effective sharpening of these tuning curves compared to the classical operational mode.

A comparison of our results to the most widely accepted, attractor neural network (ANN) description of hippocampal place cells^{10,16,17} is given. In those models, the sharpening of tuning curves is accomplished by intranetwork dynamics based on dense and topographical lateral connections between place cells. Hence they may be directly applicable to CA3 only where recurrent collaterals can be found⁶, yet the functionality of these collaterals under physiological conditions is not clear¹⁸. In our model recurrent collaterals are not necessary for sharp tuning curves but may play other roles in the dynamics of the network. We also find that, as opposed to our model, simultaneous reproduction of the rate and phase code of place cells can be problematical in ANNs. Further quantitative predictions of the two models are also compared and matched to experimental data in detail. Finally, our proposed 'one architecture–two dynamical modes' scenario is also discussed from an evolutionary perspective.

References

1. O'Keefe J and Dostrovsky J. The hippocampus as a spatial map. Preliminary evidence from unit activity in the freely moving rat. *Brain Res* 34:171–175, 1971.
2. Skaggs WE, McNaughton BL, Gothard KM, and Markus EJ. An information-theoretic approach to deciphering the hippocampal code. In *Advances in Neural Information Processing Systems* (Hanson SJ, Cowan JD, and Giles CL, eds), vol. 5, pp. 1030–1037, Morgan Kaufmann: San Mateo, CA, 1993.
3. O'Keefe J and Recce ML. Phase relationship between hippocampal place units and the EEG theta rhythm. *Hippocampus* 3:317–330, 1993.

4. Skaggs WE, McNaughton BL, Wilson MA, and Barnes CA. Theta phase precession in hippocampal neuronal populations and the compression of temporal sequences. *Hippocampus* 6:149–172, 1996.
5. Barnes CA, McNaughton BL, Mizumori SJY, Leonard BW, and Lin LH. Comparison of spatial and temporal characteristics of neuronal activity in sequential stages of hippocampal processing. *Prog Brain Res* 83:287–300, 1990.
6. Lopes da Silva FH, Witter MP, Boeijinga PH, and Lohman AH. Anatomic organization and physiology of the limbic cortex. *Physiol Rev* 70:453–511, 1990.
7. Quirk GJ, Muller RU, Kubie JL, and Ranck, Jr JB. The positional firing properties of medial entorhinal neurons: description and comparison with hippocampal place cells. *J Neurosci* 12:1945–1963, 1992.
8. McNaughton BL, Barnes CA, Meltzer J, and Sutherland RJ. Hippocampal granule cells are necessary for normal spatial learning but not for spatially-selective pyramidal cell discharge. *Exp Brain Res* 76:485–496, 1989.
9. Mizumori SJY, McNaughton BL, Barnes CA, and Fox KB. Preserved spatial coding in hippocampal CA1 pyramidal cells during reversible suppression of CA3c output: evidence for pattern completion in hippocampus. *J Neurosci* 9:3915–3928, 1989.
10. Redish AD. *Beyond the Cognitive Map: From Place Cells to Episodic Memory*, A Bradford Book. The MIT Press: Cambridge, MA, 1999.
11. Best PJ, White AM, and Minai A. Spatial information processing in the brain: the activity of hippocampal place cells. *Annu Rev Neurosci* 24:459–86, 2001.
12. Csicsvári J, Hirase H, Czurkó A, Mamiya A, and Buzsáki G. Oscillatory coupling of hippocampal pyramidal cells and interneurons in the behaving rat. *J Neurosci* 19:274–287, 1999.
13. Nuñez A, Garcia-Austt E, and Buno, Jr W. Intracellular theta-rhythm generation in identified hippocampal pyramids. *Brain Res* 416:289–300, 1987.
14. Kamondi A, Acsády L, Wang XJ, and Buzsáki Gy. Theta oscillation in somata and dendrites of hippocampal pyramidal cells in vivo: activity-dependent phase-precession of action potentials. *Hippocampus* 8:244–261, 1998.
15. Lengyel M, Szatmáry Z, and Érdi P. A detuned oscillator model of place unit phase precession in the rat hippocampus. *Soc Neurosci Abst* 25:1387, 1999.
16. Tsodyks MV, Skaggs WE, Sejnowski TJ, and McNaughton BL. Population dynamics and theta phase precession of hippocampal place cell firing: a spiking neuron model. *Hippocampus* 6:271–280, 1996.
17. Tsodyks M. Attractor neural network models of spatial maps in the hippocampus. *Hippocampus* 9:481–489, 1999.
18. Christian EP and Dudek FE. Characteristics of local excitatory circuits studied with glutamate microapplication in the CA3 area of rat hippocampal slices. *J Neuophysiol* 59:90–109, 1988.