

1 Linking microbial diversity to macrobiota diversity in eelgrass 2 meadows

3 Deirdre Loughnan^{1,2}, Laura Parfrey², Margot Hessing-Lewis¹, Mary O'Connor²

4 October 31, 2025

5 ¹ Hakai Institute

6 ² Zoology, Faculty of Science, University of British Columbia.

7
8 Corresponding Author: Deirdre Loughnan deirdre.loughnan@hakai.org
9

10 Introduction

11 • Climate change is altering the structure of ecological communities worldwide—in most systems
12 we see a decline in diversity, but effects are not uniform—limited understanding of underlying
13 mechanisms and key environmental variables that contribute to community resilience versus those
14 that are reshaping communities

- 15 1. To date considerable efforts to understand how diversity is changing globally
16 – But assessments of diversity often focus on specific guilds or taxonomic groups
17 – Rarely do they span organismal scales, such as between micro and macro scales, even
18 within a single system
19 – Increasingly clear that interactions between microbiota and macrobiota play an impor-
20 tant role in shaping ecological communities and should be incorporated into assessments
21 of biodiversity
- 22 2. Considerable research has now demonstrated the myriad of ways that microbial diversity
23 shapes ecological communities and ecosystem services
24 – shape the environment—nutrient cycling—metabolizing organic carbon and nitrogen
25 – link organisms across trophic levels
26 – species interactions and symbiosis (McRoy and Goering 1974; Kirchman et al 1984—
27 Sanders-Smith2020)
28 – microbial diversity is product of: host sp identity, space, and local envirt
29 – but still limited understanding of the complexity of interactions and the environmental
30 factors that shape them
- 31 3. Eelgrass communities are an excellent system for furthering our understanding of micro-
32 macro relationships across scales (split into 2 paragraphs)
33 (a) General relationships between eelgrass and diversity:
34 – Farjalla et al 2012: community assembly differs across scales—envirt filters and
35 interactions shape animals but microbes are stochastic

- But in marine ecosystems hosts more prone to horizontal transmission—Turon et al 2018; Russel 2019—Sanders-Smith2020; global scale find that microbial community mirror that of environment (Fahimipour2017—Sanders-Smith2020); but at finer scales see selection by host through exudates
- Seagrass release methanol and cellulose—food for microbes—and phenols which deters colonization:
- Crump et al 2018—Sanders-Smith2020: microbes consume toxic byproducts—methanol and ethanol—and produce enzymes that limit epiphyte overgrowth
- Previous studies found evidence that in seagrass animal-microbial systems covary Yeager et al 2019, Bengtsson et al. 2017
- (b) Summary of seagrass work around bc/Calvert
 - Several studies have studied eelgrass community diversity in meadows in coastal BC: meadows that are farther apart are more dissimilar—driven by environment—temp and DO explained variation more than habitat traits (LAI)—but taxa specific (Stark2020)
 - Sanders-Smith2020: differences in microbial community structure among regions but not water column; differences in diversity associated with new (lowest richness higher dominance) and old leaves
 - Segovia2020: 2 years of data: lower alpha diversity on seagrass than water, most abundant microeuk = diatoms, copepods, gregarines, and rotifers; Bray-curtis and Jaccard both showed more similar communities on seagrass than other biofilms and seawater—found spatial structuring in core taxa and a strong effect of temperature
 - Meadows can differ in their diversity spatially, but less is known how they change over time—further our understanding of how these processes covary and what potential drivers may shape community responses is unclear
- 4. Here we explore how changes in microbial diversity effect macrobial communities in both time and space—exploring the potential for microbes to mediate environmental conditions and underlie diversity changes across a community.
 - two four-year time-series of microbial diversity based on DNA sequences for the prokaryote and microeukaryote communities swabbed from eelgrass and paired with diversity of epifauna communities present on the eelgrass.
 - Use a joint Bayesian model to explore how diversity in micro and macro communities change over time and whether species interactions are correlated with environment cues, such as temperature or differences in habitat structure.
 - In addition to estimating overall trends in diversity in eelgrass communities, we also tested for trends across sites.

Materials and Methods

1. General intro to study area

- Both microbial and macro data collection was conducted around Calvert Island in BC
- Hakai Lúxvbálís conservancy: low human impacted *Zostera marina* meadows
- Meadows = monoculture of *Zostera marina* with macroalgae and microalgal
- 5 regions each with 2 sites, except choked which had 3 sites (Choked, GPS: 51.66825, -128.11855 and Pruth, GPS: 51.395, -128.069), Goose (GPS: 51.92652, -128.45317), McMullin (GPS: 52.06178, -128.41239) and Triquet (GPS: 51.80869, -128.24746) islands (Fig. 1)

2. Epifauna surveys:

- Epifauna surveys and collection of seagrass habitat traits were collected from 2014 to 2017
- Using SCUBA sampled six 0.25 m by 0.25 m quadrats established in a 15m by 30m array placed in the middle of the meadow
- Collected eelgrass, detritus, macroalgae and epifaunal invertebrates
- Prior to collecting eelgrass traits, rinsed blades to remove epifaunal invertebrates and passed through 500 μm sieve.
- Invertebrates preserved 95% EtOH and identified upon returning from the field using light microscopy (10x magnification)
- Species were identified to the lowest taxonomic level possible using manuals by Light and Smith manual (Carlton 2007) and Kozloff (1996)
- Additional domain expertise of the natural history of our surveys species was used to remove species known to predominantly reside in eelgrass sediment ($n = 17$ species)
- For additional detail of the applied methods, see....

3. Microbial surveys

- Collection and sequencing of microbial communities was conducted from 2015 to 2018
- Collected one shoot just outside each quadrat sealed underwater in a sterile plastic bag at the location of collection.
- Swabs were taken from the second oldest blade, from an area with minimal to no epibiota and 10 cm in length, rinsing the area for 10 seconds with filtered seawater (0.22 μm) to remove attached organisms, and swabbed using Puritan sterile swabs for 10 seconds.
- Swabs were placed in cryotube and placed in a -80°C freezer within 6 hours of sampling and until DNA extraction
- Extracted DNA from swabs using Qiagen Powersoil—htp 96 well DNA extraction Kit (Carlsbad, CA): V4 region of 18S rRNA gene was targeted for amplification using primers E572F: 5'-AYGGTATCTRATCRTCCTTYG-3' and PCR, library construction, and Illumina MiSeq amplicon sequencing were carried out at the Integrated Microbiome Resource Facility, Centre for Comparative Genomics and Evolutionary Bioinformatics at Dalhousie University Halifax, Canada using standard protocols (Comeau et al. 2011)
- 16S DNA extraction: MoBio PowerSoilVR -htp 96 well DNA extraction kit (Carlsbad, CA), V4 region of the 16S rRNA gene in Bacteria and Archaea was targeted for amplification using redesigned versions of the primers 515f/806r as described by Lemay et al. 2018a.
- Removed OTUs with fewer than 2 reads for that sample to minimize the impact of barcode switching, and OTUs with fewer than 250 total reads and samples with low read counts (<1,000 reads/sample)
- For additional detail, see Sanders-Smith et al. 2020 and Segovia et al. 2020 for methods for sequence data analysis.

4. Climate and habitat traits

- At time of epifauna sampling collected data on bed area, meadow depth, eelgrass leaf area index, biomass of eelgrass and epiphytes'
- Measured shoot leaf area and biomass: measured leaf length, width and blade number of five haphazardly selected representative shoots from each quadrat, multiplying the average blade area (m^2) per shoot by the shoot density (shoots per m^2)
- LAI: structural complexity of bed and amount of habitat available for colonization by epiphytes and invertebrates (Enriquez et al. 2019, Green and Short 2003): $\text{LAI} = \text{mean blade length} \times \text{width} \times \text{number of blades per shoot}$; multiplying number of shoots in quadrat

- 126 • Additional shoot selected for microepiphyte biomass, scraping biofilm with glass slide filter-
127 ing onto GF/C filters, dried and weighed
- 128 • Eelgrass and macroalgae biomass: biomass was separated and dried in a desiccator oven at
129 60°C for 48 hours.
- 130 • Our was to use temperature data recorded as close to our sites and at as temporally fine
131 scale as possible
- 132 • Combined available historic climate data for each site, pooling across the Hakai CTD and
133 sensor network data (CITE website)
- 134 • For a given site and year, the frequency of temperature measurements varied considerably,
135 as such we averaged the temperature observations across the summer months for each site
136 (with summer defined as May to August in a given year).

Statistical Analysis

Microbe sub-model

The sub-model describes the processes that shape the changes in microbial community diversity over time for 1 to n sites in our dataset. We use hierarchical modeling to partition sources of variation including measurement error, and site-level differences. In particular, we assume that a observed change in diversity for site i , Y_{micro_i} follows a normal distribution.

$$y_{\text{micro}} = \alpha_{\text{grandMicro}} + \alpha_{\text{site}_i} + \beta_{\text{year}_i} \times \text{year} + \sigma^2 \quad (1)$$

$$\alpha_{\text{microSite}} = \alpha_{\text{grandMicro}} + \alpha_{\text{site}_i} \quad (2)$$

$$(3)$$

The latent parameter $\alpha_{\text{grandMicro}}$ represents a value that is independent of site, α_{site_j} and site-level offsets from that diversity value, σ^2 is measurement error. Of these parameters, $\alpha_{\text{microSite}}$ is shared by the macro community sub-model.

Macro sub-model

The macro-community sub-model describes the processes that determine the community diversity for 1 to n sites in relation to the microbial diversity. We predict that macro community diversity will be shaped by habitat structure, including the biomass of macroalgae, bed area, depth, and the distance from the bed's edge. The diversity of the microbial community, however, may further mitigate the effects of temperature, salinity, and biomass, such that such that an observation of micro diversity at a given site i under set g of temperatures, salinity, and seagrass biomass ($\text{temp}_j, \text{salinity}_j, \text{biomass}_j$) contributes to observations of macro diversity, $Y_{\text{macro}_{i,j}}$ (which we z -scored to allow direct comparison of cues).

$$\beta_{\text{temp}} = \alpha_{\text{temp},j} + \beta_{\text{micro.temp}} \cdot \alpha_{\text{microSite},j} \quad (4)$$

$$\beta_{\text{LAI}} = \alpha_{\text{LAI},j} + \beta_{\text{micro.LAI}} \cdot \alpha_{\text{microSite},j} \quad (5)$$

$$y_{\text{macro}} = \alpha_{\text{grandMacro}} + \alpha_{\text{site}_j} + \beta_{\text{year}_j} \cdot \text{year} + \beta_{\text{temp}_j} \cdot \text{temp} + \quad (6)$$

$$\beta_{\text{LAI}_j} \cdot \text{LAI} + \beta_{\text{depth}_j} \cdot \text{depth} \quad (7)$$

Beta diversity

$$y_{\text{beta}} \sim \text{Normal}(\mu, \sigma_e^2) y_{\text{micro}} = \alpha_{\text{grandMicro}} + \alpha_{\text{site}_i} + \beta_{\text{year}_i} \times \text{year} + \sigma^2 \quad (8)$$

$$\alpha_{\text{microSite}} = \alpha_{\text{grandMicro}} + \alpha_{\text{site}_i} \quad (9)$$

$$(10)$$

Results

- Summary of community composition and characteristics:
 - Identified 3933 unique taxa of prokaryote and 1143 microeukaryote species.
 - Across all sites, sampled 58373 invertebrates from 107 taxa.

- 160 – Across the four years of sampling, the number of individual epifauna sampled per 0.25 m^2
161 of seagrass meadow varied considerably, from 122 individuals sampled in 2015 in Goose SE
162 to 9393 individuals sampled in 2014 at our Triquet S site.
- 163 – Across all sites, the five most abundant taxa (*Caprella.californica*, *Caprella.laeviscula*,
164 *Lacuna.vincta*", *Porcellidium*, *Gammaridea*) represented 60% of invertebrates collected,
165 although abundance was not consistent across all sites or years.
- 166 – Many species were not present at each site: only 11 taxa were present at each site, but 43%
167 of species were present at five or more sites.
- 168 • Changes in alpha diversity over time:
 - 169 – In modelling the relationship between prokaryote diversity and epifauna: we found a slight
170 decline in prokaryote diversity over time (-0.17, 90% UI: -0.29, -0.02), but an increase in
171 epifauna diversity over time (0.31, 90% UI: 0.06, 0.55). These estimates remained stable
172 when the model was run with the last three years of data (results not shown).
 - 173 – In trying to better understand the pathways through which microbial diversity may influence
174 epifauna diversity, we found no evidence that the observed changes in prokaryote diversity
175 interacting with changes in local temperature ($\beta_{micro.temp}$) or differences in eelgrass leaf
176 area ($\beta_{micro.LAI}$) to drive the observed increase in epifauna diversity.
 - 177 – In modelling of the relationship between microeukaryote diversity and epifauna diversity, we
178 observed no change in microeukaryote diversity over time and similarly that temperature,
179 eelgrass leaf area had no influence on how species are interacting across these organismal
180 scales. and depth are not driving the overall increase in epifauna diversity we observe.
- 181 • In addition to finding these overall trends in diversity, we found some variation in the magnitude
182 and direction in which diversity is changing at our various sites
 - 183 – The negative relationship estimated for prokaryote diversity across sites was driven by the
184 declines observed at four of our study sites, while the remaining five communities showed
185 no change
 - 186 – While we observed no overall change in microeukaryote diversity, our model does estimate
187 some changes at the site-level, with two of our most southern sites (Pruth pocket and
188 Choked inner) increasing in diversity, but the nearby Choked sandspit experienced a decline
189 in diversity.
 - 190 – Finally, we observed increases in epifauna diversity at four of our sites, while the remaining
191 five sites showed no change.
 - 192 – Contrary to our expectations, none of these shifts appear to be related to differences in
193 site-level temperature or eelgrass LAI ??

194 Disucssion

- 195 1. Summary of primary findings: some changes in diversity, changes in site level diversity—other
196 alternate drivers not currently identified
- 197 2. Compare the trends in eelgrass diversity we observed to meadows globally
- 198 3. Our dataset relative to other existing diversity datasets
- 199 4. Site-level differences and how contrast with stable temporal trends
- 200 5. The advantages of our method and the broad application to linking biodiversity across scales

Figures




Figure 1: Site-level changes in Shannon diversity over time for the a) prokaryote communities, b) microeukaryote communities and c) epifauna communities. In general, our model estimated a decline in the prokaryote diversity over time, while epifauna diversity increased, but the microeukaryote community remained stable.




Figure 2: We found no relationship between prokaryote diversity and epifauna responses to a) temperature and b) leaf area index, or between microeukaryote diversity and c) temperature and d) leaf area index, with the exception of McMullins North, for which diversity declines as LAI increased in both models.

Supporting Information

Table 1: Summary output from a joint Bayesian model linking changes in Shannon diversity of prokaryote (16S) and macroepifauna diversity in relations to environmental conditions and habitat structure, with partial pooling for site. The model includes standardized environmental cues that are z -scored to allow comparisons to be made across parameters. In addition to the mean model estimates, both the 50%, and 90% uncertainty intervals are also reported.

	mean	5%	25%	75%	95%	n_eff	Rhat
mu_grand	4.26	3.98	4.15	4.37	4.53	480.52	1.01
sigmaSt	0.26	0.01	0.08	0.39	0.65	147.52	1.01
muyear	-0.17	-0.29	-0.23	-0.12	-0.02	521.55	1.00
sigmayear	0.13	0.02	0.06	0.16	0.30	437.79	1.01
sigma_microy	0.66	0.59	0.63	0.68	0.73	829.62	1.00
mu_grandM	1.11	0.51	0.91	1.31	1.70	99.17	1.03
muyear2	0.31	0.06	0.23	0.40	0.55	375.42	1.01
sigmayear2	0.32	0.14	0.21	0.39	0.63	522.80	1.00
mutempSt	-1.76	-18.78	-4.77	2.12	12.50	317.34	1.01
sigmatempSt	0.22	0.02	0.08	0.29	0.62	895.19	1.00
muLAIS	1.01	-16.22	-1.86	4.08	16.57	223.97	1.01
sigmaLAIS	0.21	0.02	0.09	0.29	0.56	799.72	1.00
muDepth	-0.00	-0.54	-0.19	0.19	0.50	233.80	1.02
sigmaMacroSt	0.78	0.40	0.55	0.93	1.47	210.95	1.01
sigmaMacro_y	0.26	0.23	0.25	0.27	0.30	1522.14	1.00
betaMicroxtemp	0.41	-2.98	-0.50	1.13	4.39	318.45	1.01
betaMicroxLAI	-0.21	-3.87	-0.93	0.45	3.81	228.06	1.01

Table 2: Summary output from a joint Bayesian model linking changes in Shannon diversity of microeukaryotes (18S) and macroepifauna diversity in relations to environmental conditions and habitat structure, with partial pooling for site. The model includes standardized environmental cues that are z -scored to allow comparisons to be made across parameters. In addition to the mean model estimates, both the 50%, and 90% uncertainty intervals are also reported.

	mean	5%	25%	75%	95%	n_eff	Rhat
mu_grand	1.29	1.02	1.18	1.40	1.56	1501.40	1.01
sigmaSt	0.23	0.01	0.03	0.38	0.63	222.83	1.01
muyear	0.00	-0.18	-0.06	0.07	0.18	1261.63	1.00
sigmayear	0.21	0.04	0.12	0.27	0.46	399.19	1.00
sigma_microy	0.55	0.50	0.53	0.58	0.62	3821.57	1.00
mu_grandM	1.13	0.65	0.96	1.30	1.60	1229.93	1.00
muyear2	0.31	0.07	0.23	0.41	0.54	1321.86	1.00
sigmayear2	0.30	0.13	0.20	0.36	0.58	2231.48	1.00
mutempSt	-0.40	-12.30	-1.93	0.47	13.00	718.51	1.00
sigmatempSt	0.21	0.01	0.07	0.28	0.57	3029.63	1.00
muLAIST	-1.27	-15.19	-2.14	-0.13	11.49	617.68	1.01
sigmaLAIST	0.18	0.01	0.07	0.25	0.48	3237.04	1.00
muDepth	-0.03	-0.46	-0.20	0.13	0.39	1562.86	1.00
sigmaMacroSt	0.72	0.39	0.52	0.84	1.25	2240.52	1.00
sigmaMacro_y	0.26	0.23	0.25	0.27	0.30	4348.93	1.00
betaMicroxtemp	0.31	-10.17	-0.34	1.50	9.73	721.66	1.00
betaMicroxLAI	1.11	-8.76	0.20	1.81	12.01	630.40	1.01

Table 3: Summary output from a joint Bayesian model linking changes in Shannon diversity of prokaryote (16S) and macroepifauna diversity in relations to environmental conditions and habitat structure, with partial pooling for site, but with the first year of epifauna sampling removed. The model includes standardized environmental cues that are z -scored to allow comparisons to be made across parameters. In addition to the mean model estimates, both the 50%, and 90% uncertainty intervals are also reported.

	mean	5%	25%	75%	95%	n_eff	Rhat
mu_grand	4.26	3.98	4.15	4.37	4.53	345.39	1.01
sigmaSt	0.15	0.01	0.01	0.24	0.65	66.29	1.06
muyear	-0.17	-0.29	-0.24	-0.12	-0.02	284.53	1.01
sigmayear	0.14	0.02	0.07	0.18	0.30	319.64	1.00
sigma_microy	0.67	0.59	0.63	0.69	0.73	484.31	1.02
mu_grandM	1.10	0.51	0.90	1.30	1.70	195.85	1.01
muyear2	0.27	0.06	0.19	0.37	0.55	290.07	1.01
sigmayear2	0.32	0.14	0.22	0.40	0.63	446.85	1.01
mutempSt	-1.83	-18.78	-7.75	3.91	12.50	129.28	1.03
sigmatempSt	0.24	0.02	0.10	0.32	0.62	672.56	1.01
muLAIST	-0.54	-16.22	-2.11	1.56	16.57	110.81	1.03
sigmaLAIST	0.05	0.02	0.02	0.06	0.56	929.16	1.00
muDepth	-0.04	-0.54	-0.21	0.14	0.50	301.77	1.01
sigmaMacroSt	0.74	0.40	0.54	0.87	1.47	473.02	1.00
sigmaMacro_y	0.27	0.23	0.25	0.28	0.30	813.06	1.00
betaMicroxtemp	0.43	-2.98	-0.93	1.81	4.39	130.36	1.03
betaMicroxLAI	0.14	-3.87	-0.35	0.51	3.81	110.69	1.03

Table 4: Summary output from a joint Bayesian model linking changes in Shannon diversity of microeukaryotes (18S) and macroepifauna diversity in relations to environmental conditions and habitat structure, with partial pooling for site, but with the first year of epifauna sampling removed. The model includes standardized environmental cues that are z -scored to allow comparisons to be made across parameters. In addition to the mean model estimates, both the 50%, and 90% uncertainty intervals are also reported.

	mean	5%	25%	75%	95%	n_eff	Rhat
mu_grand	1.29	1.02	1.20	1.39	1.56	1346.62	1.00
sigmaSt	0.06	0.01	0.00	0.04	0.63	80.01	1.05
muyear	-0.02	-0.18	-0.09	0.06	0.18	533.42	1.01
sigmayear	0.28	0.04	0.19	0.34	0.46	556.43	1.00
sigma_microy	0.56	0.50	0.53	0.58	0.62	1798.76	1.00
mu_grandM	1.10	0.65	0.93	1.28	1.60	263.80	1.01
muyear2	0.29	0.07	0.20	0.37	0.54	536.10	1.01
sigmayear2	0.32	0.13	0.21	0.38	0.58	786.20	1.01
mutempSt	-1.73	-12.30	-8.38	4.35	13.00	502.29	1.01
sigmatempSt	0.23	0.01	0.09	0.31	0.57	1061.62	1.00
muLAIS	-0.11	-15.19	-2.70	2.39	11.49	525.11	1.01
sigmaLAIS	0.04	0.01	0.02	0.06	0.48	1884.35	1.00
muDepth	-0.08	-0.46	-0.23	0.10	0.39	326.30	1.01
sigmaMacroSt	0.73	0.39	0.52	0.85	1.25	486.49	1.01
sigmaMacro_y	0.27	0.23	0.25	0.28	0.30	1720.49	1.00
betaMicroxtemp	1.35	-10.17	-3.36	6.54	9.73	489.23	1.01
betaMicroxLAI	0.15	-8.76	-1.80	2.10	12.01	517.12	1.01