The Journal of Open Source Software

ODINN.jl: Scientific machine learning glacier modelling

Jordi Bolibar [1,2¶], **Facundo Sapienza** [3,4¶], **Alban Gossard**[1], **Lisa Girod**[1], **Mathieu le Séac'h**[1], **Lucille Gimenes**[1], **Vivek Gajadhar**[2], **Fabien Maussion**[5,6], **Ching-Yao Lai**[3], **Bert Wouters**[2], **and Fernando Pérez** [4]

1 Univ. Grenoble Alpes, CNRS, IRD, G-INP, Institut des Géosciences de l'Environnement, Grenoble, France **2** Faculty of Civil Engineering and Geosciences, Delft University of Technology, Delft, The Netherlands **3** Department of Geophysics, Stanford University, Stanford, United States **4** Department of Statistics, University of California, Berkeley, United States **5** Bristol Glaciology Centre, School of Geographical Sciences, University of Bristol, Bristol, UK **6** Department of Atmospheric and Cryospheric Sciences, University of Innsbruck, Innsbruck, Austria ¶ Corresponding author

DOI: 10.xxxxxx/draft

Software

- Review
- Repository
- Archive

Editor: Open Journals

Reviewers:

- @openjournals

Submitted: 01 January 1970
Published: unpublished

License
Authors of papers retain copyright and release the work under a Creative Commons Attribution 4.0 International License (CC BY 4.0).

DRAFT

Summary

`ODINN.jl` is a glacier model leveraging scientific machine learning (SciML) methods to perform forward and inverse simulations of large-scale glacier evolution, here meaning tens to thousands of glaciers. It can simulate both surface mass balance and ice flow dynamics through a modular architecture which enables the user to easily modify model components.

The key feature of `ODINN.jl` is its differentiability, which enables hybrid modelling that combines mechanistic and data-driven components. Since the whole ecosystem is differentiable (where differentiable means the ability to compute model derivatives with respect to parameters (Shen et al., 2023)), we can optimize almost any model component, providing a framework to tackle a wide range of scientific problems (Bolibar et al., 2023). `ODINN.jl` can optimize, separately or together, in a steady-state (time-independent simulation) or transient (time-dependent simulation) way the following:

- The initial or intermediate state of glaciers (i.e. their ice thickness) or the equivalent ice surface velocities.
- Model parameters (e.g. the Glen coefficient `A` related to ice viscosity in a 2D Shallow Ice Approximation (Hutter, 1983)), in a gridded or scalar format. This can be done for multiple time steps where observations (e.g. ice surface velocities) are available.
- The parameters of a statistical regressor (e.g. a neural network), used to parametrize a subpart or one or more coefficients of an ice flow mechanistic model. This enables the exploration of empirical laws describing physical processes of glaciers, leveraging Universal Differential Equations (UDEs, Rackauckas et al. (2021)).

For this, it is necessary to differentiate (that is, computing gradients or derivatives) through complex code, including numerical solvers, which is a non-trivial task (Sapienza et al., 2024). We use reverse mode automatic differentiation (AD) based on the adjoint method to achieve this. We have two strategies for computing both the adjoint and the required vector-jacobian products (VJPs): (1) hand-derived adjoint equations, evaluated via AD using `Enzyme.jl` (Moses et al., 2021), as well as fully manual implementations of the spatially discrete and spatially continuous VJPs; and (2) automatic adjoints using `SciMLSensitivity.jl` (Rackauckas et al., 2019), available with different AD back-ends for the VJPs computation. These two approaches are complementary, with the manual adjoints being ideal for high-performance tasks by providing more control on the implementation, and serving as a ground truth for benchmarking and testing automatic adjoint methods from `SciMLSensitivity.jl`.

Beyond all these inverse modelling capabilities, `ODINN.jl` can also act as a more conventional forward glacier model, simulating glaciers in parallel, and easily customizing different model parametrizations and choices within the simulation. Its high modularity, combined with the easy access to a vast collection of datasets coming from the Open Global Glacier Model (OGGM, Maussion et al. (2019)), makes it very easy to run simulations, even on a single laptop: a machine with 8 CPU cores and 32 GB of RAM can run simulations of up to ten glaciers. Multiple ice flow dynamics models can be easily swapped, thanks to a modular architecture (see Software design). Models based on partial differential equations (PDEs) are solved using `DifferentialEquations.jl` (Rackauckas & Nie, 2017), which provides access to a large amount of numerical solvers. For now, we have implemented a 2D Shallow Ice Approximation (SIA, Hutter (1983)), but in the future we plan to incorporate other models, such as the Depth-Integrated Viscous Approximation (DIVA, Goldberg (2011)). Validation of numerical forward simulations is evaluated in the test suite based on exact analytical solutions of the SIA equation (Bueler et al., 2005). Surface mass balance is computed with a temperature-index model with a single degree-day factor (DDF). Temperature-index models with two DDFs might be added in the future. Machine learning-based models from the `MassBalanceMachine` (Sjursen et al., 2025) are partially available via `MassBalanceMachine.jl`. Models can be loaded and evaluated, but coupling them to the ice flow simulation is ongoing work.

Statement of need

`ODINN.jl` addresses the need for a glacier model that combines the physical interpretability of mechanistic approaches with the flexibility and data-assimilation capabilities of data-driven methods (Bolibar et al., 2023). By integrating both paradigms, it enables targeted inverse methods to learn parametrizations of glacier processes, capturing unknown physics while preserving the physically grounded structure of glacier dynamics through differential equations. `ODINN.jl` is aimed at users working on glacier modelling from a single glacier to whole regions, Earth system modellers, or researchers interested in using and exploring methods based on differentiable programming in the context of geoscientific modelling.

While purely mechanistic and purely data-driven glacier models already exist (e.g. Gagliardini et al. (2013), Maussion et al. (2019), Rounce et al. (2023), Bolibar et al. (2022)), they often do not fully exploit growing glacier observations, including surface velocity, ice thickness, topography, surface mass balance, and climate reanalyses. Existing empirical laws are not always directly linked to these observables, so calibration remains challenging. Differentiable programming and functional inversion offer a path forward by enabling new empirical relationships from proxies and testing hypotheses about poorly constrained processes such as basal sliding, creep, and calving.

Improving the representation of these processes is crucial for reliable projections of glacier evolution and impacts on freshwater availability and sea-level rise (IPCC, 2021). To this end, `ODINN.jl` provides a unified modelling ecosystem supporting advanced inverse calibration and efficient modular forward simulations for large-scale glacier studies.

Developing such a framework places strong demands on scientific software. Inefficient code and irreproducible implementations can restrict progress, emphasizing open-source, community-driven tools that follow modern research software engineering (RSE) practices (Combemale et al., 2023). These tools should support modular design for rapid prototyping and extension of workflows (Nyenah et al., 2024). Julia provides two key advantages: it addresses the two-language problem by combining high-level expressiveness with high-performance (Bezanson et al., 2017), and it enables source-code differentiability for modular gradient-based inverse optimization and hybrid modelling, including UDEs. With `ODINN.jl`, we aim to create a robust, future-proof framework bridging physical understanding and data-driven discovery. Its modular architecture, testing, and continuous integration (CI) ensure reproducibility and reliability, while its open design enables collaboration across glaciology and Earth system modelling.

DRAFT




Figure 1: Overview of the ODINN.jl ecosystem.

State of the field

The field of large-scale glacier modelling has considerably grown over the last decade, with most models being (or becoming) open-source. While global glacier models such as OGGM (Maussion et al., 2019), GloGEM (Huss & Hock, 2015) and PyGEM (Rounce et al., 2023) target global-scale efficient simulations of past and future glacier changes and sea-level rise contributions, `ODINN.jl` has a focus on inverse modelling for glacier physical processes and catchment-scale glacio-hydrological simulations. This scale is beyond the practical reach of high-order ice flow models such as Elmer/Ice (Gagliardini et al., 2013), which are usually limited to a small number of glaciers due to their computational cost. Moreover, `ODINN.jl` has the potential to implement more robust glacier physics compared to global models focusing on simplicity and lightweight parametrizations. IGM (Jouvet, 2023), with its neural network-driven ice flow emulator approach has some parallelisms to `ODINN.jl`, but the model's focus has mainly been on speeding up simulations for large-scale or paleo simulations. Alternatively, the new ice sheet model `DJuice.jl` (Moses et al., 2026) – a Julia translation of the well-known ISSM model – is perhaps the most similar model to `ODINN.jl`. They both leverage Julia's differentiability capabilities, and aim to exploit them to infer parametrizations and physical processes related to ice flow. Nonetheless, while `DJuice.jl` is an ice sheet model, `ODINN.jl` is highly specialized to simulate glaciers (i.e. anything outside Greenland and Antarctica), thanks to its compatibility with OGGM's data preprocessing.

DRAFT

Software design

`ODINN.jl` is an ecosystem composed of multiple packages, each one handling a specific task:

- `Sleipnir.jl`: Handles all the basic types, functions and datasets, common through the whole ecosystem, as well as data management tasks.
- `Muninn.jl`: Handles surface mass balance processes, via different types of models.
- `Huginn.jl`: Handles ice flow dynamics, by solving the ice flow partial differential equations (PDEs) using numerical methods. It can accommodate multiple types of ice flow models.
- `ODINN.jl`: Acts as the interface to the whole ecosystem, and provides the necessary tools to differentiate and optimize any model component. It can be seen as the SciML layer, enabling different types of inverse methods, using hybrid models combining differential equations with data-driven models.

Splitting large Julia (Bezanson et al., 2017) packages into smaller, focused subpackages is a good practice that enhances maintainability, usability, and collaboration. Modular design simplifies debugging, testing, and updates by isolating functionalities, while users benefit from faster precompilation and reduced memory overhead by loading only the subpackages they need. This approach also lowers the barrier for new contributors, fosters clearer dependency management, and ensures scalability as projects grow, ultimately creating a robust and adaptable software ecosystem. The ODINN ecosystem extends beyond this suite of Julia packages, by leveraging the data preprocessing tools of OGGM. This is done via the auxiliary Python library `Gungnir`, which is responsible for generating all the necessary data to initialize and run the model, such as glacier outlines from the Randolph Glacier Inventory (RGI Consortium (2023), RGI), digital elevation models (DEMs), ice thickness observations from GlaThiDa (GlaThiDa Consortium, 2020), ice surface velocities from different studies (Millan et al., 2022), and different sources of climate reanalyses and projections (Eyring et al., 2016; Lange, 2019). This implies that `ODINN.jl`, like OGGM, is virtually capable of simulating any of the ~274,000 glaciers on Earth (RGI Consortium, 2023).

`ODINN.jl` provides a high-level user-friendly interface, enabling the user to swap and replace most elements of a glacier simulation in a modular fashion. The main elements of a simulation, such as the `Parameters`, a `Model`, and a `Simulation` (i.e. a `Prediction` or an `Inversion`), are all objects that can be easily modified and combined. In a few lines of code, the user can automatically retrieve all necessary information for most glaciers on Earth, compose a `Model` based on a specific combination of surface mass balance and ice flow models, and incorporate data-driven models (e.g. a neural network) to parametrize specific physical processes of any of these components. Because each glacier evolves independently, both forward and inverse simulations run in parallel using multiprocessing. This is what makes it possible to run simulations and inversions for tens to thousands of the ~274,000 glaciers on Earth. These design choices come with trade-offs. Parallelism only happens across glaciers, not within a single glacier, so a large glacier does not run faster on more CPU cores. The current 2D SIA ice flow model is fast and differentiable, but does not include longitudinal stresses; a Depth-Integrated Viscous Approximation (DIVA), which is closer to a full Stokes solution, is planned next. Graphics Processing Unit (GPU) compatibility is still not ready, due to the difficulties of making GPU architectures compatible with automatic differentiation (AD). Nonetheless, it is planned for future versions.

Research impact statement

`ODINN.jl` has evolved through the last five years with code contributions during three postdoctoral positions, one PhD, and three master's internships. It has so far been used to explore the use of UDEs to invert hidden empirical laws in a synthetic glacier setup, where a prescribed rheological law was successfully recovered using a neural network (Bolibar et al., 2023). This proof-of-concept then served as a backbone to create the current architecture of

DRAFT

`ODINN.jl`, finalized with the recent 2.0 release. The main changes and scientific goals of this large software development investment are the capacity to now apply these methods to real remote sensing data for tens to thousands of glaciers, through transient inversions initialized from observed glacier states. This enables the exploration of new glacier basal sliding laws directly from heterogeneous observations, which remains a long-standing problem in glaciology (Minchew & Joughin, 2020). The development of the differentiable programming methods in `ODINN.jl` also served as a catalyst to write a review paper, together with other key players in this community, on differentiable programming for differential equations (Sapienza et al., 2024).

Additionally, `ODINN.jl` is currently being used in a newly funded 4-year project, to simulate past and future glacier changes in several catchments in the Andes and the Alps. These model outputs will then be combined with a hydrological model to investigate the impacts of glacier retreat on the hydrological regimes and drought mitigation under different climate change scenarios.

With these two research venues, `ODINN.jl` will continue to be developed and used to pursue both fundamental research on glacier modelling, and applied research to assess the impact of glacier retreat on freshwater availability and drought mitigation.

AI usage disclosure

Generative AI tools (Claude Sonnet 5.5, Claude Opus 5.5, GLM 5.3 hosted by Mistral, and Mistral Medium 3.5) have been used to generate docstrings and documentation, and to assist in the coding of tests, debugging and the implementation of small new features over the existing architecture. They were also used for small corrections to the paper text during the review process. All AI-assisted outputs were reviewed, edited and validated by the authors, who made all core design decisions.

Acknowledgements

We acknowledge the help of Chris Rackauckas for the debugging and discussion of issues related to the SciML Julia ecosystem, Redouane Lguensat for scientific discussions on the first prototype of the model, and Julien le Sommer for scientific discussions around differentiable programming. We thank all the developers of the SciML Julia ecosystem who work in each one of the core libraries used within `ODINN.jl`. JB acknowledges financial support from the Nederlandse Organisatie voor Wetenschappelijk Onderzoek, Stichting voor de Technische Wetenschappen (Vidi grant 016.Vidi.171.063), a TU Delft Climate Action grant and a PEPR TRACCS grant from the Agence Nationale de la Recherce as part of France 2030 (reference ANR-25-EXTR-0006). FS and CYL were supported by NSF via grant number OPP-2441132 and the Alfred P. Sloan Foundation under grant number FG-2024-21649. FS and FP acknowledge funding from the National Science Foundation (EarthCube programme under awards 1928406 and 1928374). AG acknowledges funding from the MIAI cluster and Agence Nationale de la Recherche (ANR) in the context of France 2030 (grant ANR-23-IACL-0006).

References

Bezanson, J., Edelman, A., Karpinski, S., & Shah, V. B. (2017). Julia: A Fresh Approach to Numerical Computing. *SIAM Review*, *59*(1), 65–98. https://doi.org/10.1137/141000671

Bolibar, J., Rabatel, A., Gouttevin, I., Zekollari, H., & Galiez, C. (2022). Nonlinear sensitivity of glacier mass balance to future climate change unveiled by deep learning. *Nature Communications*, *13*(1), 409. https://doi.org/10.1038/s41467-022-28033-0

DRAFT

Bolibar, J., Sapienza, F., Maussion, F., Lguensat, R., Wouters, B., & Pérez, F. (2023). Universal differential equations for glacier ice flow modelling. *Geoscientific Model Development*, *16*(22), 6671–6687. https://doi.org/10.5194/gmd-16-6671-2023

Bueler, E., Lingle, C. S., Kallen-Brown, J. A., Covey, D. N., & Bowman, L. N. (2005). Exact solutions and verification of numerical models for isothermal ice sheets. *Journal of Glaciology*, *51*(173), 291–306. https://doi.org/10.3189/172756505781829449

Combemale, B., Gray, J., & Rumpe, B. (2023). Research software engineering and the importance of scientific models. *Software and Systems Modeling*, *22*(4), 1081–1083. https://doi.org/10.1007/s10270-023-01119-z

Eyring, V., Bony, S., Meehl, G. A., Senior, C. A., Stevens, B., Stouffer, R. J., & Taylor, K. E. (2016). Overview of the Coupled Model Intercomparison Project Phase 6 (CMIP6) experimental design and organization. *Geoscientific Model Development*, *9*(5), 1937–1958. https://doi.org/10.5194/gmd-9-1937-2016

Gagliardini, O., Zwinger, T., Gillet-Chaulet, F., Durand, G., Favier, L., Fleurian, B. de, Greve, R., Malinen, M., Martín, C., Råback, P., Ruokolainen, J., Sacchettini, M., Schäfer, M., Seddik, H., & Thies, J. (2013). Capabilities and performance of Elmer/Ice, a new-generation ice sheet model. *Geoscientific Model Development*, *6*(4), 1299–1318. https://doi.org/10.5194/gmd-6-1299-2013

GlaThiDa Consortium. (2020). *Glacier Thickness Database 3.1.0*. World Glacier Monitoring Service, Zurich, Switzerland.

Goldberg, D. N. (2011). A variationally derived, depth-integrated approximation to a higher-order glaciological flow model. *Journal of Glaciology*, *57*(201), 157–170. https://doi.org/10.3189/002214311795306763

Huss, M., & Hock, R. (2015). A new model for global glacier change and sea-level rise. *Frontiers in Earth Science*, *3*. https://doi.org/10.3389/feart.2015.00054

Hutter, K. (1983). *Theoretical Glaciology*. Springer Netherlands. https://doi.org/10.1007/978-94-015-1167-4

IPCC. (2021). *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change: Vols. In Press*. Cambridge University Press. https://doi.org/10.1017/9781009157896

Jouvet, G. (2023). Inversion of a Stokes glacier flow model emulated by deep learning. *Journal of Glaciology*, *69*(273), 13–26. https://doi.org/10.1017/jog.2022.41

Lange, S. (2019). *WFDE5 over land merged with ERA5 over the ocean (W5E5)*. GFZ Data Services. https://doi.org/10.5880/PIK.2019.023

Maussion, F., Butenko, A., Champollion, N., Dusch, M., Eis, J., Fourteau, K., Gregor, P., Jarosch, A. H., Landmann, J., Oesterle, F., Recinos, B., Rothenpieler, T., Vlug, A., Wild, C. T., & Marzeion, B. (2019). The Open Global Glacier Model (OGGM) v1.1. *Geoscientific Model Development*, *12*(3), 909–931. https://doi.org/10.5194/gmd-12-909-2019

Millan, R., Mouginot, J., Rabatel, A., & Morlighem, M. (2022). Ice velocity and thickness of the world's glaciers. *Nature Geoscience*, *15*(2), 124–129. https://doi.org/10.1038/s41561-021-00885-z

Minchew, B., & Joughin, I. (2020). Toward a universal glacier slip law. *Science*, *368*(6486), 29–30. https://doi.org/10.1126/science.abb3566

Moses, W. S., Cheng, G., Churavy, V., Gelbrecht, M., Klöwer, M., Kump, J., Morlighem, M., Williamson, S., Apte, D., Berg, P., Giordano, M., Hill, C., Loose, N., Montoison, A., Narayanan, S. H. K., Pal, A., Schanen, M., Silvestri, S., Wagner, G. L., & Heimbach, P. (2026). DJ4Earth: Differentiable, and Performance-portable Earth System Modeling

DRAFT

via Program Transformations. *Journal of Advances in Modeling Earth Systems*, *18*(5), e2025MS005615. https://doi.org/10.1029/2025MS005615

Moses, W. S., Churavy, V., Paehler, L., Hückelheim, J., Narayanan, S. H. K., Schanen, M., & Doerfert, J. (2021). Reverse-mode automatic differentiation and optimization of GPU kernels via enzyme. *Proceedings of the International Conference for High Performance Computing, Networking, Storage and Analysis*. https://doi.org/10.1145/3458817.3476165

Nyenah, E., Döll, P., Katz, D. S., & Reinecke, R. (2024). Software sustainability of global impact models. *Geoscientific Model Development Discussions*, *2024*, 1–29. https://doi.org/10.5194/gmd-2024-97

Rackauckas, C., Innes, M., Ma, Y., Bettencourt, J., White, L., & Dixit, V. (2019). DiffEqFlux.jl - A Julia Library for Neural Differential Equations. *arXiv:1902.02376 [Cs, Stat]*. http://arxiv.org/abs/1902.02376

Rackauckas, C., Ma, Y., Martensen, J., Warner, C., Zubov, K., Supekar, R., Skinner, D., Ramadhan, A., & Edelman, A. (2021). *Universal Differential Equations for Scientific Machine Learning*. arXiv. https://doi.org/10.48550/arXiv.2001.04385

Rackauckas, C., & Nie, Q. (2017). DifferentialEquations.jl – A Performant and Feature-Rich Ecosystem for Solving Differential Equations in Julia. *Journal of Open Research Software*, *5*, 15. https://doi.org/10.5334/jors.151

RGI Consortium. (2023). *Randolph Glacier Inventory - A Dataset of Global Glacier Outlines, Version 7*. National Snow; Ice Data Center. https://doi.org/10.5067/F6JMOVY5NAVZ

Rounce, D. R., Hock, R., Maussion, F., Hugonnet, R., Kochtitzky, W., Huss, M., Berthier, E., Brinkerhoff, D., Compagno, L., Copland, L., Farinotti, D., Menounos, B., & McNabb, R. W. (2023). Global glacier change in the 21st century: Every increase in temperature matters. *Science*, *379*(6627), 78–83. https://doi.org/10.1126/science.abo1324

Sapienza, F., Bolibar, J., Schäfer, F., Groenke, B., Pal, A., Boussange, V., Heimbach, P., Hooker, G., Pérez, F., Persson, P.-O., & Rackauckas, C. (2024). *Differentiable Programming for Differential Equations: A Review*. arXiv. http://arxiv.org/abs/2406.09699

Shen, C., Appling, A. P., Gentine, P., Bandai, T., Gupta, H., Tartakovsky, A., Baity-Jesi, M., Fenicia, F., Kifer, D., Li, L., Liu, X., Ren, W., Zheng, Y., Harman, C. J., Clark, M., Farthing, M., Feng, D., Kumar, P., Aboelyazeed, D., … Lawson, K. (2023). Differentiable modelling to unify machine learning and physical models for geosciences. *Nature Reviews Earth & Environment*, 1–16. https://doi.org/10.1038/s43017-023-00450-9

Sjursen, K. H., Bolibar, J., Meer, M. van der, Andreassen, L. M., Biesheuvel, J. P., Dunse, T., Huss, M., Maussion, F., Rounce, D. R., & Tober, B. (2025). Machine learning improves seasonal mass balance prediction for unmonitored glaciers. *EGUsphere*, 1–39. https://doi.org/10.5194/egusphere-2025-1206

DRAFT