

Analysis of Linear Programming

Part 3e: Matrix Form, Dual Simplex and Revised Simplex

IE 311 – Operations Research I

- So far we have learned the (primal) simplex method as the only solution procedure for linear programs.
- We now discuss different versions of the simplex method:
- first we derive the **matrix form** of the (primal) simplex method;
- then we discuss the **optimality conditions** in linear programming.
- Finally, combining the above, we reach the **Dual Simplex** and **Revised Simplex** methods.

- Let us consider an LP in standard form (assume that c is an n -row vector, b is a m -column vector and A is an $m \times n$ matrix):

$$\max x_0 = cx$$

$$\text{s.t.} \quad Ax = b$$

$$x \geq 0.$$

- Suppose that we are given a basic solution where the basic variables are indexed by the set BV .
- Question: How can we construct the simplex tableau via matrix operations?

- Recall the LP in standard form:

$$\max x_0 = cx$$

$$\text{s.t.} \quad Ax = b$$

$$x \geq 0.$$

- Let us partition the variables as basic, denoted by x_{BV} , and nonbasic, denoted by x_{NBV} (also partition c and A accordingly).

$$\max x_0 = c_{BV}x_{BV} + c_{NBV}x_{NBV}$$

$$\text{s.t.} \quad Bx_{BV} + Nx_{NBV} = b$$

$$x_{BV}, x_{NBV} \geq 0.$$

- Since B is a basis matrix (B^{-1} exists), we have $x_{BV} = B^{-1}(b - Nx_{NBV})$.

- If we plug in $x_{BV} = B^{-1}(b - Nx_{NBV})$ into the following LP,

$$\max x_0 = c_{BV}x_{BV} + c_{NBV}x_{NBV}$$

$$\text{s.t.} \quad Bx_{BV} + Nx_{NBV} = b$$

$$x_{BV}, x_{NBV} \geq 0.$$

- we obtain,

$$\max x_0 \quad + (c_{BV}B^{-1}N - c_{NBV})x_{NBV} = c_{BV}B^{-1}b$$

$$\text{s.t.} \quad x_{BV} + (B^{-1}N)x_{NBV} = B^{-1}b$$

$$x_{BV}, x_{NBV} \geq 0.$$

- or in tableau form:

Row	BV		RHS
0	x_0	$x_0 + (c_{BV}B^{-1}N - c_{NBV})x_{NBV}$	$= c_{BV}B^{-1}b$
$1, \dots, m$	x_{BV}	$x_{BV} + (B^{-1}N)x_{NBV}$	$= B^{-1}b$

- Given a basis matrix B (corresponding to the columns in BV), we obtain the following simplex tableau:

Row	BV		RHS
0	x_0	$x_0 + (c_{BV}B^{-1}N - c_{NBV})x_{NBV}$	$= c_{BV}B^{-1}b$
$1, \dots, m$	x_{BV}	$x_{BV} + (B^{-1}N)x_{NBV}$	$= B^{-1}b$

- If we have $B^{-1}b \geq 0$, then BV gives a **basic feasible solution**.
- The row vector $\bar{c} := c_{BV}B^{-1}N - c_{NBV}$ is called *the reduced cost*.
- Reduced cost of variable j is computed as $\bar{c}_j := c_{BV}B^{-1}A^j - c_j$, where A^j is the corresponding column of the matrix A .
- Note that the current tableau is **optimal if all reduced costs are nonnegative**.

- Consider the following LP in standard form:

$$\max x_0 = 3x_1 - 2x_2$$

$$\text{s.t.} \quad x_1 + x_2 + x_3 = 2$$

$$x_1 + x_4 = 1$$

$$x_1, x_2, x_3, x_4 \geq 0.$$

- Construct the simplex tableau via matrix operations for
 - $BV = \{1,2\}$, $NBV = \{3,4\}$
 - $BV = \{2,1\}$, $NBV = \{3,4\}$
 - $BV = \{1,3\}$, $NBV = \{2,4\}$
 - $BV = \{2,4\}$, $NBV = \{1,3\}$.

Matrix Form of the Simplex Tableau: Example (continued)

$$B = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}, N = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, c_{\text{BV}} = [3 \quad -2], c_{\text{NBV}} = [0 \quad 0], b = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, B^{-1} = \begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix},$$

$$B^{-1}N = \begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix}, c_{\text{BV}}B^{-1}N - c_{\text{NBV}} = [-2 \quad 5], B^{-1}b = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, c_{\text{BV}}B^{-1}b = 1.$$

Row	BV				RHS
0	x_0	x_0	$-2x_3$	$+5x_4$	$= 1$
1	x_1	x_1		$+x_4$	$= 1$
2	x_2	x_2	$+x_3$	$-x_4$	$= 1$

Matrix Form of the Simplex Tableau: Example (continued)

$$B = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, N = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, c_{BV} = [-2 \quad 3], c_{NBV} = [0 \quad 0], b = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, B^{-1} = \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix},$$

$$B^{-1}N = \begin{bmatrix} 1 & -1 \\ 0 & 1 \end{bmatrix}, c_{BV}B^{-1}N - c_{NBV} = [-2 \quad 5], B^{-1}b = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, c_{BV}B^{-1}b = 1.$$

Row	BV				RHS	
0	x_0	x_0	$-2x_3$	$+5x_4$	$= 1$	
1	x_2		x_2	$+x_3$	$-x_4$	$= 1$
2	x_1		x_1		$+x_4$	$= 1$

Matrix Form of the Simplex Tableau: Example (continued)

$$B = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}, N = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, c_{\text{BV}} = [3 \quad 0], c_{\text{NBV}} = [-2 \quad 0], b = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, B^{-1} = \begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix},$$

$$B^{-1}N = \begin{bmatrix} 0 & 1 \\ 1 & -1 \end{bmatrix}, c_{\text{BV}}B^{-1}N - c_{\text{NBV}} = [2 \quad 3], B^{-1}b = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, c_{\text{BV}}B^{-1}b = 3.$$

Row	BV					RHS
0	x_0	x_0	$+2x_2$	$+3x_4$	$= 3$	
1	x_1	x_1		$+x_4$	$= 1$	
2	x_3		x_2	$+x_3$	$-x_4$	$= 1$

Matrix Form of the Simplex Tableau: Example (continued)

$$B = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, N = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}, c_{\text{BV}} = [-2 \quad 0], c_{\text{NBV}} = [3 \quad 0], b = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, B^{-1} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix},$$

$$B^{-1}N = \begin{bmatrix} 1 & 1 \\ 1 & 0 \end{bmatrix}, c_{\text{BV}}B^{-1}N - c_{\text{NBV}} = [-5 \quad -2], B^{-1}b = \begin{bmatrix} 2 \\ 1 \end{bmatrix}, c_{\text{BV}}B^{-1}b = -4.$$

Row	BV				RHS
0	x_0	x_0	$-5x_1$	$-2x_3$	$= -4$
1	x_2		$x_1 + x_2$	$+x_3$	$= 2$
2	x_4		x_1	$+x_4$	$= 1$

Exercise: Try to construct the simplex tableau via matrix operations for the following selections. Comment on which ones lead to basic solutions, basic feasible solutions or none.

- $BV = \{3,1\}$, $NBV = \{2,4\}$
- $BV = \{3,4\}$, $NBV = \{1,2\}$
- $BV = \{1,4\}$, $NBV = \{2,3\}$
- $BV = \{2,3\}$, $NBV = \{1,3\}$.

- Consider the following primal-dual pair (here, y is a row-vector):

$$\max \quad cx$$

$$\text{s.t.} \quad Ax = b$$

$$x \geq 0.$$

$$\min \quad yb$$

$$\text{s.t.} \quad yA \geq c$$

$$y \text{ urs.}$$

- Given BV, we can rewrite as follows:

$$\max \quad c_{\text{BV}}x_{\text{BV}} + c_{\text{NBV}}x_{\text{NBV}}$$

$$\text{s.t.} \quad Bx_{\text{BV}} + Nx_{\text{NBV}} = b$$

$$x_{\text{BV}}, x_{\text{NBV}} \geq 0.$$

$$\min \quad yb$$

$$\text{s.t.} \quad yB \geq c_{\text{BV}}, \quad yN \geq c_{\text{NBV}}$$

$$y \text{ urs.}$$

- Given BV, we can rewrite primal-dual pair as follows:

$$\begin{array}{ll} \max & c_{BV}x_{BV} + c_{NBV}x_{NBV} \\ \text{s.t.} & Bx_{BV} + Nx_{NBV} = b \\ & x_{BV}, x_{NBV} \geq 0. \end{array} \quad \begin{array}{ll} \min & yb \\ \text{s.t.} & yB \geq c_{BV}, yN \geq c_{NBV} \\ & y \text{ urs.} \end{array}$$

- Let us *predict* a dual solution as $y_{BV} := c_{BV}B^{-1}$.
- When initialized with a BFS, the simplex method guarantees that
 - $(x_{BV}, x_{NBV}) := (B^{-1}b, 0)$ is feasible at every iteration
 - complementary slackness conditions hold between (x_{BV}, x_{NBV}) and y_{BV}
 - we terminate when the dual solution $y_{BV} := c_{BV}B^{-1}$ becomes feasible.

- CLAIM:** The simplex method guarantees that complementary slackness conditions $(yA - c)_j x_j = 0$ (or equivalently $\bar{c}_j x_j = 0$) hold between $(x_{BV}, x_{NBV}) := (B^{-1}b, 0)$ and $y_{BV} := c_{BV}B^{-1}$.

Row	BV		RHS
0	x_0	$x_0 + (c_{BV}B^{-1}N - c_{NBV})x_{NBV}$	$= c_{BV}B^{-1}b$
$1, \dots, m$	x_{BV}	$x_{BV} + (B^{-1}N)x_{NBV}$	$= B^{-1}b$

- PROOF:**
 - If x_j is basic, then its reduced cost $\bar{c}_j$ is 0.
 - If x_j is nonbasic, then its value is 0.

- **CLAIM:** The simplex method terminates when the dual solution $y_{BV} := c_{BV}B^{-1}$ becomes feasible.

Row	BV		RHS
0	x_0	$x_0 + (c_{BV}B^{-1}N - c_{NBV})x_{NBV}$	$= c_{BV}B^{-1}b$
$1, \dots, m$	x_{BV}	$x_{BV} + (B^{-1}N)x_{NBV}$	$= B^{-1}b$

- **PROOF:** At termination, we have $\bar{c} := c_{BV}B^{-1}N - c_{NBV} \geq 0$. Since

- $yB = c_{BV}B^{-1}B \geq c_{BV}$, and $yN = c_{BV}B^{-1}N \geq c_{NBV}$,

$y_{BV} := c_{BV}B^{-1}$ is a feasible solution to the dual problem:

$$\min yb$$

$$\text{s.t. } yB \geq c_{BV}, \quad yN \geq c_{NBV}$$

$$y \text{ urs.}$$

- CLAIM:** Strong duality *holds* between $(x_{\text{BV}}, x_{\text{NBV}}) := (B^{-1}b, 0)$ and $y_{\text{BV}} := c_{\text{BV}}B^{-1}$ at every iteration.

Row	BV		RHS
0	x_0	$x_0 + (c_{\text{BV}}B^{-1}N - c_{\text{NBV}})x_{\text{NBV}}$	$= c_{\text{BV}}B^{-1}b$
$1, \dots, m$	x_{BV}	$x_{\text{BV}} + (B^{-1}N)x_{\text{NBV}}$	$= B^{-1}b$

• PROOF:

$$cx = c_{\text{BV}}x_{\text{BV}} = c_{\text{BV}}(B^{-1}b) = (c_{\text{BV}}B^{-1})b = y_{\text{BV}}b = yb.$$

- Consider the standard form of a well-structured LP (only $\leq$ constraints with nonnegative RHS):

$$\max x_0 = cx$$

$$\text{s.t.} \quad Ax + I\tilde{x} = b$$

$$x \geq 0, \tilde{x} \geq 0.$$

- Then,
 - values of the dual variables are equal to the reduced cost of the slack variables since $\bar{c}_{\text{slack}} := c_{\text{BV}}B^{-1}I - c_{\text{slack}} = c_{\text{BV}}B^{-1} = y_{\text{BV}}$ (note $c_{\text{slack}} = 0$)
 - the basis inverse can be retrieved from the columns under the slack variables since $B^{-1}I = B^{-1}$.
- See the example above!

- Consider the following primal-dual pair (here, y is a row-vector):

$$\max \quad cx$$

$$\min \quad yb$$

$$\text{s.t.} \quad Ax = b$$

$$\text{s.t.} \quad yA \geq c$$

$$x \geq 0.$$

$$y \text{ urs.}$$

- A primal-dual pair (x, y) is **optimal** if and only if
 - x is primal feasible, that is, $Ax = b, x \geq 0$
 - y is dual feasible, that is, $yA \geq c$
 - (x, y) satisfies complementary slackness, that is, $(yA - c)_j x_j = 0$.
- Complementary slackness is equivalent to strong duality, that is, $yb = cx$.

- **(Primal) Simplex Method:**
 - keeps **primal feasibility** and complementary slackness intact at each iteration, and
 - terminates when **dual feasibility** is achieved.
- There is an alternative method for LPs with appropriate structure, the **Dual Simplex Method**.
- **Dual Simplex Method:**
 - keeps **dual feasibility** and complementary slackness intact at each iteration, and
 - terminates when **primal feasibility** is achieved.

The simplex algorithm proceeds as follows:

- **Step 1:** Convert the LP to standard form.
- **Step 2:** Obtain a BFS (if possible) from the standard form.
- **Step 3:** Determine whether the current BFS is optimal.
- **Step 4:** If not optimal, determine which nonbasic variable should become basic and which basic variable should become nonbasic, to obtain a better objective value (or detect unboundedness).
- **Step 5:** Use EROs to obtain that new BFS, then return to Step 3.

The dual simplex algorithm proceeds as follows:

- **Step 1:** Convert the LP to standard form.
- **Step 2:** Obtain a **dual BFS** from the standard form.
- **Step 3:** Determine whether the current dual BFS is **primal feasible**.
- **Step 4:** If not primal feasible, determine which basic variable should become nonbasic and which nonbasic variable should become basic, to obtain a new dual BFS with a worse objective value (or detect infeasibility).
- **Step 5:** Use EROs to obtain that new dual BFS, then return to Step 3.


- Solve the following problem using the dual simplex method:

$$\min x_0 = x_1 + x_2$$

$$\text{s.t.} \quad x_1 + x_2 \geq 4$$

$$x_1 + 3x_2 \geq 6$$

$$x_1, x_2 \geq 0.$$



- In Step 1, we transform the problem into the following standard form of a maximization LP (note the negative RHS vector):

$$-\max x_0 = -x_1 - x_2$$

$$\text{s.t.} \quad -x_1 - x_2 + x_3 = -4$$


$$-x_1 - 3x_2 + x_4 = -6$$

Dual Simplex Example: Step 2

- In Step 2, we obtain an initial dual BFS:

Row	BV					RHS
0	x'_0	x'_0	$+x_1$	$+x_2$		$= 0$
1	x_3		$-x_1$	$-x_2$	$+x_3$	$= -4$
2	x_4		$-x_1$	$-3x_2$	$+x_4$	$= -6$

- Note:** The slack variables x_3 and x_4 are chosen as basic variables.
- Reduced costs** of the slack variables correspond to the **dual variables** (in this iteration, the origin is a dual BFS).
- RHS column** in this tableau is not nonnegative.



- In Step 4, we first decide which basic variable to make nonbasic (**look at RHS**):

Row	BV					RHS
0	x'_0	x'_0	$+x_1$	$+x_2$		$= 0$
1	x_3		$-x_1$	$-x_2$	$+x_3$	$= -4$
2	x_4		$-x_1$	$-3x_2$	$+x_4$	$= -6$

- Idea:** Intuitively, choosing the most negative RHS might decrease the infeasibility the most (although any basic variable with negative RHS can be chosen).
- In this case, we can choose the basic variable x_4 to leave the basis, hence, make it nonbasic.**

- In Step 4, we next decide which nonbasic variable to make basic (**look at the row of the leaving variable and row-0**):

Row	BV					RHS
0	x'_0	x'_0	$+x_1$	$+x_2$		$= 0$
1	x_3		$-x_1$	$-x_2$	$+x_3$	$= -4$
2	x_4		$-x_1$	$-3x_2$		$+x_4 = -6$

- Idea:** What is the adjacent BFS obtained by making x_4 nonbasic?
- We need to perform the so-called **minimum ratio test**:
 - Obtain the negative of the entrywise ratio of row-0 and the row of the leaving variable among the columns that have negative entry in the row of the leaving variable.
 - Choose the nonbasic variable that gives the minimum ratio as the entering variable (if there is no such variable, dual is unbounded and primal is infeasible):


$$\text{Minimum ratio test: } \min \left(\frac{0}{-1}, \frac{0}{-3} \right) = 0$$

- In Step 5, we use EROs to find a new BFS with a worse objective function value (**pivoting**):

Row	BV					RHS
0	x'_0	x'_0	$+x_1$	$+x_2$		$= 0$
1	x_3		$-x_1$	$-x_2$	$+x_3$	$= -4$
2	x_4		$-x_1$	$-3x_2$	$+x_4$	$= -6$

- Idea:** How can we make the column of the entering variable a unit vector by pivoting?

Row	BV					RHS
0	x'_0	x'_0	$+\frac{2}{3}x_1$		$+\frac{1}{3}x_4$	$= -2$
1	x_3		$-\frac{2}{3}x_1$	$+x_3$	$-\frac{1}{3}x_4$	$= -2$
2	x_2		$\frac{1}{3}x_1$	$+x_2$	$-\frac{1}{3}x_4$	$= 2$



- In Step 3, we check whether the current BFS is primal feasible (**look at RHS**):

Row	BV					RHS
0	x'_0	x'_0	$+\frac{2}{3}x_1$		$+\frac{1}{3}x_4$	$= -2$
1	x_3		$-\frac{2}{3}x_1$	$+x_3$	$-\frac{1}{3}x_4$	$= -2$
2	x_2		$\frac{1}{3}x_1$	$+x_2$	$-\frac{1}{3}x_4$	$= 2$

- Idea:** If we move to an adjacent dual BFS, can we decrease the *infeasibility* (make the RHS less negative while keeping reduced costs nonnegative)?
- Note that the value of x_3 is negative.**
- Therefore, the current dual BFS is NOT primal feasible.

- In Step 4, we first decide which basic variable to make nonbasic (**look at RHS**):

Row	BV					RHS
0	x'_0	x'_0	$+\frac{2}{3}x_1$		$+\frac{1}{3}x_4$	$= -2$
1	x_3		$-\frac{2}{3}x_1$	$+x_3$	$-\frac{1}{3}x_4$	$= -2$
2	x_2		$\frac{1}{3}x_1$	$+x_2$	$-\frac{1}{3}x_4$	$= 2$

- Idea:** Intuitively, choosing the most negative RHS might decrease the infeasibility the most (although any basic variable with negative RHS can be chosen).
- In this case, there is only one candidate: the basic variable x_3 leaves the basis.**

- In Step 4, we next decide which nonbasic variable to make basic (**look at the row of the leaving variable and row-0**):

Row	BV					RHS
0	x'_0	x'_0	$+\frac{2}{3}x_1$		$+\frac{1}{3}x_4$	$= -2$
1	x_3		$-\frac{2}{3}x_1$	$+x_3$	$-\frac{1}{3}x_4$	$= -2$
2	x_2		$\frac{1}{3}x_1$	$+x_2$	$-\frac{1}{3}x_4$	$= 2$

- Idea:** What is the adjacent BFS obtained by making x_3 nonbasic?
- We perform the so-called **minimum ratio test**:

$$\text{There is a tie since } \min \left\{ -\frac{2/3}{-2/3}, -\frac{1/3}{-1/3} \right\} = 1.$$

- We choose x_1 to enter the basis (we could have also chosen x_4).
- Note: The next dual BFS will be degenerate


Dual Simplex Example: Step 5 (Again)

- In Step 5, we use EROs to find a new BFS with a worse objective function value (**pivoting**):

Row	BV				RHS
0	x'_0	x'_0	$+\frac{2}{3}x_1$	$+\frac{1}{3}x_4$	$= -2$
1	x_3		$-\frac{2}{3}x_1$	$+x_3$	$-\frac{1}{3}x_4$ $= -2$
2	x_2		$\frac{1}{3}x_1$	$+x_2$	$-\frac{1}{3}x_4$ $= 2$

- Idea:** How can we make the column of the entering variable a unit vector by pivoting?

Row	BV				RHS
0	x'_0	x'_0		$+x_3$	$= -4$
1	x_1		x_1	$-\frac{3}{2}x_3$	$+\frac{1}{2}x_4$ $= 3$
2	x_2		$+x_2$	$+\frac{1}{2}x_3$	$-\frac{1}{2}x_4$ $= 1$



- In Step 3, we check whether the current BFS is primal feasible (**look at RHS**):

Row	BV				RHS
0	x'_0	x'_0	$+x_3$		$= -4$
1	x_1	x_1	$-\frac{3}{2}x_3$	$+\frac{1}{2}x_4$	$= 3$
2	x_2	$+x_2$	$+\frac{1}{2}x_3$	$-\frac{1}{2}x_4$	$= 1$

- Note that the values of x_1 and x_2 are nonnegative, hence, we obtain a primal feasible solution.**
- Therefore, the current BFS is **OPTIMAL** with objective value 4.
- Optimal primal solution: $x_1 = 3$, $x_2 = 1$.

- Consider the following LP:

$$\max x_0 = cx$$

$$\text{s.t.} \quad Ax = b$$

$$x \geq 0.$$

- Given a basis BV, we can rewrite the LP as follows:

$$\max x_0 \quad + (c_{\text{BV}}B^{-1}N - c_{\text{NBV}})x_{\text{NBV}} = c_{\text{BV}}B^{-1}b$$

$$\text{s.t.} \quad x_{\text{BV}} + (B^{-1}N)x_{\text{NBV}} = B^{-1}b$$

$$x_{\text{BV}}, x_{\text{NBV}} \geq 0.$$

- We will now see a version of the simplex method (called the *Revised Simplex Method*) which does NOT use tableaus at all.

The revised simplex algorithm proceeds as follows:

- **Step 1** Convert the LP to standard form.
- **Step 2** Obtain an initial feasible basis BV.
- **Step 3** Given BV, compute basis matrix inverse B^{-1} , primal solution $B^{-1}b$, reduced cost $\bar{c} := c_{BV}B^{-1}N - c_{NBV}$ and objective value $c_{BV}B^{-1}b$. Determine whether the current BFS is optimal (check if there exists $\bar{c}_j < 0$).
- **Step 4** If the current BFS is not optimal, then determine which nonbasic variable (say j^E) should become a basic variable. Compute the column of the entering variable $B^{-1}A^{j^E}$. Use minimum ratio test to decide which basic variable (say j^L) should become a nonbasic variable (or determine if the problem is unbounded).
- **Step 5** Update $BV = BV \cup \{j^E\} \setminus \{j^L\}$. Go back to step 3.

- Solve the following problem using the revised simplex method:

$$\max x_0 = 5x_1 + 4x_2$$

$$\text{s.t.} \quad 3x_1 + 2x_2 \leq 12$$

$$x_1 + 6x_2 \leq 12$$

$$x_1, x_2 \geq 0.$$

- In Step 1, we transform the problem into the following standard form

$$\max x_0 = 5x_1 + 4x_2$$

$$\text{s.t.} \quad 3x_1 + 2x_2 + x_3 = 12$$

$$x_1 + 6x_2 + x_4 = 12$$

$$x_1, x_2, x_3, x_4 \geq 0.$$

- Here is the matrix form of the LP:

$$A = \begin{bmatrix} 3 & 2 & 1 & 0 \\ 1 & 6 & 0 & 1 \end{bmatrix}, c = [5 \quad 4 \quad 0 \quad 0], b = \begin{bmatrix} 12 \\ 12 \end{bmatrix}.$$

- In Step 2, we should obtain an initial feasible BFS.
- Note:** The slack variables x_3 and x_4 can be chosen as basic variables since the RHS is nonnegative.
- Hence, $BV = \{3,4\}$.

- Recall the matrix form:

$$A = \begin{bmatrix} 3 & 2 & 1 & 0 \\ 1 & 6 & 0 & 1 \end{bmatrix}, c = [5 \quad 4 \quad 0 \quad 0], b = \begin{bmatrix} 12 \\ 12 \end{bmatrix}.$$

- We compute the following matrices for $BV = \{3,4\}$:

$$B = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, N = \begin{bmatrix} 3 & 2 \\ 1 & 6 \end{bmatrix}, c_{BV} = [0 \quad 0], c_{NBV} = [5 \quad 4], B^{-1} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix},$$

$$B^{-1}N = \begin{bmatrix} 3 & 2 \\ 1 & 6 \end{bmatrix}, B^{-1}b = \begin{bmatrix} 12 \\ 12 \end{bmatrix}, \bar{c} = c_{BV}B^{-1}N - c_{NBV} = [-5 \quad -4], c_{BV}B^{-1}b = 0.$$

- Reduced cost $\bar{c}$ is NOT nonnegative. Current BV is not proven to be optimal.

- Recall the the following matrices for $BV = \{3,4\}$:

$$B^{-1}N = \begin{bmatrix} 3 & 2 \\ 1 & 6 \end{bmatrix}, B^{-1}b = \begin{bmatrix} 12 \\ 12 \end{bmatrix}, \bar{c} = c_{BV}B^{-1}N - c_{NBV} = [-5 \quad -4], c_{BV}B^{-1}b = 0.$$

- We choose x_1 as the **entering variable** (most negative reduced cost).
- We compute the column of the entering variable:

$$B^{-1}A^j = \begin{bmatrix} 3 \\ 1 \end{bmatrix}.$$

- We choose x_3 as the **leaving variable** (minimum ratio test).
- We update $BV = \{3,4\} \cup \{1\} \setminus \{3\} = \{1,4\}$.

- Recall the matrix form:

$$A = \begin{bmatrix} 3 & 2 & 1 & 0 \\ 1 & 6 & 0 & 1 \end{bmatrix}, c = [5 \quad 4 \quad 0 \quad 0], b = \begin{bmatrix} 12 \\ 12 \end{bmatrix}.$$

- We compute the following matrices for $BV = \{1, 4\}$:

$$B = \begin{bmatrix} 3 & 0 \\ 1 & 1 \end{bmatrix}, N = \begin{bmatrix} 2 & 1 \\ 6 & 0 \end{bmatrix}, c_{BV} = [5 \quad 0], c_{NBV} = [4 \quad 0], B^{-1} = \begin{bmatrix} \frac{1}{3} & 0 \\ -\frac{1}{3} & 1 \end{bmatrix},$$

$$B^{-1}N = \begin{bmatrix} \frac{2}{3} & \frac{1}{3} \\ \frac{16}{3} & -\frac{1}{3} \end{bmatrix}, B^{-1}b = \begin{bmatrix} 4 \\ 8 \end{bmatrix}, \bar{c} = c_{BV}B^{-1}N - c_{NBV} = \left[-\frac{2}{3} \quad \frac{5}{3}\right], c_{BV}B^{-1}b = 20.$$

- Reduced cost $\bar{c}$ is NOT nonnegative. Current BV is not proven to be optimal.

- Recall the the following matrices for $BV = \{1,4\}$:

$$B^{-1}N = \begin{bmatrix} \frac{2}{3} & \frac{1}{3} \\ \frac{16}{3} & -\frac{1}{3} \end{bmatrix}, B^{-1}b = \begin{bmatrix} 4 \\ 8 \end{bmatrix}, \bar{c} = c_{BV}B^{-1}N - c_{NBV} = \begin{bmatrix} -\frac{2}{3} & \frac{5}{3} \end{bmatrix}, c_{BV}B^{-1}b = 20.$$

- We choose x_2 as the **entering variable** (most negative reduced cost).
- We compute the column of the entering variable:

$$B^{-1}A^j = \begin{bmatrix} \frac{2}{3} \\ \frac{16}{3} \end{bmatrix}.$$

- We choose x_4 as the **leaving variable** (minimum ratio test).
- We update $BV = \{1,4\} \cup \{2\} \setminus \{4\} = \{1,2\}$.

- Recall the matrix form:

$$A = \begin{bmatrix} 3 & 2 & 1 & 0 \\ 1 & 6 & 0 & 1 \end{bmatrix}, c = [5 \quad 4 \quad 0 \quad 0], b = \begin{bmatrix} 12 \\ 12 \end{bmatrix}.$$

- We compute the following matrices for $BV = \{1,2\}$:

$$B = \begin{bmatrix} 3 & 2 \\ 1 & 6 \end{bmatrix}, N = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}, c_{BV} = [5 \quad 4], c_{NBV} = [0 \quad 0], B^{-1} = \begin{bmatrix} \frac{3}{8} & -\frac{1}{8} \\ -\frac{1}{16} & \frac{3}{16} \end{bmatrix},$$

$$B^{-1}N = \begin{bmatrix} \frac{3}{8} & -\frac{1}{8} \\ -\frac{1}{16} & \frac{3}{16} \end{bmatrix}, B^{-1}b = \begin{bmatrix} 3 \\ \frac{3}{2} \end{bmatrix}, \bar{c} = c_{BV}B^{-1}N - c_{NBV} = \left[\frac{13}{8} \quad \frac{1}{8}\right], c_{BV}B^{-1}b = 21.$$

- Reduced cost $\bar{c}$ is nonnegative. Hence, current BV is OPTIMAL.
- Optimal **primal** solution: $x_1 = 3, x_2 = 3/2$.
- Optimal **dual** solution: $y_1 = 13/8, y_2 = 1/8$.

- **Exercise:** Solve the following problem using the revised simplex method. Double check the iterations using the tableau implementation in Python.

$$\begin{aligned} \max x_0 &= 2x_1 + 3x_2 \\ \text{s.t.} \quad &2x_1 + x_2 \leq 10 \\ &x_1 + x_2 \leq 6 \\ &-x_1 + x_2 \leq 4 \\ &x_1, x_2 \geq 0. \end{aligned}$$

- As presented, the revised simplex method is virtually the same as the simplex method.
- However, commercial solvers such as Gurobi always use the **revised** simplex method, for two main reasons:
- The basis-matrix inversion can be implemented efficiently using the previous basis inverse (e.g. the product form of the inverse).
- Elementary row operations in the plain simplex method accumulate round-off errors; the revised method has ways to control these (e.g. occasionally re-inverting the basis matrix from scratch).
- The implementation details of the revised simplex method are advanced topics beyond the scope of this course.

- At each iteration of simplex, we can update B^{-1} using the following:
- Let k be the index of entering variable and r be that of the leaving variable. $B_{new}^{-1} = EB_{old}^{-1}$,

where matrix E is an identity matrix except that its r^{th} column is replaced by the vector

$$\eta = \begin{bmatrix} \eta_1 \\ \eta_2 \\ \vdots \\ \eta_m \end{bmatrix}, \text{ where}$$

$$\eta_i = \begin{cases} \frac{-a'_{ik}}{a'_{rk}} & \text{if } i \neq r \\ \frac{1}{a'_{rk}} & \text{if } i = r \end{cases}$$

$$\begin{array}{ll}\max & 3x_1 + 5x_2 \\ \text{s.t.} & x_1 \leq 4 \\ & 2x_2 \leq 12 \\ & 3x_1 + 2x_2 \leq 18 \\ & x_1, x_2 \geq 0\end{array}$$

$$\begin{array}{ll}\max & z - 3x_1 - 5x_2 + 0x_3 + 0x_4 + 0x_5 = 0 \\ \text{s.t.} & x_1 + x_3 = 4 \\ & 2x_2 + x_4 = 12 \\ & 3x_1 + 2x_2 + x_5 = 18 \\ & x_1, x_2, x_3, x_4, x_5 \geq 0\end{array}$$

$$\max \quad 3x_1 + 5x_2$$

$$\text{s.t.} \quad x_1 \leq 4$$

$$2x_2 \leq 12$$

$$3x_1 + 2x_2 \leq 18$$

$$x_1, x_2 \geq 0$$

z	x_1	x_2	x_3	x_4	x_5	RHS
1	-3	-5	0	0	0	0
0	1	0	1	0	0	4
0	0	2	0	1	0	12
0	3	2	0	0	1	18

$$\max \quad 3x_1 + 5x_2$$

$$\text{s.t.} \quad x_1 \leq 4$$

$$2x_2 \leq 12$$

$$3x_1 + 2x_2 \leq 18$$

$$x_1, x_2 \geq 0$$

z	x_1	x_2	x_3	x_4	x_5	RHS
1	-3	0	0	$\frac{5}{2}$	0	30
0	1	0	1	0	0	4
0	0	1	0	$\frac{1}{2}$	0	6
0	3	0	0	-1	1	6

$$\begin{array}{ll}
 \max & 3x_1 + 5x_2 \\
 \text{s.t.} & x_1 \leq 4 \\
 & 2x_2 \leq 12 \\
 & 3x_1 + 2x_2 \leq 18 \\
 & x_1, x_2 \geq 0
 \end{array}$$

z	x_1	x_2	x_3	x_4	x_5	RHS
1	0	0	0	$\frac{3}{2}$	1	36
0	0	0	1	$\frac{1}{3}$	$-\frac{1}{3}$	2
0	0	1	0	$\frac{1}{2}$	0	6
0	1	0	0	$-\frac{1}{3}$	$\frac{1}{3}$	2