

Modelling Nonlinear optics with the Bloch-Messiah reduction

Oliver Thomas, Dara McCutcheon, Will McCutcheon

Quantum Engineering CDT
University of Bristol

August 23, 2018

Overview

Modelling
Nonlinear
optics with
the
Bloch-Messiah
reduction

Oliver
Thomas, Dara
McCutcheon,
Will
McCutcheon

- What is nonlinear optics?
- Why do we care about it?
- Gaussian optics
- What I have been doing
- Outlook

Motivation quantum nonlinear optics

Modelling
Nonlinear
optics with
the
Bloch-Messiah
reduction

Oliver
Thomas, Dara
McCutcheon,
Will
McCutcheon

The good

Spontaneous Parametric processes, SPDC, SFWM

- Heralded single photon sources
- Entangled photon pair generation (polarisation, spatial)

Kerr processes

- Self-Phase modulation (SPM), generating Bannana states (CV)
- Cross-Phase modulation (XPM) for sensing

The bad

Spontaneous parametric processes

- Generating more than two photons \rightarrow bad for quantum computing
- Understanding filtering

All Kerr nonlinear processes

- SPM \rightarrow Spectral broadening
- XPM \rightarrow Unwanted phase shifts on single photons due to propagation of the pump

What do we mean by nonlinear optics?

Modelling
Nonlinear
optics with
the
Bloch-Messiah
reduction

Oliver
Thomas, Dara
McCutcheon,
Will
McCutcheon

- Roughly processes that conserve energy but do not conserve photon number.

$$\vec{P} = \chi^{(1)} \vec{E}_1 + \chi^{(2)} \vec{E}_1 \vec{E}_2 + \chi^{(3)} \vec{E}_1 \vec{E}_2 \vec{E}_3 + \dots \quad (1)$$

Here we are going to talk about squeezing, i.e SPDC or SFWM, Hamiltonians are then of the form,

$$\hat{H} = A \hat{a}_S^\dagger \hat{a}_I^\dagger \hat{a}_P + h.c. \quad (2)$$

$$\hat{H} = A \hat{a}_S^\dagger \hat{a}_I^\dagger \hat{a}_P \hat{a}_P + h.c. \quad (3)$$

Gaussian Optics

Modelling
Nonlinear
optics with
the
Bloch-Messiah
reduction

Oliver
Thomas, Dara
McCutcheon,
Will
McCutcheon

- Using the undepleted pump approximation we can write the Hamiltonians as terms which are at most quadratic in creation and annihilation operators.

$$\hat{U} = \exp \left[-\frac{i}{\hbar} \left(\underset{\text{Power}}{P} \int d\omega_1 \int d\omega_2 \underset{\text{JSA}}{f(\omega_1, \omega_2)} \underset{\text{Signal \& Idler}}{\hat{a}_s^\dagger(\omega_1) \hat{a}_i^\dagger(\omega_2)} + h.c. \right) \right] \quad (4)$$

- These are Gaussian transforms, they take Gaussian states to Gaussian states ¹

$$\begin{bmatrix} \vec{b} \\ \vec{b}^\dagger \end{bmatrix} = \mathbf{M} \begin{bmatrix} \vec{a} \\ \vec{a}^\dagger \end{bmatrix} \quad (5)$$

¹These are linear symplectic transforms which conveniently can be written as a matrix

Types of Gaussian transformations

Modelling
Nonlinear
optics with
the
Bloch-Messiah
reduction

Oliver
Thomas, Dara
McCutcheon,
Will
McCutcheon




Figure: Two source HOM dip




Figure: Type-1 Fusion gate

Schmidt decomposition

We can re-write the Hamiltonian using a Schmidt-decomposition as,

$$P'F(\omega_1, \omega_2) = \sum_k r_k \psi_k(\omega_1) \phi_k(\omega_2) \quad (6)$$

Where r_k is the Schmidt number, ψ & ϕ are unitaries.

To solve this numerically we discretize the function and the Schmidt-decomposition is then the Singular value decomposition (SVD) of the JSA (F).

$$P'\mathbf{F}_{(\omega_1, \omega_2)} = \sum_k r_k \mathbf{U}_{(\omega_1, k)} \mathbf{V}_{(k, \omega_2)}^\dagger \quad (7)$$

- with $\psi_k(\omega_1)$ is the k -th row and ω_1 -th column of $\mathbf{U}_{(\omega_1, k)}$,
- with $\phi_k(\omega_2)$ is the ω_2 -th row and k -th column of $\mathbf{V}_{(k, \omega_2)}^\dagger$

Joint Spectral Amplitudes (JSAs)

Modelling
Nonlinear
optics with
the
Bloch-Messiah
reduction



Oliver
Thomas, Dara
McCutcheon,
Will
McCutcheon

¹Moving to the rotating frame...

Seperable JSAs Schmidt modes

Modelling
Nonlinear
optics with
the
Bloch-Messiah
reduction

Oliver
Thomas, Dara
McCutcheon,
Will
McCutcheon



(a) Signal (red) and Idler (blue)

Non-separable JSAs



Modelling
Nonlinear
optics with
the
Bloch-Messiah
reduction

Oliver
Thomas, Dara
McCutcheon,
Will
McCutcheon

Non-separable JSAs Schimdt modes

Modelling
Nonlinear
optics with
the
Bloch-Messiah
reduction

Oliver
Thomas, Dara
McCutcheon,
Will
McCutcheon



(a) Signal (red) and Idler (blue)

Reducing the size of the state-space

Modelling
Nonlinear
optics with
the
Bloch-Messiah
reduction

Oliver
Thomas, Dara
McCutcheon,
Will
McCutcheon

- The Schmidt decomposition lets us represent the system in a finite number of broadband modes ($\psi_k(\omega_1), \phi_k(\omega_2)$)
-

Correlations in a HOM dip

Modelling
Nonlinear
optics with
the
Bloch-Messiah
reduction

Oliver
Thomas, Dara
McCutcheon,
Will
McCutcheon





Figure: Two source HOM dip

Two squeezers JSA

Modelling
Nonlinear
optics with
the
Bloch-Messiah
reduction

Oliver
Thomas, Dara
McCutcheon,
Will
McCutcheon



G(4) correlation function

Modelling
Nonlinear
optics with
the
Bloch-Messiah
reduction

Oliver
Thomas, Dara
McCutcheon,
Will
McCutcheon

$$G^{(4)} = \frac{\langle \hat{a}_1^\dagger \hat{a}_2^\dagger \hat{a}_3^\dagger \hat{a}_4^\dagger \hat{a}_1 \hat{a}_2 \hat{a}_3 \hat{a}_4 \rangle}{\langle \hat{a}_1^\dagger \hat{a}_1 \rangle \langle \hat{a}_2^\dagger \hat{a}_2 \rangle \langle \hat{a}_3^\dagger \hat{a}_3 \rangle \langle \hat{a}_4^\dagger \hat{a}_4 \rangle} \quad (8)$$

Where,

$$a_i = \sum_j a_i(\omega_j) \quad (9)$$

Meaning we sum over all of the spectral modes of the spatial modes (1,2,3,4) separately. We end up with,

$$G^{(4)} = 1 - \left(\frac{2 | \cosh(r) |^2}{| \cosh(r) |^2 + | \sinh(r) |^2} \sin(\theta) \cos(\theta) \right)^2 \quad (10)$$

$G(4)$ correlation function

Modelling
Nonlinear
optics with
the
Bloch-Messiah
reduction

Oliver

Thomas, Dara
McCutcheon,
Will
McCutcheon

Summary

Modelling
Nonlinear
optics with
the
Bloch-Messiah
reduction

Oliver
Thomas, Dara
McCutcheon,
Will
McCutcheon