STAT 590 Survival Analysis -Final Project

A clinical Trial in the Treatment of Carcinoma of the Oropharynx

Andres Gonzalez Yushan Zhao

May 8, 2023




California State University - Long Beach
Departement of Mathematics and Statistics
1250 Bellflower Boulevard
Long Beach, California 90840

Table of contents


1	Exploratory Data Analysis		
	1.1	Age	1
	1.2	Gender	4
	1.3	Time and Status	6
	1.4	Kaplan Meier	6

1.1 Age

Age Distribution




Age Distribution by Grade Differentiation




Source: Clinical Trial in the Treatment of Carcinoma of the Oropharynx Dataset

Age Distribution by Condition




Age Distribution by Site




Source: Clinical Trial in the Treatment of Carcinoma of the Oropharynx Dataset

Age Distribution by T_Stage




Age Distribution by T_Stage



Source: Clinical Trial in the Treatment of Carcinoma of the Oropharynx Dataset

1.2 Gender



Condition

195 Participants



Source: Clinical Trial in the Treatment of Carcinoma of the Oropharynx Dataset


Stacked Distribution of Grades by Gender

195 Participants



1.3 Time and Status

Scatter Plot of Time and Age by Status



Source: Clinical Trial in the Treatment of Carcinoma of the Oropharynx Dataset

1.4 Kaplan Meier


```
pharynx <- pharynx %>%
  mutate(
    Status = as.numeric(Status),
    TX = as_factor(Tx),
    SEX = as_factor(Sex),
    T_STAGE = as_factor(T_Stage)
    )

pharynx_surv <- Surv(pharynx$Time, pharynx$Status)

# Fit a Kaplan-Meier curve, stratified by treatment
pharynx_km_tx <- survfit2(pharynx_surv ~ TX, data = pharynx)

# Plot the Kaplan-Meier curve, stratified by treatment
ggsurvfit(pharynx_km_tx, linewidth = 1) +
    add_confidence_interval() +
    add_censor_mark() +</pre>
```

Kaplan-Meier Survival Curve by Treatment



```
pharynx_surv <- Surv(pharynx$Time, pharynx$Status)

# Fit a Kaplan-Meier curve, stratified by treatment
pharynx_km_tx <- survfit2(pharynx_surv ~ Sex, data = pharynx)

# Plot the Kaplan-Meier curve, stratified by treatment</pre>
```

```
ggsurvfit(pharynx_km_tx, linewidth = 1) +
 add_confidence_interval() +
 add_censor_mark() +
 add_pvalue(location = "annotation") +
 labs(y = "Percentage Survival",
       title = "Kaplan-Meier Survival Curve by Treatment") +
 scale_y_continuous(label = scales::percent,
                     breaks = seq(0,1, by = 0.2),
                     expand = c(0.015,0)) +
 scale_color_manual(values = c("#B83377", "#2EA9D8")) +
 scale_fill_manual(values = c("#B83377", "#2EA9D8")) +
 theme(
   text = element_text(family = "montse"),
   title = element_text(face = "bold"),
   plot.title.position = "plot",
   legend.position = "top"
```

Kaplan-Meier Survival Curve by Treatment

