

# ME 599/699 Robot Modeling & Control

## Fall 2021

### Homogenous Coordinate Transformations


Hasan A. Poonawala

Department of Mechanical Engineering  
University of Kentucky

Email: [hasan.poonawala@uky.edu](mailto:hasan.poonawala@uky.edu)

Web: <https://www.engr.uky.edu/~hap>

# Same Vector Space, Different Bases



A basis and an origin together form a **coordinate frame** or **reference frame**.

# Change Of Basis

The coordinate  $(1, 0)$  will produce different points under different bases.

# Change Of Basis

The coordinate  $(1, 0)$  will produce different points under different bases.

When we use a different basis, the coordinates assigned to a point must change, in order to correctly regenerate that point using the new basis.

# Change Of Basis

The coordinate  $(1, 0)$  will produce different points under different bases.

When we use a different basis, the coordinates assigned to a point must change, in order to correctly regenerate that point using the new basis.

Let  $A, B, C, \dots$  be different coordinate frames.

A point  $p$  then has coordinates  $p^A, p^B, p^C \dots$  corresponding to each basis.

# Change Of Vector Space Basis

Given  $p^A$ , what is  $p^B$ , or  $p^C$ ?

Answer:

$$p^B = \left(T_B^A\right)^{-1} p^A,$$

where

$$T_B^A = \begin{bmatrix} (e_B^1)^A & (e_B^2)^A & \cdots & (e_B^n)^A \end{bmatrix},$$

and  $(e_B^i)^A$  is the coordinates in frame  $A$  of the  $i^{\text{th}}$  basis vector of frame  $B$ .

## Example


**Problem:** Find  $p^B$  if  $p^A = (1, 1)$ .

**Solution:** From the diagram,

$$\begin{aligned}e_B^1 &= e_A^1 \\e_B^2 &= e_A^1 + e_A^2 \\ \Rightarrow T_B^A &= \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}\end{aligned}$$

Apply the formula:

$$\begin{aligned}p^B &= \left(T_B^A\right)^{-1} p^A \\ &= T_A^B p^A = \begin{bmatrix} 0 \\ 1 \end{bmatrix}\end{aligned}$$



The columns of  $T_B^A$  tell us how to draw the basis of  $B$  in  $A$

# Change Of Vector Space Basis

## Full derivation:

The vector  $e_B^i$  has coordinates  $(e_B^i)^A = (T_{1i}, T_{2i}, \dots, T_{ni})$  in frame  $A$ .



# Change Of Vector Space Basis

## Full derivation:

The vector  $e_B^i$  has coordinates  $(e_B^i)^A = (T_{1i}, T_{2i}, \dots, T_{ni})$  in frame  $A$ .

Let the coordinates of  $p$  in frame  $A$  be  $p^A = (\alpha_1^A, \alpha_2^A, \dots, \alpha_n^A)$ , so that the point  $p$  can be expressed as

$$p \iff \sum_j^n \alpha_j^A e_A^j$$

# Change Of Vector Space Basis

## Full derivation:

The vector  $e_B^i$  has coordinates  $(e_B^i)^A = (T_{1i}, T_{2i}, \dots, T_{ni})$  in frame  $A$ .

Let the coordinates of  $p$  in frame  $A$  be  $p^A = (\alpha_1^A, \alpha_2^A, \dots, \alpha_n^A)$ , so that the point  $p$  can be expressed as

$$p \iff \sum_j^n \alpha_j^A e_A^j$$

Note that  $p$  is an abstract point equivalent to the coordinate-given combination of the basis  $\{e_A^1, e_A^2, \dots, e_A^n\}$ .

# Change Of Vector Space Basis

## Full derivation:

The vector  $e_B^i$  has coordinates  $(e_B^i)^A = (T_{1i}, T_{2i}, \dots, T_{ni})$  in frame  $A$ .

Let the coordinates of  $p$  in frame  $A$  be  $p^A = (\alpha_1^A, \alpha_2^A, \dots, \alpha_n^A)$ , so that the point  $p$  can be expressed as

$$p \iff \sum_j^n \alpha_j^A e_A^j$$

Note that  $p$  is an abstract point equivalent to the coordinate-given combination of the basis  $\{e_A^1, e_A^2, \dots, e_A^n\}$ .

Similarly, if  $p^B = (\beta_1^B, \beta_2^B, \dots, \beta_n^B)$ , then

$$p \iff \sum_i^n \beta_i^B e_B^i$$

# Change Of Vector Space Basis

So, we can write

$$e_B^i = \sum_j^n T_{ji} e_A^j; \quad p \iff \sum_i^n \beta_i^B e_B^i; \quad p \iff \sum_j^n \alpha_j^A e_A^j \quad (1)$$

# Change Of Vector Space Basis

So, we can write

$$e_B^i = \sum_j^n T_{ji} e_A^j; \quad p \iff \sum_i^n \beta_i^B e_B^i; \quad p \iff \sum_j^n \alpha_j^A e_A^j \quad (1)$$

Combining the first and second equation in (1), we get

$$\begin{aligned} p &\iff \sum_i^n \beta_i^B e_B^i = \sum_i^n \beta_i^B \left( \sum_j^n T_{ji} e_A^j \right) \\ &\iff \sum_j^n \left( \sum_i^n \left( \beta_i^B T_{ji} \right) \right) e_A^j \end{aligned} \quad (2)$$

Comparing (2) to the third equation in (1), we get

$$\alpha_j^A = \sum_i^n \left( \beta_i^B T_{ji} \right).$$

# Change Of Vector Space Basis

The expression

$$\alpha_j^A = \sum_i^n \left( \beta_i^B T_{ji} \right)$$

represents a linear transformation of the coordinates of a point.

# Change Of Vector Space Basis

The expression

$$\alpha_j^A = \sum_i^n \left( \beta_i^B T_{ji} \right)$$

represents a linear transformation of the coordinates of a point.

$$p \iff \begin{bmatrix} e_A^1 & e_A^2 & \cdots & e_A^n \end{bmatrix} \begin{bmatrix} \alpha_1^A \\ \alpha_2^A \\ \vdots \\ \alpha_n^A \end{bmatrix} = \begin{bmatrix} e_B^1 & e_B^2 & \cdots & e_B^n \end{bmatrix} \begin{bmatrix} \beta_1^B \\ \beta_2^B \\ \vdots \\ \beta_n^B \end{bmatrix}$$

# Change Of Vector Space Basis

The expression

$$\alpha_j^A = \sum_i^n \left( \beta_i^B T_{ji} \right)$$

represents a linear transformation of the coordinates of a point.

$$p \iff \begin{bmatrix} e_A^1 & e_A^2 & \cdots & e_A^n \end{bmatrix} \begin{bmatrix} \alpha_1^A \\ \alpha_2^A \\ \vdots \\ \alpha_n^A \end{bmatrix} = \begin{bmatrix} e_B^1 & e_B^2 & \cdots & e_B^n \end{bmatrix} \begin{bmatrix} \beta_1^B \\ \beta_2^B \\ \vdots \\ \beta_n^B \end{bmatrix}$$

The coordinates of  $e_B^i$  in frame  $A$  give:

$$e_B^1 = \begin{bmatrix} e_A^1 & \cdots & e_A^n \end{bmatrix} \begin{bmatrix} T_{11} \\ \vdots \\ T_{n1} \end{bmatrix}, e_B^2 = \begin{bmatrix} e_A^1 & \cdots & e_A^n \end{bmatrix} \begin{bmatrix} T_{12} \\ \vdots \\ T_{n2} \end{bmatrix}, \dots$$



# Change Of Vector Space Basis

We can collect these expressions for point  $e_B^i$  as

$$\begin{bmatrix} e_B^1 & e_B^2 & \cdots & e_B^n \end{bmatrix} = \begin{bmatrix} e_A^1 & e_A^2 & \cdots & e_A^n \end{bmatrix} \begin{bmatrix} T_{11} & T_{12} & \cdots & T_{1n} \\ T_{21} & T_{22} & \cdots & T_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ T_{n1} & T_{n2} & \cdots & T_{nn} \end{bmatrix},$$

So that

$$\begin{bmatrix} e_B^1 & e_B^2 & \cdots & e_B^n \end{bmatrix} \begin{bmatrix} \beta_1^B \\ \beta_2^B \\ \vdots \\ \beta_n^B \end{bmatrix} = \begin{bmatrix} e_A^1 & e_A^2 & \cdots & e_A^n \end{bmatrix} \begin{bmatrix} T_{11} & T_{12} & \cdots & T_{1n} \\ T_{21} & T_{22} & \cdots & T_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ T_{n1} & T_{n2} & \cdots & T_{nn} \end{bmatrix} \begin{bmatrix} \beta_1^B \\ \beta_2^B \\ \vdots \\ \beta_n^B \end{bmatrix}$$

# Change Of Vector Space Basis

Since

$$p = [e_A^1 \quad e_A^2 \quad \cdots \quad e_A^n] \begin{bmatrix} \alpha_1^A \\ \alpha_2^A \\ \vdots \\ \alpha_n^A \end{bmatrix},$$

we find that transforming coordinates is a linear operation represented by matrix operations:


$$\begin{bmatrix} \alpha_1^A \\ \alpha_2^A \\ \vdots \\ \alpha_n^A \end{bmatrix} = \begin{bmatrix} T_{11} & T_{12} & \cdots & T_{1n} \\ T_{21} & T_{22} & \cdots & T_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ T_{n1} & T_{n2} & \cdots & T_{nn} \end{bmatrix} \begin{bmatrix} \beta_1^B \\ \beta_2^B \\ \vdots \\ \beta_n^B \end{bmatrix}$$

More compactly:  $p^B = (T_B^A)^{-1} p^A$ , where [► to example](#)

$$T_B^A = \left[ (e_B^1)^A \quad (e_B^2)^A \quad \cdots \quad (e_B^n)^A \right].$$


# Change Of Origin

Suppose points  $p$ ,  $q$  have coordinates  $p^A$ ,  $q^A$  in a frame  $A$ . Consider frame  $B$  whose origin is at  $p$ , with the same basis elements for its vector space as the frame  $A$ . What is  $q^B$ ?



## Change Of Origin


Suppose points  $p$ ,  $q$  have coordinates  $p^A$ ,  $q^A$  in a frame  $A$ . Consider frame  $B$  whose origin is at  $p$ , with the same basis elements for its vector space as the frame  $A$ . What is  $q^B$ ?



The coordinates of  $q$  in frame  $B$  is the same as coordinates of the vector  $v = q - p$  in the basis common to both frame  $A$  and  $B$ .

## Change Of Origin

Suppose points  $p$ ,  $q$  have coordinates  $p^A$ ,  $q^A$  in a frame  $A$ . Consider frame  $B$  whose origin is at  $p$ , with the same basis elements for its vector space as the frame  $A$ . What is  $q^B$ ?




The coordinates of  $q$  in frame  $B$  is the same as coordinates of the vector  $v = q - p$  in the basis common to both frame  $A$  and  $B$ .

Precisely because vectors are free, the coordinates of  $v$  in frame  $B$  will be the same as that in frame  $A$ .

# Change Of Origin

Suppose points  $p$ ,  $q$  have coordinates  $p^A$ ,  $q^A$  in a frame  $A$ . Consider frame  $B$  whose origin is at  $p$ , with the same basis elements for its vector space as the frame  $A$ . What is  $q^B$ ?



The coordinates of  $q$  in frame  $B$  is the same as coordinates of the vector  $v = q - p$  in the basis common to both frame  $A$  and  $B$ .

Precisely because vectors are free, the coordinates of  $v$  in frame  $B$  will be the same as that in frame  $A$ . So,  $q^B = q^A - p^A$ .

In general,  $q^B = q^A - (\text{coordinates of origin of } B \text{ in } A)$

# Change Of Frames

Combining previous discussions, we get that to map coordinates from one frame to another we :

1. express the coordinates of the basis vectors of one frame in the other (through, say, matrix  $T_B^A$ ),
2. express the coordinates of the origin of one frame in another (through, say coordinate vector  $o_B^A$ ),
3. use the map

$$p^B = \left(T_B^A\right)^{-1} (p^A - o_B^A)$$

# Checkpoint

- ▶ We relate points by picking an origin and using a vector space



# Checkpoint

- ▶ We relate points by picking an origin and using a vector space
- ▶ Coordinates of vectors, given a basis, become coordinates of points

# Checkpoint

- ▶ We relate points by picking an origin and using a vector space
- ▶ Coordinates of vectors, given a basis, become coordinates of points
- ▶ These coordinates correspond to a frame: origin + basis

# Checkpoint

- ▶ We relate points by picking an origin and using a vector space
- ▶ Coordinates of vectors, given a basis, become coordinates of points
- ▶ These coordinates correspond to a frame: origin + basis
- ▶ All coordinates are  $n$ -tuples, we can't say anything about the basis from the coordinates.

# Checkpoint

- ▶ We relate points by picking an origin and using a vector space
- ▶ Coordinates of vectors, given a basis, become coordinates of points
- ▶ These coordinates correspond to a frame: origin + basis
- ▶ All coordinates are  $n$ -tuples, we can't say anything about the basis from the coordinates.
- ▶ If we know the bases and origins, we can transform coordinates from one frame to another.

# Checkpoint

- ▶ We relate points by picking an origin and using a vector space
- ▶ Coordinates of vectors, given a basis, become coordinates of points
- ▶ These coordinates correspond to a frame: origin + basis
- ▶ All coordinates are  $n$ -tuples, we can't say anything about the basis from the coordinates.
- ▶ If we know the bases and origins, we can transform coordinates from one frame to another.

# Checkpoint

- ▶ We relate points by picking an origin and using a vector space
- ▶ Coordinates of vectors, given a basis, become coordinates of points
- ▶ These coordinates correspond to a frame: origin + basis
- ▶ All coordinates are  $n$ -tuples, we can't say anything about the basis from the coordinates.
- ▶ If we know the bases and origins, we can transform coordinates from one frame to another.

If all bases for the plane give us two numbers, what's special about a basis where the two elements are at 90 degrees , and have the same 'length'?

# Norms and Distances

Let's reconsider our earlier example:

$$T_B^A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}; \quad q^A = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \implies q^B = \left(T_B^A\right)^{-1} q^A = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

# Norms and Distances

Let's reconsider our earlier example:

$$T_B^A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}; \quad q^A = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \implies q^B = \left(T_B^A\right)^{-1} q^A = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

REMEMBER: We're talking about the same two points in Euclidean space.



# Norms and Distances

Let's reconsider our earlier example:

$$T_B^A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}; \quad q^A = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \implies q^B = \left(T_B^A\right)^{-1} q^A = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

REMEMBER: We're talking about the same two points in Euclidean space.

$$\|q^A\|_A = \sqrt{2}. \quad \|q^B\|_B = 1.$$

# Norms and Distances

Let's reconsider our earlier example:

$$T_B^A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}; \quad q^A = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \implies q^B = \left(T_B^A\right)^{-1} q^A = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

REMEMBER: We're talking about the same two points in Euclidean space.

$\|q^A\|_A = \sqrt{2}$ .  $\|q^B\|_B = 1$ . What gives?

# Norms and Distances

Let's reconsider our earlier example:

$$T_B^A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}; \quad q^A = \begin{bmatrix} 1 \\ 1 \end{bmatrix} \implies q^B = (T_B^A)^{-1} q^A = \begin{bmatrix} 0 \\ 1 \end{bmatrix}.$$

**REMEMBER:** We're talking about the same two points in Euclidean space.

$\|q^A\|_A = \sqrt{2}$ .  $\|q^B\|_B = 1$ . What gives?

Note that  $\|q^B\|_B = \|(T_B^A)^{-1} q^A\|_A$ .

**Q:** What kinds of matrices preserve the norms of the vectors they act upon?

# Special Orthogonal Group in Three Dimensions

if  $T_B^A \in SO(3)$ , then we'd have norm-preservation.

## Definition ( $SO(3)$ )

The Special Orthogonal Group  $SO(3)$  is the set of matrices  $R \in \mathbb{R}^{3 \times 3}$  such that

$$R^T R = Id, \text{ and } \det R = 1$$

.

$SO(3)$  is known as the orientation group **and** the rotation group.

## Example

Problem: Find  $p^B$  if  $p^A = (1, 1)$ .

Solution: From the diagram,


$$e_B^1 = e_A^1$$

$$e_B^2 = e_A^1 + e_A^2$$

$$\Rightarrow T_B^A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$

Apply the formula:

$$\begin{aligned} p^B &= (T_B^A)^{-1} p^A \\ &= T_A^B p^A = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \end{aligned}$$



The columns of  $T_B^A$  tell us how to draw the basis of  $B$  in  $A$

[▶ to defn](#)

## Example

Problem: Find  $p^B$  if  $p^A = (1, 1)$ .

Solution: From the diagram,


$$e_B^1 = e_A^1$$

$$e_B^2 = e_A^1 + e_A^2$$

$$\Rightarrow T_B^A = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}$$

Apply the formula:

$$\begin{aligned} p^B &= (T_B^A)^{-1} p^A \\ &= T_A^B p^A = \begin{bmatrix} 0 \\ 1 \end{bmatrix} \end{aligned}$$



Not norm-preserving.

$$(T_A^B)^T T_A^B = \begin{bmatrix} 1 & -1 \\ -1 & 2 \end{bmatrix}$$

## Example

Problem: Find  $p^B$  if  $p^A = (1, 1)$ .

Solution: From the diagram,

$$e_B^1 = \frac{1}{\sqrt{2}} e_A^1 - \frac{1}{\sqrt{2}} e_A^2$$


$$e_B^2 = e_A^1 + e_A^2$$

$$\Rightarrow T_B^A = \begin{bmatrix} \frac{1}{\sqrt{2}} & 1 \\ -\frac{1}{\sqrt{2}} & 1 \end{bmatrix}$$

Apply the formula:

$$p^B = \left(T_B^A\right)^{-1} p^A$$

$$= T_A^B p^A = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$



The columns of  $T_B^A$  tell us how to draw the basis of  $B$  in  $A$

► to defn

## Example

Problem: Find  $p^B$  if  $p^A = (1, 1)$ .

Solution: From the diagram,

$$e_B^1 = \frac{1}{\sqrt{2}} e_A^1 - \frac{1}{\sqrt{2}} e_A^2$$


$$e_B^2 = e_A^1 + e_A^2$$

$$\Rightarrow T_B^A = \begin{bmatrix} \frac{1}{\sqrt{2}} & 1 \\ -\frac{1}{\sqrt{2}} & 1 \end{bmatrix}$$

Apply the formula:

$$p^B = (T_B^A)^{-1} p^A$$

$$= T_A^B p^A = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$



Not norm-preserving.

$$(T_A^B)^T T_A^B = \begin{bmatrix} 0.75 & -0.25 \\ -0.25 & 0.75 \end{bmatrix}$$



## Example

Problem: Find  $p^B$  if  $p^A = (1, 1)$ .

Solution: From the diagram,

$$e_B^1 = \frac{1}{\sqrt{2}} e_A^1 - \frac{1}{\sqrt{2}} e_A^2$$


$$e_B^2 = \frac{1}{\sqrt{2}} e_A^1 + \frac{1}{\sqrt{2}} e_A^2$$

$$\Rightarrow T_B^A = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix}$$

Apply the formula:

$$p^B = \left(T_B^A\right)^{-1} p^A$$

$$= T_A^B p^A = \begin{bmatrix} 0 \\ \sqrt{2} \end{bmatrix}$$



## Example

Problem: Find  $p^B$  if  $p^A = (1, 1)$ .

Solution: From the diagram,

$$e_B^1 = \frac{1}{\sqrt{2}} e_A^1 - \frac{1}{\sqrt{2}} e_A^2$$


$$e_B^2 = \frac{1}{\sqrt{2}} e_A^1 + \frac{1}{\sqrt{2}} e_A^2$$

$$\Rightarrow T_B^A = \begin{bmatrix} \frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \\ -\frac{1}{\sqrt{2}} & \frac{1}{\sqrt{2}} \end{bmatrix}$$

Apply the formula:

$$p^B = \left(T_B^A\right)^{-1} p^A$$

$$= T_A^B p^A = \begin{bmatrix} 0 \\ \sqrt{2} \end{bmatrix}$$



norm-preserving!

$$\left(T_A^B\right)^T T_A^B = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

# Orthonormal Vectors

We have seen that

$$T_B^A = \begin{bmatrix} (e_B^1)^A & (e_B^2)^A & \cdots & (e_B^n)^A \end{bmatrix}.$$

Therefore,

$$(T_B^A)^T T_B^A = \begin{bmatrix} \left((e_B^1)^A\right)^T \\ \left((e_B^2)^A\right)^T \\ \vdots \\ \left((e_B^n)^A\right)^T \end{bmatrix} \begin{bmatrix} (e_B^1)^A & (e_B^2)^A & \cdots & (e_B^n)^A \end{bmatrix}$$

# Orthonormal Vectors

$$\begin{aligned} (T_B^A)^T T_B^A &= \begin{bmatrix} ((e_B^1)^A)^T (e_B^1)^A & ((e_B^1)^A)^T (e_B^2)^A & \cdots & ((e_B^1)^A)^T (e_B^n)^A \\ ((e_B^2)^A)^T (e_B^1)^A & ((e_B^2)^A)^T (e_B^2)^A & \cdots & ((e_B^2)^A)^T (e_B^n)^A \\ \vdots & \vdots & \ddots & \vdots \\ ((e_B^n)^A)^T (e_B^1)^A & ((e_B^n)^A)^T (e_B^2)^A & \cdots & ((e_B^n)^A)^T (e_B^n)^A \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 & \cdots & 0 \\ 0 & 1 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 1 \end{bmatrix} \end{aligned}$$

Effectively, the coordinates of basis vectors of  $B$  in frame  $A$  are unit length and perpendicular to each other.

# Checkpoint

- ▶ Frames are origin+basis

# Checkpoint

- ▶ Frames are origin+basis
- ▶ Frames define vector coordinates for points in Euclidean space, relative to the frame

# Checkpoint

- ▶ Frames are origin+basis
- ▶ Frames define vector coordinates for points in Euclidean space, relative to the frame
- ▶ Can transform vector coordinates of a point in different frames using an affine map

# Checkpoint

- ▶ Frames are origin+basis
- ▶ Frames define vector coordinates for points in Euclidean space, relative to the frame
- ▶ Can transform vector coordinates of a point in different frames using an affine map
- ▶ To preserve distance, the linear part of the affine map must be in  $SO(3)$



# Checkpoint


- ▶ Frames are origin+basis
- ▶ Frames define vector coordinates for points in Euclidean space, relative to the frame
- ▶ Can transform vector coordinates of a point in different frames using an affine map
- ▶ To preserve distance, the linear part of the affine map must be in  $SO(3)$
- ▶  $T_B^A \in SO(3)$  when basis vectors are all unit length, mutually perpendicular.

# Checkpoint

- ▶ Frames are origin+basis
- ▶ Frames define vector coordinates for points in Euclidean space, relative to the frame
- ▶ Can transform vector coordinates of a point in different frames using an affine map
- ▶ To preserve distance, the linear part of the affine map must be in  $SO(3)$
- ▶  $T_B^A \in SO(3)$  when basis vectors are all unit length, mutually perpendicular.
- ▶ The coordinate transformation is then
$$p^B = (R_B^A)^{-1} (p^A - o_B^A)$$


▶ mobile robot

# Coordinate Transformation Vs Rigid Motion




Consider a robot with a center, a camera in 'front', and two wheels to the side.

# Coordinate Transformation Vs Rigid Motion




Whenever we move the robot, the distances between these points don't change.

# Coordinate Transformation Vs Rigid Motion




As the robot moves, we can take a snapshot of these points, and they each define a coordinate frame for Euclidean space.

# Coordinate Transformation Vs Rigid Motion



Q1: How would the robot compare observations of either purple point over time? A1: Coordinate transformations

# Coordinate Transformation Vs Rigid Motion



Q1: How would the robot compare observations of either purple point over time? A1: Coordinate transformations

Q2: How do we keep track of all the points on the robots?

A2: Coordinate transformations, **but reinterpreted.**

► rigid motion

# Coordinate Transformation Vs Rigid Motion

We know how to express all points in the robot's frame in any other frame: Use a distance-preserving coordinate transformation.



# Coordinate Transformation Vs Rigid Motion

We know how to express all points in the robot's frame in any other frame: Use a distance-preserving coordinate transformation.

This **transformation itself** becomes a representative for all points on the robot.

# Coordinate Transformation Vs Rigid Motion

We know how to express all points in the robot's frame in any other frame: Use a distance-preserving coordinate transformation.

This **transformation itself** becomes a representative for all points on the robot.

We have seen that

$$p^B = \left(T_B^A\right)^{-1} \left(p^A - o_B^A\right) \quad (3)$$

Let  $d = o_B^A$  and  $R = T_B^A$ .

# Coordinate Transformation Vs Rigid Motion

We know how to express all points in the robot's frame in any other frame: Use a distance-preserving coordinate transformation.

This **transformation itself** becomes a representative for all points on the robot.

We have seen that

$$p^B = \left(T_B^A\right)^{-1} \left(p^A - o_B^A\right) \quad (3)$$

Let  $d = o_B^A$  and  $R = T_B^A$ . From (3), we can derive

$$p^B = R^{-1} \left(p^A - d\right) \quad (4)$$

$$p^A = R p^B + d. \quad (5)$$

# Coordinate Transformation Vs Rigid Motion

We know how to express all points in the robot's frame in any other frame: Use a distance-preserving coordinate transformation.

This **transformation itself** becomes a representative for all points on the robot.

We have seen that

$$p^B = \left(T_B^A\right)^{-1} \left(p^A - o_B^A\right) \quad (3)$$

Let  $d = o_B^A$  and  $R = T_B^A$ . From (3), we can derive

$$p^B = R^{-1} \left(p^A - d\right) \quad (4)$$

$$p^A = R p^B + d. \quad (5)$$

# Rigid Body Pose

We refer to the pair  $(d, R)$  as the pose – relative to frame  $A$  – of the rigid body to which frame  $B$  is attached.

# Rigid Body Pose

We refer to the pair  $(d, R)$  as the pose – relative to frame  $A$  – of the rigid body to which frame  $B$  is attached.

Let's reinterpret the two affine transformations associated with  $(d, R)$ . Consider vector  $v$  in frame  $A$ :

$$u_1 = R^{-1}(v - d) \quad \text{(Change of Basis)} \quad (6)$$


$$u_2 = R v + d. \quad \text{(Rigid motion)} \quad (7)$$

# Rigid Body Pose


Let's reinterpret the two affine transformations associated with  $(d, R)$ . Consider vector  $v$  in frame  $A$ :

$$u_1 = R^{-1}(v - d) \quad (\text{Change of Basis}) \quad (6)$$

$$u_2 = R v + d. \quad (\text{Rigid motion}) \quad (7)$$




# Rigid Body Pose



- ▶ If we view  $u_1$  as coordinates in frame  $B$ , we've changed coordinates of  $v$  from world to body frame.
- ▶ If we view  $u_2$  as coordinates in frame  $A$ , we've moved the point  $o_A \oplus v$  relative to frame  $A$ .




# Rigid Body Pose



The pair  $(d, R) \in \mathbb{R}^3 \times \text{SO}(3)$  tells us how to move points in frame A to achieve the same coordinates in frame B.


# Rigid Body Pose



The pair  $(d, R) \in \mathbb{R}^3 \times \text{SO}(3)$  tells us how to move points in frame  $A$  to achieve the same coordinates in frame  $B$ .

$d$  is a vector, the coordinates of origin of frame  $B$ , and  $R$  is a matrix containing the coordinates of axes of  $B$ , both relative to  $A$

# Rigid Body Pose




The pair  $(d, R) \in \mathbb{R}^3 \times \text{SO}(3)$  tells us how to move points in frame  $A$  to achieve the same coordinates in frame  $B$ .

$d$  is a vector, the coordinates of origin of frame  $B$ , and  $R$  is a matrix containing the coordinates of axes of  $B$ , both relative to  $A$

Move in frame  $A$  = reorient by  $R$  and then move by  $d$  :  $Rv + d$

# Example



# Special Euclidean Group SE(3)

Coordinates of points in 3D Euclidean space =  $p^A \in \mathbb{R}^3$

Coordinates of cartesian frames in 3D Euclidean space =  
 $(d, R) \in \mathbb{R}^3 \times \text{SO}(3)$

Points: Euclidean Space :: Cartesian Frames : Special Euclidean Group

# Special Euclidean Group SE(3)

Coordinates of points in 3D Euclidean space =  $p^A \in \mathbb{R}^3$

Coordinates of cartesian frames in 3D Euclidean space =  
 $(d, R) \in \mathbb{R}^3 \times \text{SO}(3)$

Points: Euclidean Space :: Cartesian Frames : Special Euclidean Group

Affine Space : Euclidean Space :: G-Torsor : Special Euclidean Group

# Special Euclidean Group SE(3)

Coordinates of points in 3D Euclidean space =  $p^A \in \mathbb{R}^3$

Coordinates of cartesian frames in 3D Euclidean space =  
 $(d, R) \in \mathbb{R}^3 \times \text{SO}(3)$

Points: Euclidean Space :: Cartesian Frames : Special Euclidean Group

Affine Space : Euclidean Space :: G-Torsor : Special Euclidean Group

G-Torsor: A group  $G$  with an action that maps a group element to another group element

# Special Euclidean Group SE(3)

Coordinates of points in 3D Euclidean space =  $p^A \in \mathbb{R}^3$

Coordinates of cartesian frames in 3D Euclidean space =  
 $(d, R) \in \mathbb{R}^3 \times \text{SO}(3)$

Points: Euclidean Space :: Cartesian Frames : Special Euclidean Group

Affine Space : Euclidean Space :: G-Torsor : Special Euclidean Group

G-Torsor: A group  $G$  with an action that maps a group element to another group element

Again, no coordinate frame is unique.

For a  $G$ -Torsor, we don't define origin+basis (not a vector space).



# Special Euclidean Group SE(3)

Coordinates of points in 3D Euclidean space =  $p^A \in \mathbb{R}^3$

Coordinates of cartesian frames in 3D Euclidean space =  
 $(d, R) \in \mathbb{R}^3 \times \text{SO}(3)$

Points: Euclidean Space :: Cartesian Frames : Special Euclidean Group

Affine Space : Euclidean Space ::  $G$ -Torsor : Special Euclidean Group

$G$ -Torsor: A group  $G$  with an action that maps a group element to another group element

Again, no coordinate frame is unique.

For a  $G$ -Torsor, we don't define origin+basis (not a vector space).

Instead, we define an identity element (it's a group): the reference coordinate frame.

# Homogenous Transformations

We can convert the affine map between two Euclidean spaces of dimension 3 into a linear map between two subsets of  $\mathbb{R}^4$ .

Define a homogenization  $h: \mathbb{R}^3 \mapsto \mathbb{R}^4$  as  $h(p^A) = \begin{bmatrix} p^A \\ 1 \end{bmatrix}$ .

If  $p^A = Rp^B + d$ , then

$$h(p^A) = \begin{bmatrix} R & d \\ 0 & 1 \end{bmatrix} h(p^B). \quad (6)$$

The matrix  $\begin{bmatrix} R & d \\ 0 & 1 \end{bmatrix}$  represents a homogenous transformation, and forms a group.

# Checkpoint

- ▶ The coordinate transformation is  $p^B = (R_B^A)^{-1} (p^A - o_B^A)$
- ▶ Norm-preserving coordinate transformation = rigid motion of points within the same coordinate frame.
- ▶ Set of rigid body poses/rigid motions forms a group:  $SE(3)$
- ▶ After choosing a reference frame, we assign coordinates – aka rigid body pose –  $(d, R)$  to frame (Torsor structure)

## Back to $SO(3)$

- ▶ Rigid bodies correspond to cartesian frames

## Back to $SO(3)$

- ▶ Rigid bodies correspond to cartesian frames
- ▶ Cartesian frames have a position and orientation

## Back to $SO(3)$

- ▶ Rigid bodies correspond to cartesian frames
- ▶ Cartesian frames have a position and orientation
- ▶ Orientations are also a  $G$ -Torsor

## Back to $SO(3)$

- ▶ Rigid bodies correspond to cartesian frames
- ▶ Cartesian frames have a position and orientation
- ▶ Orientations are also a  $G$ -Torsor
- ▶ Rotations ( $\sim$  vector) help us get from one orientation ( $\sim$  point) to another

## Back to $SO(3)$

- ▶ Rigid bodies correspond to cartesian frames
- ▶ Cartesian frames have a position and orientation
- ▶ Orientations are also a  $G$ -Torsor
- ▶ Rotations ( $\sim$  vector) help us get from one orientation ( $\sim$  point) to another
- ▶ The rotation relative to a reference that produces an orientation becomes the 'orientation' coordinate of that frame.



## Back to SO(3)

- ▶ Rigid bodies correspond to cartesian frames
- ▶ Cartesian frames have a position and orientation
- ▶ Orientations are also a  $G$ -Torsor
- ▶ Rotations ( $\sim$  vector) help us get from one orientation ( $\sim$  point) to another
- ▶ The rotation relative to a reference that produces an orientation becomes the 'orientation' coordinate of that frame.
- ▶ We've called this matrix  $T_B^A$ ,  $R_B^A$ ,  $R$ ,  $T$

## Back to $SO(3)$

- ▶ Rigid bodies correspond to cartesian frames
- ▶ Cartesian frames have a position and orientation
- ▶ Orientations are also a  $G$ -Torsor
- ▶ Rotations ( $\sim$  vector) help us get from one orientation ( $\sim$  point) to another
- ▶ The rotation relative to a reference that produces an orientation becomes the 'orientation' coordinate of that frame.
- ▶ We've called this matrix  $T_B^A$ ,  $R_B^A$ ,  $R$ ,  $T$
- ▶ The  $G$ -Torsor nature is why  $SO(3)$  is called both the rotation group and the orientation group.

## Back to $SO(3)$

- ▶ Rigid bodies correspond to cartesian frames
- ▶ Cartesian frames have a position and orientation
- ▶ Orientations are also a  $G$ -Torsor
- ▶ Rotations ( $\sim$  vector) help us get from one orientation ( $\sim$  point) to another
- ▶ The rotation relative to a reference that produces an orientation becomes the 'orientation' coordinate of that frame.
- ▶ We've called this matrix  $T_B^A$ ,  $R_B^A$ ,  $R$ ,  $T$
- ▶ The  $G$ -Torsor nature is why  $SO(3)$  is called both the rotation group and the orientation group.
- ▶ Assigning coordinates to an orientation is the same as defining the rotation that generates that frame.


# Basic Rotations

Consider three frames rotated about each one of the world frame axes by an angle  $\theta$ .

# Basic Rotations

Consider three frames rotated about each one of the world frame axes by an angle  $\theta$ . Each rotation is given by

$$R_{x,\theta} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos(\theta) & -\sin(\theta) \\ 0 & \sin(\theta) & \cos(\theta) \end{bmatrix}$$




# Basic Rotations

Consider three frames rotated about each one of the world frame axes by an angle  $\theta$ . Each rotation is given by

$$R_{x,\theta} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos(\theta) & -\sin(\theta) \\ 0 & \sin(\theta) & \cos(\theta) \end{bmatrix}$$

$$R_{y,\theta} = \begin{bmatrix} \cos(\theta) & 0 & \sin(\theta) \\ 0 & 1 & 0 \\ -\sin(\theta) & 0 & \cos(\theta) \end{bmatrix}$$






# Basic Rotations

Consider three frames rotated about each one of the world frame axes by an angle  $\theta$ . Each rotation is given by

$$R_{x,\theta} = \begin{bmatrix} 1 & 0 & 0 \\ 0 & \cos(\theta) & -\sin(\theta) \\ 0 & \sin(\theta) & \cos(\theta) \end{bmatrix}$$

$$R_{y,\theta} = \begin{bmatrix} \cos(\theta) & 0 & \sin(\theta) \\ 0 & 1 & 0 \\ -\sin(\theta) & 0 & \cos(\theta) \end{bmatrix}$$

$$R_{z,\theta} = \begin{bmatrix} \cos(\theta) & -\sin(\theta) & 0 \\ \sin(\theta) & \cos(\theta) & 0 \\ 0 & 0 & 1 \end{bmatrix}$$



# General Rotations

We can construct a general rotation using a sequence of basic rotations. (Compare to Euclidean space)



# General Rotations

We can construct a general rotation using a sequence of basic rotations. (Compare to Euclidean space)

So, orientation coordinates can be derived by sequences of basic rotations.

# General Rotations

We can construct a general rotation using a sequence of basic rotations. (Compare to Euclidean space)

So, orientation coordinates can be derived by sequences of basic rotations.

For Euclidean vector spaces, the order of a sequence of (vector space) operations didn't matter:  $v + w = w + v$ .

# General Rotations

We can construct a general rotation using a sequence of basic rotations. (Compare to Euclidean space)

So, orientation coordinates can be derived by sequences of basic rotations.

For Euclidean vector spaces, the order of a sequence of (vector space) operations didn't matter:  $v + w = w + v$ .

For rotations, they do. In general,  $R_1 R_2 \neq R_2 R_1$ .

# Change-of-Basis For Orientations

Suppose we define an orientation  $B$  relative to a orientation  $A$  through a rotation  $R_B^A$ .

# Change-of-Basis For Orientations

Suppose we define an orientation  $B$  relative to a orientation  $A$  through a rotation  $R_B^A$ .

Now, someone decides to change the identity element to be orientation  $C$ , with coordinate  $R_C^A$  (in frame  $A$ ).

# Change-of-Basis For Orientations

Suppose we define an orientation  $B$  relative to a orientation  $A$  through a rotation  $R_B^A$ .

Now, someone decides to change the identity element to be orientation  $C$ , with coordinate  $R_C^A$  (in frame  $A$ ).

Which rotation  $R_B^C$  below correctly defines orientation of  $B$  relative to orientation  $C$ ?

1.  $R_B^C = R_B^A R_A^C$
2.  $R_B^C = R_B^A R_C^A$
3.  $R_B^C = R_A^C R_B^A$
4.  $R_B^C = R_C^A R_B^A$

# Change-of-Basis For Orientations

Suppose we define an orientation  $B$  relative to a orientation  $A$  through a rotation  $R_B^A$ .

Now, someone decides to change the identity element to be orientation  $C$ , with coordinate  $R_C^A$  (in frame  $A$ ).

Which rotation  $R_B^C$  below correctly defines orientation of  $B$  relative to orientation  $C$ ?

1.  $R_B^C = R_B^A R_A^C$
2.  $R_B^C = R_B^A R_C^A$
3.  $R_B^C = R_A^C R_B^A$
4.  $R_B^C = R_C^A R_B^A$

How would you pick the right transformation? Why did we not consider  $R_A^B$ ?

# Change-of-Basis For Orientations

- ▶ For example, imagine you, a driver, and a passenger in a car. Your orientation frames are aligned: Forward:  $x$ , upwards:  $z$ .
- ▶ When the car stops, the passenger opens the door spins to their right ( $R_C^A = R_{z,-90^\circ}$ )
- ▶ You lean back in your driver's seat ( $R_B^A = R_{y,-20^\circ}$ )
- ▶ What is your orientation according to the passenger?
  1.  $R_B^C = R_B^A R_A^C$
  2.  $R_B^C = R_B^A R_C^A$
  3.  $R_B^C = R_A^C R_B^A$
  4.  $R_B^C = R_C^A R_B^A$



# Change-of-Basis For Orientations

Rotation matrix  $R_B^A$  gives the coordinates of the basis vectors of frames  $B$  in  $A$ .

# Change-of-Basis For Orientations

Rotation matrix  $R_B^A$  gives the coordinates of the basis vectors of frames  $B$  in  $A$ .

We want to change the frame of these coordinates to frame  $C$ .

# Change-of-Basis For Orientations

Rotation matrix  $R_B^A$  gives the coordinates of the basis vectors of frames  $B$  in  $A$ .

We want to change the frame of these coordinates to frame  $C$ .

To change the coordinates of vectors from  $A$  to  $C$ , we must pre-multiply by  $(R_C^A)^{-1} = R_A^C$ . So,

$$R_B^C = R_A^C R_B^A$$

# Change-of-Basis For Orientations

**Alternatively,** The rigid motion in  $A$  corresponding to moving to frame  $B$  is  $R_B^A$ ; the rigid motion in frame  $C$  corresponding to moving to frame  $A$  is  $R_A^C$ .

The combined rigid motion in  $C$  is achieved by first moving by  $R_B^A$  **in  $C$** , then moving the result by  $R_A^C$ .

Therefore,

$$R_B^C = R_A^C R_B^A$$

.

# Change-of-Basis For Orientations

**Alternatively,** The rigid motion in  $A$  corresponding to moving to frame  $B$  is  $R_B^A$ ; the rigid motion in frame  $C$  corresponding to moving to frame  $A$  is  $R_A^C$ .

The combined rigid motion in  $C$  is achieved by first moving by  $R_B^A$  **in  $C$** , then moving the result by  $R_A^C$ .

Therefore,

$$R_B^C = R_A^C R_B^A$$

.

# Transforming Transforms

- ▶ We have looked at transforming points (or orientations) between frames, either as

# Transforming Transforms

- ▶ We have looked at transforming points (or orientations) between frames, either as
  - ▶ a rigid motion (coordinate stay in the same frame)

# Transforming Transforms

- ▶ We have looked at transforming points (or orientations) between frames, either as
  - ▶ a rigid motion (coordinates stay in the same frame)
  - ▶ a change of basis (coordinates are in a new frame)



# Transforming Transforms

- ▶ We have looked at transforming points (or orientations) between frames, either as
  - ▶ a rigid motion (coordinate stay in the same frame)
  - ▶ a change of basis (coordinates are in e new frame)
- ▶ Only two frames are involved

# Transforming Transforms

- ▶ We have looked at transforming points (or orientations) between frames, either as
  - ▶ a rigid motion (coordinate stay in the same frame)
  - ▶ a change of basis (coordinates are in e new frame)
- ▶ Only two frames are involved
- ▶ We now look at transforming **transformations** between frames.

# Transforming Transforms

- ▶ We have looked at transforming points (or orientations) between frames, either as
  - ▶ a rigid motion (coordinate stay in the same frame)
  - ▶ a change of basis (coordinates are in e new frame)
- ▶ Only two frames are involved
- ▶ We now look at transforming **transformations** between frames.

# Transforming Transforms

- ▶ We have looked at transforming points (or orientations) between frames, either as
  - ▶ a rigid motion (coordinate stay in the same frame)
  - ▶ a change of basis (coordinates are in e new frame)
- ▶ Only two frames are involved
- ▶ We now look at transforming **transformations** between frames.

Instead of orientation  $R_B^A$  in frame  $A$ , what if we define rotation  $R^A$  in frame  $A$ .

How do we represent this rotation in frame  $C$ ?

# Change-of-Basis For Rotations

- ▶ The rotation  $R^A$  is relative to frame  $A$ .
- ▶ A general orientation  $P$  has coordinates  $R_P^A$  in frame  $A$
- ▶ Rotating this orientation results in a new orientation  $R^A R_P^A$  in frame  $A$ :

$$R_P^A \mapsto R^A R_P^A$$

- ▶ But, note that  $R_P^A = R_C^A R_P^C$
- ▶ Therefore :

$$R_C^A R_P^C \mapsto R^A R_C^A R_P^C, \text{ or}$$


$$R_P^C \mapsto \left(R_C^A\right)^{-1} R^A R_C^A R_P^C, \text{ or}$$

- ▶ Therefore, a rotation  $R^A$  in frame  $A$  becomes a rotation



$$R^C = \left(R_C^A\right)^{-1} R^A R_C^A$$

in frame  $C$ .

# Extrinsic vs Intrinsic Rotations




# Extrinsic vs Intrinsic Rotations



Rotate about  $z$

# Extrinsic vs Intrinsic Rotations





# Extrinsic vs Intrinsic Rotations

- ▶ A first rigid motion corresponding to rotation  $R_1$  relative to a frame  $A$  produces frame  $B$
- ▶ A second rigid motion rotation  $R_2$  can be applied relative to either  $A$  or  $B$ .
- ▶ When applied relative to  $B$ , the second rotation is an intrinsic rotation.  $R = R_1 R_2$ .
- ▶ When applied relative to  $A$ , the second rotation is an extrinsic rotation.  $R = R_2 R_1$ .