Vrite your name here Surname	Other	names
Pearson Edexcel nternational Advanced Level	Centre Number	Candidate Number
Further Pu	Ire	
Mathemate Advanced/Advanced	tics F3	
Mathemat	tics F3	Paper Reference WFM03/01

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information

- The total mark for this paper is 75.
- The marks for each question are shown in brackets
 use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over ▶



P51515A

©2018 Pearson Education Ltd.

1/1/1/1/1/1/1/



1. Solve the equation				
$15 \operatorname{sech}^2 x + 7 \tanh x = 13$				
Give your answers in terms of simplified natural logarithms.				

		Leave blank
Question 1 continued		
		Q1
	(Total 6 marks)	
	(IUIAI U IIIAI KS)	



DO NOT WRITE IN THIS AREA

(a) Find the eigenvalues and corresponding normalised eigenvectors of the matrix ${\bf A}.$

(7)

(b) Write down a matrix **P** and a diagonal matrix **D** such that $P^{T}AP = D$.

(2)

AND THE PROPERTY OF A SECTION ASSESSMENT ASSESSMENT ASSESSMENT OF A SECTION ASSESSMENT ASSESSMENT ASSESSMENT ASSESSMENT ASSESSMENT ASSESSMENT ASSESSME

			2		
×					
×					
×					
		>			
	Į	ŗ	9		,
S	8	5	₹		7
۶	Š	2	Š	2	
S	Š	2	Š	2	
5	Š	5	Š	5	
8	3	2	Š	2	
5	3	5			
8	3	2			
8	3	2			
>					
>					
\ \ \ \		>			
>		2			
>		}			
>					
>					
\ \ \ \ \					
\ \ \ \ \					
>					
\ \ \ \ \		>			
\ \ \ \ \					
\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\					
\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\					
\ \ \ \ \					
\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	à				
\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	à				
\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	à				
\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	à				
\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\					
\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	à				
\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	à				
	à				
\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\\	à				
	à				
	à				
	à				

	Leav
uestion 2 continued	



ied		

Question 2 continued		
		1
		Q2
	(Total 9 marks)	



3. Given that

$$y = \arctan\left(\frac{\sin x}{\cos x - 1}\right)$$
 $x \neq 2n\pi$, $n \in \mathbb{Z}$

Show that

$$\frac{\mathrm{d}y}{\mathrm{d}x} = k$$

where k is a constant to be found.

(6)

Question 3 continued	b.
destion 5 continued	



Question 3 continued	

Question 3 continued	Leave blank
Question 5 continued	
	Q3
(Total 6 marks)	
(======================================	



4. The hyperbola H has equation

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$$

The line l is a normal to H at the point $P(a \sec \theta, b \tan \theta)$, $0 < \theta < \frac{\pi}{2}$

(a) Using calculus, show that an equation for l is

$$ax\sin\theta + by = (a^2 + b^2)\tan\theta$$
(5)

The line l meets the x-axis at the point Q, and the point M is the midpoint of PQ.

(b) Find the coordinates of M.

(3)

(c) Hence find the cartesian equation of the locus of M as θ varies, giving your answer in the form $y^2 = f(x)$.

(4)

uestion 4 continued	



Question 4 continued	
guestion 4 continued	

Question 4 continued		blank
		Q 4
	(T-4-140 1)	
	(Total 12 marks)	



5.

$$\mathbf{M} = \begin{pmatrix} 4 & -5 & 0 \\ k & 2 & 0 \\ -3 & -5 & k \end{pmatrix}, \text{ where } k \text{ is a real constant, } k \neq 0, \ k \neq -\frac{8}{5}$$

(a) Find, in terms of k, the inverse of the matrix M.

(5)

A transformation $T: \mathbb{R}^3 \to \mathbb{R}^3$ is represented by the matrix

$$\begin{pmatrix} 4 & -5 & 0 \\ -1 & 2 & 0 \\ -3 & -5 & -1 \end{pmatrix}$$

The transformation T maps the plane Π_1 onto the plane Π_2

Given that the plane Π_2 has equation 2x - z = 4

(b) find a cartesian equation of the plane Π_1

1	•		
	h		
	₹,		
L	_	,	

	Leave blank
Question 5 continued	



Question 5 continued	

Question 5 continued		blank
Question 3 continued		
		Q5
	(Total 11 manular)	
	(Total 11 marks)	



(5)

6. The curve C has parametric equations

$$x = \theta - \tanh \theta$$
, $y = \operatorname{sech} \theta$, $0 \le \theta \le \ln 3$

- (a) Find
 - (i) $\frac{\mathrm{d}x}{\mathrm{d}\theta}$
 - (ii) $\frac{\mathrm{d}y}{\mathrm{d}\theta}$

The curve C is rotated through 2π radians about the x-axis.

(b) Find the exact area of the curved surface formed, giving your answer as a multiple of π .

	Leave
Question 6 continued	



Question 6 continued		

	blank
Question 6 continued	
	Q6
	20
(Total 7 marks)	



7. The plane Π_1 has equation x + y + z = 3 and the plane Π_2 has equation 2x + 3y - z = 4

The planes $\Pi_{\rm l}$ and $\Pi_{\rm l}$ intersect in the line L.

(a) Find a cartesian equation for the line L.

(6)

The plane Π_3 has equation

$$\mathbf{r.} \begin{pmatrix} 5 \\ -4 \\ 4 \end{pmatrix} = 12$$

The line L meets the plane Π_3 at the point A.

(b) Find the coordinates of A.

(3)

(c) Find the acute angle between \overrightarrow{OA} and the line L, where O is the origin. Give your answer in degrees to one decimal place.

(3)

	Leave
	blank
Question 7 continued	



Question 7 continued	
Question / continueu	

	blank
Question 7 continued	
	07
	Q7
(Total 12 marks)	
(Total 12 Illal K3)	+



8.

$$I_n = \int \frac{x^n}{\sqrt{(x^2 + k^2)}} dx$$
 where k is a constant and $n \in \mathbb{Z}^+$

(a) Show that, for $n \ge 2$

$$I_n = \frac{x^{n-1}}{n} (x^2 + k^2)^{\frac{1}{2}} - \frac{(n-1)}{n} k^2 I_{n-2}$$

(7)

(b) Hence find the exact value of

$$\int_0^1 \frac{x^5}{\sqrt{(x^2+1)}} \, \mathrm{d}x$$

(5)

	I
uestion 8 continued	



uestion 8 continued		

	Le
	bl
Question 8 continued	
	1



END		
	(Total 12 marks) TOTAL FOR PAPER: 75 MARKS	
	/TD / 144	
		Q8