Surname	Other n.	ames
Pearson Edexcel International Advanced Level	Centre Number	Candidate Number
Mechanica Advanced/Advance		
Advanced/Advance	u Jubjiului v	
Tuesday 25 October 2016 - Time: 1 hour 30 minutes		Paper Reference WME01/01

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer **all** questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- Whenever a numerical value of g is required, take g = 9.8 m s⁻², and give your answer to either two significant figures or three significant figures.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information

- The total mark for this paper is 75.
- The marks for each question are shown in brackets
 use this as a quide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

PEARSON

Turn over ▶



1.	Two particles, P and Q , have masses $2m$ and $3m$ respectively. They are moving tow each other, in opposite directions, along the same straight line, on a smooth horizon plane. The particles collide. Immediately before they collide the speed of P is $2u$ and speed of Q is u . In the collision the magnitude of the impulse exerted on P by Q is S	ontal d the
	(a) Find the speed of <i>P</i> immediately after the collision.	(3)
	(b) State whether the direction of motion of P has been reversed by the collision.	(1)
	(c) Find the speed of Q immediately after the collision.	(3)

Question 1 continued	Leave blank
	-
	_
	-
	_
	-
	_
	-
	_
	-
	_
	-
	_
	_
	_
	_
	_
	_
	_
	_
	-
	-
	Q1
(Total 7 marks)	



2.	[In this question i and j are perpendicular unit vectors in a horizontal plane.]
	Three forces, $(-10\mathbf{i} + a\mathbf{j})$ N, $(b\mathbf{i} - 5\mathbf{j})$ N and $(2a\mathbf{i} + 7\mathbf{j})$ N, where a and b are constants, act on a particle P of mass 3 kg. The acceleration of P is $(3\mathbf{i} + 4\mathbf{j})$ m s ⁻²
	(a) Find the value of a and the value of b. (5)
	At time $t = 0$ seconds the speed of P is u m s ⁻¹ and at time $t = 4$ seconds the velocity of P is $(20\mathbf{i} + 20\mathbf{j})$ m s ⁻¹
	(b) Find the value of u. (4)



Question 2 continued		bla
		Q2
	(Total 9 marks)	



3.




Figure 1

A plank AB has length 8 m and mass 12 kg. The plank rests on two supports. One support is at C, where AC = 3 m and the other support is at D, where AD = x metres. A block of mass 3 kg is placed on the plank at B, as shown in Figure 1. The plank rests in equilibrium in a horizontal position. The magnitude of the force exerted on the plank by the support at D is twice the magnitude of the force exerted on the plank by the support at C. The plank is modelled as a uniform rod and the block is modelled as a particle.

Question 3 continued	blank
	Q3
(Total 7 marks)	



4. [In this question **i** and **j** are horizontal unit vectors due east and due north respectively and position vectors are given relative to a fixed origin O]

A particle P is moving with velocity $(\mathbf{i} - 2\mathbf{j})$ km h⁻¹. At time t = 0 hours, the position vector of P is $(-5\mathbf{i} + 9\mathbf{j})$ km. At time t hours, the position vector of P is \mathbf{p} km.

(a) Find an expression for \mathbf{p} in terms of t.

(2)

The point A has position vector $(3\mathbf{i} + 2\mathbf{j})$ km.

(b) Find the position vector of P when P is due west of A.

(4)

Another particle Q is moving with velocity $[(2b-1)\mathbf{i} + (5-2b)\mathbf{j}] \text{ km h}^{-1}$ where b is a constant.

Given that the particles are moving along parallel lines,

(c) find the value of b.

(4)

8

Question 4 continued		blank
		Q4
	(Total 10 marks)	



(9)

5.




Figure 2

A particle P of mass 0.5 kg is at rest on a rough plane which is inclined to the horizontal at 30° . The particle is held in equilibrium by a force of magnitude $8\,\mathrm{N}$, acting at an angle of 40° to the plane, as shown in Figure 2. The line of action of the force lies in the vertical plane containing P and a line of greatest slope of the plane. The coefficient of friction between P and the plane is μ . Given that P is on the point of sliding up the plane, find the value of μ .

10

Question 5 continued	Leave blank
Question 5 continued	
	1



Question 5 continued		

Question 5 continued		blank
Zaronon e communes		
		Q5
	/m	
	(Total 9 marks)	



ac ar	celeration $0.5 \mathrm{ms^{-2}}$. At the instant when <i>B</i> is 200 m lad the speed of <i>B</i> is $44 \mathrm{ms^{-1}}$. Find the speed of <i>B</i> wh	behind A , the speed of A is 35 m en it overtakes A .

Question 6 continued	Leave
	Q6
(Total 9 marks)	



7.	A train moves on a straight horizontal track between two stations <i>A</i> and <i>B</i> . The train starts from rest at <i>A</i> and moves with constant acceleration $1 \mathrm{m}\mathrm{s}^{-2}$ until it reaches a speed of $V \mathrm{m}\mathrm{s}^{-1}$. The train maintains this speed of $V \mathrm{m}\mathrm{s}^{-1}$ for the next <i>T</i> seconds before slowing down with constant deceleration $0.5 \mathrm{m}\mathrm{s}^{-2}$, coming to rest at <i>B</i> . The journey from <i>A</i> to <i>B</i> takes $180 \mathrm{s}$ and the distance between the stations is $4800 \mathrm{m}$.
	(2)
	(b) Show that $T = 180 - 3V$. (2)
	(c) Find the value of V . (7)

	I t
Question 7 continued	



I		

Question 7 continued	Leave blank
	05
	Q7
(Total 11 marks)	



8.




Figure 3

Two particles P and Q have masses $2 \, \text{kg}$ and $3 \, \text{kg}$ respectively. The particles are attached to the ends of a light inextensible string. The string passes over a smooth light pulley which is fixed at the top of a rough plane. The plane is inclined to horizontal ground at an angle α , where $\tan \alpha = \frac{3}{4}$. Initially P is held at rest on the inclined plane with the part of the string from P to the pulley parallel to a line of greatest slope of the plane. The particle Q hangs freely below the pulley at a height of $0.5 \, \text{m}$ above the ground, as shown in Figure 3. The coefficient of friction between P and the plane is μ . The system is released from rest, with the string taut, and Q strikes the ground before P reaches the pulley. The speed of Q at the instant when it strikes the ground is $1.4 \, \text{m s}^{-1}$.

(a) For the motion before Q strikes the ground, find the tension in the string.

(5)

(b) Find the value of μ .

(8)

	Leave
	blank
Question 8 continued	
Question o continueu	



Question 8 continued		

	Leave
	blank
Question 8 continued	



Question 8 continued	Leave
	00
(Total 12 manks)	Q8
TOTAL FOR PAPER: 75 MARKS END	