/rite your name here Surname	Oth	er names
Pearson Edexcel International Idvanced Level	Centre Number	Candidate Number
Further Pu	Iro	
Mathemat Advanced/Advanced	tics F3	
Mathemat	tics F3	Paper Reference WFM03/01

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information

- The total mark for this paper is 75.
- The marks for each question are shown in brackets
 use this as a guide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

Turn over 🕨





Leave blank

1.	The curve C has equation
	$y = 9\cosh x + 3\sinh x + 7x$
	Use differentiation to find the exact x coordinate of the stationary point of C , giving your
	answer as a natural logarithm. (6)

(6)

2. An ellipse has equation

$$\frac{x^2}{25} + \frac{y^2}{4} = 1$$

The point *P* lies on the ellipse and has coordinates $(5\cos\theta, 2\sin\theta)$, $0 < \theta < \frac{\pi}{2}$

The line L is a normal to the ellipse at the point P.

(a) Show that an equation for L is

$$5x \sin \theta - 2y \cos \theta = 21 \sin \theta \cos \theta \tag{5}$$


Given that the line L crosses the y-axis at the point Q and that M is the midpoint of PQ,

(b) find the exact area of triangle OPM, where O is the origin, giving your answer as a multiple of $\sin 2\theta$

	blank
Question 2 continued	



3. Without using a calculator, find



(b) $\int_{-1}^{4} \frac{1}{\sqrt{4x^2 - 12x + 34}} \, dx$, giving your answer in the form $p \ln(q + r\sqrt{2})$,

where p, q and r are rational numbers to be found.

(7)

	Leave
	blank
Question 3 continued	



4. $\mathbf{M} = \begin{pmatrix} 1 & k & 0 \\ -1 & 1 & 1 \\ 1 & k & 3 \end{pmatrix}, \text{ where } k \text{ is a constant}$

(a) Find \mathbf{M}^{-1} in terms of k.

(5)

(4)

Hence, given that k = 0

(b) find the matrix N such that

$$\mathbf{MN} = \begin{pmatrix} 3 & 5 & 6 \\ 4 & -1 & 1 \\ 3 & 2 & -3 \end{pmatrix}$$

	1
uestion 4 continued	



- 5. Given that $y = \operatorname{artanh}(\cos x)$
 - (a) show that

$$\frac{\mathrm{d}y}{\mathrm{d}x} = -\mathrm{cosec}\,x$$

(b) Hence find the exact value of

$$\int_0^{\frac{\pi}{6}} \cos x \, \operatorname{artanh}(\cos x) \, \mathrm{d}x$$

giving your answer in the form $a \ln(b + c\sqrt{3}) + d\pi$, where a, b, c and d are rational numbers to be found.

(5)

(2)

	Leave
	blank
Question 5 continued	



Leave blank

6.	The coordinates of the points A , B and C relative to a fixed origin O are $(1, 2, 3)$, $(-1, 3, 4)$ and $(2, 1, 6)$ respectively. The plane Π contains the points A , B and C .	
	(a) Find a cartesian equation of the plane Π . (5)	
	The point D has coordinates $(k, 4, 14)$ where k is a positive constant.	
	Given that the volume of the tetrahedron ABCD is 6 cubic units,	
	(b) find the value of k .	
	(4)	

	Leave
	blank
Question 6 continued	
	1



7. The curve C has parametric equations

$$x = 3t^4, \quad y = 4t^3, \qquad 0 \leqslant t \leqslant 1$$

The curve C is rotated through 2π radians about the x-axis. The area of the curved surface generated is S.

(a) Show that

$$S = k\pi \int_0^1 t^5 (t^2 + 1)^{\frac{1}{2}} dt$$

where k is a constant to be found.

(4)

(b) Use the substitution $u^2 = t^2 + 1$ to find the value of S, giving your answer in the form $p\pi(11\sqrt{2} - 4)$ where p is a rational number to be found.

(7)

	Leave
	blank
Question 7 continued	



$$I_n = \int_0^{\ln 2} \tanh^{2n} x \, \mathrm{d}x, \quad n \geqslant 0$$

(a) Show that, for $n \ge 1$

$$I_n = I_{n-1} - \frac{1}{2n-1} \left(\frac{3}{5}\right)^{2n-1}$$

(5)

(b) Hence show that

$$\int_0^{\ln 2} \tanh^4 x \, \mathrm{d}x = p + \ln 2$$

where p is a rational number to be found.

(5)

	1
uestion 8 continued	
	_

