Surname	Other nar	nes
Pearson Edexcel International Advanced Level	Centre Number	Candidate Number
Core Math Advanced Subsidiar		s C12
Tuesday 12 January 2016 – Time: 2 hours 30 minutes	•	Paper Reference WMA01/01

Candidates may use any calculator allowed by the regulations of the Joint Council for Qualifications. Calculators must not have the facility for symbolic algebra manipulation, differentiation and integration, or have retrievable mathematical formulae stored in them.

Instructions

- Use **black** ink or ball-point pen.
- If pencil is used for diagrams/sketches/graphs it must be dark (HB or B). Coloured pencils and highlighter pens must not be used.
- **Fill in the boxes** at the top of this page with your name, centre number and candidate number.
- Answer all questions and ensure that your answers to parts of questions are clearly labelled.
- Answer the questions in the spaces provided
 there may be more space than you need.
- You should show sufficient working to make your methods clear. Answers without working may not gain full credit.
- When a calculator is used, the answer should be given to an appropriate degree of accuracy.

Information

- The total mark for this paper is 125.
- The marks for each question are shown in brackets
 use this as a quide as to how much time to spend on each question.

Advice

- Read each question carefully before you start to answer it.
- Try to answer every question.
- Check your answers if you have time at the end.

P 4 6 9 5 7 A 0 1 5 2

Turn over ▶



(2)

(3)

1. A sequence of numbers $u_1, u_2, u_3, ...$ satisfies $u_{n+1} = 2u_n - 6, \quad n \geqslant 1$ Given that $u_1 = 2$ (a) find the value of u_3 (b) evaluate $\sum_{i=1}^4 u_i$

	Question 1 continued	Lea bla
		01
(Total 5 marks)		V1



- 2. (i) Given that $\frac{49}{\sqrt{7}} = 7^a$, find the value of a.
 - (ii) Show that $\frac{10}{\sqrt{18} 4} = 15\sqrt{2} + 20$

You must show all stages of your working.

(3)

(2)

		blank
Question 2 continued		
		03
		Q2
	(Total 5 marks)	



3.	Find, using calculus and showing each step of your working,	
	$\int_{1}^{4} \left(6x - 3 - \frac{2}{\sqrt{x}}\right) \mathrm{d}x$	
		(5)
_		

	Leave
	blank
Question 3 continued	
	•
	•
	.
	.
	Q3
(Total 5 marks)	.
(Total S marks)	



	Find the first term and the common difference of this sequence.	
	Find the first term and the common difference of this sequence.	
_		



	blank
Question 4 continued	
	Q4
(Total 6 marks)	



5. (a) Sketch the graph of $y = \sin 2x$, $0 \le x \le \frac{3\pi}{2}$

Show the coordinates of the points where your graph crosses the *x*-axis.

(2)

The table below gives corresponding values of x and y, for $y = \sin 2x$. The values of y are rounded to 3 decimal places where necessary.

x	0	$\frac{\pi}{12}$	$\frac{\pi}{6}$	$\frac{\pi}{4}$
y	0	0.5	0.866	1

(b) Use the trapezium rule with all the values of *y* from the table to find an approximate value for

$$\int_0^{\frac{\pi}{4}} \sin 2x \, dx \tag{3}$$

Question 5 continued	blank
	Q5
(Total 5 marks)	



6. $f(x) = x^3 + x^2 - 12x - 18$ (a) Use the factor theorem to show that $(x + 3)$ is a factor of $f(x)$. (2) (b) Factorise $f(x)$. (2) (c) Hence find exact values for all the solutions of the equation $f(x) = 0$ (3)			
 (b) Factorise f(x). (c) Hence find exact values for all the solutions of the equation f(x) = 0 	6.	$f(x) = x^3 + x^2 - 12x - 18$	
(b) Factorise f(x).(c) Hence find exact values for all the solutions of the equation f(x) = 0		(a) Use the factor theorem to show that $(x + 3)$ is a factor of $f(x)$.	(2)
(c) Hence find exact values for all the solutions of the equation $f(x) = 0$			(2)
(c) Hence find exact values for all the solutions of the equation $f(x) = 0$		(b) Factorise $f(x)$.	(2)
			(2)
		(c) Hence find exact values for all the solutions of the equation $f(x) = 0$	(3)

	Leave
	blank
Question 6 continued	
	1



uestion 6 continued	

Question 6 continued	blank
	Q6
(Total	7 marks)
(200	



7. (a) Find the first 4 terms, in ascending powers of x , of the binomial expansion	
of $(1 + kx)^8$, where k is a non-zero constant. Give each term in its simplest form.	(4)
Given that the coefficient of x^3 in this expansion is 1512	
(b) find the value of k .	
	(3)



estion 7 continued	



uestion 7 continued

estion 7 continued	



8.	(a) Given that $7 \sin x = 3 \cos x$, find the exact value of $\tan x$.	(1)
	(b) Hence solve for $0 \leqslant \theta < 360^{\circ}$	
	$7\sin(2\theta + 30^\circ) = 3\cos(2\theta + 30^\circ)$	
	giving your answers to one decimal place.	
	(Solutions based entirely on graphical or numerical methods are not acceptable.)	(5)

(Total 6 marks)	Question 8 continued	blank
(Total 6 marks)		Q8
	(Total 6 marks)	



9. The resident population of a city is 130 000 at the end of Year 1

A model predicts that the resident population of the city will increase by 2% each year, with the populations at the end of each year forming a geometric sequence.

(a) Show that the predicted resident population at the end of Year 2 is 132 600

(1)

(b) Write down the value of the common ratio of the geometric sequence.

(1)

The model predicts that Year N will be the first year which will end with the resident population of the city exceeding 260 000

(c) Show that

$$N > \frac{\log_{10} 2}{\log_{10} 1.02} + 1$$

(4)

(d) Find the value of N.

(1)

	Leav
	blanl
Question 9 continued	



estion 9 continued	

Question 9 continued		Leav blan
		Ω^{0}
		Q9
	(Total 7 marks)	



10. The curve C has equation

$$y = 12x^{\frac{5}{4}} - \frac{5}{18}x^2 - 1000, \qquad x > 0$$

(a) Find $\frac{dy}{dx}$

(2)

(b) Hence find the coordinates of the stationary point on C.

(5)

(c) Use $\frac{d^2y}{dx^2}$ to determine the nature of this stationary point.

(3)

THE DIVINE ANDA

	Leave
	blank
Question 10 continued	
Question to commute	



	Leave blank
Question 10 continued	

		Leave blank
Question 10 continued		
		Q10
	(Tatal 10 mayles)	
	(Total 10 marks)	



11.




Figure 1

Figure 1 shows a triangle XYZ with XY = 10 cm, YZ = 16 cm and ZX = 12 cm.

(a) Find the size of the angle *YXZ*, giving your answer in radians to 3 significant figures.

(3)

The point A lies on the line XY and the point B lies on the line XZ and AX = BX = 5 cm. AB is the arc of a circle with centre X.

The shaded region S, shown in Figure 1, is bounded by the lines BZ, ZY, YA and the arc AB.

Find

(b) the perimeter of the shaded region to 3 significant figures,

(4)

(c) the area of the shaded region to 3 significant figures.

(4)

	L
	b
Question 11 continued	



Overtion 11 continued	Leave blank
Question 11 continued	

Question 11 continued	Leave blank
	Q11
(Total 11 marks)	



12.

$$f(x) = \frac{(4+3\sqrt{x})^2}{x}, \qquad x > 0$$

(a) Show that $f(x) = Ax^{-1} + Bx^{k} + C$, where A, B, C and k are constants to be determined.

(4)

(b) Hence find f'(x).

(2)

(c) Find an equation of the tangent to the curve y = f(x) at the point where x = 4

(4)

	Leave
	blank
Question 12 continued	



Question 12 continued

	Leave
Question 12 continued	blank
Question 12 Continued	
	Q12
(Total 10 marks)	



Leave blank

(a) Show that k satisfies the inequality	
$11k^2 - 30k - 9 > 0$	
(b) Find the range of possible values for k .	

	Leave
	blank
Question 13 continued	



estion 13 continued	

Question 13 continued		eave lank
	Q1	13
	(Total 8 marks)	
	()	j



14. (i) Given that

$$\log_a x + \log_a 3 = \log_a 27 - 1$$
, where a is a positive constant

find, in its simplest form, an expression for x in terms of a.

(4)

(ii) Solve the equation

$$(\log_5 y)^2 - 7(\log_5 y) + 12 = 0$$

showing each step of your working.

(4)

Question 14 continued	blank
	Q14
(Total 8 marks)	



Leave blank

15. The points A and B have coordinates $(-8, -8)$ and $(12, 2)$ respectively. AB is the diameter of a circle C .	
(a) Find an equation for the circle <i>C</i> .	
(6))
The point $(4, 8)$ also lies on C .	
(b) Find an equation of the tangent to C at the point (4, 8), giving your answer in the form $ax + by + c = 0$	
4)
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_
	_



	Leave
Question 15 continued	blank
Question to continuou	



estion 15 continued	

Question 15 continued	Leave
Question 13 continued	
	015
	Q15
(Total 10 marks)	






Figure 2

The straight line *l* with equation $y = \frac{1}{2}x + 1$ cuts the curve *C*, with equation $y = x^2 - 4x + 3$, at the points *P* and *Q*, as shown in Figure 2

(a) Use algebra to find the coordinates of the points P and Q.

(5)

The curve C crosses the x-axis at the points T and S.

(b) Write down the coordinates of the points T and S.

(2)

The finite region R is shown shaded in Figure 2. This region R is bounded by the line segment PQ, the line segment TS, and the arcs PT and SQ of the curve.

(c) Use integration to find the exact area of the shaded region R.

(8)

	Leave
	blank
Question 16 continued	
Question to continuou	



estion 16 continued	

	Leave
	blank
Question 16 continued	



Question 16 continued	blank
	Q16
(Total 15 marks)	
TOTAL FOR PAPER: 125 MARKS	
END	