
A Expression-matched controls
TP53I3 rank 20/21; empirical P = 0.095; 10/20 controls had nominal P < 0.05
Control genes
TP53I3
KCNAB2
TP53I3
RUNDC1
FAM89B
ANKS6
ARHGEF6
GALNT4
AC093019.2
SMCR5
MUL1
TAS2R14
SNHG18
LINC01560
ZNF418
AC096642.1
TUBB6
AL139095.4
AC005070.3
ZFX-AS1
MFSD4A
AL138824.1
0.75
1
2
4
Odds ratio, aMASH versus ctrl (95% CI)
B Disease and alcohol contrasts
GSE256398; fold change versus Healthy
BH q >= 0.05
BH q < 0.05
Alcohol hepatitis (n = 5)
7.80x; BH q = 0.0355
MASH fibrosis (n = 4)
10.45x; BH q = 0.0355
Alcohol cirrhosis (n = 4)
6.63x; BH q = 0.245
MASH cirrhosis (n = 4)
2.15x; BH q = 0.245
MASLD F0 (n = 3)
2.35x; BH q = 0.245
1
2
4
8
Fold change versus Healthy (log scale)
C MASL-referenced sensitivity
Ratio of CP10K + 0.01; 95% CI
eMASH
aMASH
Matched DecontX
Matched raw
Primary model
1
2
3
4
5
Ratio versus MASL (log scale)