

(A) Hepatocyte TP53I3 by group
GSE256398; one point per donor (n = 17)
TP53I3 expression (CP10K)
0.20
0.15
0.10
0.05
0.00
Healthy
MASLD F0
MASH fibrosis
MASH cirrhosis
(B) Fibrosis-strata trend
Jonckheere-Terpstra P = 0.0389
Median TP53I3 expression (CP10K)
0.20
0.15
0.10
0.05
0.00
Healthy
MASLD F0
MASH fibrosis
MASH cirrhosis
(C) Positive fraction by cell type
GSE256398; 111,435 nuclei
Other cell types
Hepatocyte
T_NK
HSC_Stellate
Neutrophil
B_Plasma
Endothelial
Hepatocyte
Unassigned
Cholangiocyte
0.14% (n= 7,174)
1.01% (n= 8,591)
1.17% (n= 7,697)
1.24% (n= 647)
1.49% (n= 7,036)
2.46% (n=72,421)
5.08% (n= 1,162)
6.46% (n= 6,707)
0.0
2.5
5.0
7.5
% TP53I3-positive cells
(D) Fold change versus Healthy
BH-adjusted three-comparison family
BH-adjusted P < 0.05
Not significant
MASH Fibrosis
MASLD F0
MASH Cirrhosis
10.45x BH P = 0.043
2.35x BH P = 0.245
2.15x BH P = 0.245
5
10
15
Fold change (median pseudobulk)