

Formula Name KETAMINE HYDROGEL 5% GEL

Initials AP **Quantity** 60 **MLS** **Schedule** 3 **Days to Exp** 14 **Date of Formula** 9/9/2025
Cost \$11.99 **Cost Per Unit** \$0.20 **AWP** \$0.00 **AWP Per Unit** \$0.00 **Last Review** 1/6/2016

Chemical Ingredients

Chemical Name	Form	Qty	Units	QS	Est Cost	Ha	Note
KETAMINE HCL USP	PWD	3.4500	GMS	☐	\$7.51		SOLUBILITY: WATER (FS) ALCOHOL (S) MAXIMUM SOLUBILITY IS 200MG/ML WHITE CRYSTALLINE POWDER WITH A CHARACTERISTIC COLOR SOLUTIONS MAY BE CLEAR TO SLIGHTLY GREEN USED IN ANESTHESIA MAY CAUSE AMNESIA
PROPYLENE GLYCOL USP	LIQU	5.0000	MLS	☐	\$0.07		SOLUBILITY:N/A
HYDROGEL	GEL	60.0000	GMS	☒	\$4.42		

Therapy**Description**

Clear thin hydrated gel

Classification(s)Non-Preserved Aqueous**Disease State(s)****Instructions****A. Calculations/Special Considerations:**

Ketamine 1mg = Ketamine HCl 1.15 mg

B. Equipment Required:

Beakers; weigh boats; mortar& pestle, #2 rubber spatulas, metal spatula, , #2 60mL luer lock syringe, luer-luer adaptor, luer syringe tip cap, oral syringe,

For batches 240 mL or less: 60 mL syringes, luer lock adapters.

For batches 240+ mL: large mixing bowl w/spade-shaped attachment or unguator

C. Method of Preparation:

1. Calculate required amounts and fill out compounding and control usage logs
2. Measure out ingredients
3. Triturate Ketamine in mortar.
4. Wet step 3 with propylene glycol (for 240 mL or less: use syringe to syringe method; for 240+ mL: use mixer on lowest speed)
5. Remove plunger from a 60mL syringe, cap the tip, and stand syringe upright in a 30mL beaker, then pour mixture into syringe
6. In new syringe fill with Hydrogel base add plunger loosely, invert and tap Hydrogel base to the bottom, once tip is clear of base, fully insert plunger and expel extra air
7. Attach syringe from step 5 to Hydrogel base containing syringe with luer lock adapter and mix via syringe to syringe transfer until evenly dispersed (complete 40 transfers)
8. Package and label with barcode label and affix "external use only" label
9. Tape labels
10. Have compound certified pharmacist perform final verification

D. Labeling:

Label with lot number and expiration barcode and ingredient(s) label.

External use only auxillary sticker

Refrigerate sticker

Add prepack label if compound lot made to stock for multiple Rxs

Add Rx auxillary label if compound lot made to order for a unique Rx

_____ (place label here) _____

Appropriate References Should Be
Consulted. Safety And Efficacy Of
These Formulas Are The
Responsibility Of The Compounding
Pharmacist

E. Description of Finished Product:

Clear thin hydrated gel

F. Quality Control Procedures

Physical appearance of compound verified by compounding certified pharmacist

Volume of final product is within 5% of intended batch size

Pharmacist signature: _____

G. Packaging Container:

May bulk store in 60mL luer-lock syringe with UV protective bag

May dispense in UV protective oral syringe, metered dose pump, or uno-dose applicator

H. Storage Requirements

Store at refrigerated temperature (36°–46 °F)

Maximum BUD - 14 days

I. Formula Source & BUD Testing:

Formulation - OPPC

BUD - USP 795

INGREDIENT WEIGHT #1